





Tightening Targets



Courtesy Martin Haigh (SXE)

Updated Reasons for Concern



(Smith et al. 2009 PNAS)


Timothy M. Lenton*†, Hermann Held[‡], Elmar Kriegler^{‡§}, Jim W. Hall[¶], Wolfgang Lucht[‡], Stefan Rahmstorf[‡], and Hans Joachim Schellnhuber†‡||**

*School of Environmental Sciences, University of East Anglia, and Tyndall Centre for Climate Change Research, Norwich NR4 7TJ, United Kingdom; *Potsdam Institute for Climate Impact Research, P.O. Box 60 12 03, 14412 Potsdam, Germany; *Department of Engineering and Public Policy, Carnegie Mellon University, Pittsburgh, PA 15213-3890; [¶]School of Civil Engineering and Geosciences, Newcastle University, and Tyndall Centre for Climate Change Research, Newcastle NE1 7RU, United Kingdom; and Environmental Change Institute, Oxford University, and Tyndall Centre for Climate Change Research, Oxford OX1 3OY, United Kingdom


**This contribution is part of the special series of Inaugural Articles by members of the National Academy of Sciences elected on May 3, 2005.

Edited by William C. Clark, Harvard University, Cambridge, MA, and approved November 21, 2007 (received for review June 8, 2007)

PNAS










February 12, 2008 1786 – 1793 PNAS vol. 105

1995-2004 mean temperatures



Global temperature has increased by 1-1.5C in 9 years and...





...is expected to continue to increase more aggressively in the near future



Fig. 1.1 Arctic sea ice extent. Satellite imagery of sea ice extent in September








Fig 1.6 - Permafrost coverage in the northern hemisphere. Source: UNEP




Fig 1.4 Simulated future Arctic temperature trends. Regional heating of the Arctic following rapid sea ice loss events. Following such events, heating extends up to 1500km inland from the sea. Source: Steve Deyo, @University Corporation of Atmospheric Research




Fig 1.4 Simulated future Arctic temperature trends. Regional heating of the Arctic following rapid sea ice loss events. Following such events, heating extends up to 1500km inland from the sea. Source: Steve Deyo, @University Corporation of Atmospheric Research






Fig 1.5 - Carbon content. Volumes of total carbon content estimated in billion tonnes. Sources: Schuur et al., UNEP, CDIAC.




A Miserable Alternative: Surviving in a + 5°C World

Stabilizing, at last,...




... carrying capacity estimates below 1 billion people.

Business as Usual, Scramble, Blueprints



Shell Energy Scenarios to 2050 (2008), with extrapolations to 2011 for MIT Climate Review

Business as UsualScramble Blueprints Better Blueprints?




"India Magazia" aure is sughiy dia 2016 pedadiliy care Is day adhin 275 maning (PTZ-P manasaris speri)



Coal is not clean





Nor are the fleets



Natural Gas versus Coal and Oil




A perfect storm to disrupt business as usual

- 1. Enormous new supply of Natural Gas
- Huge volume of installed, highly efficient and available natural gas power generating capacity
 - 100 to 400 million tonnes of displacable CO_2 5% to 19% of total 2,000 million tonnes of coal emissions (plus SOx, NOx, Mercury)
 - 1.2 to 4.6 TCF (5% to 20%) of increased demand for Natural Gas

Switch the capacity curves

Figure 9. Monthly Capacity Factors in 2007 for Study Group Coal and NGCC Plants




Source: Calculated by CRS from the EIA-906/920 and EIA-860 databases.


Notes: For information on the study groups of coal and NGCC plants, see footnote 11. NGCC= natural gas combined cycle.

Switch the capacity curves

Figure 9. Monthly Capacity Factors in 2007 for Study Group Coal and NGCC Plants



Renewable Portfolio standards adopted by states




- Renewable Portfolio Standard
- Alternative Energy Portfolio Standard
- Renewable or Alternative Energy Goal

http://www.pewclimate.org

Today

Estimated U.S. Energy Use in 2007: ~101.5 Quads






Source: LLNL 2008. Data is based on DOE/EIA-0384(2007), June 2008. If this information or a reproduction of it is used, credit must be given to the Lawrence Livermore National Laboratory and the Department of Energy, under whose auspices the work was performed. Distributed electricity represents only retail electricity sales and does not include self-generation. EIA reports flows for non-thermal resources (i.e., hydro, wind and solar) in BTU-equivalent values by assuming a typical fossil fuel plant "heat rate." The efficiency of electricity production is calculated as the total retail electricity delivered divided by the primary energy input into electricity generation. End use efficiency is estimated as 80% for the residential, commercial and industrial sectors, and as 25% for the transportation sector. Totals may not equal sum of components due to independent rounding. LLNL-MI-410527


5-10 years from now


~101.5 Quads

Lawrence Livermore National Laboratory



Source: LLNL 2008. Data is based on DOE/EIA-0384(2007), June 2008. If this information or a reproduction of it is used, credit must be given to the Lawrence Livermore National Laboratory and the Department of Energy, under whose auspices the work was performed. Distributed electricity represents only retail electricity sales and does not include self-generation. EIA reports flows for non-thermal resources (i.e., hydro, wind and solar) in BTU-equivalent values by assuming a typical fossil fuel plant "heat rate." The efficiency of electricity production is calculated as the total retail electricity delivered divided by the primary energy input into electricity generation. End use efficiency is estimated as 80% for the residential, commercial and industrial sectors, and as 25% for the transportation sector. Totals may not equal sum of components due to independent rounding. LLNL-MI-410527






Wall Street Journal: Blowout Occurs at Pennsylvania Gas Well

DEP orders Texas company to cease gas drilling in Pennsylvania

Bloomberg: Drilling Rules Change May Harm Industry, Exxon Says



Blast at Abandoned W. Va. Coal Mine Injures 7
Flames Shoot 70 Feet in Air as Workers Drilling Gas Well Hit Pocket of Methane in Old Coal Mine Chamber

Surface impacts on:


- 1.Surface water
- 2.Shallow Groundwater
- 3.Biodiversity and Ecology
- **4.Local Communities**

The Jonah Gas Field


Located in Wyoming's Upper Green River Valley, Jonah contains an estimated 7 to 10 TCF. In the last 10 years, the field turned from sagebrush into one of the nation's richest gas fields with 500 wells. A new proposal would add 3,100 more wells.

The image was acquired September 3, 2003, covers an area of 10.7 x 13.5 km NASA/GSFC/METI/ERSDAC/ JAROS, and U.S./Japan ASTER Science Team









IN INTERNATIONAL VOY CONTANT PRODUCTION
PRESENTS A FILM OF JOSE FOR
"EASTAND"
EDITED OF MATTHEY SANCIES EXECUTIVE PRODUCEDS DESMA VINSES AND HONIES SENAY
PRODUCED OF TRISS ADDRESSES OF THE SOURCE OF TRISS ADDRESSES OF THE ATTER PARK CITY
CHECK LISTINGS OF TEMPLE THEATER, PARK CITY
CHECK LISTINGS OF OTHER SCREENINGS CHECK LISTINGS FOR OTHER SCREENINGS

JANUARY 24TH...

SUNDANCE



Align Goals

Regulators

Reduced Risks to as low as reasonably practical

Industry

Regulatory Efficiency

Reduce costs

NGOs

Public Assurance that industry is

responsible

"show me"


Simple model



Control

Recover

Program Timeline



Lead the change

