


Impacts to Air from Marcellus Shale Gas Extraction Operations OGAP Conference _ November 19, 2010

Conrad (Dan) Volz, DrPH, MPH
Assistant Professor, University of Pittsburgh, Graduate School
of Public Health
Director—Center for Healthy Environments and Communities
(CHEC)

Potential Shale Gas Extraction Air Pollution Impacts

1. How organic compounds in the shale layer enter air and become Hazardous Air Pollutants

- ▶ good evidence that flowback and produced water from shale layers themselves contain organic compounds that could volatilize into the environment when brought to the surface




1. How organic compounds in the shale layer can enter air and become Hazardous Air Pollutants

- ▶ Gas is of thermogenic or biogenic origin and stored as sorbed hydrocarbons, as free gas in fracture and intergranular porosity, and as gas dissolved in kerogen and bitumen (Schettler and Parmely, 1990; Martini et al., 1998).
- ▶ Kerogen and bitumen are extremely large molecular weight and a diverse group of organic compounds.




1. How organic compounds in the shale layer can enter air and become Hazardous Air Pollutants

- ▶ The [USGS factsheet 2009-3032](#) states clearly that hydrofrac water “in close contact with the rock during the course of the stimulation treatment, and when recovered may contain a variety of formation materials, including brines, heavy metals, radionuclides, and organics that can make wastewater treatment difficult and expensive” to dispose of.




Organic Compounds Come From the Shale Layer Itself

- ▶ Certainly gas shales contain numerous organic hydrocarbons; we know, for example, that the Marcellus contains from 3–12% organic carbon (OC), the Barnett: 4.5% OC, and the Fayetteville: 4–9.8% OC (Arthur et al, 2008).
- ▶ Volatile hydrocarbons occur naturally in produced water and that produced water from gas–condensate–producing platforms contains higher concentrations of organic compounds than from oil–producing platforms (Veil et al., 2004).



Organic Compounds Come From the Shale Layer Itself


- ▶ Produced waters from gas production have higher contents of low molecular-weight aromatic hydrocarbons such as benzene, toluene, ethylbenzene, and xylene than those from oil operations.
- ▶ Produced water contains: aliphatic and aromatic carboxylic acids, phenols, and aliphatic and aromatic hydrocarbons. They are not easily removed from produced water and are generally discharged directly into fracing ponds.



Organic Chemicals in Flowback and Produced Water May Also Come From Chemical Additives

- ▶ Chemicals added to produced water or put into a producing well – such as corrosion and scale inhibitors, scale solvents, biocides, antifreeze, and oil and grease, and impurities in the chemicals used.
- ▶ Further, some paraffin's and aromatics have moderate solubility in water; as long as oil-gas and water flow upward together these can become dissolved in water. The longer the transit time (as in deep Marcellus wells) the more hydrocarbon can dissolve into water. This paper reports finding toluene, ethylbenzene, [phenol](#), [naphthalene](#) and [2,4-dimethylphenol](#) in produced water and states that [bis\(2-ethyl-hexyl\) phthalate](#), [di-n-butyl phthalate](#), [fluorine](#) and [diethyl phthalate](#) have been found in produced water by the EPA.

Fang CS, 1990,
[Petroleum drilling and production operations in the Gulf of Mexico. Estuaries](#), Vol 13, No 1, pp. 89–97.



How do organic compounds in gas extraction waters enter air?

First a review of Henry's Law

Henry's Law states that the solubility of a gas in a liquid is a function of the partial pressure of the gas above that liquid.

The concentration of the gas in the liquid is proportional to its concentration in the atmosphere with the Henry's Law Constant describing the relationship.

Every organic compound has a unique Henry's Law constant, m (dimensionless), or K_h (pressure-mass per volume) at a specific temperature and pressure (remember $PV=nRT$) so this constant is critical to predicting volatilization of organic chemical from the water phase into the air phase.

The equations corresponding to the above two constants are;

$$C_{air} = m * C_{water}$$

$$P_v = K_h * C_{water}$$

where;

C_{air} : concentration in headspace, (mass/volume)

C_{water} : concentration in water, (mass/volume)

P_v : partial pressure in vapor phase


K_h : Henry's Law constant

m : dimensionless Henry's Law constant




How do organic compounds in gas extraction waters enter air?

- ▶ Flowback or produced water that returns to the surface and goes into a frac pond–pit or impoundment will offgas (become a vapor in air) its organic compounds into the air.
- ▶ Each organic compound enters air according to its Henry's Law constant, its concentration in the water and its partial pressure in air.
- ▶ This is dependant on T and P–all things being equal more volatilization will occur on hot sunny days with low humidity.
- ▶ This conceptually becomes an air pollution problem, and the organic compounds are now termed Hazardous Air Pollutants (HAP's).
- ▶ We have little data now on species of organic chemicals in air as a result of this process–research needs to be done.
- ▶ Indications from other shale plays are that there are contributions from the processes to air– but there is controversy regarding if these levels can produce health effects.




How do organic compounds in gas extraction waters enter air?

- ▶ Additionally, separators, condensers, cryo plants and compressors can leak causing some volatile organic compounds to enter air. Incomplete combustion in flaring also adds VOC's to air.
- ▶ CHEC is right now doing UV-DOAS spectrophotometry of gas extraction processes in the Marcellus to determine the concentrations and species of organic chemicals that may be given off by these processes.




How volatile organic compounds can act as precursor chemicals for the formation of ozone


- ▶ Ozone is a secondary pollutant that is formed in polluted areas by atmospheric reactions involving two main types of precursor pollutants volatile organic compounds (VOC's) and nitrogen oxides (NO_x).
- ▶ Carbon monoxide (CO) from incomplete combustion of fuels is also an important precursor for ozone formation.





How volatile organic compounds can act as precursor chemicals for the formation of ozone

- ▶ The formation of ozone and other oxidation products (like peroxyacyl nitrates and hydrogen peroxide), including oxidation products of the precursor chemicals, is an extremely complex reaction that depends on the intensity and wavelength of sunlight, atmospheric mixing and interactions with cloud and other aerosol particulates, the concentrations of the VOC's and NO_x in the air, and the rates of all the chemical reactions.







Existing Ozone Problems in 4 state area-pr-Marcellus Extraction




Existing NO2 Concentrations Over 4 State Area—Means of 1–Hour Values Over 1998–2008 (influence of Marcellus Shale would not be significant yet—data given to get approximation of NO2 that is available to react with volatile organics from gas operations)



Emitters of NO₂ in 2002 by Tonnage Category (Does not include mobile sources of NO₂ but does give major sources of NO₂ that are available to react with volatile organics from gas operations—to form ozone)



Sublette County, Wyoming– Robert Field, PhD

- ▶ In 2005 wintertime ozone was found to be a new phenomenon in Sublette County, Wyoming.
 - ▶ Ozone episodes are associated with **specific meteorological conditions and local oil and gas development.**
 - ▶ The Wyoming Department of Environmental Quality has developed a wide range of monitoring and modeling studies.
 - ▶ The University of Wyoming helped determine the scope of ozone episodes through large-scale passive sampling surveys performed through collaboration with stakeholders, including local, county, state, and federal authorities along with energy companies and citizens.
 - ▶ A mobile laboratory has also provided important information at locations not covered by established monitoring sites.
 - ▶ **Modeling done by Dr Field clearly shows daily ozone levels building with solar radiation in the Sublette County gas fields.**
- 


TCEQ Barnett Shale Air Monitoring of One Location

Detected in parts per billion by volume (ppbv)

Ethane ;ethylene ;propane;

propylene;dichlorodifluoromethane; methyl
chloride; isobutane; n-butane; isopentane; n-
pentane; 2-methylpentane ;


n-hexane; chloroform;methylcyclopentane ;
benzene; carbon tetrachloride; 3-
methylhexane; 2,2,4-trimethylpentane; 2-
chloropentane; toluene



PA DEP Preliminary/Snapshot Study – Mr. Nick Lazor, Chief, Air Quality Division

- ▶ PA DEP released a report on a five-week air quality study conducted near Marcellus Shale natural gas operations in Greene and Washington counties.
- ▶ DEP's assessment focused on concentrations of volatile organic compounds, including benzene, toluene and xylene, which are typically found in petroleum products. The department also sampled for other pollutants including carbon monoxide and nitrogen dioxide near natural gas extraction and processing sites.
- ▶ The agency gathered samples to provide background data at its monitoring station in Florence, a section of Hanover Township, Washington County.
- ▶ The air monitoring surveys near natural gas operations were conducted at a wastewater impoundment, tank farm and two compressor stations.
- ▶ Those surveys detected the main constituents of natural gas—including methane, ethane, propane and butane—as well as low levels of associated compounds, including benzene and n-hexane, which were detected infrequently at the tank farm and at a compressor station.
- ▶ Higher concentrations of the main constituents of natural gas were detected mainly near the compressor stations.
- ▶ Methyl mercaptan, a gas which has a penetrating and unpleasant odor similar to rotten cabbage or rotten eggs, was also detected at concentrations that generally produce odors at each location where samples were taken.

▶



VOLZ NOTE – THE METHODOLOGY OF THE DEP STUDY DOES NOT ALLOW FOR VALID CONCLUSIONS REGARDING PERSONAL EXPOSURE TO AIR TOXICS OR FOR ASSESSMENT OF STANDARD ATTAINMENT– THIS WAS A PRELIMINARY STUDY TO UNDERSTAND THE RELATIVE CONCENTRATIONS OF SELECTED SPECIES OF AIR CONTAMINANTS.

IT WAS INTENDED TO BE A SURVEY STUDY

