

Table I-22
AGE-ADJUSTED U.S. DEATH RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

Both Sexes

	All Races		White			Black		
	Rate ^b 2000-2003	APC ^c 1994-2003	Rate ^b 2000-2003	APC ^c 1994-2003	Rate ^b 2000-2003	APC ^c 1994-2003	Rate ^b 2000-2003	APC ^c 1994-2003
All Sites	194.5	-1.1*	192.4	-1.0*	241.7	-1.6*		
Lung and Bronchus	55.1	-0.9*	55.3	-0.7*	62.5	-1.6*		
Colon and Rectum	19.8	-1.9*	19.3	-2.0*	27.3	-1.0*		
Breast	14.6	-2.5*	14.3	-2.6*	22.4	-3.2*		
Prostate ^e	10.7	-3.5*	10.3	0.2	20.4	-1.4*		
Pancreas	10.5	0.0	9.8	-3.5*	13.8	-0.8*		
Non-Hodgkin Lymphoma	7.7	-1.9*	8.1	-1.9*	8.4	-3.0*		
Leukemia	7.5	-0.6*	7.8	-0.5*	7.2	-1.2*		
Ovary ^e	5.1	-0.5*	5.2	-0.5	6.7	-1.2*		
Liver & IBD ^f	4.8	1.8*	4.8	-0.9*	6.3	1.1*		
Brain and ONS ^f	4.5	-1.0*	4.5	-0.1	6.1	-4.3*		
Esophagus	4.4	0.4*	4.4	1.8*	5.3	-1.7*		
Urinary Bladder	4.3	-0.3*	4.3	1.4*	4.4	-0.4		
Stomach	4.3	-3.1*	4.3	-0.2	4.3	0.4		
Kidney and Renal Pelvis	4.2	-0.3	3.8	-3.2*	4.1	-0.5		
Myeloma	3.8	-0.7*	3.5	-0.6*	3.9	-4.2*		
	Asian/Pacific Islander		American Indian/Alaska Native			Hispanic ^d		
	Rate ^b 2000-2003	APC ^c 1994-2003	Rate ^b 2000-2003	APC ^c 1994-2003	Rate ^b 2000-2003	APC ^c 1994-2003	Rate ^b 2000-2003	APC ^c 1994-2003
All Sites	117.0	-1.8*	127.0	-1.4*	131.0	-0.9*		
Lung and Bronchus	27.2	-1.3*	33.1	-2.0*	23.9	-1.3*		
Colon and Rectum	12.6	-1.7*	13.0	-1.9	13.8	-0.5*		
Liver & IBD ^f	10.7	-0.5	7.5	-2.0	9.0	-2.3*		
Stomach	8.3	-4.2*	7.1	-3.9*	8.5	-2.9*		
Pancreas	7.2	-1.1*	6.0	-0.7	8.3	0.1		
Breast	7.0	-0.8	5.9	1.0	7.6	2.0*		
Non-Hodgkin Lymphoma	4.8	-1.9*	5.1	-1.6	6.8	-1.8*		
Prostate ^e	4.6	-6.1*	4.7	0.1	5.9	-2.1*		
Leukemia	3.9	-2.3*	4.4	-1.9	5.2	-0.6		
Ovary ^e	2.8	0.8	3.9	0.0	3.6	0.3		
Oral Cavity and Pharynx	2.4	-1.6*	3.1	1.8	3.5	0.0		
Brain and ONS ^f	1.9	-0.2	2.9	0.0	3.2	0.1		
Urinary Bladder	1.8	0.5	2.9	1.1	2.9	0.5		
Kidney and Renal Pelvis	1.8	-1.9	2.2	-1.1	2.4	0.3		
Esophagus	1.8	-3.7*	2.0	-0.3	2.4	-1.5*		

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 1992-2002 age-adjusted rates for the race/ethnic group.

^b Mortality data used in calculating the rates are analyzed from a public use file provided by the National Center for Health Statistics (NCHS). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). The rates shown for sex-specific cancer sites are calculated using the population for both sexes combined.

^c The APC is the Annual Percent Change over the time interval. Mortality data used in calculating the trends are analyzed from a public use file provided by the National Center for Health Statistics (NCHS). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. The 2000-2003 Hispanic death rates do not include deaths from Minnesota, New Hampshire and North Dakota. The 1994-2003 Hispanic mortality trends do not include deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

^e The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^f IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* The APC is significantly different from zero ($p < .05$).