



## Complete Summary

---

### GUIDELINE TITLE

Prevention of rotavirus gastroenteritis among infants and children.  
Recommendations of the Advisory Committee on Immunization Practices (ACIP).

### BIBLIOGRAPHIC SOURCE(S)

Parashar UD, Alexander JP, Glass RI, Advisory Committee on Immunization Practices (ACIP), Centers for Disease Control and Prevention. Prevention of rotavirus gastroenteritis among infants and children. Recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR Recomm Rep 2006 Aug 11;55(RR-12):1-13. [85 references] [PubMed](#)

### GUIDELINE STATUS

This is the current release of the guideline.

### \*\* REGULATORY ALERT \*\*

### FDA WARNING/REGULATORY ALERT

**Note from the National Guideline Clearinghouse:** This guideline references a drug(s) for which important revised regulatory and/or warning information has been released.

- [June 15, 2007, RotaTeq \(Rotavirus, Live, Oral, Pentavalent Vaccine\)](#): Changes to the ADVERSE REACTIONS and POST-MARKETING sections of the product's prescribing information. The ADVERSE REACTIONS section was updated to include six cases of Kawasaki disease that were observed during the Phase 3 clinical trial.
- [February 13, 2007, Rotavirus, Live, Oral, Pentavalent Vaccine \(RotaTeq\)](#): FDA Public Health Notification regarding 28 post-marketing reports of intussusception following administration of Rotavirus, Live, Oral, Pentavalent vaccine (RotaTeq).

### COMPLETE SUMMARY CONTENT

\*\* REGULATORY ALERT \*\*

SCOPE

METHODOLOGY - including Rating Scheme and Cost Analysis

RECOMMENDATIONS

EVIDENCE SUPPORTING THE RECOMMENDATIONS

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

CONTRAINDICATIONS

QUALIFYING STATEMENTS

## SCOPE

### **DISEASE/CONDITION(S)**

Rotavirus gastroenteritis

### **GUIDELINE CATEGORY**

Prevention

### **CLINICAL SPECIALTY**

Family Practice  
Infectious Diseases  
Pediatrics  
Preventive Medicine

### **INTENDED USERS**

Advanced Practice Nurses  
Health Care Providers  
Nurses  
Physician Assistants  
Physicians  
Public Health Departments

### **GUIDELINE OBJECTIVE(S)**

To provide recommendations on the use of rotavirus vaccine in infants and children

### **TARGET POPULATION**

Infants from 6 to 36 weeks

### **INTERVENTIONS AND PRACTICES CONSIDERED**

Immunization with live, oral, human-bovine reassortant rotavirus vaccine (RotaTeq®)

### **MAJOR OUTCOMES CONSIDERED**

- Incidence of rotavirus gastroenteritis
- Adverse effects

## METHODOLOGY

### METHODS USED TO COLLECT/SELECT EVIDENCE

Searches of Electronic Databases

### DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

Not stated

### NUMBER OF SOURCE DOCUMENTS

Not stated

### METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

### RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

#### Level of Evidence

- I. Evidence from randomized controlled trials
- II. Evidence from other epidemiologic studies
- III. Opinions of authorities

#### Strength of Evidence

- A. Good evidence to support recommendation
- B. Fair evidence to support recommendation
- C. Insufficient evidence

### METHODS USED TO ANALYZE THE EVIDENCE

Review

### DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

### METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

### RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

## **COST ANALYSIS**

In an analysis that used estimates of current rotavirus disease burden, vaccine efficacy, vaccine coverage rates, and health costs, investigators estimated that a national rotavirus vaccination program in which 3 doses of RotaTeq® are administered at ages 2, 4, and 6 months would result in 255,000 fewer physician visits, 137,000 fewer emergency department (ED) visits, 44,000 fewer hospitalizations, and 13 fewer deaths per year in children aged <5 years. From the health-care perspective alone, vaccination is likely to be cost-saving at a total cost per child (including administration costs) of up to \$66 per child (approximately \$12 per vaccine dose). A higher-priced vaccine would be increasingly unlikely to be cost-saving, and at a cost of more than \$143 per child (approximately \$38 per dose), a rotavirus vaccination program would most likely have a net cost to the health-care system. From the societal perspective, vaccination is likely to be cost-saving at a total cost per child of up to \$156 per child (approximately \$42 per dose). A higher-priced vaccine would be increasingly unlikely to be cost-saving, and at a cost of more than \$268 per child (approximately \$79 per dose), a rotavirus vaccination program would most likely have a net cost to society.

## **METHOD OF GUIDELINE VALIDATION**

Peer Review

## **DESCRIPTION OF METHOD OF GUIDELINE VALIDATION**

Not stated

## **RECOMMENDATIONS**

### **MAJOR RECOMMENDATIONS**

The levels of evidence (I–III) and strength of evidence (A–C) supporting the recommendations are defined at the end of the "Major Recommendations" field.

### **Recommendations for the Use of Rotavirus Vaccine**

#### **Routine Administration**

Advisory Committee on Immunization Practices (ACIP) recommends routine vaccination of U.S. infants with 3 doses of rotavirus vaccine administered orally at ages 2, 4, and 6 months (see Table below titled "Recommendations and quality of evidence for recommendations for use of rotavirus vaccine"). The first dose should be administered between ages 6–12 weeks. Subsequent doses should be administered at 4–10-week intervals, and all 3 doses of vaccine should be administered by age 32 weeks. Vaccination should not be initiated for infants aged >12 weeks because of insufficient data on safety of the first dose of rotavirus vaccine in older infants. Vaccine should not be administered after age 32 weeks because of insufficient data on the safety and efficacy of rotavirus vaccine in infants after this age. For infants in whom the first dose of rotavirus vaccine is inadvertently administered off label at age  $\geq 13$  weeks, the rest of the rotavirus

vaccination series should be completed as per the schedule because timing of the first dose should not affect the safety and efficacy of the second and third dose. Infants who have had rotavirus gastroenteritis before receiving the full course of rotavirus vaccinations should still initiate or complete the 3-dose schedule because the initial infection frequently provides only partial immunity.

Infants who are being breastfed can receive rotavirus vaccine. The efficacy of rotavirus vaccine is similar among breastfed and nonbreastfed infants. Like other vaccines, rotavirus vaccine can be administered to infants with transient, mild illnesses, and with or without low-grade fever.

### **Simultaneous Administration**

Rotavirus vaccine can be administered together with diphtheria, tetanus, and pertussis vaccine (DTaP), *Haemophilus influenzae* type b conjugate (Hib) vaccine, inactivated poliovirus vaccine (IPV), hepatitis B vaccine, and pneumococcal conjugate vaccine. Available evidence suggests that the rotavirus vaccine does not interfere with the immune response to the Hib vaccine, IPV, hepatitis B vaccine, and pneumococcal conjugate vaccine, and the diphtheria and tetanus antigens in DTaP. Because validation of the pertussis assays is still under review, insufficient immunogenicity data are available to confirm lack of interference of immune responses when rotavirus vaccine is concomitantly administered with childhood vaccines to prevent pertussis.

### **Contraindications**

Rotavirus vaccine should not be administered to infants who have severe hypersensitivity to any component of the vaccine or who have experienced a serious allergic reaction to a previous dose of rotavirus vaccine.

### **Precautions**

#### *Altered Immunocompetence*

Practitioners should consider the potential risks and benefits of administering rotavirus vaccine to infants with known or suspected altered immunocompetence. Children and adults who are immunocompromised because of congenital immunodeficiency, hematopoietic transplantation, or solid organ transplantation sometimes experience severe, prolonged, and even fatal rotavirus gastroenteritis. However, no safety or efficacy data are available for the administration of rotavirus vaccine to infants who are potentially immunocompromised, including

- Infants with blood dyscrasias, leukemia, lymphomas of any type, or other malignant neoplasms affecting the bone marrow or lymphatic system
- Infants on immunosuppressive therapy (including high-dose systemic corticosteroids)
- Infants with primary and acquired immunodeficiency states, including human immunodeficiency virus/acquired immunodeficiency syndrome (HIV/AIDS) or other clinical manifestations of infection with HIV; cellular immune deficiencies; and hypogammaglobulinemic and dysgammaglobulinemic states. Data are insufficient from the clinical trials to support administration of

rotavirus vaccine to infants with indeterminate HIV status who are born to mothers with HIV/AIDS.

- Infants who have received a blood transfusion or blood products, including immunoglobulins, within 42 days. In general, rotavirus vaccine should be deferred for 42 days following receipt of an antibody-containing product if possible. However, if the 42-day deferral would cause the first dose of rotavirus vaccine to be scheduled for age  $\geq 13$  weeks, a shorter deferral interval should be used to ensure the first dose is administered before age 13 weeks.

### *Acute Gastroenteritis*

In usual circumstances, rotavirus vaccine should not be administered to infants with acute, moderate-to-severe gastroenteritis until the condition improves. However, infants with mild acute gastroenteritis can be vaccinated, particularly if the delay in vaccination might be substantial and might make the child ineligible to receive vaccine (e.g., aged  $\geq 13$  weeks before vaccination is initiated).

Rotavirus vaccine has not been studied among infants with concurrent acute gastroenteritis. In these infants, the immunogenicity and efficacy of rotavirus vaccine can theoretically be compromised. For example, infants who receive oral poliovirus vaccine (OPV) during an episode of acute gastroenteritis in some instances have diminished poliovirus antibody responses to OPV.

### *Moderate-to-Severe Illness*

Infants with moderate-to-severe illness should be vaccinated as soon as they have recovered from the acute phase of the illness. This precaution avoids superimposing adverse effects of the vaccine on the underlying illness or mistakenly attributing a manifestation of the underlying illness to the vaccine.

### *Preexisting Chronic Gastrointestinal Disease*

Practitioners should consider the potential risks for and benefits of administering rotavirus vaccine to infants with preexisting chronic gastrointestinal disease. Infants with preexisting chronic gastrointestinal conditions who are not undergoing immunosuppressive therapy should benefit from rotavirus vaccine vaccination, and the benefits outweigh the theoretical risks. However, the safety and efficacy of rotavirus vaccine have not been established for infants with these preexisting conditions (e.g., congenital malabsorption syndromes, Hirschsprung's disease, short-gut syndrome, or persistent vomiting of unknown cause).

### *Intussusception*

Following administration of a previously licensed rotavirus vaccine, RRV-TV, an increased risk for intussusception was observed. Available prelicensure data from a trial of 70,000 infants indicated no evidence of an association between intussusception and the current vaccine. However, additional postlicensure surveillance data are required to confirm that the vaccine is not associated with intussusception at a lower rate than would have been detected in prelicensure trials. In addition, data suggest that infants with a history of intussusception

might be at higher risk for a repeat episode than other infants. Therefore, until postlicensure data on safety of rotavirus vaccine are available, the risks for and the benefits of vaccination should be considered when vaccinating infants with a previous episode of intussusception.

### **Special Situations**

#### *Premature Infants (<37 weeks' gestation)*

Practitioners should consider the potential risks for and benefits of vaccinating premature infants against rotavirus. Limited data suggest that premature infants are at increased risk for hospitalization from viral gastroenteritis during their first year of life. In clinical trials, the safety and efficacy of rotavirus vaccine appears to be similar for premature and term infants, although a relatively small number of preterm infants have been evaluated. The lower level of maternal antibody to rotaviruses in very low birthweight, premature infants theoretically could increase the risk for adverse reactions from rotavirus vaccine. ACIP supports vaccination of prematurely born infants if they are at least aged 6 weeks, are being or have been discharged from the hospital nursery, and are clinically stable. Until further data are available, ACIP considers that the benefits of rotavirus vaccine vaccination of premature infants outweigh the theoretical risks.

#### *Exposure of Immunocompromised Persons to Vaccinated Infants*

Infants living in households with persons who have or are suspected of having an immunodeficiency disorder or impaired immune status can be vaccinated. The majority of experts believe the protection of the immunocompromised household member afforded by vaccination of young children in the household outweighs the small risk for transmitting vaccine virus to the immunocompromised household member and any subsequent theoretical risk for vaccine virus-associated disease. To minimize potential virus transmission, all members of the household should employ measures such as good hand washing after contact with the feces of the vaccinated infant (e.g., after changing a diaper).

#### *Exposure of Pregnant Women to Vaccinated Infants*

Infants living in households with pregnant women can be vaccinated. The majority of women of childbearing age would have pre-existing immunity to rotavirus and so the risk for infection and disease from potential exposure to the attenuated vaccine virus strain is low. In addition, no evidence exist that rotavirus infection or disease during pregnancy poses any risk to the fetus. Furthermore, vaccination of young children would avoid potential exposure of the pregnant women to wild virus if the unvaccinated infant suffers from rotavirus gastroenteritis.

#### *Regurgitation of Vaccine*

The practitioner should not readminister a dose of rotavirus vaccine to an infant who regurgitates, spits out, or vomits during or after administration of vaccine. The infant can receive the remaining recommended doses of rotavirus vaccine at appropriate intervals. Data are limited regarding the safety of administering a dose of rotavirus vaccine higher than the recommended dose and on the efficacy

of administering a partial dose. Additional data on safety and efficacy are needed to evaluate the benefits of and risks for readministration.

#### *Hospitalization After Vaccination*

If a recently vaccinated child is hospitalized for any reason, no precautions other than routine universal precautions need be taken to prevent the spread of vaccine virus in the hospital setting.

### **Recommendations and Quality of Evidence for Recommendations for Use of Rotavirus Vaccine**

	<b>Level of Evidence</b>	<b>Strength of Evidence</b>
<b>Recommendations</b>		
Routine vaccination at ages 2, 4, and 6 months	I	A
Administer to breastfed infants	I	A
Co-administer with DTaP, Hib vaccine, IPV, hepatitis B vaccine, and pneumococcal conjugate vaccine	I	A
Administer to infants with mild illness	I	B
<b>Contraindications</b>		
Serious allergy to a vaccine component or a previous vaccine dose	III	B
<b>Precautions</b>		
Altered immunocompetence	III	C
Moderate-to-severe illness, including acute gastroenteritis	III	C
Chronic gastrointestinal disease	III	C
History of intussusception	III	C
<b>Special Situations</b>		
Premature infants (aged <37 weeks)	I	B
Infants living in households with immunocompromised persons	III	C
Infants living in households with pregnant women	III	C
Regurgitation of vaccine	III	C
Children hospitalized after vaccination	III	C

#### **Definitions:**

##### **Level of Evidence**

- I. Evidence from randomized controlled trials
- II. Evidence from other epidemiologic studies
- III. Opinions of authorities

##### **Strength of Evidence**

- A. Good evidence to support recommendation
- B. Fair evidence to support recommendation



C. Insufficient evidence

## **CLINICAL ALGORITHM(S)**

None provided

## **EVIDENCE SUPPORTING THE RECOMMENDATIONS**

### **TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS**

The type of supporting evidence is graded and identified for select recommendations (see "Major Recommendations").

## **BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS**

### **POTENTIAL BENEFITS**

Prevention of rotavirus gastroenteritis and associated morbidity and mortality among infants and children

### **POTENTIAL HARMS**

A discussion of potential harms post-vaccination, including intussusception and other adverse events, can be found in the original guideline document.

## **CONTRAINDICATIONS**

### **CONTRAINDICATIONS**

Refer to "Major Recommendations" field for a description of contraindications to vaccine administration.

## **QUALIFYING STATEMENTS**

### **QUALIFYING STATEMENTS**

References to non-Centers for Disease Control (CDC) sites on the Internet are provided as a service to Morbidity and Mortality Weekly Report (MMWR) readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in MMWR were current as of the date of publication.

## **IMPLEMENTATION OF THE GUIDELINE**

### **DESCRIPTION OF IMPLEMENTATION STRATEGY**

An implementation strategy was not provided.

## INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

### IOM CARE NEED

Staying Healthy

### IOM DOMAIN

Effectiveness

## IDENTIFYING INFORMATION AND AVAILABILITY

### BIBLIOGRAPHIC SOURCE(S)

Parashar UD, Alexander JP, Glass RI, Advisory Committee on Immunization Practices (ACIP), Centers for Disease Control and Prevention. Prevention of rotavirus gastroenteritis among infants and children. Recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR Recomm Rep 2006 Aug 11;55(RR-12):1-13. [85 references] [PubMed](#)

### ADAPTATION

Not applicable: The guideline was not adapted from another source.

### DATE RELEASED

2006 Aug 11

### GUIDELINE DEVELOPER(S)

Centers for Disease Control and Prevention - Federal Government Agency [U.S.]

### SOURCE(S) OF FUNDING

United States Government

### GUIDELINE COMMITTEE

Advisory Committee on Immunization Practices Rotavirus Vaccine Work Group  
Advisory Committee on Immunization Practices

### COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

*Advisory Committee on Immunization Practices Rotavirus Vaccine Work Group*  
*Members:* John Treanor, MD (*Chair*) Rochester, New York; James Alexander, MD, Atlanta, Georgia; Umesh Parashar, MBBS, Atlanta, Georgia; Marc-Alain Widdowson, MA, Atlanta, Georgia; Rosemary Tiernan, Bethesda, Maryland; Hector Izurieta, MD, Rockville Maryland; Geoffrey Evans, MD, Rockville, Maryland; Roger

Glass, MD, Atlanta, Georgia; Edgar Marcuse, MD, Seattle, Washington; Trudy Murphy, MD, Atlanta, Georgia; Paul Gargiullo, PhD, Atlanta, Georgia; Larry Pickering, Atlanta, Georgia; Jane Seward, MD, Atlanta, Georgia; Penina Haber, Atlanta, Georgia; Greg Wallace, MD, Atlanta, Georgia; Penelope Dennehy, MD, Providence, Rhode Island; Samuel Katz, MD, Durham, North Carolina; John Modlin, MD, Lebanon, New Hampshire; and Paul Offit, MD, Philadelphia, Pennsylvania

*Advisory Committee on Immunization Practices Members:* Jon S. Abramson, MD (*Chairman*) Wake Forest University School of Medicine, Winston-Salem, North Carolina; Larry K. Pickering, MD (*Executive Secretary*) CDC, Atlanta, Georgia; Ban Mishu Allos, MD, Vanderbilt University School of Medicine, Nashville, Tennessee; Robert L. Beck, Palmyra, Virginia; Judith Campbell, MD, Baylor College of Medicine, Houston, Texas; Reginald Finger, MD, Colorado Springs, Colorado; Janet R. Gilsdorf, MD, University of Michigan, Ann Arbor, Michigan; Harry Hull, MD, Minnesota Department of Health, Minneapolis, Minnesota; Tracy Lieu, MD, Harvard Pilgrim Health Care and Harvard Medical School, Boston, Massachusetts; Edgar K. Marcuse, MD, Children's Hospital and Regional Medical Center, Seattle, Washington; Dale L. Morse, MD, New York State Department of Health, Albany, New York; Julia Morita, MD, Chicago Department of Public Health, Chicago, Illinois; Gregory A. Poland, MD, Mayo Clinic and Foundation, Rochester, Minnesota; Patricia Stinchfield, Children's Hospitals and Clinics, St. Paul, Minnesota; John J. Treanor, MD, University of Rochester, Rochester, New York; and Robin J. Womeodu, MD, University of Tennessee Health Science Center, Memphis, Tennessee

*Ex-Officio Members:* James Cheek, MD, Indian Health Service, Albuquerque, New Mexico; Wayne Hachey, DO, Department of Defense, Falls Church, Virginia; Geoffrey S. Evans, MD, Health Resources and Services Administration, Rockville, Maryland; Bruce Gellin, MD, National Vaccine Program Office, Washington, D.C.; Linda Murphy, Centers for Medicare and Medicaid Services, Baltimore, Maryland; George T. Curlin, MD, National Institutes of Health, Bethesda, Maryland; Norman Baylor, PhD, U.S. Food and Drug Administration, Rockville, Maryland; and Kristin Lee Nichol, MD, Department of Veterans Affairs, Minneapolis, Minnesota

*Liaison Representatives:* Johathan Temte, MD, Madison, Wisconsin, Doug Campos-Outcalt, MD, Phoenix, Arizona, American Academy of Family Physicians; Keith Powell, MD, Akron, Ohio, Carol Baker, MD, Houston, Texas, American Academy of Pediatrics; Andrea Gelzer, MD, Hartford, Connecticut, America's Health Insurance Plans; James C. Turner, MD, Charlottesville, Virginia, American College Health Association; Stanley Gall, MD, Louisville, Kentucky, American College of Obstetricians and Gynecologists; Kathleen M. Neuzil, MD, Seattle, Washington, American College of Physicians; Litjen Tan, PhD, Chicago, Illinois, American Medical Association; Stephan L. Foster, PharmD, Memphis, Tennessee, American Pharmacists Association; Paul W., McKinney, MD, Louisville, Kentucky, Association of Teachers of Preventive Medicine; Clement Lewin, PhD, Orange, Connecticut, Biotechnology Industry Organization; Monica Naus, MD, Vancouver, British Columbia, Canada, Canadian National Advisory Committee on Immunization; Steve Gordon, MD, Cleveland, Ohio, Healthcare Infection Control Practices Advisory Committee; Samuel L. Katz, MD, Durham, North Carolina, Infectious Diseases Society of America; David Salisbury, MD, London, England, United Kingdom, London Department of Health; Nancy Bennett, MD, Rochester,

New York, Jeffrey Duchin, MD, Seattle, Washington, National Association of County and City Health Officials; David A. Neumann, PhD, Alexandria, Virginia, National Coalition for Adult Immunization; William Schaffner, MD, Nashville, Tennessee, National Foundation for Infectious Diseases; Romeo S. Rodriguez, Mexico, National Immunization Council and Child Health Program, Mexico; Patricia Whitley-Williams, MD, New Brunswick, New Jersey, Peter Paradiso, PhD, Collegeville, Pennsylvania, National Medical Association; and Amy B. Middleman, MD, Houston, Texas, Society for Adolescent Medicine

## **FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST**

Not stated

## **GUIDELINE STATUS**

This is the current release of the guideline.

## **GUIDELINE AVAILABILITY**

Electronic copies: Available from the Centers for Disease Control and Prevention (CDC) Web site.

- [HTML Format](#)
- [Portable Document Format \(PDF\)](#)

Print copies: Available from the Centers for Disease Control and Prevention, MMWR, Atlanta, GA 30333. Additional copies can be purchased from the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402-9325; (202) 783-3238.

## **AVAILABILITY OF COMPANION DOCUMENTS**

None available

## **PATIENT RESOURCES**

None available

## **NGC STATUS**

This NGC summary was completed by ECRI on August 21, 2006. This summary was updated by ECRI on February 19, 2007 following the FDA advisory on Rotavirus, Live, Oral, Pentavalent vaccine (RotaTeq). This summary was updated by ECRI Institute on July 9, 2007 following the FDA advisory on RotaTeq (Rotavirus, Live, Oral, Pentavalent) vaccine.

## **COPYRIGHT STATEMENT**

No copyright restrictions apply.

## DISCLAIMER

### NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 10/6/2008

