

BLACK NEEDLERUSH

Juncus roemerianus Scheele

Plant Symbol = JURO

Contributed by: USDA NRCS Plant Materials
Program and USDA NRCS National Plant Data
Center:



© Larry Allain
Larry Allain, NWRC, GS, USDI

Alternate Names

black grass, Roemers rush, black rush, black
needlegrass.

Uses

Dense stands of black needlerush form deep fibrous
root systems, which provide very good shoreline
protection, filter suspended solids, uptake nutrients,
and facilitate substrate oxidation. With its range of
salinity tolerances, black needlerush is used in tidal
estuary restoration along the Atlantic and Gulf coasts.
Seed and vegetative parts of black needlerush are
utilized by waterfowl, muskrats, nutria, rice rat,
marsh rabbit and non-game birds. More than 60 bird
species use black needlerush dominated marsh at
least seasonally.

Ninety percent of the biomass of marsh plants, such
as black needlerush, is not consumed by herbivores.
Instead, marsh plant biomass is decomposed to
microbial biomass. This microbial biomass is
available to primary consumers which initiate food
webs leading to commercially important fishes and
crustaceans.

Black needlerush is commonly planted in constructed
wetlands, which are used for the treatment of dilute
organic wastes.

Researchers evaluated three marsh plant species,
black needlerush, common reed (*Phragmites
australis*) and bulltongue arrowhead (*Sagittaria
lancifolia*) for tolerance of soil contaminated with
diesel oil. Bulltongue arrowhead exhibited the
highest level of tolerance, common reed the least
tolerance and black needlerush was intermediate.

Status

Please consult the PLANTS Web site and your State
Department of Natural Resources for this plant's
current status (e.g. threatened or endangered species,
state noxious status, and wetland indicator values).

Description

General: Black needlerush is a moderate growing,
bunch forming, grass-like perennial. It is coarse and
rigid. Where soil salinity is low, black needle grass is
robust with leaves reaching over 2.2 m in height. But
in high saline areas the plants are dwarfed often less
than 0.3 m tall. Leaves are terete, stiff and pungent.
Inflorescence appear laterally; involucrel bract terete
and erect. Perianth usually brownish, 3-3.5 mm long;
sepals longer and are more pointed than petals. It is
composed of two types of plants based on flower
morphology: one form produces perfect (bisexual
flowers) and another form produces pistillate
(unisexual) flowers. Seeds are dark and 0.6 mm long
(May to October). Black needlerush is found growing
in brackish marshes in dense zonal stands.

Distribution: Black needlerush is one of the dominate
species in the marshes on the southern Atlantic and
Gulf coasts. It dominates 20.7% of the marsh in the
south Atlantic states and 7.3% of Gulf coast marshes.
But, it covers more marsh area on the Gulf coast than
on the Atlantic coast. Its distribution is continuous
from Maryland to Florida and westward to
southwestern Texas. It is usually restricted to coastal

marches and estuaries, but it may extend 10 to 15 miles inland along river estuaries.

For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

Habitat: Black needlerush occupies the edge of ditches and shorelines of bays, back bays and tributaries of tidal systems in coastal systems. Black needlerush grows on soils ranging from almost pure sand to fine silt loam and clay mixtures. It also, grows in highly organic soils including peat soils. It has a high tolerance to anaerobic conditions, high tolerance to CaCO₃ and tolerates pH ranges from 4.0-7.0

Adaptation

Black needlerush dominated plant communities are classified into three generalized categories based on elevation and salinity influences: 1) Saline marsh, which experiences little dilution of tidal waters; 2) Brackish marsh, where tidal waters are routinely diluted, 3) Intermediate marsh, which is transitional between brackish and freshwater marsh. The number of species in association with black needlerush tends to increase as water salinity decreases. However, *Spartina* species are thought to be more tolerant of flooding and salinity than black needlerush. In freshwater habitats black needlerush's growth may be restricted by organisms that feed on its rhizomes.

Salt marsh plant communities are characterized by striking zonal patterns across elevation gradients. In a study conducted in Georgia, black needlerush dominated the high elevation marsh and smooth cordgrass (*Spartina alterniflora*) dominated the middle and low elevation marsh. Black needlerush did not occur naturally in the cordgrass zone and performed poorly when transplanted there. The poor performance of black needlerush occurred whether or not cordgrass neighbors were removed, which indicates the poor performance was caused by physical stress. In contrast, although smooth cordgrass occurred naturally at low densities in the black needlerush zone, it performed well there only if the black needlerush neighbors were removed. The excellent performance of smooth cordgrass only where black needlerush neighbors were removed indicates that performance is limited by competition from black needlerush.

Establishment

In the wild, black needlerush establishes from seed, rhizomes or vegetative divisions from storm derived floatsum. Pistillate-flowered plants produce more

seed and seed with higher viability than perfect-flowered plants. Seed germination requires light and seeds will not germinate if covered. Seeds remain viable for over a year. Seeds will germinate when submerged or floating on water. Black needlerush seedlings are found mainly in areas with no or little associated vegetation. Seedlings are seldom found in mature stands of black needlerush or other marsh species. Black needlerush maintains itself in established stands though rhizome growth.

The percentage germination of seed collected in Mississippi averaged 75 for pistillate flowered plants and 60 for perfect flowered plants. Black needlerush seeds collected from the Mississippi Gulf coast exhibited high percentage germination without a cold-wet stratification. Nursery and greenhouse production are effectively accomplished with seed or vegetative divisions. Seed can be stratified in saturated soil-less potting media with high percentage of fine organics. At the USDA Natural Resources Conservation Service (NRCS) Cape May Plant Materials Center (PMC), New Jersey, seed and media were thoroughly moistened, placed in zip lock bags and stored at approximately 36 °F for 60 days. Materials were direct seeded into greenhouse starter trays with diurnal temperatures of 80° F days and 50° F nights and 14 hour photoperiods. Seedling emergence was limited and rate of growth slow.

For greenhouse forced materials, vegetative divisions use similar environmental photoperiods and diurnal temperatures listed above.

Tissue culture and plant regeneration protocols have been developed for blank needlerush.

Management

The effects of fire on blank needlerush dominated marsh vary with water depth and soil moisture. On flooded sites and sites with exposed but saturated soil, fires will not harm the underground reproductive structures. When the marsh soil is dry, fire can consume the rhizomes and the entire stand. In Mississippi, burning increased the biomass produced in a black needlerush dominated marsh, but black needlerush recovered more slowly than *Spartina* species. Three years after burning black needlerush biomass was less than before burning. In Maryland, the biomass and stem densities of several marsh species increased after burning, but burning did not change either the biomass or stem density of black needlerush. In Florida, black needlerush biomass was 47% less one year after burning. In another Florida study, winter burning increased the frequency of common three-square (*Schoenoplectus pungens* var.

pungens) when it occurred as a competing subdominant with black needlerush. But, in nearly pure stands of black needlerush burning did not change the species composition. In general, the effect of burning on black needlerush was either a decrease in biomass and frequency or no change in these parameters.

Pests and Potential Problems

In the mid-Atlantic, several major nursery pests may present themselves and required IPM strategies. These pests can include pythium, brown patch, fungus gnats and algae. This is based upon professional experiences at the NRCS Cape May PMC, New Jersey and diagnosis of pests by Rutgers University Turf Grass Department staff.

Consult your local university extension service agent for pesticide recommendations.

Cultivars, Improved, and Selected Materials (and area of origin)

No improved varieties are known, but numerous wetland nurseries carry local or regional ecotypes. Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be listed under the subheading "Department of Agriculture."

References

Britton, L. and A. Brown, 1970. *An illustrated flora of the northern United States and Canada*. Dover Press.

Brown, M.L. and R.G. Brown. 1984. *Herbaceous plants of Maryland*. Port City Press.

Eleuterius, L.N. 1975. *The life history of the salt marsh rush, Juncus roemerianus*. Bull. Torrey Pines Bot. Club. 102: 135-140.

Eleuterius, L.N. 1976. *The distribution of Juncus roemerianus in the salt marshes of North America*. Chesapeake Science 17: 289-292.

Eleuterius, L.N. 1976. *Vegetative morphology and anatomy of the salt marsh rush, Juncus roemerianus*. Gulf Research Reports 5: 1-10.

Eleuterius, L.N. 1984. *Autecology of the black needlerush Juncus roemerianus*. Gulf Research Reports 7: 27-34.

Fell, J.W. and I.L. Hunter. 1979. *Fungi associated with the decomposition of the black rush Juncus roemerianus in south Florida*. Mycologia 71: 323-342.

Fernald, M.L. 1950. *Grays manual of botany*. 8th Edition. American Book Co.

Flores, C. and D. Bonds. *Evaluation of vegetative response to fire exclusion and prescribe burning on Blackwater National Wildlife Refuge and Fishing Bay Wildlife Management Area*. Accessed: 070215. <http://www.fws.gov/blackwater/fire_research.pdf>

Gleason, H.A and A.Cronquist. 1963. *Manual of vascular plants of the United States and adjacent Canada*. Van Norstrand Reinhold Co.

Hill, D.T., W.E. Payne, J.W. Jones and S.R. Kown. 1997. *Ammonia effects on the biomass production of five constructed wetland plant species*. Bioresource Technology 62: 109-133.

Hackney, C.T. and A.A. De LaCruz. 1981. *Effects of fire on two brackish marsh communities" management implications*. Wetlands 1: 75-86.

Lin, Q, & I.A. Mendelsohn. 2006. *Determination of diesel oil tolerance of coastal dominant marsh plants for restoration and remediation of the oil-impacted habitats*. <http://www.osradp.lsu.edu/maySymp06/MaySymp06_Lin.pdf>

Meyers, K.E. 1956. *Management of needlerush marsh at the Chassahowitzka Refuge*. Proc. Annual Conference Southeast Assoc. Game and Fish Comm. 9: 175-177.

Odum, W.E., J.C. Zieman and E.J. Heald. 1973. *Importance of various plant detritus to estuaries*. Pp. 91-114. In: *Proceedings of the Coastal Marsh and Estuary Management Symposium*. Ed. R.H. Chabrech, Louisiana State Univ., Baton Rouge.

Pennings, S.C., M. Grant and M.D. Bertness. 2005. *Plant zonation in low-latitude salt marshes: disentangling the roles of flooding, salinity and competition*. Journal of Ecology 93: 159-167.

Stout, J.P. 1984. *The ecology of irregularly flooded salt marches of the Northeast Gulf of Mexico; a community profile*. Biol. Rep. 85. U.S. Dept. of Interior, Fish & Wildlife Service.

USDA-Forest Service. *Fire effect information system*
<<http://www.fs.fed.us/database/feis/>>. Accessed:
070215.

[Read about Civil Rights at the Natural Resources Conservation Service.](#)

USDA, NRCS. 2007. *The PLANTS database*
(<http://plants.usda.gov>, 31 January 2007). National
Plant Data Center, Baton Rouge, LA 70874-4490
USA.

Wang, J., D.M. Seliskar and J.L. Gallagher. 2005.
*Tissue culture and plant regeneration of the sast
marsh monocots Juncus roemerianus and Juncus
gerardi.*
<[http://cat.inist.fr/?aModele=afficheN&cpsid=16953
946](http://cat.inist.fr/?aModele=afficheN&cpsid=16953946)>.

Prepared By

William Skaradek
USDA NRCS Cape May Plant Materials Center,
Cape May, New Jersey
James F. Henson
USDA NRCS National Plant Data Center, Baton
Rouge, Louisiana

Species Coordinator

William Skaradek
USDA NRCS Cape May Plant Materials Center,
Cape May, New Jersey

Document Reviewer:

Joel Douglas
USDA NRCS Central National Technology Support
Center, Fort Worth, Texas
Robert Glennon
USDA NRCS National Headquarters, Washington,
D.C.

Edited: 070227.jsp

For more information about this and other plants, please contact
your local NRCS field office or Conservation District, and visit the
PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials
Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

*The U.S. Department of Agriculture (USDA) prohibits
discrimination in all its programs and activities on the basis of
race, color, national origin, sex, religion, age, disability, political
beliefs, sexual orientation, and marital or family status. (Not all
prohibited bases apply to all programs.) Persons with disabilities
who require alternative means for communication of program
information (Braille, large print, audiotape, etc.) should contact
USDA's [TARGET Center](#) at 202-720-2600 (voice and TDD).*

*To file a complaint of discrimination write USDA, Director, Office
of Civil Rights, Room 326-W, Whitten Building, 14th and
Independence Avenue, SW, Washington, DC 20250-9410 or call
202-720-5964 (voice or TDD). USDA is an equal opportunity
provider and employer.*