

ALKALI SACATON

Sporobolus airoides (Torr.)

Torr

plant symbol = SPAI

Contributed By: USDA, NRCS, National Plant Data Center



R. Mohlenbrock
USDA, NRCS, Wetland Sciences Institute
@PLANTS

Alternative Names

giant sacaton, finetop saltgrass, hairgrass dropseed, zacaton alcalino

Uses

Ethnobotanic: The Hopi Indians use the seeds in times of famine (Whiting 1939). They were ground into flour, eaten dry or made into a mush.

Wildlife: Alkali sacaton's abundant herbage is eaten by cattle, sheep, and horses (Hitchcock 1951). This species is used as a good forage or grazing grass in lowland and in alkali regions (Gates 1937).

Status

Please consult the Plants Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Description

General: Grass Family (Poaceae). Alkali sacaton is a tough perennial two to three and a half feet tall, growing in large bunches. The culms are erect to spreading and range in height from 50 to 100 centimeters tall. The blades are elongate, flat, soon becoming involute, and usually less than four millimeters wide (Hitchcock 1951).

Distribution: Alkali sacaton ranges from South Dakota to Washington, south to Missouri, Kansas, Texas, and Mexico (Steyermark 1963). For current distribution, please consult the Plant profile page for this species on the PLANTS Web site.

Adaptation

Alkali sacaton grows on dry to moist sites with sand or gravelly soil. This species is often found growing on alkaline flats, prairies, and sandy plateaus. It is common along drainage in desert and semi-desert areas.

Establishment

Propagation by Seed: Alkali sacaton seeds should be sown in the spring in a greenhouse. Cover the seeds with a light layer of the growing media. Germination for this species should take place within two weeks. When seedlings are large enough to handle, plant them directly into their permanent positions in the summer.

Large divisions can be planted directly into their permanent positions. However, smaller divisions should be placed in individual pots in a cold frame, planting them when they are well established in the summer.

Management

Once *Sporobolus airoides* plants are well established little maintenance is required. It is best that the stands of this species is grazed in the spring and summer when growth is active.

Cultivars, Improved and Selected Materials (and area of origin)

Commonly available through commercial seed vendors.

References

Britton, N.L. & A. Brown 1970. *An illustrated flora of the northern United States and Canada*. Dover Publications, New York, New York.

Gates, F.C. 1937. *Grasses in Kansas*. Kansas State Printing Plant, Topeka, Kansas.

Hitchcock, A.S. 1951. *Manual of the grasses of the United States*. Misc. Publ. No. 200. U.S. Department of Agricultural, Washington, D.C.

Hitchcock, A.S. 1971. *Manual of the grasses of the United States*. Dover Publications, New York, New York.

Kearney, T.H.; R.H. Peebles; J. T. Howell; & E. McClintock 1960. *Arizona flora*. 2nd ed. University of California press, Berkeley, California.

Munz, P.A. 1974. *A flora of southern California*. University of California Press, Berkeley, California.

Steyermark, J.A. 1963. *Flora of Missouri*. The Iowa State University Press, Ames, Iowa.

The Great Plains Flora Association 1986. *Flora of the Great Plains*. University Press of Kansas, Lawrence, Kansas.

Whiting, A.F. 1939. *Ethnobotany of the Hopi*. North Arizona Society of Science and Art.

Prepared By

Jammie Favorite

formerly USDA, NRCS, National Plant Data Center
Baton Rouge, Louisiana

Species Coordinator

Lincoln M. Moore

USDA, NRCS, National Plant Data Center
Baton Rouge, Louisiana

Edited: 10jan02 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>>.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.