

NORTHERN SWEETGRASS

Hierochloa hirta (Schrank)
Borbás ssp. arctica (J. Presl) G.
Weim.
 plant symbol = HIHIA

Contributed by: USDA NRCS Rose Lake Plant Materials Center



Uses

Cultural: The cultural uses of sweetgrass include ceremonial incense, perfume, hair wash, bedding, basketry, dermatological aid, cold remedy, cough medicine, eyewash, febrifuge, respiratory aid, analgesic, insecticide, veterinary aid, decoration and adornment.

Because of the sweet, vanilla-like fragrance that develops once the plant has been harvested and begins to dry the use of sweetgrass as incense and fragrance is fairly ubiquitous among Native American tribes. The longer leaves are often braided and burned for religious and peace ceremonies, to invoke good power, and for various other rituals of cultural significance. In addition, sweetgrass has been utilized for several medicinal and cosmetic purposes. Whether alone, as adornment and perfume, or soaked in water for tea and other solutions the therapeutic applications of sweetgrass are scattered throughout the tribes. Another traditional use of this plant, particularly among the people of the Great Lakes and Northeast, is in handicrafts. Sweetgrass is often employed to craft or decorate baskets, bowls, trays and mats.


Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status (e.g. threatened or endangered species, state noxious status, and wetland indicator values).

Description

Sweetgrass is a native perennial grass. The culms are semi-erect, up to 30 inches tall and arise from slender, creeping rhizomes. Leaves are few, rough-edged and have shiny, hairless undersides. Often it has a reddish-purple color near the base of the plant. The highly prized longer leaves that grow on sterile shoots reach 18+ inches in length. The inflorescence is an open, pyramid-shaped, golden brown panicle with slender branches. Spikelets have 3 florets with awnless lemmas; glumes are thin, translucent and nearly equal in length. The fruit is a caryopsis.

Known Distribution



05/04/01

Adaptation

Sweetgrass is circumpolar, being native to both the western and eastern hemispheres. In North America its extensive range traverses the northern regions from Alaska to Newfoundland, moving down to New England, across the Great Lakes region and the upper Midwest to Oregon, and into the Southwest. In the Great Lakes, Northeastern and Midwestern regions sweetgrass usually inhabits moist ground on shores (fresh or brackish), meadows, low prairies, at the edges of woods, bogs and marshes. Normally, it is not found in pure stands but among other grasses and shrubs in mid-successional communities.

Establishment

Sweetgrass spreads vigorously by often-deep creeping rhizomes. In the spring these rhizomes produce inconspicuous fruiting stems with sparse, short leaves. Longer leaves develop later from separate sterile basal shoots. Although sweetgrass can reproduce by seed, it is mostly infertile, producing few seedheads that contain few seeds. This species is extremely cold hardy. It will go dormant in cold weather and resprout once ground temperatures reach 40 °F. For the Great Lakes, Northeastern and Midwestern regions flowering begins in the spring.

Sweetgrass development from seed is very slow. This coupled with the infertile nature of the species explains why plant division is the most successful method of reproducing sweetgrass. Dividing a plant is accomplished by separating the individual propagules that have developed from the rhizomes of a spreading plant. Each propagule can then be placed in a container for further separation or future planting. Newly separated plants will do best if placed in the shade for 2–3 weeks while their roots establish. After this, transplant at 1-foot spacings into areas of partial shade to full sun.

Management

As sweetgrass is not drought tolerant, keep the soil moist but not saturated. Fertilize with a balanced lawn-starter 2-3 times during the establishment year and years following harvests. Organic fertilizers may also be used at 5 lbs. per 100 square feet. Sweetgrass should be weeded at least every other year.

Longer leaves of the sterile shoots may be harvested several times during the year, however, the mid-season growth is considered to be superior. These leaves are gathered by grasping the shoots firmly near the ground and pulling until they break from the rootstock an inch or two below the surface of the soil.



Prepared By & Species Coordinator:

Tony Bush, Agronomist
USDA NRCS Rose Lake Plant Materials Center
East Lansing, Michigan

04May2001 JLK

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.