

SEER Incidence* and US Death Rates,# 1998-2002 5-Year Relative Survival Rates,^ 1995-2001 All Cancer Combined, by Race and Sex





Figure 1-9

* Incidence rates are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia) and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 # Death rates are from the NCHS public use data file for the total US and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 ^ Survival rates are from the SEER 9 areas. Relative survival rates are expressed as percents.

Figure I-11

5-Year Relative Survival Rates SEER Program, 1995-2001 Both Sexes, by Race




Source: SEER 9 areas.


SEER Cancer Incidence and US Death Rates, 1998-2002 By Cancer Site and Race

Incidence Cancer Site Mortality


Lung and Bronchus




Colon and Rectum



Breast (Female)



Prostate



Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia) and NCHS public use data file for the total US. Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

^^ Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics excludes cases from Detroit, Hawaii, Alaska Native Registry and Rural Georgia. Mortality data for Hispanics excludes cases from Connecticut, Maine, Maryland, Minnesota, Oklahoma, New Hampshire, New York, North Dakota, and Vermont.