

Table I-14

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site, Race and Sex

13 SEER Areas, 2000-2002

Site	All Races		Whites		Blacks	
	Males	Females	Males	Females	Males	Females
All Sites	45.67	38.09	46.11	39.45	42.50	31.88
Invasive and In Situ	46.91	41.44	47.48	42.99	42.79	33.99
Oral cavity and Pharynx	1.38	0.68	1.40	0.70	1.29	0.50
Esophagus	0.76	0.25	0.78	0.24	0.77	0.34
Stomach	1.22	0.75	1.06	0.61	1.27	0.97
Colon and Rectum	5.84	5.51	5.84	5.48	4.99	5.41
Invasive and In Situ	6.13	5.74	6.13	5.70	5.25	5.66
Liver and Intrahepatic bile duct	0.89	0.43	0.71	0.36	0.95	0.35
Pancreas	1.26	1.27	1.28	1.23	1.16	1.46
Larynx	0.62	0.14	0.62	0.15	0.85	0.19
Invasive and In Situ	0.66	0.15	0.67	0.16	0.88	0.20
Lung and Bronchus	7.58	5.72	7.59	6.01	8.08	5.24
Melanoma of skin	1.94	1.30	2.33	1.59	0.08	0.06
Invasive and In Situ	3.13	2.14	3.73	2.59	0.10	0.09
Breast	0.12	13.22	0.12	14.04	0.16	9.98
Invasive and In Situ	0.13	15.81	0.14	16.71	0.17	11.97
Cervix uteri	-	0.74	-	0.71	-	0.87
Corpus and Uterus, NOS	-	2.61	-	2.79	-	1.87
Invasive and In Situ	-	2.64	-	2.83	-	1.89
Ovary	-	1.48	-	1.59	-	0.92
Prostate	17.93	-	17.73	-	20.66	-
Testis	0.36	-	0.43	-	0.10	-
Urinary bladder(Invasive and In Situ)	3.58	1.14	3.99	1.24	1.39	0.80
Kidney and Renal pelvis	1.56	0.91	1.64	0.94	1.45	0.91
Brain and Other nervous system	0.65	0.50	0.74	0.57	0.32	0.28
Thyroid	0.35	0.97	0.38	1.02	0.17	0.53
Hodgkin lymphoma	0.24	0.19	0.25	0.21	0.21	0.15
Non-Hodgkin lymphoma	2.18	1.82	2.32	1.94	1.25	1.06
Multiple myeloma	0.70	0.55	0.68	0.51	0.94	0.93
Leukemia	1.50	1.07	1.61	1.14	0.87	0.79
Acute Lymphocytic Leukemia	0.12	0.11	0.14	0.12	0.04	0.06
Chronic Lymphocytic Leukemia	0.53	0.33	0.59	0.35	0.26	0.20
Acute Myeloid Leukemia	0.46	0.35	0.48	0.37	0.29	0.30
Chronic Myeloid Leukemia	0.19	0.14	0.19	0.14	0.13	0.11
Kaposi Sarcoma	0.12	0.02	0.10	0.02	0.23	0.02
Mesothelioma	0.19	0.05	0.22	0.05	0.06	0.02

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia).

Note: Invasive cancer only unless specified otherwise.

Table I-15

Lifetime Risk (Percent) of Dying from Cancer by Site, Race and Sex

Total U.S., 2000-2002

Site	All Races		Whites		Blacks	
	Males	Females	Males	Females	Males	Females
All Sites	23.56	19.93	23.62	20.16	24.21	19.43
Oral cavity and Pharynx	0.38	0.19	0.37	0.19	0.50	0.17
Esophagus	0.74	0.22	0.74	0.21	0.81	0.31
Stomach	0.59	0.40	0.53	0.35	0.89	0.67
Colon and Rectum	2.36	2.23	2.36	2.20	2.46	2.51
Liver and Intrahepatic bile duct	0.65	0.37	0.61	0.35	0.69	0.38
Pancreas	1.17	1.17	1.18	1.16	1.13	1.33
Larynx	0.23	0.06	0.22	0.06	0.38	0.09
Lung and Bronchus	7.25	4.92	7.33	5.12	7.29	3.97
Melanoma of skin	0.35	0.20	0.40	0.22	0.03	0.04
Breast	0.03	2.96	0.03	2.96	0.05	3.25
Cervix uteri	-	0.26	-	0.23	-	0.46
Corpus and Uterus, NOS	-	0.51	-	0.48	-	0.75
Ovary	-	1.05	-	1.10	-	0.74
Prostate	2.97	-	2.78	-	4.71	-
Testis	0.02	-	0.02	-	0.01	-
Urinary bladder	0.75	0.32	0.80	0.32	0.39	0.33
Kidney and Renal pelvis	0.59	0.34	0.61	0.35	0.45	0.28
Brain and Other nervous system	0.49	0.39	0.53	0.42	0.23	0.21
Thyroid	0.04	0.06	0.05	0.06	0.03	0.06
Hodgkin lymphoma	0.05	0.04	0.05	0.04	0.04	0.03
Non-Hodgkin lymphoma	0.94	0.81	1.00	0.86	0.50	0.43
Multiple myeloma	0.46	0.40	0.44	0.38	0.63	0.67
Leukemia	0.97	0.72	1.02	0.75	0.64	0.54
Acute Lymphocytic Leukemia	0.05	0.04	0.05	0.04	0.03	0.02
Chronic Lymphocytic Leukemia	0.23	0.15	0.24	0.16	0.16	0.11
Acute Myeloid Leukemia	0.33	0.25	0.35	0.26	0.19	0.17
Chronic Myeloid Leukemia	0.07	0.05	0.07	0.05	0.06	0.05

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: NCHS public use data file for the total US.

Table II-8
ALL SITES (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.17	0.35	0.71	45.67	0.16	0.32	0.78	38.09
10	0.18	0.55	1.28	46.18	0.16	0.63	1.86	38.41
20	0.37	1.11	3.07	46.32	0.47	1.70	4.72	38.39
30	0.75	2.74	9.13	46.65	1.25	4.29	10.14	38.24
40	2.04	8.57	22.40	46.95	3.10	9.06	18.27	37.70
50	6.86	21.40	37.36	47.20	6.23	15.86	26.66	36.18
60	16.46	34.54	43.87	45.68	10.54	22.38	30.69	32.80

Lifetime Risk of Being Diagnosed = 45.67% Lifetime Risk of Being Diagnosed = 38.09%
 Lifetime Risk of Dying = 23.56% Lifetime Risk of Dying = 19.93%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.19	0.38	0.78	46.11	0.17	0.35	0.85	39.45
10	0.20	0.60	1.36	46.53	0.18	0.68	1.96	39.72
20	0.41	1.17	3.13	46.64	0.51	1.79	4.89	39.69
30	0.78	2.77	9.10	46.90	1.30	4.42	10.49	39.51
40	2.03	8.51	22.52	47.12	3.18	9.36	19.04	38.93
50	6.79	21.48	37.59	47.28	6.46	16.56	27.81	37.35
60	16.56	34.72	43.91	45.64	11.07	23.38	31.79	33.82

Lifetime Risk of Being Diagnosed = 46.11% Lifetime Risk of Being Diagnosed = 39.45%
 Lifetime Risk of Dying = 23.62% Lifetime Risk of Dying = 20.16%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.11	0.22	0.47	42.50	0.11	0.22	0.55	31.88
10	0.11	0.37	1.08	43.58	0.12	0.45	1.51	32.52
20	0.26	0.98	3.50	43.92	0.34	1.40	4.17	32.53
30	0.74	3.33	12.03	44.91	1.07	3.87	9.24	32.53
40	2.69	11.73	27.16	45.90	2.87	8.36	16.67	32.23
50	9.87	26.72	40.57	47.18	5.84	14.66	23.80	31.19
60	21.00	38.26	45.05	46.50	9.94	20.25	26.80	28.58

Lifetime Risk of Being Diagnosed = 42.50% Lifetime Risk of Being Diagnosed = 31.88%
 Lifetime Risk of Dying = 24.21% Lifetime Risk of Dying = 19.43%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table III-8
BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.03	0.06	0.09	0.65	0.03	0.05	0.08	0.50
10	0.03	0.05	0.09	0.63	0.02	0.04	0.07	0.47
20	0.03	0.06	0.12	0.60	0.02	0.05	0.10	0.46
30	0.04	0.10	0.19	0.58	0.03	0.08	0.14	0.44
40	0.06	0.16	0.30	0.56	0.04	0.11	0.21	0.41
50	0.10	0.25	0.42	0.51	0.07	0.16	0.29	0.37
60	0.16	0.34	0.43	0.44	0.10	0.23	0.30	0.32

Lifetime Risk of Being Diagnosed = 0.65%
 Lifetime Risk of Dying = 0.49%

Lifetime Risk of Being Diagnosed = 0.50%
 Lifetime Risk of Dying = 0.39%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.04	0.07	0.10	0.74	0.04	0.06	0.08	0.57
10	0.03	0.06	0.10	0.71	0.02	0.04	0.08	0.53
20	0.03	0.07	0.14	0.68	0.02	0.06	0.11	0.51
30	0.04	0.11	0.22	0.66	0.04	0.09	0.16	0.49
40	0.07	0.18	0.34	0.62	0.05	0.13	0.23	0.46
50	0.11	0.28	0.47	0.57	0.08	0.19	0.32	0.41
60	0.18	0.38	0.48	0.49	0.11	0.26	0.33	0.35

Lifetime Risk of Being Diagnosed = 0.74%
 Lifetime Risk of Dying = 0.53%

Lifetime Risk of Being Diagnosed = 0.57%
 Lifetime Risk of Dying = 0.42%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.02	0.04	0.05	0.32	0.03	0.05	0.06	0.28
10	0.02	0.03	0.05	0.31	0.02	0.03	0.04	0.26
20	0.01	0.04	0.08	0.30	0.01	0.03	0.05	0.24
30	0.03	0.06	0.12	0.29	0.02	0.04	0.08	0.23
40	0.04	0.10	0.16	0.27	0.02	0.06	0.12	0.22
50	0.06	0.13	0.21	0.25	0.04	0.10	0.15	0.20
60	0.07	0.17	0.21	0.21	0.06	0.12	0.17	0.17

Lifetime Risk of Being Diagnosed = 0.32%
 Lifetime Risk of Dying = 0.23%

Lifetime Risk of Being Diagnosed = 0.28%
 Lifetime Risk of Dying = 0.21%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table IV-11
BREAST CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.12	0.00	0.00	0.05	13.22
10	0.00	0.00	0.00	0.12	0.00	0.05	0.48	13.37
20	0.00	0.00	0.01	0.12	0.05	0.48	1.92	13.40
30	0.00	0.01	0.02	0.13	0.44	1.88	4.49	13.41
40	0.01	0.02	0.06	0.13	1.46	4.11	7.56	13.14
50	0.02	0.05	0.09	0.12	2.73	6.30	9.64	12.06
60	0.04	0.08	0.11	0.11	3.82	7.40	9.52	9.99

Lifetime Risk of Being Diagnosed = 0.12% Lifetime Risk of Being Diagnosed = 13.22%
 Lifetime Risk of Dying = 0.03% Lifetime Risk of Dying = 2.96%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.12	0.00	0.00	0.05	14.04
10	0.00	0.00	0.00	0.13	0.00	0.05	0.48	14.17
20	0.00	0.00	0.01	0.13	0.05	0.48	1.96	14.20
30	0.00	0.01	0.02	0.13	0.43	1.92	4.66	14.21
40	0.01	0.02	0.06	0.13	1.50	4.28	8.01	13.94
50	0.02	0.05	0.09	0.13	2.86	6.71	10.31	12.83
60	0.04	0.08	0.11	0.12	4.12	7.96	10.18	10.66

Lifetime Risk of Being Diagnosed = 0.12% Lifetime Risk of Being Diagnosed = 14.04%
 Lifetime Risk of Dying = 0.03% Lifetime Risk of Dying = 2.96%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.16	0.00	0.00	0.07	9.98
10	0.00	0.00	0.00	0.16	0.00	0.07	0.53	10.21
20	0.00	0.00	0.02	0.17	0.07	0.53	1.83	10.24
30	0.00	0.02	0.04	0.17	0.47	1.78	3.98	10.26
40	0.01	0.04	0.09	0.17	1.34	3.58	6.20	9.99
50	0.03	0.08	0.12	0.17	2.36	5.11	7.52	9.10
60	0.06	0.10	0.13	0.16	3.05	5.72	7.13	7.48

Lifetime Risk of Being Diagnosed = 0.16% Lifetime Risk of Being Diagnosed = 9.98%
 Lifetime Risk of Dying = 0.05% Lifetime Risk of Dying = 3.25%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table V-6
CERVIX UTERI CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.04	0.74
10	0.00	0.04	0.15	0.75
20	0.04	0.15	0.30	0.75
30	0.12	0.26	0.40	0.71
40	0.15	0.28	0.42	0.60
50	0.14	0.28	0.38	0.46
60	0.15	0.26	0.32	0.34

Lifetime Risk of Being Diagnosed = 0.74%
 Lifetime Risk of Dying = 0.26%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.04	0.71
10	0.00	0.04	0.17	0.72
20	0.04	0.17	0.31	0.72
30	0.13	0.27	0.40	0.68
40	0.15	0.28	0.41	0.56
50	0.13	0.26	0.36	0.42
60	0.14	0.23	0.29	0.30

Lifetime Risk of Being Diagnosed = 0.71%
 Lifetime Risk of Dying = 0.23%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.03	0.87
10	0.00	0.03	0.14	0.89
20	0.03	0.13	0.27	0.89
30	0.10	0.24	0.39	0.86
40	0.14	0.29	0.47	0.77
50	0.15	0.34	0.50	0.66
60	0.21	0.38	0.50	0.55

Lifetime Risk of Being Diagnosed = 0.87%
 Lifetime Risk of Dying = 0.46%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table VI-13
COLON AND RECTUM CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	5.84	0.00	0.00	0.01	5.51
10	0.00	0.01	0.07	5.91	0.00	0.01	0.06	5.57
20	0.01	0.07	0.28	5.95	0.01	0.06	0.25	5.58
30	0.06	0.27	0.95	6.01	0.05	0.24	0.74	5.60
40	0.22	0.90	2.39	6.05	0.19	0.70	1.78	5.59
50	0.71	2.25	4.31	6.05	0.52	1.63	3.36	5.52
60	1.66	3.89	5.43	5.77	1.16	2.98	4.73	5.25

Lifetime Risk of Being Diagnosed = 5.84%
 Lifetime Risk of Dying = 2.36%

Lifetime Risk of Being Diagnosed = 5.51%
 Lifetime Risk of Dying = 2.23%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	5.84	0.00	0.00	0.01	5.48
10	0.00	0.01	0.07	5.91	0.00	0.01	0.06	5.53
20	0.01	0.07	0.27	5.94	0.01	0.06	0.23	5.54
30	0.06	0.26	0.91	5.99	0.05	0.22	0.70	5.55
40	0.20	0.87	2.35	6.03	0.18	0.66	1.72	5.54
50	0.69	2.22	4.30	6.03	0.49	1.58	3.32	5.47
60	1.65	3.88	5.41	5.74	1.14	2.96	4.72	5.21

Lifetime Risk of Being Diagnosed = 5.84%
 Lifetime Risk of Dying = 2.36%

Lifetime Risk of Being Diagnosed = 5.48%
 Lifetime Risk of Dying = 2.20%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	4.99	0.00	0.00	0.02	5.41
10	0.00	0.01	0.08	5.13	0.00	0.02	0.08	5.53
20	0.01	0.08	0.34	5.18	0.02	0.08	0.32	5.54
30	0.07	0.34	1.17	5.30	0.06	0.31	1.00	5.57
40	0.28	1.13	2.63	5.41	0.25	0.96	2.28	5.60
50	0.91	2.53	4.25	5.51	0.73	2.11	3.82	5.58
60	1.88	3.88	5.06	5.35	1.51	3.38	4.86	5.31

Lifetime Risk of Being Diagnosed = 4.99%
 Lifetime Risk of Dying = 2.46%

Lifetime Risk of Being Diagnosed = 5.41%
 Lifetime Risk of Dying = 2.51%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table VII-6
CORPUS AND UTERUS, NOS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.61
10	0.00	0.01	0.06	2.64
20	0.01	0.06	0.25	2.64
30	0.05	0.24	0.76	2.65
40	0.19	0.72	1.50	2.62
50	0.54	1.33	2.02	2.48
60	0.83	1.55	1.97	2.04

Lifetime Risk of Being Diagnosed = 2.61%
Lifetime Risk of Dying = 0.51%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.79
10	0.00	0.01	0.06	2.81
20	0.01	0.06	0.25	2.82
30	0.05	0.24	0.80	2.82
40	0.19	0.75	1.59	2.79
50	0.57	1.43	2.17	2.65
60	0.90	1.67	2.11	2.18

Lifetime Risk of Being Diagnosed = 2.79%
Lifetime Risk of Dying = 0.48%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.87
10	0.00	0.01	0.06	1.92
20	0.01	0.06	0.19	1.92
30	0.05	0.18	0.50	1.93
40	0.13	0.46	1.12	1.91
50	0.34	1.03	1.56	1.86
60	0.75	1.33	1.61	1.65

Lifetime Risk of Being Diagnosed = 1.87%
Lifetime Risk of Dying = 0.75%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table VIII-8
ESOPHAGUS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.76	0.00	0.00	0.00	0.25
10	0.00	0.00	0.01	0.77	0.00	0.00	0.00	0.25
20	0.00	0.01	0.03	0.77	0.00	0.00	0.01	0.25
30	0.00	0.03	0.15	0.78	0.00	0.01	0.03	0.26
40	0.03	0.14	0.37	0.79	0.01	0.03	0.09	0.26
50	0.12	0.35	0.62	0.79	0.02	0.08	0.16	0.25
60	0.25	0.54	0.69	0.72	0.06	0.14	0.22	0.24

Lifetime Risk of Being Diagnosed = 0.76%
 Lifetime Risk of Dying = 0.74%

Lifetime Risk of Being Diagnosed = 0.25%
 Lifetime Risk of Dying = 0.22%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.78	0.00	0.00	0.00	0.24
10	0.00	0.00	0.01	0.79	0.00	0.00	0.00	0.25
20	0.00	0.01	0.03	0.80	0.00	0.00	0.01	0.25
30	0.01	0.03	0.15	0.80	0.00	0.01	0.03	0.25
40	0.03	0.14	0.37	0.81	0.01	0.03	0.08	0.25
50	0.12	0.35	0.63	0.81	0.02	0.07	0.15	0.25
60	0.25	0.55	0.71	0.74	0.06	0.14	0.22	0.24

Lifetime Risk of Being Diagnosed = 0.78%
 Lifetime Risk of Dying = 0.74%

Lifetime Risk of Being Diagnosed = 0.24%
 Lifetime Risk of Dying = 0.21%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.77	0.00	0.00	0.00	0.34
10	0.00	0.00	0.01	0.79	0.00	0.00	0.00	0.35
20	0.00	0.01	0.05	0.80	0.00	0.00	0.02	0.35
30	0.00	0.05	0.24	0.82	0.00	0.02	0.07	0.35
40	0.04	0.24	0.53	0.84	0.02	0.08	0.17	0.36
50	0.21	0.52	0.76	0.85	0.06	0.16	0.27	0.35
60	0.36	0.64	0.74	0.75	0.11	0.24	0.30	0.32

Lifetime Risk of Being Diagnosed = 0.77%
 Lifetime Risk of Dying = 0.81%

Lifetime Risk of Being Diagnosed = 0.34%
 Lifetime Risk of Dying = 0.31%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table IX-8
HODGKIN LYMPHOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.06	0.24	0.00	0.02	0.06	0.19
10	0.02	0.06	0.10	0.23	0.02	0.06	0.09	0.19
20	0.04	0.08	0.11	0.22	0.04	0.07	0.09	0.17
30	0.04	0.07	0.10	0.18	0.03	0.05	0.07	0.13
40	0.03	0.06	0.10	0.14	0.02	0.03	0.05	0.10
50	0.03	0.06	0.10	0.11	0.02	0.04	0.06	0.08
60	0.04	0.07	0.09	0.09	0.02	0.05	0.06	0.07

Lifetime Risk of Being Diagnosed = 0.24%
Lifetime Risk of Dying = 0.05%

Lifetime Risk of Being Diagnosed = 0.19%
Lifetime Risk of Dying = 0.04%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.06	0.25	0.00	0.02	0.07	0.21
10	0.02	0.06	0.10	0.25	0.02	0.07	0.10	0.21
20	0.04	0.08	0.12	0.23	0.04	0.08	0.10	0.18
30	0.04	0.08	0.10	0.19	0.03	0.05	0.07	0.14
40	0.04	0.07	0.11	0.16	0.02	0.04	0.06	0.10
50	0.03	0.07	0.10	0.12	0.02	0.04	0.07	0.09
60	0.04	0.08	0.10	0.10	0.03	0.05	0.07	0.07

Lifetime Risk of Being Diagnosed = 0.25%
Lifetime Risk of Dying = 0.05%

Lifetime Risk of Being Diagnosed = 0.21%
Lifetime Risk of Dying = 0.04%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.06	0.21	0.00	0.02	0.06	0.15
10	0.01	0.06	0.10	0.22	0.02	0.06	0.09	0.15
20	0.04	0.09	0.12	0.20	0.04	0.07	0.10	0.13
30	0.04	0.08	0.11	0.17	0.03	0.05	0.07	0.09
40	0.04	0.07	0.09	0.13	0.02	0.04	0.04	0.06
50	0.04	0.06	0.09	0.09	0.02	0.02	0.03	0.04
60	0.03	0.06	0.07	0.07	0.01	0.02	0.02	0.03

Lifetime Risk of Being Diagnosed = 0.21%
Lifetime Risk of Dying = 0.04%

Lifetime Risk of Being Diagnosed = 0.15%
Lifetime Risk of Dying = 0.03%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table X-9
KAPOSI SARCOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	0.12	0.00	0.00	0.00	0.02
10	0.00	0.01	0.04	0.12	0.00	0.00	0.00	0.02
20	0.01	0.04	0.07	0.12	0.00	0.00	0.00	0.02
30	0.03	0.07	0.08	0.11	0.00	0.00	0.00	0.02
40	0.03	0.04	0.05	0.08	0.00	0.00	0.00	0.02
50	0.01	0.02	0.03	0.05	0.00	0.00	0.01	0.02
60	0.01	0.02	0.03	0.04	0.00	0.01	0.01	0.02

Lifetime Risk of Being Diagnosed = 0.12%

Lifetime Risk of Being Diagnosed = 0.02%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	0.10	0.00	0.00	0.00	0.02
10	0.00	0.01	0.03	0.10	0.00	0.00	0.00	0.02
20	0.01	0.03	0.06	0.10	0.00	0.00	0.00	0.02
30	0.03	0.06	0.07	0.10	0.00	0.00	0.00	0.02
40	0.03	0.04	0.05	0.07	0.00	0.00	0.00	0.02
50	0.01	0.02	0.03	0.05	0.00	0.00	0.01	0.02
60	0.01	0.02	0.03	0.04	0.00	0.01	0.01	0.02

Lifetime Risk of Being Diagnosed = 0.10%

Lifetime Risk of Being Diagnosed = 0.02%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.04	0.23	0.00	0.00	0.00	0.02
10	0.00	0.04	0.13	0.23	0.00	0.00	0.01	0.02
20	0.04	0.13	0.19	0.23	0.00	0.01	0.01	0.02
30	0.10	0.16	0.18	0.20	0.00	0.01	0.01	0.01
40	0.06	0.09	0.10	0.11	0.01	0.01	0.01	0.01
50	0.03	0.04	0.04	0.05	0.00	0.01	0.01	0.01
60	0.02	0.02	0.02	0.02	0.00	0.00	0.00	0.00

Lifetime Risk of Being Diagnosed = 0.23%

Lifetime Risk of Being Diagnosed = 0.02%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia).

Table XI-8

KIDNEY AND RENAL PELVIS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.01	0.02	1.56	0.01	0.01	0.02	0.91
10	0.00	0.01	0.03	1.57	0.00	0.01	0.02	0.91
20	0.00	0.03	0.13	1.58	0.00	0.02	0.08	0.91
30	0.03	0.13	0.39	1.59	0.02	0.07	0.20	0.91
40	0.10	0.37	0.82	1.59	0.06	0.19	0.42	0.90
50	0.27	0.75	1.25	1.55	0.13	0.37	0.64	0.86
60	0.51	1.05	1.33	1.37	0.25	0.53	0.72	0.76

Lifetime Risk of Being Diagnosed = 1.56%
Lifetime Risk of Dying = 0.59%

Lifetime Risk of Being Diagnosed = 0.91%
Lifetime Risk of Dying = 0.34%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.01	0.02	1.64	0.01	0.01	0.02	0.94
10	0.00	0.01	0.03	1.65	0.00	0.01	0.02	0.94
20	0.00	0.03	0.13	1.66	0.00	0.02	0.08	0.94
30	0.03	0.13	0.39	1.67	0.02	0.07	0.21	0.94
40	0.11	0.37	0.86	1.67	0.06	0.20	0.44	0.93
50	0.28	0.78	1.31	1.61	0.14	0.38	0.66	0.89
60	0.53	1.11	1.39	1.43	0.26	0.54	0.74	0.78

Lifetime Risk of Being Diagnosed = 1.64%
Lifetime Risk of Dying = 0.61%

Lifetime Risk of Being Diagnosed = 0.94%
Lifetime Risk of Dying = 0.35%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.02	0.02	1.45	0.02	0.03	0.03	0.91
10	0.00	0.01	0.04	1.48	0.00	0.01	0.04	0.91
20	0.01	0.04	0.16	1.49	0.01	0.03	0.10	0.91
30	0.03	0.16	0.48	1.52	0.02	0.09	0.24	0.91
40	0.13	0.46	0.91	1.54	0.06	0.22	0.47	0.90
50	0.36	0.84	1.25	1.52	0.16	0.42	0.72	0.87
60	0.56	1.03	1.29	1.35	0.29	0.61	0.76	0.77

Lifetime Risk of Being Diagnosed = 1.45%
Lifetime Risk of Dying = 0.45%

Lifetime Risk of Being Diagnosed = 0.91%
Lifetime Risk of Dying = 0.28%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XII-8
LARYNX CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.62	0.00	0.00	0.00	0.14
10	0.00	0.00	0.00	0.62	0.00	0.00	0.00	0.14
20	0.00	0.00	0.03	0.63	0.00	0.00	0.01	0.14
30	0.00	0.03	0.14	0.64	0.00	0.01	0.03	0.14
40	0.03	0.14	0.35	0.64	0.01	0.03	0.08	0.14
50	0.12	0.33	0.53	0.63	0.03	0.07	0.12	0.14
60	0.23	0.45	0.54	0.56	0.05	0.10	0.11	0.12

Lifetime Risk of Being Diagnosed = 0.62%
 Lifetime Risk of Dying = 0.23%

Lifetime Risk of Being Diagnosed = 0.14%
 Lifetime Risk of Dying = 0.06%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.62	0.00	0.00	0.00	0.15
10	0.00	0.00	0.00	0.62	0.00	0.00	0.00	0.15
20	0.00	0.00	0.03	0.63	0.00	0.00	0.01	0.15
30	0.00	0.03	0.14	0.63	0.00	0.01	0.03	0.15
40	0.03	0.14	0.35	0.64	0.01	0.03	0.08	0.15
50	0.11	0.33	0.53	0.63	0.03	0.08	0.12	0.14
60	0.23	0.45	0.54	0.56	0.05	0.10	0.12	0.12

Lifetime Risk of Being Diagnosed = 0.62%
 Lifetime Risk of Dying = 0.22%

Lifetime Risk of Being Diagnosed = 0.15%
 Lifetime Risk of Dying = 0.06%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.85	0.00	0.00	0.00	0.19
10	0.00	0.00	0.01	0.87	0.00	0.00	0.00	0.20
20	0.00	0.01	0.07	0.88	0.00	0.00	0.02	0.20
30	0.01	0.07	0.28	0.90	0.00	0.02	0.06	0.20
40	0.06	0.28	0.58	0.93	0.02	0.06	0.13	0.20
50	0.23	0.56	0.83	0.93	0.04	0.11	0.17	0.19
60	0.38	0.69	0.80	0.81	0.08	0.13	0.15	0.16

Lifetime Risk of Being Diagnosed = 0.85%
 Lifetime Risk of Dying = 0.38%

Lifetime Risk of Being Diagnosed = 0.19%
 Lifetime Risk of Dying = 0.09%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XIII-14

LEUKEMIA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.06	0.09	0.12	1.50	0.05	0.08	0.10	1.07
10	0.03	0.06	0.10	1.46	0.03	0.05	0.08	1.03
20	0.03	0.07	0.13	1.44	0.02	0.05	0.10	1.01
30	0.04	0.10	0.25	1.43	0.03	0.07	0.17	0.99
40	0.07	0.22	0.53	1.42	0.05	0.14	0.32	0.97
50	0.16	0.48	0.95	1.40	0.09	0.28	0.57	0.94
60	0.35	0.85	1.24	1.34	0.19	0.50	0.80	0.89

Lifetime Risk of Being Diagnosed = 1.50%
Lifetime Risk of Dying = 0.97%

Lifetime Risk of Being Diagnosed = 1.07%
Lifetime Risk of Dying = 0.72%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.06	0.10	0.13	1.61	0.06	0.09	0.11	1.14
10	0.04	0.06	0.10	1.57	0.03	0.05	0.08	1.09
20	0.03	0.07	0.13	1.54	0.02	0.05	0.10	1.07
30	0.04	0.11	0.27	1.53	0.03	0.08	0.17	1.05
40	0.07	0.23	0.56	1.51	0.05	0.14	0.34	1.03
50	0.17	0.51	1.01	1.49	0.10	0.30	0.61	1.00
60	0.37	0.91	1.32	1.42	0.21	0.54	0.85	0.94

Lifetime Risk of Being Diagnosed = 1.61%
Lifetime Risk of Dying = 1.02%

Lifetime Risk of Being Diagnosed = 1.14%
Lifetime Risk of Dying = 0.75%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.03	0.04	0.06	0.87	0.03	0.05	0.06	0.79
10	0.02	0.04	0.07	0.87	0.02	0.04	0.06	0.78
20	0.02	0.06	0.12	0.86	0.02	0.04	0.08	0.76
30	0.04	0.10	0.22	0.86	0.02	0.06	0.13	0.75
40	0.07	0.18	0.42	0.85	0.04	0.11	0.26	0.74
50	0.12	0.38	0.64	0.84	0.07	0.24	0.47	0.73
60	0.30	0.60	0.79	0.83	0.18	0.43	0.64	0.72

Lifetime Risk of Being Diagnosed = 0.87%
Lifetime Risk of Dying = 0.64%

Lifetime Risk of Being Diagnosed = 0.79%
Lifetime Risk of Dying = 0.54%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XIV-8
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	0.89	0.00	0.01	0.01	0.43
10	0.00	0.00	0.01	0.90	0.00	0.00	0.01	0.43
20	0.00	0.01	0.08	0.91	0.00	0.01	0.02	0.43
30	0.01	0.08	0.24	0.92	0.00	0.02	0.06	0.43
40	0.07	0.24	0.48	0.92	0.02	0.06	0.17	0.43
50	0.18	0.43	0.71	0.88	0.04	0.16	0.30	0.42
60	0.28	0.57	0.73	0.76	0.12	0.27	0.37	0.39

Lifetime Risk of Being Diagnosed = 0.89%
 Lifetime Risk of Dying = 0.65%

Lifetime Risk of Being Diagnosed = 0.43%
 Lifetime Risk of Dying = 0.37%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.71	0.00	0.01	0.01	0.36
10	0.00	0.00	0.01	0.72	0.00	0.00	0.01	0.35
20	0.00	0.01	0.06	0.72	0.00	0.00	0.02	0.35
30	0.01	0.06	0.19	0.73	0.00	0.02	0.05	0.35
40	0.05	0.18	0.37	0.74	0.01	0.05	0.13	0.35
50	0.13	0.33	0.56	0.71	0.03	0.12	0.24	0.35
60	0.21	0.46	0.59	0.61	0.09	0.22	0.30	0.33

Lifetime Risk of Being Diagnosed = 0.71%
 Lifetime Risk of Dying = 0.61%

Lifetime Risk of Being Diagnosed = 0.36%
 Lifetime Risk of Dying = 0.35%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.95	0.00	0.00	0.00	0.35
10	0.00	0.00	0.02	0.97	0.00	0.00	0.01	0.36
20	0.00	0.02	0.12	0.98	0.00	0.01	0.02	0.36
30	0.01	0.12	0.39	1.00	0.00	0.02	0.08	0.36
40	0.11	0.39	0.66	1.02	0.02	0.08	0.17	0.36
50	0.30	0.59	0.84	0.98	0.06	0.16	0.27	0.36
60	0.33	0.62	0.76	0.79	0.11	0.23	0.30	0.32

Lifetime Risk of Being Diagnosed = 0.95%
 Lifetime Risk of Dying = 0.69%

Lifetime Risk of Being Diagnosed = 0.35%
 Lifetime Risk of Dying = 0.38%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XV-13
LUNG AND BRONCHUS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	7.58	0.00	0.00	0.01	5.72
10	0.00	0.01	0.03	7.69	0.00	0.01	0.03	5.78
20	0.00	0.03	0.21	7.73	0.00	0.03	0.19	5.80
30	0.03	0.21	1.01	7.82	0.03	0.19	0.82	5.82
40	0.19	1.00	3.19	7.92	0.16	0.80	2.37	5.84
50	0.84	3.11	6.14	8.01	0.65	2.25	4.36	5.79
60	2.45	5.72	7.47	7.74	1.68	3.89	5.18	5.39

Lifetime Risk of Being Diagnosed = 7.58%
Lifetime Risk of Dying = 7.25%

Lifetime Risk of Being Diagnosed = 5.72%
Lifetime Risk of Dying = 4.92%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	7.59	0.00	0.00	0.01	6.01
10	0.00	0.01	0.03	7.68	0.00	0.01	0.03	6.07
20	0.00	0.03	0.19	7.72	0.00	0.03	0.19	6.08
30	0.02	0.19	0.94	7.79	0.03	0.19	0.84	6.10
40	0.17	0.93	3.12	7.89	0.16	0.82	2.49	6.12
50	0.79	3.05	6.13	7.98	0.67	2.37	4.62	6.07
60	2.42	5.74	7.46	7.71	1.78	4.13	5.44	5.65

Lifetime Risk of Being Diagnosed = 7.59%
Lifetime Risk of Dying = 7.33%

Lifetime Risk of Being Diagnosed = 6.01%
Lifetime Risk of Dying = 5.12%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	8.08	0.00	0.00	0.00	5.24
10	0.00	0.00	0.04	8.30	0.00	0.00	0.04	5.35
20	0.00	0.04	0.40	8.38	0.00	0.03	0.28	5.37
30	0.04	0.40	1.79	8.60	0.03	0.28	1.11	5.41
40	0.38	1.81	4.68	8.85	0.25	1.09	2.70	5.46
50	1.54	4.62	7.62	9.10	0.87	2.55	4.41	5.44
60	3.58	7.06	8.62	8.79	1.83	3.86	4.80	4.99

Lifetime Risk of Being Diagnosed = 8.08%
Lifetime Risk of Dying = 7.29%

Lifetime Risk of Being Diagnosed = 5.24%
Lifetime Risk of Dying = 3.97%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XVI-8
MELANOMA OF SKIN (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.04	1.94	0.00	0.01	0.08	1.30
10	0.01	0.04	0.13	1.96	0.01	0.08	0.21	1.31
20	0.03	0.12	0.30	1.97	0.07	0.20	0.38	1.30
30	0.09	0.27	0.59	1.96	0.13	0.31	0.53	1.24
40	0.19	0.51	0.97	1.90	0.18	0.40	0.65	1.12
50	0.34	0.81	1.36	1.78	0.22	0.47	0.73	0.95
60	0.51	1.10	1.48	1.56	0.26	0.53	0.72	0.76

Lifetime Risk of Being Diagnosed = 1.94%
 Lifetime Risk of Dying = 0.35%

Lifetime Risk of Being Diagnosed = 1.30%
 Lifetime Risk of Dying = 0.20%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.05	2.33	0.00	0.02	0.11	1.59
10	0.01	0.05	0.15	2.35	0.02	0.11	0.27	1.60
20	0.04	0.15	0.37	2.36	0.09	0.26	0.48	1.59
30	0.11	0.33	0.72	2.34	0.17	0.39	0.66	1.51
40	0.23	0.62	1.17	2.27	0.23	0.50	0.81	1.35
50	0.41	0.98	1.63	2.11	0.28	0.59	0.90	1.15
60	0.62	1.32	1.75	1.83	0.32	0.65	0.86	0.91

Lifetime Risk of Being Diagnosed = 2.33%
 Lifetime Risk of Dying = 0.40%

Lifetime Risk of Being Diagnosed = 1.59%
 Lifetime Risk of Dying = 0.22%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.06
10	0.00	0.00	0.01	0.09	0.00	0.00	0.01	0.06
20	0.00	0.01	0.01	0.09	0.00	0.01	0.01	0.06
30	0.01	0.01	0.02	0.09	0.01	0.01	0.02	0.05
40	0.00	0.02	0.05	0.09	0.01	0.01	0.02	0.05
50	0.01	0.05	0.07	0.09	0.01	0.02	0.03	0.05
60	0.04	0.07	0.08	0.08	0.01	0.03	0.04	0.04

Lifetime Risk of Being Diagnosed = 0.08%
 Lifetime Risk of Dying = 0.03%

Lifetime Risk of Being Diagnosed = 0.06%
 Lifetime Risk of Dying = 0.04%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XVII-7
MESOTHELIOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.19	0.00	0.00	0.00	0.05
10	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.05
20	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.05
30	0.00	0.00	0.02	0.20	0.00	0.00	0.01	0.05
40	0.00	0.01	0.06	0.20	0.00	0.01	0.02	0.05
50	0.01	0.06	0.14	0.21	0.01	0.01	0.03	0.05
60	0.05	0.14	0.20	0.21	0.01	0.03	0.04	0.04

Lifetime Risk of Being Diagnosed = 0.19%

Lifetime Risk of Being Diagnosed = 0.05%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.22	0.00	0.00	0.00	0.05
10	0.00	0.00	0.00	0.23	0.00	0.00	0.00	0.05
20	0.00	0.00	0.00	0.23	0.00	0.00	0.00	0.05
30	0.00	0.00	0.02	0.23	0.00	0.00	0.01	0.05
40	0.00	0.02	0.07	0.23	0.00	0.01	0.02	0.05
50	0.01	0.06	0.17	0.23	0.01	0.02	0.03	0.05
60	0.05	0.16	0.23	0.24	0.01	0.03	0.05	0.05

Lifetime Risk of Being Diagnosed = 0.22%

Lifetime Risk of Being Diagnosed = 0.05%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.02
10	0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.02
20	0.00	0.00	0.00	0.07	0.00	0.00	0.00	0.02
30	0.00	0.00	0.01	0.07	0.00	0.00	0.00	0.02
40	0.00	0.01	0.02	0.07	0.00	0.00	0.01	0.02
50	0.01	0.02	0.04	0.07	0.00	0.01	0.02	0.02
60	0.01	0.04	0.07	0.07	0.01	0.02	0.02	0.02

Lifetime Risk of Being Diagnosed = 0.06%

Lifetime Risk of Being Diagnosed = 0.02%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia).

Table XVIII-8
MULTIPLE MYELOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.70	0.00	0.00	0.00	0.55
10	0.00	0.00	0.01	0.71	0.00	0.00	0.00	0.56
20	0.00	0.01	0.03	0.71	0.00	0.00	0.02	0.56
30	0.01	0.03	0.12	0.72	0.00	0.02	0.08	0.56
40	0.03	0.11	0.29	0.72	0.02	0.08	0.21	0.56
50	0.09	0.28	0.53	0.72	0.07	0.19	0.38	0.56
60	0.20	0.47	0.65	0.68	0.13	0.33	0.49	0.51

Lifetime Risk of Being Diagnosed = 0.70%
 Lifetime Risk of Dying = 0.46%

Lifetime Risk of Being Diagnosed = 0.55%
 Lifetime Risk of Dying = 0.40%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.68	0.00	0.00	0.00	0.51
10	0.00	0.00	0.01	0.69	0.00	0.00	0.00	0.52
20	0.00	0.01	0.03	0.70	0.00	0.00	0.02	0.52
30	0.01	0.03	0.11	0.70	0.00	0.02	0.08	0.52
40	0.02	0.10	0.28	0.71	0.02	0.07	0.18	0.52
50	0.08	0.27	0.52	0.71	0.06	0.17	0.35	0.51
60	0.20	0.46	0.63	0.67	0.12	0.30	0.45	0.47

Lifetime Risk of Being Diagnosed = 0.68%
 Lifetime Risk of Dying = 0.44%

Lifetime Risk of Being Diagnosed = 0.51%
 Lifetime Risk of Dying = 0.38%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	0.94	0.00	0.00	0.00	0.93
10	0.00	0.00	0.01	0.96	0.00	0.00	0.01	0.95
20	0.00	0.01	0.06	0.97	0.00	0.01	0.05	0.95
30	0.01	0.06	0.21	1.00	0.01	0.05	0.18	0.96
40	0.05	0.20	0.50	1.02	0.04	0.17	0.40	0.96
50	0.16	0.48	0.81	1.04	0.14	0.38	0.71	0.96
60	0.37	0.75	0.97	1.01	0.26	0.61	0.86	0.90

Lifetime Risk of Being Diagnosed = 0.94%
 Lifetime Risk of Dying = 0.63%

Lifetime Risk of Being Diagnosed = 0.93%
 Lifetime Risk of Dying = 0.67%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XIX-12
NON-HODGKIN LYMPHOMA

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.03	0.06	2.18	0.01	0.02	0.04	1.82
10	0.02	0.05	0.13	2.21	0.01	0.04	0.08	1.84
20	0.03	0.11	0.27	2.20	0.03	0.07	0.17	1.83
30	0.08	0.24	0.54	2.19	0.05	0.14	0.36	1.81
40	0.16	0.47	0.97	2.15	0.10	0.31	0.71	1.78
50	0.31	0.84	1.52	2.06	0.22	0.62	1.18	1.72
60	0.56	1.30	1.79	1.88	0.42	1.01	1.47	1.57

Lifetime Risk of Being Diagnosed = 2.18%
Lifetime Risk of Dying = 0.94%

Lifetime Risk of Being Diagnosed = 1.82%
Lifetime Risk of Dying = 0.81%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.01	0.03	0.06	2.32	0.01	0.02	0.04	1.94
10	0.02	0.06	0.13	2.34	0.01	0.03	0.08	1.95
20	0.04	0.11	0.28	2.33	0.02	0.07	0.17	1.94
30	0.08	0.24	0.56	2.32	0.05	0.14	0.37	1.93
40	0.17	0.49	1.03	2.28	0.10	0.32	0.76	1.90
50	0.33	0.89	1.63	2.19	0.23	0.68	1.27	1.83
60	0.60	1.39	1.90	1.99	0.47	1.09	1.57	1.67

Lifetime Risk of Being Diagnosed = 2.32%
Lifetime Risk of Dying = 1.00%

Lifetime Risk of Being Diagnosed = 1.94%
Lifetime Risk of Dying = 0.86%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.05	1.25	0.00	0.02	0.05	1.06
10	0.01	0.04	0.15	1.28	0.01	0.05	0.10	1.08
20	0.03	0.14	0.30	1.28	0.04	0.09	0.19	1.07
30	0.11	0.28	0.52	1.28	0.05	0.15	0.33	1.04
40	0.18	0.43	0.75	1.21	0.10	0.28	0.50	1.00
50	0.27	0.62	0.93	1.11	0.19	0.42	0.71	0.94
60	0.40	0.77	0.94	0.97	0.26	0.57	0.77	0.83

Lifetime Risk of Being Diagnosed = 1.25%
Lifetime Risk of Dying = 0.50%

Lifetime Risk of Being Diagnosed = 1.06%
Lifetime Risk of Dying = 0.43%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XX-8

ORAL CAVITY AND PHARYNX CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.38	0.00	0.00	0.01	0.68
10	0.00	0.01	0.04	1.40	0.00	0.01	0.03	0.69
20	0.01	0.04	0.16	1.40	0.01	0.03	0.08	0.69
30	0.03	0.16	0.44	1.41	0.02	0.07	0.17	0.68
40	0.13	0.42	0.82	1.40	0.05	0.15	0.30	0.67
50	0.30	0.72	1.08	1.32	0.10	0.26	0.46	0.63
60	0.45	0.84	1.05	1.10	0.16	0.37	0.52	0.56

Lifetime Risk of Being Diagnosed = 1.38%
Lifetime Risk of Dying = 0.38%

Lifetime Risk of Being Diagnosed = 0.68%
Lifetime Risk of Dying = 0.19%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.40	0.00	0.01	0.02	0.70
10	0.00	0.01	0.04	1.42	0.00	0.01	0.03	0.70
20	0.01	0.04	0.16	1.42	0.01	0.03	0.07	0.70
30	0.03	0.15	0.44	1.43	0.02	0.06	0.16	0.69
40	0.13	0.42	0.82	1.42	0.04	0.14	0.30	0.68
50	0.30	0.72	1.09	1.34	0.10	0.26	0.47	0.65
60	0.45	0.85	1.07	1.12	0.17	0.39	0.54	0.58

Lifetime Risk of Being Diagnosed = 1.40%
Lifetime Risk of Dying = 0.37%

Lifetime Risk of Being Diagnosed = 0.70%
Lifetime Risk of Dying = 0.19%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.01	1.29	0.00	0.00	0.01	0.50
10	0.01	0.01	0.04	1.33	0.00	0.01	0.03	0.51
20	0.01	0.04	0.19	1.33	0.01	0.03	0.10	0.51
30	0.03	0.19	0.55	1.36	0.02	0.10	0.21	0.51
40	0.17	0.54	0.99	1.38	0.08	0.18	0.32	0.49
50	0.40	0.88	1.20	1.30	0.11	0.25	0.38	0.44
60	0.56	0.92	1.02	1.04	0.15	0.28	0.35	0.35

Lifetime Risk of Being Diagnosed = 1.29%
Lifetime Risk of Dying = 0.50%

Lifetime Risk of Being Diagnosed = 0.50%
Lifetime Risk of Dying = 0.17%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XXI-6
OVARY CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.03	1.48
10	0.01	0.03	0.07	1.49
20	0.02	0.06	0.19	1.48
30	0.04	0.17	0.43	1.47
40	0.13	0.39	0.74	1.44
50	0.26	0.63	1.02	1.34
60	0.38	0.79	1.06	1.13

Lifetime Risk of Being Diagnosed = 1.48%
 Lifetime Risk of Dying = 1.05%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.03	1.59
10	0.01	0.03	0.08	1.60
20	0.02	0.06	0.19	1.59
30	0.05	0.18	0.46	1.58
40	0.13	0.41	0.80	1.55
50	0.29	0.68	1.11	1.44
60	0.42	0.86	1.14	1.21

Lifetime Risk of Being Diagnosed = 1.59%
 Lifetime Risk of Dying = 1.10%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.02	0.92
10	0.01	0.01	0.05	0.94
20	0.01	0.04	0.12	0.93
30	0.03	0.11	0.25	0.93
40	0.08	0.22	0.47	0.91
50	0.15	0.41	0.66	0.87
60	0.28	0.55	0.73	0.79

Lifetime Risk of Being Diagnosed = 0.92%
 Lifetime Risk of Dying = 0.74%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XXII-8
PANCREAS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.26	0.00	0.00	0.00	1.27
10	0.00	0.00	0.01	1.28	0.00	0.00	0.01	1.28
20	0.00	0.01	0.05	1.29	0.00	0.01	0.04	1.29
30	0.01	0.05	0.20	1.30	0.01	0.04	0.14	1.29
40	0.05	0.20	0.53	1.31	0.03	0.13	0.40	1.29
50	0.16	0.50	0.95	1.31	0.11	0.38	0.81	1.29
60	0.37	0.85	1.17	1.24	0.28	0.74	1.12	1.24

Lifetime Risk of Being Diagnosed = 1.26%
 Lifetime Risk of Dying = 1.17%

Lifetime Risk of Being Diagnosed = 1.27%
 Lifetime Risk of Dying = 1.17%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.28	0.00	0.00	0.00	1.23
10	0.00	0.00	0.01	1.30	0.00	0.00	0.01	1.24
20	0.00	0.01	0.05	1.30	0.00	0.01	0.04	1.25
30	0.01	0.05	0.20	1.32	0.01	0.03	0.13	1.25
40	0.05	0.20	0.53	1.33	0.03	0.13	0.38	1.25
50	0.16	0.49	0.96	1.32	0.10	0.36	0.79	1.25
60	0.36	0.86	1.19	1.25	0.27	0.72	1.09	1.19

Lifetime Risk of Being Diagnosed = 1.28%
 Lifetime Risk of Dying = 1.18%

Lifetime Risk of Being Diagnosed = 1.23%
 Lifetime Risk of Dying = 1.16%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.16	0.00	0.00	0.00	1.46
10	0.00	0.00	0.02	1.19	0.00	0.00	0.01	1.49
20	0.00	0.02	0.09	1.20	0.00	0.01	0.05	1.49
30	0.02	0.09	0.29	1.23	0.01	0.05	0.19	1.50
40	0.07	0.28	0.70	1.25	0.05	0.18	0.56	1.52
50	0.22	0.67	1.08	1.27	0.14	0.53	1.01	1.54
60	0.52	0.99	1.18	1.21	0.43	0.95	1.36	1.52

Lifetime Risk of Being Diagnosed = 1.16%
 Lifetime Risk of Dying = 1.13%

Lifetime Risk of Being Diagnosed = 1.46%
 Lifetime Risk of Dying = 1.33%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XXIII-6
PROSTATE CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	17.93
10	0.00	0.00	0.01	18.17
20	0.00	0.01	0.29	18.28
30	0.01	0.30	2.63	18.50
40	0.29	2.66	8.94	18.80
50	2.46	8.98	15.85	19.22
60	7.19	14.77	17.94	18.48

Lifetime Risk of Being Diagnosed = 17.93%
 Lifetime Risk of Dying = 2.97%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	17.73
10	0.00	0.00	0.01	17.93
20	0.00	0.01	0.26	18.02
30	0.01	0.27	2.54	18.22
40	0.26	2.57	8.85	18.49
50	2.39	8.89	15.67	18.87
60	7.13	14.57	17.58	18.08

Lifetime Risk of Being Diagnosed = 17.73%
 Lifetime Risk of Dying = 2.78%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	20.66
10	0.00	0.00	0.02	21.23
20	0.00	0.02	0.67	21.43
30	0.02	0.69	4.62	22.00
40	0.68	4.75	13.12	22.71
50	4.39	13.42	20.79	23.79
60	10.97	19.91	22.96	23.55

Lifetime Risk of Being Diagnosed = 20.66%
 Lifetime Risk of Dying = 4.71%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XXIV-8
STOMACH CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.22	0.00	0.00	0.00	0.75
10	0.00	0.00	0.02	1.23	0.00	0.00	0.02	0.76
20	0.00	0.02	0.06	1.24	0.00	0.01	0.04	0.76
30	0.01	0.06	0.19	1.25	0.01	0.04	0.11	0.76
40	0.05	0.17	0.46	1.26	0.03	0.09	0.23	0.76
50	0.13	0.43	0.86	1.25	0.07	0.21	0.44	0.74
60	0.32	0.79	1.13	1.21	0.15	0.39	0.63	0.71

Lifetime Risk of Being Diagnosed = 1.22%
 Lifetime Risk of Dying = 0.59%

Lifetime Risk of Being Diagnosed = 0.75%
 Lifetime Risk of Dying = 0.40%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.06	0.00	0.00	0.00	0.61
10	0.00	0.00	0.02	1.07	0.00	0.00	0.01	0.62
20	0.00	0.02	0.05	1.08	0.00	0.01	0.04	0.62
30	0.01	0.05	0.16	1.08	0.01	0.04	0.09	0.62
40	0.04	0.15	0.41	1.09	0.02	0.08	0.19	0.61
50	0.12	0.38	0.75	1.08	0.06	0.17	0.35	0.60
60	0.29	0.68	0.97	1.04	0.12	0.31	0.51	0.57

Lifetime Risk of Being Diagnosed = 1.06%
 Lifetime Risk of Dying = 0.53%

Lifetime Risk of Being Diagnosed = 0.61%
 Lifetime Risk of Dying = 0.35%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.27	0.00	0.00	0.00	0.97
10	0.00	0.01	0.02	1.30	0.00	0.00	0.01	0.99
20	0.01	0.02	0.09	1.31	0.00	0.01	0.05	0.99
30	0.02	0.09	0.27	1.34	0.01	0.04	0.14	1.00
40	0.07	0.26	0.61	1.37	0.03	0.13	0.32	1.00
50	0.20	0.58	1.09	1.39	0.10	0.30	0.58	1.01
60	0.44	1.03	1.30	1.38	0.21	0.52	0.85	0.99

Lifetime Risk of Being Diagnosed = 1.27%
 Lifetime Risk of Dying = 0.89%

Lifetime Risk of Being Diagnosed = 0.97%
 Lifetime Risk of Dying = 0.67%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XXV-6
TESTIS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.02	0.12	0.36
10	0.02	0.12	0.24	0.36
20	0.10	0.22	0.29	0.34
30	0.12	0.19	0.22	0.24
40	0.08	0.11	0.12	0.13
50	0.03	0.04	0.05	0.05
60	0.01	0.02	0.02	0.02

Lifetime Risk of Being Diagnosed = 0.36%
 Lifetime Risk of Dying = 0.02%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.03	0.15	0.43
10	0.03	0.15	0.29	0.43
20	0.12	0.26	0.35	0.41
30	0.14	0.23	0.26	0.29
40	0.09	0.13	0.14	0.15
50	0.04	0.05	0.06	0.06
60	0.02	0.02	0.03	0.03

Lifetime Risk of Being Diagnosed = 0.43%
 Lifetime Risk of Dying = 0.02%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.03	0.10
10	0.01	0.03	0.06	0.10
20	0.03	0.06	0.08	0.09
30	0.03	0.05	0.06	0.07
40	0.02	0.03	0.03	0.04
50	0.01	0.02	0.02	0.02
60	0.01	0.01	0.01	0.01

Lifetime Risk of Being Diagnosed = 0.10%
 Lifetime Risk of Dying = 0.01%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XXVI-8
THYROID CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.02	0.35	0.00	0.02	0.12	0.97
10	0.00	0.02	0.06	0.36	0.02	0.12	0.28	0.98
20	0.02	0.06	0.11	0.36	0.10	0.26	0.45	0.96
30	0.04	0.09	0.17	0.34	0.16	0.35	0.53	0.86
40	0.05	0.13	0.22	0.31	0.19	0.37	0.53	0.71
50	0.08	0.17	0.24	0.27	0.19	0.35	0.47	0.53
60	0.10	0.17	0.20	0.20	0.17	0.30	0.35	0.36

Lifetime Risk of Being Diagnosed = 0.35%
 Lifetime Risk of Dying = 0.04%

Lifetime Risk of Being Diagnosed = 0.97%
 Lifetime Risk of Dying = 0.06%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.01	0.02	0.38	0.00	0.02	0.14	1.02
10	0.00	0.02	0.07	0.38	0.02	0.14	0.31	1.03
20	0.02	0.06	0.12	0.38	0.11	0.29	0.49	1.00
30	0.04	0.10	0.18	0.36	0.17	0.38	0.56	0.90
40	0.06	0.14	0.23	0.32	0.20	0.39	0.55	0.73
50	0.08	0.18	0.25	0.27	0.19	0.36	0.48	0.54
60	0.10	0.18	0.20	0.21	0.18	0.30	0.35	0.36

Lifetime Risk of Being Diagnosed = 0.38%
 Lifetime Risk of Dying = 0.05%

Lifetime Risk of Being Diagnosed = 1.02%
 Lifetime Risk of Dying = 0.06%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	0.17	0.00	0.00	0.03	0.53
10	0.00	0.01	0.03	0.18	0.00	0.03	0.10	0.54
20	0.01	0.02	0.04	0.18	0.03	0.10	0.19	0.53
30	0.02	0.04	0.09	0.18	0.07	0.16	0.28	0.51
40	0.02	0.07	0.11	0.16	0.09	0.21	0.31	0.45
50	0.05	0.09	0.14	0.15	0.12	0.23	0.32	0.37
60	0.04	0.10	0.11	0.11	0.12	0.21	0.26	0.27

Lifetime Risk of Being Diagnosed = 0.17%
 Lifetime Risk of Dying = 0.03%

Lifetime Risk of Being Diagnosed = 0.53%
 Lifetime Risk of Dying = 0.06%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.

Table XXVII-8

URINARY BLADDER CANCER (Invasive and In Situ)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (13 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

2000-2002

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	3.58	0.00	0.00	0.00	1.14
10	0.00	0.00	0.02	3.63	0.00	0.00	0.01	1.15
20	0.00	0.02	0.11	3.65	0.00	0.01	0.04	1.15
30	0.02	0.10	0.41	3.69	0.01	0.04	0.13	1.16
40	0.09	0.40	1.23	3.73	0.03	0.12	0.36	1.16
50	0.33	1.19	2.55	3.78	0.10	0.33	0.70	1.15
60	0.93	2.40	3.48	3.72	0.25	0.64	1.01	1.10

Lifetime Risk of Being Diagnosed = 3.58%
Lifetime Risk of Dying = 0.75%

Lifetime Risk of Being Diagnosed = 1.14%
Lifetime Risk of Dying = 0.32%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	3.99	0.00	0.00	0.00	1.24
10	0.00	0.01	0.02	4.03	0.00	0.00	0.01	1.25
20	0.00	0.02	0.12	4.06	0.00	0.01	0.04	1.25
30	0.02	0.11	0.46	4.09	0.01	0.04	0.15	1.26
40	0.09	0.45	1.39	4.14	0.03	0.14	0.40	1.26
50	0.36	1.34	2.86	4.18	0.11	0.38	0.78	1.24
60	1.05	2.68	3.84	4.10	0.28	0.70	1.09	1.18

Lifetime Risk of Being Diagnosed = 3.99%
Lifetime Risk of Dying = 0.80%

Lifetime Risk of Being Diagnosed = 1.24%
Lifetime Risk of Dying = 0.32%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.39	0.00	0.00	0.00	0.80
10	0.00	0.00	0.02	1.43	0.00	0.00	0.01	0.82
20	0.00	0.02	0.07	1.44	0.00	0.00	0.02	0.82
30	0.02	0.07	0.27	1.48	0.00	0.02	0.09	0.83
40	0.06	0.26	0.62	1.51	0.02	0.09	0.26	0.84
50	0.22	0.60	1.14	1.56	0.07	0.25	0.53	0.85
60	0.45	1.07	1.45	1.56	0.19	0.50	0.77	0.85

Lifetime Risk of Being Diagnosed = 1.39%
Lifetime Risk of Dying = 0.39%

Lifetime Risk of Being Diagnosed = 0.80%
Lifetime Risk of Dying = 0.33%

Devcan Version 6.0, April 2005, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia). Mortality data are from the NCHS public use data file for the total US.