

Table 7.25A Tobacco Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH			
	Heavy Use ¹	Binge Use But Not Heavy Use ¹	Use But Not Binge Use ¹	No Use
Any Tobacco ²	11,259	19,345	17,309	23,585
Cigarettes	9,722	16,601	14,365	20,449
Smokeless Tobacco	1,667	2,054	1,314	2,751
Cigars	3,039	4,319	2,937	2,456
Pipes	277	453	520	566

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all Heavy Alcohol Users are also Binge Alcohol Users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 7.25B Tobacco Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH			
	Heavy Use ¹	Binge Use But Not Heavy Use ¹	Use But Not Binge Use ¹	No Use
Any Tobacco ²	71.0	51.0	26.2	20.5
Cigarettes	61.3	43.8	21.8	17.7
Smokeless Tobacco	10.5	5.4	2.0	2.4
Cigars	19.2	11.4	4.4	2.1
Pipes	1.7	1.2	0.8	0.5

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all Heavy Alcohol Users are also Binge Alcohol Users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 7.26A Tobacco Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH			
	Heavy Use ¹	Binge Use But Not Heavy Use ¹	Use But Not Binge Use ¹	No Use
Any Tobacco ²	511	1,124	547	1,585
Cigarettes	449	987	458	1,316
Smokeless Tobacco	101	155	44	188
Cigars	214	407	143	354
Pipes	26	38	16	58

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all Heavy Alcohol Users are also Binge Alcohol Users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 7.26B Tobacco Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH			
	Heavy Use ¹	Binge Use But Not Heavy Use ¹	Use But Not Binge Use ¹	No Use
Any Tobacco ²	81.1	55.9	31.7	7.8
Cigarettes	71.2	49.1	26.6	6.5
Smokeless Tobacco	16.1	7.7	2.5	0.9
Cigars	33.9	20.2	8.3	1.7
Pipes	4.1	1.9	0.9	0.3

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all Heavy Alcohol Users are also Binge Alcohol Users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 7.27A Tobacco Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH			
	Heavy Use ¹	Binge Use But Not Heavy Use ¹	Use But Not Binge Use ¹	No Use
Any Tobacco ²	3,654	4,979	2,375	3,042
Cigarettes	3,293	4,483	2,120	2,754
Smokeless Tobacco	659	475	132	237
Cigars	1,133	1,258	473	554
Pipes	88	142	45	58

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all Heavy Alcohol Users are also Binge Alcohol Users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 7.27B Tobacco Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH			
	Heavy Use ¹	Binge Use But Not Heavy Use ¹	Use But Not Binge Use ¹	No Use
Any Tobacco ²	79.0	61.7	39.2	24.8
Cigarettes	71.2	55.5	35.0	22.5
Smokeless Tobacco	14.2	5.9	2.2	1.9
Cigars	24.5	15.6	7.8	4.5
Pipes	1.9	1.8	0.7	0.5

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all Heavy Alcohol Users are also Binge Alcohol Users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 7.28A Tobacco Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH			
	Heavy Use ¹	Binge Use But Not Heavy Use ¹	Use But Not Binge Use ¹	No Use
Any Tobacco ²	7,095	13,242	14,388	18,959
Cigarettes	5,981	11,131	11,786	16,379
Smokeless Tobacco	907	1,425	1,138	2,327
Cigars	1,693	2,654	2,322	1,547
Pipes	163	274	460	451

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all Heavy Alcohol Users are also Binge Alcohol Users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 7.28B Tobacco Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH			
	Heavy Use ¹	Binge Use But Not Heavy Use ¹	Use But Not Binge Use ¹	No Use
Any Tobacco ²	66.9	47.6	24.7	22.9
Cigarettes	56.4	40.0	20.2	19.8
Smokeless Tobacco	8.6	5.1	2.0	2.8
Cigars	16.0	9.5	4.0	1.9
Pipes	1.5	1.0	0.8	0.5

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all Heavy Alcohol Users are also Binge Alcohol Users.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.