

**REFERENCE COPY**

Do Not Remove from the Library


U. S. Fish and Wildlife Service

Biological Report 82(11.29) National Wetlands Research Center TR EL-82-4  
March 1985

700 Cajun Dome Boulevard  
Lafayette, Louisiana 70506

**Species Profiles: Life Histories and  
Environmental Requirements of Coastal Fishes  
and Invertebrates (Gulf of Mexico)**

**SHEEPSHEAD**



Fish and Wildlife Service  
U.S. Department of the Interior

Coastal Ecology Group  
Waterways Experiment Station  
U.S. Army Corps of Engineers

This is one of the first reports to be published in the new "Biological Report" series. This technical report series, published by the Research and Development branch of the U.S. Fish and Wildlife Service, replaces the "FWS/OBS" series published from 1976 to September 1984. The Biological Report series is designed for the rapid publication of reports with an application orientation, and it continues the focus of the FWS/OBS series on resource management issues and fish and wildlife needs.

Biological Report 82(11.29)  
TR EL-82-4  
March 1985

Species Profiles: Life Histories and Environmental Requirements  
of Coastal Fishes and Invertebrates (Gulf of Mexico)

SHEEPSHEAD

by

Cecil A. Jennings  
P.O. Drawer LW  
Department of Wildlife & Fisheries  
Mississippi State University  
Mississippi State, MS 39762

Project Manager  
Larry Shanks  
Project Officer  
John Parsons  
National Coastal Ecosystems Team  
U.S. Fish and Wildlife Service  
1010 Gause Boulevard  
Slidell, LA 70458

Performed for

Coastal Ecology Group  
U.S. Army Corps of Engineers  
Waterways Experiment Station  
Vicksburg, MS 39180

and

National Coastal Ecosystems Team  
Division of Biological Services  
Research and Development  
Fish and Wildlife Service  
U.S. Department of the Interior  
Washington, DC 20240

This series should be referenced as follows:

U.S. Fish and Wildlife Service. 1983-19\_\_\_. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates. U.S. Fish Wildl. Serv. Biol. Rep. 82(11). U.S. Army Corps of Engineers, TR EL-82-4.

This profile should be cited as follows:

Jennings, C.A. 1985. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Gulf of Mexico) -- sheepshead. U.S. Fish Wildl. Serv. Biol. Rep. 82(11.29). U.S. Army Corps of Engineers, TR EL-82-4. 10 pp.

## PREFACE

This species profile is one of a series on coastal aquatic organisms, principally fish, of sport, commercial, or ecological importance. The profiles are designed to provide coastal managers, engineers, and biologists with a brief comprehensive sketch of the biological characteristics and environmental requirements of the species and to describe how populations of the species may be expected to react to environmental changes caused by coastal development. Each profile has sections on taxonomy, life history, ecological role, environmental requirements, and economic importance, if applicable. A three-ring binder is used for this series so that new profiles can be added as they are prepared. This project is jointly planned and financed by the U.S. Army Corps of Engineers and the U.S. Fish and Wildlife Service.

Suggestions or questions regarding this report should be directed to one of the following addresses.

Information Transfer Specialist  
National Coastal Ecosystems Team  
U.S. Fish and Wildlife Service  
NASA-Slidell Computer Complex  
1010 Gause Boulevard  
Slidell, LA 70458

or

U.S. Army Engineer Waterways Experiment Station  
Attention: WESER-C  
Post Office Box 631  
Vicksburg, MS 39180

## CONTENTS

	<u>Page</u>
PREFACE.....	iii
CONVERSION TABLE.....	v
ACKNOWLEDGMENTS.....	vi
NOMENCLATURE/TAXONOMY/RANGE.....	1
MORPHOLOGY/IDENTIFICATION AIDS.....	1
REASON FOR INCLUSION IN THE SERIES.....	3
LIFE HISTORY.....	3
Spawning.....	3
Larvae.....	3
Juveniles.....	4
Migration.....	4
GROWTH CHARACTERISTICS.....	5
FISHERY.....	5
ECOLOGICAL ROLE.....	5
DISEASE AND PARASITES.....	6
ENVIRONMENTAL REQUIREMENTS.....	6
Temperature and Salinity.....	6
Other Environmental Factors.....	6
LITERATURE CITED.....	8

## CONVERSION TABLE

### Metric to U.S. Customary

<u>Multiply</u>	<u>By</u>	<u>To Obtain</u>
millimeters (mm)	0.03937	inches
centimeters (cm)	0.3937	inches
meters (m)	3.281	feet
kilometers (km)	0.6214	miles
square meters (m <sup>2</sup> )	10.76	square feet
square kilometers (km <sup>2</sup> )	0.3861	square miles
hectares (ha)	2.471	acres
liters (l)	0.2642	gallons
cubic meters (m <sup>3</sup> )	35.31	cubic feet
cubic meters	0.0008110	acre-feet
milligrams (mg)	0.00003527	ounces
grams (g)	0.03527	ounces
kilograms (kg)	2.205	pounds
metric tons (t)	2205.0	pounds
metric tons	1.102	short tons
kilocalories (kcal)	3.968	British thermal units
Celsius degrees	1.8(°C) + 32	Fahrenheit degrees

### U.S. Customary to Metric

inches	25.40	millimeters
inches	2.54	centimeters
feet (ft)	0.3048	meters
fathoms	1.829	meters
miles (mi)	1.609	kilometers
nautical miles (nmi)	1.852	kilometers
square feet (ft <sup>2</sup> )	0.0929	square meters
acres	0.4047	hectares
square miles (mi <sup>2</sup> )	2.590	square kilometers
gallons (gal)	3.785	liters
cubic feet (ft <sup>3</sup> )	0.02831	cubic meters
acre-feet	1233.0	cubic meters
ounces (oz)	28.35	grams
pounds (lb)	0.4536	kilograms
short tons (ton)	0.9072	metric tons
British thermal units (Btu)	0.2520	kilocalories
Fahrenheit degrees	0.5556(°F - 32)	Celsius degrees

## ACKNOWLEDGMENTS

The author is grateful for the reviews by Bennie A. Rohr, National Marine Fisheries Service, Pascagoula, Mississippi, and by Thomas D. McIlwain, Gulf Coast Research Laboratory, Ocean Springs, Mississippi.






Figure 1. Sheepshead.

SHEEPSHEAD

NOMENCLATURE/TAXONOMY/RANGE

Scientific name.....Archosargus probatocephalus  
 Preferred Common Name.....Sheepshead  
 Other Common Names.....English: Sheepshead bream, sheepshead porgie, convict fish; French: Rondeau mouton; Spanish: Sargo chopá, pargo  
 Class.....Osteichthyes  
 Order.....Perciformes  
 Family.....Sparidae

Geographic range: The sheepshead (Figure 1) is common along the Atlantic and Gulf of Mexico coasts (Figure 2) of the United States, from Cape Cod to Texas (Bigelow and Schroeder 1953). It has been reported from Nova Scotia (Gilhen et

al. 1976) to Rio de Janeiro (Randall et al. 1978). The sheepshead is divided into three subspecies: A. p. probatocephalus, the northern form, is distributed from Nova Scotia to Cedar Key, Florida; A. p. oviceps ranges in the Gulf of Mexico from St. Marks, Florida to the Campeche Bank, Mexico; and A. p. aries, the southern form, ranges from Belize to Bahia de Sepetiba, Brazil (Caldwell 1965; Randall et al. 1978; Burgess 1980).

MORPHOLOGY/IDENTIFICATION AIDS

Morphological descriptions and identification aids for the sheepshead in this report were largely extracted




Figure 2. Distribution of sheephead along the coast of the Gulf of Mexico.

from Hildebrand and Schroeder (1927), Hoese and Moore (1977), and Johnson (1978). The primary counts of fin rays are as follows: Dorsal-XII (spiny rays), 10-12 (soft rays); and anal-III, 10-11 respectively. There are 44-49 scales in the lateral line and 8 scales above and 15 below the lateral line at the insertion of the dorsal fin. The body is deep and compressed and the back is elevated above the short and deep head. The snout is short, 2.1 to 2.6 times in head length; the eye is 2.75 to 4.55 in head length. The mouth is moderate and nearly horizontal, 2.7 to 3.3 times in head length. The incisor-like anterior teeth and broad posterior molars are strong. Gill rakers are short, with six to seven on the lower limb of the first arch. Scales are finely serrate. The dorsal fin has strong spines; the spinous portion is longer than the soft portion. The caudal fin has a shallow fork. The second spine of the anal fin is enlarged. The pectoral fins are long, 2.7 to 3.7 times in standard length (SL).

The adult sheepshead is gray to greenish-yellow and has five to seven vertical black crossbars along the body. Crossbars are more distinct in young specimens. The dorsal, anal, and ventral fins are mostly dusky to black; the caudal and pectoral fins are greenish.

#### REASON FOR INCLUSION IN THE SERIES

The sheepshead is a relatively popular sport fish in much of its range (McClane 1964) and usually can be caught throughout the year in the Gulf of Mexico and its estuaries (Perret 1971). It has some commercial value for food (Hildebrand and Schroeder 1927; Perlmutter 1961; Burgess 1980), but its acceptance as a food fish varies among coastal states.

In Alabama, and probably the other gulf states as well, more sheepshead

could be caught without endangering their abundance (Swingle 1977).

Information on the biology and population dynamics of the sheepshead is scarce.

#### LIFE HISTORY

##### Spawning

Sheepshead probably spawn in spring in mid-Atlantic coastal waters (Jordan and Evermann 1896; Hildebrand and Cable 1938; Springer and Woodburn 1960) and in the Mississippi Sound (Christmas and Waller 1973). Spawning probably takes place offshore because larvae and ripe adults are scarce in spring in shallow beach waters and estuaries (Hildebrand and Cable 1938; Springer and Woodburn 1960).

Sheepshead tend to congregate in nearshore waters of the gulf in March, April, and May (Swingle 1977). Adults migrate to offshore waters to spawn in spring and later return to nearshore waters and estuaries.

Sheepshead eggs are about 0.8 mm in diameter (Rathburn 1892), are buoyant, and hatch in about 40 hours at water temperatures near 25°C (Rathburn 1892; Smith 1907).

##### Larvae

The data on larvae in this section are largely taken from Mook (1977). His findings are the result of his examinations of specimens from the collection of the Florida Department of Natural Resources Marine Laboratory.

Yolk-sac larvae are about 2.0 to 4.5 mm long (Johnson 1978) and have a median fin fold (Mook 1977). The caudal fin is the first of the median fins to form, becoming apparent in larvae 3 mm long as ray elements in the post-ventral part of the fin fold. The caudal and anal fins are well developed

in larvae 4-mm long. The pectorals, the first paired fins to develop, appear as small fin buds in 2-mm long larvae and have developed ray elements in larvae 4 mm long. Yolk-sac larvae of sheepshead have about 24 myomeres (successive muscle segments).

Pigmentation in yolk-sac larvae is limited to one small melanophore at the angle of the jaw and three melanophores along the ventral side of the body posterior to the vent. Some individuals may have one or two small melanophores on the yolk sac. Most melanophores disappear in larvae 6 mm long (Mook 1977).

The morphological development of larval sheepshead (5 to 25 mm long) is similar to that of other perciform fishes, but they attain adult characteristics at a smaller size than do other sparid fishes (Mook 1977). At 6 mm, body depth/length ratio is 3.4, head length/body length ratio is 3.0, and the snout length/head length ratio is 4.2 (Hildebrand and Cable 1938; Johnson 1978). The caudal fin is well developed (Hildebrand and Cable 1938), and the dorsal fin has 7 spines and 12 soft rays.

Pigmentation of the body of larval sheepshead is brownish with a median ventral line. There are three black spots -- one behind the isthmus, one below the base of the pectoral fin, and one a short distance anterior to the anal fin (Hildebrand and Cable 1938). Two dark specks are located at the base of the anal fin as well (Hildebrand and Cable 1938, Johnson 1978).

Most evidence suggests that sheepshead eggs hatch in offshore waters and that the larvae or postlarvae move inshore along beaches, and into estuaries (Hildebrand and Cable 1938; McClane 1964; Burgess 1980). Ostracods constitute much of the diet of sheepshead less than 30 mm long (Hildebrand and Cable 1938).

## Juveniles

The dorsal and anal spines of juvenile sheepshead (25 to 30 mm long) are similar to those of adults (Hildebrand and Cable 1938; Johnson 1978). The caudal fin is forked and the lateral line is fully developed. The pectoral fins become larger and oblique instead of rounded as in smaller fish (Hildebrand and Cable 1938). Juvenile sheepshead are most abundant in grass flats and over mud bottoms (Hildebrand and Cable 1938; McClane 1964; Burgess 1980). In late summer, when juveniles are about 40 mm long, they begin leaving the grass flats (Hildebrand and Cable 1938; Johnson 1978) and congregate with adults around stone jetties, breakwaters, piers, and wrecks (Hildebrand and Cable 1938; LaMonte 1952; McClane 1964; Mook 1977; Burgess 1980; Juneau and Pollard 1981).

Juvenile sheepshead collected on grass flats had eaten primarily copepods and filamentous algae (Hildebrand and Cable 1938). Sheepshead that inhabited areas with structures (e.g., pilings and jetties) fed primarily on mollusks and crustaceans (Hildebrand and Cable 1938; LaMonte 1952; Randall et al. 1978).

## Migration

Sheepshead are not a true migratory species (Gilhen et al. 1976) -- their movements are primarily offshore and inshore. In Texas and Louisiana they move offshore with the onset of cool weather in late fall (Gunter 1945; Kelly 1965). In winter, high concentrations of sheepshead have been reported at depths of 7 to 12 fathoms near oil platforms and artificial reefs off the Mississippi River Delta and the Mississippi and Alabama coasts (Bennie A. Rohr, National Marine Fisheries Service, Pascagoula, Mississippi; pers. comm. 1983). Sheepshead overwinter in offshore gulf waters and spawn in spring. After spawning, they

apparently return to nearshore waters.

#### GROWTH CHARACTERISTICS

The sheepshead grows and matures slowly (Hildebrand and Cable 1938; Johnson 1978). No published data were found on growth beyond age group 0 or on size at maturity. Springer and Woodburn (1960) reported that the average total lengths (TL) for young-of-the-year in the Tampa Bay area were 21, 29, and 42 mm long in June, July, and August, respectively.

The heaviest sheepshead recorded weighed 66 kg (Hildebrand and Schroeder 1927; Perlmutter 1961). Large sheepshead in the Chesapeake Bay area ranged from 11 to 33 kg (Hildebrand and Schroeder 1927). Near Beaufort, North Carolina, most sheepshead caught by anglers were 280-380 mm long and weighed 0.5 - 1 kg. Individuals up to 500 mm and 2.3 kg sometimes were caught (Hildebrand and Cable 1938). A sheepshead weighing about 5.4 kg is considered large along the Mississippi coast; they are larger in Louisiana (Bennie A. Rohr, National Marine Fisheries Service, Pascagoula, Mississippi; pers. comm. 1983).

#### FISHERY

In 1964, sheepshead constituted 15% of the brackish water fish caught in the Mobile Delta by sport fishermen (Swingle et al. 1966). In 1975, about 64% of the sheepshead landed in Alabama were taken by sport fishermen (Wade 1977). In 1979, 1.8 million sheepshead were caught in the Gulf of Mexico by sport fishermen (National Marine Fisheries Service 1981). The sport fishery for sheepshead probably produces a major portion of the total landings of sheepshead in the Gulf States.

Commercially, most sheepshead are accidentally caught by shrimp trawlers

and discarded (Swingle 1977; Benson 1982), but some are taken intentionally for the market in March, April, and May. Sheepshead are also caught with bottom longlines, seines, and trammel nets (Randall et al. 1978; Tarver and Savoie 1976; Juneau 1975; Christmas and Waller 1973). In some years, sheepshead compose up to 12% of the finfish biomass taken in 12-m trawls off the Mississippi and Alabama coasts during the National Marine Fisheries Service spring groundfish survey (Bennie A. Rohr, National Marine Fisheries Service, Pascagoula, Mississippi; pers. comm. 1983).

Commercial catches of sheepshead were largest in Florida and Alabama in 1981, and in Texas and Louisiana in 1982 (Table 1) (National Marine Fisheries Service 1983). From 1971 to 1982, annual commercial landings of sheepshead in the northern Gulf of Mexico (including Florida's east coast) was about 453.6 tons (t) (National Marine Fisheries Service 1983). The northern gulf landings (Table 1) were 554 t valued at \$246,500 in 1981 and 558 t valued at \$225,300 in 1982 (National Marine Fisheries Service 1983).

Sheepshead landings declined in the Gulf States from about 1 million lb in 1939 to a low of 572,000 lb in 1965 (Gunter 1945; Lyles 1967; Christmas and Waller 1973). This decline was attributed more to reduced fishing intensity than to a reduction in available stocks. Landing statistics from 1971 to 1982 support this conclusion (Perret 1971; Christmas and Waller 1973).

#### ECOLOGICAL ROLE

Sheepshead are omnivorous, and often eat plant material as well as invertebrates and small vertebrates that inhabit shallow, inshore brackish waters (Hildebrand and Cable 1938; Gunter 1945; Darnell 1961; Johnson

Table 1. Commercial sheepshead landings in the Gulf States, 1981-82 (National Marine Fisheries Service 1983).

State	1981		1982	
	Weight (t)	Value (thousands of dollars)	Weight (t)	Value (thousands of dollars)
Florida	253	\$139.3	77	\$ 43.3
Alabama	111	36.0	132	43.8
Mississippi	80	24.4	68	21.7
Louisiana	59	27.7	135	36.9
Texas	51	19.1	146	79.6
Gulf Total	554	246.5	558	225.3

1978; Kelly 1965; Odum et al. 1982; Overstreet and Heard 1982). The diet of larval, juvenile, and adult sheepshead was summarized by Benson (1982). Sheepshead larvae usually eat copepods, amphipods, and other zooplankton. Small juveniles eat copepods, mysids, polychaetes, and chironomid larvae; and large juveniles and adults eat blue crabs, barnacles, young oysters, clams, other mollusks and crustaceans, and small fish -- primarily young Atlantic croakers (Odum et al. 1982). The role of the sheepshead in the trophic dynamics of estuaries has not been well documented.

#### DISEASE AND PARASITES

Sheepshead are hosts of ciliates (Overstreet and Howse 1977), nematodes (Norris and Overstreet 1975), trematodes (Sparks 1957; Hendrix and Overstreet 1977; Deardorff and Overstreet 1981), and isopods (Overstreet 1978), none of which are known to endanger populations of the species.

#### ENVIRONMENTAL REQUIREMENTS

##### Temperature and Salinity

Sheepshead have been collected from water at temperatures of 5°

(Christmas and Waller 1973; Perret 1971) to 35.1°C (Johnson 1978). Juveniles and adults were collected in gulf waters with a temperature range of 8.0° to 29.6°C. Dunham (1972) and Tarver and Savoie (1976) collected sheepshead from the Lake Pontchartrain-Lake Maurepas estuarine complex in water temperatures of 10° to 34.9°C.

Sheepshead are euryhaline (Gunter 1956) and have been collected in water at salinities of 5.3 to 25.0 ppt (Perret and Caillouet 1974), 2.9 to 22.5 ppt (Dunham 1972), 0.1 to 9.3 ppt (Juneau 1975), 0.0 to 30.0 ppt (Perret 1971), 0.0 to 9.9 ppt (Tarver and Savoie 1976), and 0.0 to 1.5 ppt (Kelly 1965).

No information on lethal upper and lower thermal limits for larvae, juveniles, or adults has been reported. In the Lake Pontchartrain-Lake Maurepas estuarine complex, sheepshead were most abundant at temperatures of 30° to 35°C and salinities of 5 to 10 ppt (Tarver and Savoie 1976).

##### Other Environmental Factors

Young juveniles commonly live in grass flats over mud bottoms, whereas adults and older juveniles usually congregate on the bottom or along the shore near rocks, pilings, breakwaters,

jetties, and piers.

Sheepshead were killed during severe oxygen depletion in semi-open

and closed canals in coastal Louisiana (Adkins and Bowman 1976), but the minimal dissolved oxygen tolerances for sheepshead are unknown.

## LITERATURE CITED


- Adkins, G., and P. Bowman. 1976. A study of the fauna in dredged canals of coastal Louisiana. La. Wildl. Fish. Comm. Tech. Bull. No. 18. 72 pp.
- Benson, N. G., ed. 1982. Life history requirements of selected finfish and shellfish in Mississippi Sound and adjacent areas. U.S. Fish. Wildl. Serv. FWS/OBS-81/51. 97 pp.
- Bigelow, H. B., and W. C. Schroeder. 1953. Fishes of the Gulf of Maine. U.S. Fish. Wildl. Serv. Fish Bull. 53:1-577.
- Burgess, G. H. 1980. Sheepshead. Page 754 in D. S. Lee, ed. Atlas of North American freshwater fishes. Publ. No. 1980-12. North Carolina State Mus. Nat. Hist., Raleigh. 854 pp.
- Caldwell, D. K. 1965. Systematics and variation in the sparid fish Archosargus probatocephalus. Bull. South Calif. Acad. Sci. 64(2):89-100.
- Christmas, J. Y., and R. S. Waller. 1973. Estuarine vertebrates, Mississippi. Page 320-406 in J.Y. Christmas, ed. Cooperative Gulf of Mexico estuarine inventory and study, Mississippi. Gulf Coast Res. Lab., Ocean Springs, Miss.
- Darnell, R. M. 1961. Trophic spectrum of an estuarine community based on studies of Lake Pontchartrain, Louisiana. Ecology 42(3): 553-568.
- Deardorff, T. L., and R. M. Overstreet. 1981. Review of Hysterothylacium and Iheringascaris (both previously = Thynnascaris) (Nematoda:Anisakidae) from the Northern Gulf of Mexico. Proc. Biol. Soc. Wash. 93(4): 1035-1079.
- Dunham, F. 1972. A study of commercially important estuarine dependent industrial fishes. La. Wildl. Fish. Comm. Tech. Bull. No. 4. 63 pp.
- Gilhen, J., C. G. Gruchy, and D. E. McAllister. 1976. The sheepshead, Archosargus probatocephalus, and the feather blenny, Hypsoblennius hentzi, two additions to the Canadian Atlantic ichthyofauna. Can. Field Nat. 90:42-46.
- Gunter, G. 1945. Studies on marine fishes of Texas. Inst. Mar. Sci. Univ. Tex. 1(1):1-190.
- Gunter, G. 1956. A revised list of euryhaline fishes of North America. Am. Midl. Nat. 56(2):345-354.
- Hendrix, S. S., and R. M. Overstreet. 1977. Marine aspidogastriids (Trematoda) from fishes in the northern Gulf of Mexico. J. Parasitol. 63(5):810-817.
- Hildebrand, S. F., and L. E. Cable. 1938. Further notes on the development and life history of some teleosts at Beaufort, N.C. Bull. U.S. Bur. Fish. 48:505-642.



- Hildebrand, S. F., and W. C. Schroeder. 1927. Fishes of Chesapeake Bay. Bull. U.S. Bur. Fish. 43(1): 366 pp.
- Hoese, H. D., and R. H. Moore. 1977. Fishes of the Gulf of Mexico: Texas, Louisiana, and adjacent waters. Texas A&M University Press, College Station. 327 pp.
- Johnson, D. G. 1978. Development of fishes in the mid-Atlantic Bight: an atlas of egg, larval, and juvenile stages. Vol. 4. Carangidae through Ehippidae. U.S. Fish Wildl. Serv. FWS/OBS-78/12. 311 pp.
- Jordan, D. S., and B. W. Evermann. 1896. The fishes of North and Middle America. U.S. Natl. Mus. Bull. 47 (in 4 parts):1-3313.
- Juneau, C. L. 1975. An inventory and study of the Vermilion Bay-Atchafalaya Bay Complex. Phase II, Biology. La. Wildl. Fish. Comm. Tech. Bull. No. 13. 21-74.
- Juneau, C. L., and J. F. Pollard. 1981. A survey of the recreational shrimp and finfish harvest of the Vermilion Bay area and their impact on commercial fishery resources. La. Dep. Wildl. Fish. Tech. Bull. No. 33. 40 pp.
- Kelly, J. R. 1965. A taxonomic survey of the fishes of Delta National Wildlife Refuge with emphasis upon distribution and abundance. M.S. Thesis. Louisiana State University, Baton Rouge. 133 pp.
- LaMonte, F. 1952. Marine game fishes of the world. Doubleday Nature Guide, Doubleday and Co., Inc., Garden City, N.Y. 190 pp.
- Lyles, C. H. 1967. Fishery statistics of the United States, 1965. U.S. Fish Wildl. Serv., Fish. Stat. Dig. 59. 363 pp.
- McClane, A. J. ed. 1964. McClane's standard fishing encyclopedia. Holt, Rinehart, and Winston, New York. 1057 pp.
- Mook, D. 1977. Larval and osteological development of the sheepshead Archosargus probatocephalus (Pisces:Sparidae). Copeia 1977(1):126-133.
- National Marine Fisheries Service. 1981. Current fishery statistics No. 8100, 1980. U.S. Dep. Commer., Natl. Oceanogr. Atmos. Admin.
- National Marine Fisheries Service. 1983. Landings data: Jan. 1971 through Dec. 1981. 19 pp.
- Norris, D. E., and R. M. Overstreet. 1975. Thynnascaris reliquens N. and T. habena (Linton 1900) (Nematoda:Ascaridoidea) from fishes in the northern Gulf of Mexico and eastern U.S. seaboard. J. Parasitol. 61(2):330-336.
- Odum, W. E., C. C. McIvor, and T. J. Smith III. 1982. The ecology of the mangroves of south Florida: a community profile. U.S. Fish Wildl. Serv. FWS/OBS-81/24. 144 pp.
- Overstreet, R. M. 1978. Marine maladies? Worms, germs, and other symbionts from the northern Gulf of Mexico. Miss.-Ala. Publ. Ocean Springs, Miss. MASGP-78-021. 140 pp.
- Overstreet, R. M., and R. W. Heard. 1982. Food contents of six commercial fishes from Mississippi Sound. Gulf Res. Rep. 7(2):137-149.
- Overstreet, R. M., and H. D. Howse. 1977. Some parasites and diseases of estuarine fishes in polluted

- habitats of Mississippi. Annu. N.Y. Acad. Sci. 298:427-426.
- Perlmutter, A. 1961. Guide to marine fishes. Bramhall House, New York. 431 pp.
- Perret, W. S. 1971. Cooperative Gulf of Mexico estuarine inventory study. Phase 4, Biology. Pages 31-69. La. Wildl. Fish. Comm. 171 pp.
- Perret, W. S., and C. W. Caillouet, Jr. 1974. Abundance and size of fishes taken by trawling in Vermilion Bay, Louisiana. Bull. Mar. Sci. 24(10):52-74.
- Randall, J. E., B. Bishop, and R. Vergaro. 1978. SPARID Archo. Page 12 in E. Fischer, ed. FAO species identification sheets for fishery purposes. Western Central Atlantic (fishing area 31). Vol. 5. Food Agric. Organ. U.N., Rome, Italy.
- Rathburn, R. 1892. Successful hatching of sheepshead eggs on Fish Hawk. Rep. U.S. Comm. Fish and Fish. 1888-1889:59.
- Smith, H. M. 1907. The fishes of North Carolina. North Carolina Geol. Econ. Surv. 2:1-453.
- Sparks, A. K. 1957. Some digenetic trematodes of fishes of Grand Isle, Louisiana. La. Acad. Sci. 20:71-82.
- Springer, V. G., and K. D. Woodburn. 1960. An ecological study of the fishes of the Tampa Bay area. Fla. Board Conserv. Mar. Lab. Prof. Pap. Ser. 1. 104 pp.
- Swingle, H. A. 1977. Coastal fishery resources of Alabama. Ala. Mar. Res. Bull. 12:31-58.
- Swingle, W. E., S. L. Spencer, and T. M. Scott, Jr. 1966. Statistics of the sport fishery of the Mobile Delta during the period of July 1, 1963 to June 30, 1964. Proc. Southeast. Assoc. Game Fish. Comm. 19:439-446.
- Tarver, J. W., and L. B. Savoie. 1976. An inventory and study of the Lake Pontchartrain-Lake Maurepas estuarine complex. Phase II, Biology. La. Wildl. Fish. Comm. Tech. Bull. 19:7-99.
- Wade, C. W. 1977. Survey of the Alabama marine recreational fishery. Ala. Mar. Res. Bull. 12:1-22.

<b>REPORT DOCUMENTATION PAGE</b>	<b>1. REPORT NO.</b> Biological Report 82(11.29)*	<b>2.</b>	<b>3. Recipient's Accession No.</b>
<b>4. Title and Subtitle</b> Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Gulf of Mexico) -- Sheepshead		<b>5. Report Date</b> March 1985	
<b>7. Author(s)</b> Cecil A. Jennings		<b>8. Performing Organization Rept. No.</b>	
<b>9. Performing Organization Name and Address</b> MS Cooperative Fish and Wildlife Research Unit P.O. Drawer BX Mississippi State University Mississippi State, MS 39762		<b>10. Project/Task/Work Unit No.</b>	
<b>12. Sponsoring Organization Name and Address</b> National Coastal Ecosystems Team Division of Biological Services Fish and Wildlife Service U.S. Department of Interior Washington, DC 20240		<b>11. Contract(C) or Grant(G) No.</b> (C) (G)	
<b>15. Supplementary Notes</b> *U.S. Army Corps of Engineers Report No. TR EL-82-4		<b>13. Type of Report &amp; Period Covered</b>  <b>14.</b>	
<b>16. Abstract (Limit: 200 words)</b> Species profiles are literature summaries of the taxonomy, morphology, range, life history, and environmental requirements of coastal aquatic species. They are prepared to assist in environmental impact statements. The sheepshead, <i>Archosargus probatocephalus</i> , contributes moderately to the multi-million dollar commercial and sport fishery in the northern Gulf of Mexico. Commercial landings for 1982 were 558 t valued at \$225,300; more than 1.8 million sheepshead were caught by anglers in 1979. Juveniles and adults move offshore as temperatures decline in fall. Spawning probably takes place in offshore waters during the spring. Larvae move inshore to grass flats in estuaries where they feed on soft plants and animals. Juveniles and adults live near structures on the bottom, such as pilings, or along the shore; they are omnivorous, but primarily eat mollusks and crustaceans. Sheepshead can tolerate wide ranges of temperatures and salinities. Little is known about the actual time of inshore and offshore movements, growth rate, age composition, role in trophic dynamics of estuaries, and age at maturity.			
<b>17. Document Analysis a. Descriptors</b> Estuaries                      Spawning Fishes                              Feeding Growth  <b>b. Identifiers/Open-Ended Terms</b> Sheepshead                              Spawning <i>Archosargus probatocephalus</i> Habitat requirements Salinity requirements                      Life history  <b>c. COSATI Field/Group</b>			
<b>18. Availability Statement</b> Unlimited	<b>19. Security Class (This Report)</b> Unclassified	<b>21. No. of Pages</b> 16	
	<b>20. Security Class (This Page)</b> Unclassified	<b>22. Price</b>	



**REGION 1**

Regional Director  
 U.S. Fish and Wildlife Service  
 Lloyd Five Hundred Building, Suite 1692  
 500 N.E. Multnomah Street  
 Portland, Oregon 97232

**REGION 2**

Regional Director  
 U.S. Fish and Wildlife Service  
 P.O. Box 1306  
 Albuquerque, New Mexico 87103

**REGION 3**

Regional Director  
 U.S. Fish and Wildlife Service  
 Federal Building, Fort Snelling  
 Twin Cities, Minnesota 55111

**REGION 4**

Regional Director  
 U.S. Fish and Wildlife Service  
 Richard B. Russell Building  
 75 Spring Street, S.W.  
 Atlanta, Georgia 30303

**REGION 5**

Regional Director  
 U.S. Fish and Wildlife Service  
 One Gateway Center  
 Newton Corner, Massachusetts 02158

**REGION 6**

Regional Director  
 U.S. Fish and Wildlife Service  
 P.O. Box 25486  
 Denver Federal Center  
 Denver, Colorado 80225

**REGION 7**

Regional Director  
 U.S. Fish and Wildlife Service  
 1011 E. Tudor Road  
 Anchorage, Alaska 99503



# DEPARTMENT OF THE INTERIOR

## U.S. FISH AND WILDLIFE SERVICE



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.