THE IHS PRIMARY Care provider



A journal for health professionals working with American Indians and Alaska Natives

September 1997

Published by the IHS Clinical Support Center

Volume 22, Number 9

The Accreditation Association for Ambulatory Health Care: An Option for Ambulatory Health Centers

Robert W. Boyce, RPh, PP, FASHP, Director of Pharmacy, and Facility Accreditation Coordinator; James E. Edge, RPh, MPH, Service Unit Director; Mark J. Nurre, MD, Clinical Director; and Donnie Lee, MD, Quality Improvement Coordinator; all from the Western Oregon Service Unit, Chemawa Indian Health Center, Salem, Oregon.

Introduction

In February 1997, the Chemawa Indian Health Center of the Western Oregon Service Unit became the first facility in the Indian Health Service to be accredited by the Accreditation Association for Ambulatory Health Care (AAAHC). Achievement of accreditation by the AAAHC was a result of many months of work in evaluating various accrediting organizations and, then, requesting a special exception from the Indian Health Service (IHS) to seek an alternative method of accreditation other than the Joint Commission on Accreditation of Healthcare Organizations (JCAHO).

This is not the first time that the Chemawa Indian Health Center has blazed the trail for Ambulatory Health Centers in the IHS. Chemawa was one of the first Ambulatory Health Centers in the IHS to be accredited by the JCAHO. This accreditation occurred 16 years ago, in 1981, and we have remained JCAHOaccredited since. Following our last JCAHO survey in 1993, we were awarded accreditation with commendation.

Why Seek Alternative Accreditation?

Since the 1993 JCAHO survey, many of our staff questioned the need to continue to be accredited by JCAHO. We felt that the JCAHO (which is primarily hospital-based in focus) really did not meet our needs as an ambulatory health center, and that there must be a more appropriate accrediting body that focused on ambulatory care rather than hospital care. Additionally, we were looking for an organization that had "deemed status" from the Health Care Financing Administration, which is a requirement of IHS.* Concurrently, Service Unit Directors in the Portland Area were expressing concern about the burden, both financial and workload, placed upon their health care programs by the JCAHO accreditation process. They were seeing JCAHO accreditation becoming more expensive (most ambulatory surveys cost in the neighborhood of \$10,000) and increasingly more complex with

* The "deemed status" of an accrediting organization allows entities that it accredits to collect Medicare reimbursement

In This Issue...

- 137 The Accreditation Association for Ambulatory Health Care: An Option for Ambulatory Health Centers
- 140 Injuries and the Ten Leading Causes of Death for Native Americans in the U.S.: Opportunities for Prevention
- 146 Letter to the Editor: Pre-Medical Students Interested in Volunteering With IHS
- 147 Secretary's Quality of Work Life Initiative
- 149 Meetings of Interest
- 150 NCME Videotapes Available
- 151 Special Announcements

regard to documentation requirements that placed a heavy burden on their health care providers. This was occurring despite the fact that the JCAHO accreditation process had established a separate ambulatory component.

In late 1995, the Chemawa Indian Health Center asked the Area Director, Portland Area IHS, for permission to explore alternatives to JCAHO, with the understanding that we would continue to maintain some type of nationally recognized accreditation to assure both ourselves and our patients that we truly do provide high quality health care. We researched a number of possible accreditation services, and selected the Accreditation Association for Ambulatory Health Care (AAAHC). At the same time, the Portland Area Chief Medical Officer was working at the IHS Headquarters level to get the IHS regulations modified to formally allow this.

In January 1996, the Portland Area Director approved the Western Oregon Service Unit to be a pilot project and undergo an AAAHC survey, stating that there would then be evaluation of the process and outcome of that survey before other service units in the Portland Area might be allowed to pursue similar accreditation. Over several months, the Western Oregon Service Unit proceeded with preparation for such a survey and contacted the AAAHC to arrange the details of the survey process. The Application for Accreditation, Presurvey Questionnaire, and supporting documents were submitted to AAAHC in September 1996. In October, AAAHC contacted the service unit with the scheduled dates of the survey, and the service unit was surveyed on February 3 and 4, 1997.

What is the AAAHC?

The Accreditation Association for Ambulatory Health Care is a private, not-for-profit organization that was formed in 1979 to assist ambulatory health care organizations in improving the quality of care they provide to their patients. It accomplishes this by setting standards, which are similar to JCAHO standards, measuring performance, providing consultation and education where needed, and, ultimately, by awarding accreditation to those organizations that are found to be in compliance with its standards. AAAHC surveyors are clinical health care professionals who practice in settings similar to those that the AAAHC surveys. All surveyors are volunteers, serving without pay because they believe in promoting high quality ambulatory health care. They conduct an on-site survey, and the results, with the facility self-assessment, are sent to the AAAHC board which renders the final decision. This board is appointed by 13 of the nation's leading health care associations (see Table 1). Accreditation may be awarded for either one or three years, depending on the level of compliance with the standards.

Table 1. Organizations that appoint members to the AAAHC Board of Directors.

- Medical Group Management Association
- Federated Ambulatory Surgery Association
- American College Health Association
- National Association of Community Health Centers
- Outpatient Ophthalmic Surgery Society
- American Academy of Facial Plastic and Reconstructive Surgery
- American Society of Outpatient Surgeons
 - American College of Occupational and Environmental Medicine
- American Academy of Dental Group Practice
- American Association of Oral and Maxillofacial Surgeons
- American Academy of Cosmetic Surgery
- Associations of Freestanding Radiation Oncology Centers
- American Society for Dermatologic Surgery.

The Standards

AAAHC has 8 Core Standards and 14 Adjunct Standards. The titles of the standards (see Table 2) are very similar to the 1992 Joint Commission Standards for Ambulatory Health Care. The core standards are a requirement that every ambulatory health center must meet. These standards are more general than those of the JCAHO, and AAAHC does not specifically tell the organization being accredited what they have to do to meet these standards. For example, AAAHC requires the organization to be able to show how it meets the standard, not what type of activity to do to meet it. The adjunct standards refer only to specific services for unique ambulatory care environments, such as dental, pharmacy, laboratory, or surgical services. Thus, a health center only has to prepare for the specific services they provide. This philosophy of how to comply with the standards is the main difference between AAAHC and JCAHO.

Findings

The following are some of the important lessons from both the preparation process and the survey itself at our service unit:

- The costs of our AAAHC survey were \$375 for the application fee and \$3,825 for the survey fee. An additional cost that we incurred was a separate survey of our laboratory by a laboratory accreditation organization, for a fee of \$1200. For this we used the Commission on Laboratory Accreditation (COLA), and that survey was conducted separately about a month after our AAAHC survey. This was necessary because AAAHC does not have deemed status for the Clinical Laboratory Improvement Act (CLIA).
- The standards of the AAAHC and the JCAHO are nearly identical. Our process of preparing for the survey, though,

Table 2. The AAAHC standards.

Core Standards

- Rights of Patients
- Governance
- Administration
- Quality of Care Provided
- Quality Management and Improvement
- Clinical Records
- Professional Improvement
- Facilities and Environment
- Diagnostic Imaging Services
- Radiation Oncology Treatment Services
- Occupational Health Services
- Other Professional and Technical Services
- Teaching and Publication Activities
- Research Activities

Adjunct Standards

- Anesthesia Services
- Surgical Services
- · Overnight Care and Services
- Dental Services
- Emergency Services
- Immediate/Urgent Care Services
- Pharmaceutical Services
- · Pathology and Medical Laboratory Services

was not as intensive as it had been previously when preparing for a JCAHO survey. The AAAHC preparation did include a review of our policies and procedures, updating of manuals and minutes, assigning individual staff members to review our compliance with the chapters in their field of expertise, and, of course, the apprehension and near panic that immediately precedes an important survey such as this.

The primary difference between AAAHC and JCAHO seemed to us to be a difference in philosophy: a difference in the way that the standards are looked at and evaluated. The JCAHO has always used the philosophy that "if it's not documented in writing, we assume it didn't happen, and the standard is not met." AAAHC's philosophy is that "if you can explain to us how you meet the standard and show us that you do meet it, you are in compliance with the standard." They state, appropriately, that a small ambulatory health program should not be held to the same documentation standards as a large hospital.

The actual survey went very well and we were pleased with both the more relaxed process of the AAAHC survey and the helpfulness of the AAAHC surveyor. (Even though more relaxed, however, we felt he was very thorough and complete in his review of all areas.) At the closeout, he complimented us on the quality of our program and stated that he found no substantial problems.

• We received the final report in 8 weeks, and it was more thorough than that done in the past by the JCAHO. We received a bound report of all of the surveyors' comments and the rating we received from the committee. We did not have any areas of noncompliance. If there had been areas of non-compliance, the AAAHC does not require the organization to respond in writing. They do expect that these areas will be corrected before the next survey.

Recommendations

We felt that our pilot project was very successful and were pleased with the process and outcome of the AAAHC survey. Quality was not compromised with AAAHC, but rather we feel it was enhanced. The AAAHC gives more freedom to improve quality and has less tendency to require a specific activity be measured to fulfill a standard. We would recommend that all Ambulatory Health Centers in Indian Country consider this organization prior to preparing for their next accreditation survey.

Injuries and the Ten Leading Causes of Death for Native Americans in the U.S.: Opportunities for Prevention

L.J. David Wallace, MSEH, Injury Prevention Specialist, Division of Unintentional Injury Prevention; David A. Sleet, PhD, Associate Director for Science, Division of Unintentional Injury Prevention; and Steven P. James, Computer Specialist, Office of Statistics and Programming; all from the National Center for Injury Prevention and Control, Centers for Disease Control and Prevention, Atlanta, Georgia.

Introduction

Identifying the leading causes of death for American Indians and Alaska Natives (also referred to in this article as Native Americans) by age-group and sex is one way to help prioritize public health needs and to determine the main causes of premature mortality. To study the effect of injuries among Native Americans and to demonstrate the need for prevention programs, the authors have highlighted the unintentional injury- and violence-related causes of death, by age group, for Native Americans for the years 1992-1994. These data should also prove useful for identifying other important causes of death among Native Americans.

Methods

These data are from Centers for Disease Control and Prevention's (CDC) National Center for Health Statistics (NCHS) and include all deaths of American Indians and Alaska Natives who resided in the United States during 1992-1994. In a few instances for the tenth leading cause of death, there were ties among two or more causes, and such instances are footnoted. The standard cause-of-death groupings used by the NCHS from the International Classification of Diseases, 9th Revision (ICD9) were used to define the causes of death, except that alcohol abuse (ICD9 code 305.0, *nondependent alcohol abuse**) was separated from *alcoholism*. We used the following ICD9 codes to define deaths due to *alcoholism*: 291, 303, 425.5, 535.3, 571.0-571.3, 790.3, and E860.[†]

Because motor vehicle crashes are such a significant primary cause of death, the authors have identified their impact separately from the rest of the unintentional injury category. The category *other unintentional injury* includes all deaths due to unintentional injury <u>except</u> those that are related to motor vehicle crashes.

Key Findings

Figures 1 through 3 show the ten leading causes of death in American Indians and Alaska Natives during the years 1992 through 1994, by age group, and by sex. Key findings include:

- Unintentional injury, of which motor vehicle-related injury is the greatest contributor, is the leading cause of death for Native Americans age 1-44 years.
- The number of American Indian and Alaska Native youth and young adults age 15-24 years who were killed by injuries and violence is 10 times greater than the combined number who died of the next six leading causes.
- When combined, unintentional injury (including injury due to motor vehicle crashes), homicide, and suicide are the leading cause of death among males, and the second leading cause among both sexes combined. Unintentional injuries are the third leading cause of death among females.
- Suicide outranks homicide as a cause of death, especially among males.
- Over four times more Native American men than women died from suicide.
- Over a third (37.3%) of all suicides occurred before the age of 25.
- More Native American men died of injuries related to motor vehicle crashes, other unintentional injuries, suicides, and homicides than from heart disease or cancer.

Discussion

American Indians and Alaska Natives are at risk for injury-related death for several reasons. First, on average this

^{*} Nondependent alcohol abuse includes drunkenness not otherwise specified (NOS), excessive drinking of alcohol NOS, alcohol "hangover," and inebriety NOS.

[†] Deaths due to alcoholism include alcoholic psychoses, alcohol dependence syndrome, alcoholic cardiomyopathy, alcoholic gastritis, alcoholic fatty liver, acute alcoholic hepatitis, alcoholic cirrhosis of the liver, unspecified alcoholic liver damage, excessive blood level of alcohol, and accidental poisoning by alcohol.

Figure 1. Ten leading causes of death in American Indians and Alaska Natives, by age group, 1992-1994.*

KOUPS	55-64 ≥65 Total	Heart Heart Heart Disease Disease 0,454	Malignant Malignant Malignant Neoplasms Neoplasms 2,558 4,471	Diabetes Cerebro- Motor 361 vascular Vehicle 945 2,002	Alcoholism Diabetes Other 230 765 Unint. Injury 1,627	Cerebro- Pneumonia Alcoholism vascular & Influenza 1,381 178 709	OtherBronchitisDiabetesUnint. InjuryEmphysema1,354140Asthma 5851	MotorOtherOrecebro-VehicleUnint. Injuryvascular137266I,286	BronchitisNephritisPneumoniaEmphysema253& InfluenzaAsthma 1351,021	Liver Alcoholism Suicide Disease 157 784 93	Pneumonia Septicemia Bronchitis & Influenza 150 Emphysema
	35-44 45-54	Alcoholism Heart 421 Disease 676	Motor Malignant Vehicle Neoplasms 329 529	Heart Alcoholism Disease 385 271	Other Motor Unint. Injury Vehicle 268 188	Malignant Other Veoplasms Unint. Injury 220 178	Homicide Diabetes	Suicide Liver 144 Disease 78	HIV Cerebro- 100 vascular 76	Liver Suicide Disease 61 70	Pneumonia Homicide & Influenza 53† 66
	25-34 3	Motor Ald Vehicle 495	Other N Unint. Injury V 360	Suicide 238 D	Homicide (215 Uni	Alcoholism Ma 164 Ne	HIV 124	Malignant S i Neoplasms 88	Heart Disease 80	Alcohol Abuse 35	Pneumonia Pn & Influenza & I
AGE GROUPS	15-24	Motor Vehicle 563	Suicide 272	Homicide 189	Other Unint. Injury 185	Malignant Neoplasms 34	Alcoholism 24	Heart Disease 24	Congenital Anomalies 13	Alcohol Abuse 12	Pneumonia & Influenza o
	10-14	Motor Vehicle 49	Other Unint. Injury 45	Homicide 22	Suicide 21	Malignant Neoplasms 17	Congenital Anomalies 7	Heart Disease 3	Pneumonia & Influenza 3	HIV 2	Septicemia 2
	5-9	Other Unint. Injury 45	Motor Vehicle 39	Malignant Neoplasms 15	Heart Disease 8	Congenital Anomalies 7	Homicide 5	Pneumonia & Influenza 4	Meningitis 3	Septicemia 3	Benign Neoplasms 1+
	1-4	Other Unint. Injury 94	Motor Vehicle 77	Homicide 31	Congenital Anomalies 30	Pneumonia & Influenza 15	Malignant Neoplasms 13	Heart Disease 12	Meningitis 6	Bronchitis Emphysema Asthma 4	Cerebro- vascular 3+
	7	SIDS 258	Congenital Anomalies 247	Prem./Resp. Distress Synd. 92	Other Unint. Injury 43	Pneumonia & Influenza 42	Placenta Cord Membranes 36	Maternal Complications 25	Perinatal Infections 22	Homicide 21	Intrauterine Hypoxia 20
	RANK	7	2	e	4	a	Q	2	8	6	10

Source: NCHS Mortality Tapes, 1992-1994. Includes all American Indians and Alaska Natives in the United States.
* Motor vehicle fatalities separated from other unintentional injury category. Number of deaths during the 3 year period for each cause is shown in the appropriate box.
† Tied with at least one other cause of death category.

Figure 2. Ten leading causes of death in American Indian and Alaska Native females, by age group, 1992-1994.*

			JS JS	w		. 4)	ε	lia 2a	ury	s na 71	
	Total	Heart Disease 2,713	Malignant Neoplasms 2,138	Diabetes 795	Cerebro- vascular 716	Motor Vehicle 644	Alcoholism 517	Pneumonia & Influenza 475	Other Unint. Injury 393	Bronchitis Emphysema Asthma 371	Liver Disease 192†
AGE GROUPS	≥65	Heart Disease 2,052	Malignant Neoplasms 1,228	Cerebro- vascular 551	Diabetes 484	Pneumonia & Influenza 359	Bronchitis Emphysema Asthma 271	Nephritis 143	Other Unint. Injury 97	Septicemia 86	Liver Disease 68
	55-64	Malignant Neoplasms 451	Heart Disease 382	Diabetes 203	Cerebro- vascular 87	Alcoholism 83	Bronchitis Emphysema Asthma 67	Liver Disease 49	Motor Vehicle 48	Other Unint. Injury 37	Pneumonia & Influenza 33
	45-54	Malignant Neoplasms 279	Heart Disease 161	Alcoholism 139	Diabetes 83	Motor Vehicle 59	Cerebro- vascular 40	Other Unint. Injury 35	Liver Disease 28	Bronchitis Emphysema Asthma 23	Suicide 19
	35-44	Alcoholism 164	Malignant Neoplasms 126	Motor Vehicle 92	Heart Disease 68	Other Unint. Injury 48	Homicide 36	Liver Disease 34	Suicide 24	Pneumonia & Influenza 21	Cerebro- vascular 20
	25-34	Motor Vehicle 152	Alcoholism 73	Other Unint. Injury 60	Homicide 53	Malignant Neoplasms 44	Suicide 38	Heart Disease 26	HIV 16	Pneumonia & Influenza 15	Alcohol Abuse 13
	15-24	Motor Vehicle 172	Suicide 44	Other Unint. Injury 37	Homicide 30	Malignant Neoplasms 16	Heart Disease 6	Congenital Anomalies <i>5</i>	Pregnancy Complications 4	Alcoholism 3	Cerebro- vascular 3†
	10-14	Motor Vehicle 18	Other Unint. Injury 11	Suicide 8	Malignant Neoplasms 7	Congenital Anomalies <i>5</i>	Homicide 2	Pneumonia & Influenza 2	Septicemia 2	Anemias 1	Heart Disease 1†
	5-9	Motor Vehicle 21	Other Unint. Injury 13	Malignant Neoplasms 7	Congenital Anomalies 3	Homicide 3	Benign Neoplasms 1	Heart Disease 1	Meningo Coccal 1	Pneumonia & Influenza 1	Speticemia 1
	1-4	Other Unint. Injury 36	Motor Vehicle 34	Congenital Anomalies 15	Homicide 14	Malignant Neoplasms 8	Pneumonia & Influenza 8	Heart Disease 3	HIV 3	Meningitis 3	Benign Neoplasms 2
	>1	Congenital Anomalies 124	SIDS 101	Prem./Resp. Distress Synd. 36	Other Unint. Injury 19	Placenta Cord Membranes 17	Pneumonia & Influenza 16	Maternal Complications 10	Homicide 9	Intrauterine Hypoxia 9	Perinatal Infections 7
	RANK	7	2	e	4	2	9	7	8	6	10

Source: NCHS Mortality Tapes, 1992-1994. Includes all American Indians and Alaska Natives in the United States.
* Motor vehicle fatalities separated from other unintentional injury category. Number of deaths during the 3 year period for each cause is shown in the appropriate box.
† Tied with at least one other cause of death category.

Figure 3. Ten leading causes of death in American Indian and Alaska Native males, by age group, 1992-1994.*

		0	ns NS	. 4)	ury	E	۵	1 -	S	e e	za
	Total	Heart Disease 3,741	Malignant Neoplasms 2,303	Motor Vehicle 1,358	Other Unint. Injury 1,234	Alcoholism 864	Suicide 640	Cerebro- vascular 570	Diabetes 559	Homicide 554	Pneumonia & Influenza 546
AGE GROUPS	≥65	Heart Disease 2,177	Malignant Neoplasms 1,330	Cerebro- vascular 394	Pneumonia & Influenza 350	Bronchitis Emphysema Asthma 314	Diabetes 281	Other Unint. Injury 169	Nephritis 110	Alcoholism 102	Motor Vehicle 69
	55-64	Heart Disease 741	Malignant Neoplasms 541	Diabetes 158	Alcoholism 147	Other Unint. Injury 103	Cerebro- vascular 91	Motor Vehicle 89	Bronchitis Emphysema Asthma 68	Pneumonia & Influenza 57	Liver Disease 44
	45-54	Heart Disease 515	Malignant Neoplasms 250	Alcoholism 246	Other Unint. Injury 143	Motor Vehicle 129	Diabetes 86	Liver Disease 50	Suicide 42	Homicide 39	Pneumonia & Influenza 36†
	35-44	Alcoholism 257	Motor Vehicle 237	Other Unint. Injury 220	Heart Disease 203	Suicide 120	Homicide 111	Malignant Neoplasms 94	HIV 89	Pneumonia & Influenza 45	Liver Disease 36
	25-34	Motor Vehicle 343	Other Unint. Injury 300	Suicide 200	Homicide 162	HIV 108	Alcoholism 91	Heart Disease 54	Malignant Neoplasms 44	Alcohol Abuse 22	Liver Disease 17
	15-24	Motor Vehicle 391	Suicide 228	Homicide 159	Other Unint. Injury 148	Alcoholism 21	Heart Disease 18	Malignant Neoplasms 18	Alcohol Abuse 10	Congenital Anomalies 8	HIV 6†
	10-14	Other Unint. Injury 34	Motor Vehicle 31	Homicide 20	Suicide 13	Malignant Neoplasms 10	Congenital Anomalies 2	Heart Disease 2	HIV 2	Bronchitis Emphysema Asthma 1	Meningitis 1†
	5-9	Other Other Unint. Injury Unint. Injury 58 32	Motor Vehicle 18	Malignant Neoplasms 8	Heart Disease 7	Congenital Anomalies 4	Meningitis 3	Pneumonia & Influenza 3	Homicide 2	Septicemia 2	Suicide 1†
	1-4	Other Unint. Injury 58	Motor Vehicle 43	Homicide 17	Congenital Anomalies 15	Heart Disease 9	Pneumonia & Influenza 7	Malignant Neoplasms <i>5</i>	Bronchitis Emphysema Asthma 3	Meningitis 3	Meningo Coccal 2†
	>1	SIDS 157	Congenital Anomalies 123	Prem./Resp. Distress Synd. 56	Pneumonia & Influenza 26	Other Unint. Injury 24	Placenta Cord Membranes 19	Maternal Complications 15	Perinatal Infections 15	Homicide 12	Intrauterine Hypoxia 11
	RANK	-	5	ю	4	Q	9	7	8	6	10

Source: NCHS Mortality Tapes, 1992-1994. Includes all American Indians and Alaska Natives in the United States. * Motor vehicle fatalities separated from other unintentional injury category. Number of deaths during the 3 year period for each cause is shown in the appropriate box. † Tied with at least one other cause of death category.

population is younger than the total U.S. population (median age: 24 years versus 33 years¹), and young persons are at higher risk for injury because of risk taking behaviors such as drinking and driving; not wearing seat belts; and, especially for Native Americans, alcohol abusive drinking styles (sporadic binge drinking).² Second, about half of all Native Americans live in rural or non-metropolitan areas,³ which means that they may have limited access to advanced emergency medical care when an injury occurs (such as from a motor vehicle crash) and their treatment may be delayed. In addition, because alcohol is not legally available in many Native American communities, to obtain alcohol, those residents who drink may drive long distances and then return on isolated two-lane highways while impaired. Third, numerous studies have shown that, for many Native American communities, alcohol plays a substantial role in injury occurrence and premature mortality.⁴⁻⁶ The role of alcohol in unintentional injuries, homicides, and suicides could not be determined from these data. However, other researchers have estimated that in states with reservations, 75% of suicides, 80% of homicides, and 65% of motor vehicle-related deaths among Native Americans involve alcohol.⁷ Although alcohol and abusive binge drinking appear to be important risk factors for injury death among Native Americans, we must note that a substantial proportion of the population, especially older adults, do not drink.^{2,6}

Preventing Motor Vehicle-Related Injuries

In terms of prevention, important steps have been taken to reduce injuries through the efforts of tribes, the Indian Health Service (IHS), and others. Effective injury prevention strategies, such as increasing occupant restraint use by passing tribal occupant restraint laws (or adopting the state law) combined with strict enforcement, and highway lighting projects to target pedestrian injuries are well documented.^{8,9} The Navajo Nation is a good example of a tribe that successfully increased occupant restraint use through this type of program.⁹ However, a 1995 survey of tribal traffic laws found that of 174 tribes reporting, 63 still did not have a seat belt law and 56 did not have a child passenger restraint law.¹⁰

Even in instances where tribes have laws, they are not always rigorously enforced. In one example, from a tribal community in New Mexico with a child passenger restraint law but very little enforcement, toddler child restraint usage was only 9% (unpublished survey data, New Mexico Department of Health, 1996). Passage of occupant restraint laws in combination with public information campaigns, low cost child restraint programs, and strict enforcement of the law by police and tribal judges are still vital to reducing motor vehiclerelated injuries among Native Americans. Enforcing tribal seat belt laws will also help police target drunk driving, because those drivers who drink are much less likely to wear seat belts than those drivers who do not drink. One recent study in Washington found that only 36% of intoxicated Native American drivers involved in a crash reported using seat belts.¹¹ In a similar study in New Mexico, only 4% of drivers involved in an alcohol-related motor vehicle crash were wearing seat belts (Philip May, PhD, personal communication, 1996).

Our data for 1992-1994 show that 36% of the Native Americans who died in motor vehicle crashes were children or young adults less than age 25. Teenage drivers are particularly at risk from dying in a car crash for at least three reasons:

- Driver inexperience: Nationally, most passenger deaths among teenagers (65%) occur in crashes in which another teen is driving.¹²
- Nighttime driving: About half of all deaths among teenagers due to motor vehicle crashes occur between 9 p.m. and 6 a.m.¹² Research has shown that cities with curfew laws prohibiting teens 13-17 years old from being out between 9 p.m. and 6 a.m. have had a 23% reduction in teen motor vehicle deaths.¹³
- Alcohol use: The combination of driver inexperience and • alcohol is a deadly mix: female teenage drivers are 54 times more likely to die in a motor vehicle crash when their blood alcohol concentrations are between 0.05 g/dl and 0.10 g/dl. Male teenage drivers are 18 times more likely to die under such circumstances.¹² Parents can enforce a zero tolerance of alcohol with their teen drivers and not allow their children to ride with others who have been drinking. Many tribes have the authority to revoke or restrict driving privileges on the reservations, enforce zero alcohol tolerance for underage drinking, and set curfew ordinances. Tribes should consider adopting components of new legislation being adopted by states on graduated licensing,14 impaired driving, and nighttime curfews to help reduce deaths related to motor vehicle crashes among teenagers.

Preventing Violence-Related Injuries

Many promising programs and resources are available for preventing suicide, family and intimate violence, and youth violence.¹⁵⁻¹⁷ The April 1997 issue of *The IHS Primary Care Provider* included articles^{18,19} on and references²⁰ to preventing violence in Native American communities, and resource contacts¹⁸ for tribes interested in suicide prevention programs, and family violence programs.

Suicide among Native Americans has the greatest impact among youth and young adults. From 1992 through 1994, 37.3% of all suicides occurred before age 25, and 67.7% before age 35. There are promising strategies to prevent suicide such as training school and community gatekeepers; educating the community; screening for suicide risk in schools and clinics; developing peer support projects; and restricting access to lethal means.^{15,20} The American Indian and Alaska Native Community Suicide Prevention Center and Network of the Jicarilla Apache Tribe, in Dulce, New Mexico, is a good example of a community-based program that has shown promising results in reducing suicides among youth and young adults.¹⁸ The Center staff provides training, technical assistance, and program support to tribes who are interested in developing their own suicide prevention programs. Restricting access to lethal means for those considered at risk for suicide may be an effective, yet underused, strategy for preventing suicide. Firearms are used in approximately 59% of suicides by Native American males, and 41% of females.²⁰ In some areas, Alaska for example, the rates are much higher.^{20,21} Friends and relatives of persons at risk for suicide, and the public at large, should be made aware of research findings indicating that having a firearm in the home is associated with approximately a five-fold increase in the risk of suicide among household residents.²²

A Multifaceted, Community-Based Approach

Preventing injuries among people living in Native American communities often requires a comprehensive, communitybased approach tailored to specific local settings and involves many partners. The Native American communities that bring the necessary partners together and work collaboratively will be the most successful at reducing injuries. Some of the key components of a successful community injury prevention program are these:

- Availability of local injury data that can be used to target high risk groups, set program priorities, and evaluate the effectiveness of intervention programs.
- Local programs to screen people for alcohol abuse and, when necessary, to provide counseling, and treatment.
- A system for referring victims of family and intimate violence to social services, law enforcement, and safe homes.
- Youth services and crisis response programs.
- Strong community support for the police and judges to aggressively enforce occupant restraint, drunk driving, and family violence laws.
- Tribal leaders who are engaged and supportive.
- Support from and participation by community members and local health professionals.

Although Figures 1 through 3 present national data, IHS Areas or service units and tribes can develop their own similar profiles of leading causes of death as a means for supporting local prevention programs. Such approaches have worked well in other settings.²³ In order to meet *Healthy People* 2000²⁴ objectives for reducing injuries and violence, these types of concerted efforts are needed, as well as directing resources toward effective or promising intervention programs that can further reduce injuries as a leading health problem among American Indians and Alaska Natives.

References

- Indian Health Service. *Trends in Indian Health*, 1996. Rockville, MD: U.S. Department of Health and Human Services; 1997.
- May PA. Overview of alcohol abuse epidemiology for American Indian populations. In: *Changing Numbers, Changing Needs: American Indian Demography and Public Health.* Washington, DC: National Academy Press; 1996.
- D'Angelo A. American Indians and Alaska Natives: defining where they reside. *IHS Primary Care Provider*. 1996;21(3):36-42.
- Becker TM, Samet JM, Wiggins CL, Key CR. Violent death in the West: suicide and homicide in New Mexico, 1958-1987. Suicide Life Threat Behav. 1990;20(4):324-334.
- Gallaher MM, Fleming DW, Berger LR, Sewell CM. Pedestrian and hypothermia deaths among Native Americans in New Mexico: between bar and home. *JAMA*. 1992;267:1345-1348.
- May PA. Alcohol abuse and alcoholism among American Indians: an overview. In: Watts TD, Wright R, eds. *Alcoholism in Minority Populations*. Springfield IL: Charles C. Thomas; 1989.
- May PA. The epidemiology of alcohol abuse among American Indians: the mythical and real properties. *Am Indian Culture Research J*. 1994;18(2):121-143.
- Centers for Disease Control and Prevention. Motor vehicle crashes and injuries in an Indian Community — Arizona. *MMWR*. 1989;38:589-591.
- Centers for Disease Control and Prevention. Safety-belt use and motor-vehicle-related injuries — Navajo Nation, 1988-1991. MMWR. 1992;41:705-708.
- Inventory of Tribal Traffic Laws. Rockville, MD: Indian Health Service; 1995.
- 11. Grossman DC, Sugarman JR, Fox C, Moran J. Motor-vehicle crashinjury risk factors among American Indians. *Accid Anal and Prev*. 1997;29(3):313-319.
- 12. *Fatality Facts, 1994: Teenagers.* Arlington VA: Insurance Institute for Highway Safety; 1994.
- Preusser DF, Zador PL, Williams AF. The effect of city curfew ordinances on teenage motor vehicle fatalities. *Accid Anal and Prev.* 1993;25(5):641-645.
- Centers for Disease Control and Prevention. Involvement by young drivers in fatal motor-vehicle crashes — United States, 1988-1995. MMWR. 1996;45:1049-1053.
- National Center for Injury Prevention and Control. *Youth Suicide Prevention Programs: A Resource Guide*. Atlanta, GA: Centers for Disease Control and Prevention; 1992.
- National Center for Injury Prevention and Control. *The Prevention of Youth Violence: a Framework for Community Action*. Atlanta, GA: Centers for Disease Control and Prevention; 1993.
- Powell KE, Hawkins DF, eds. Youth violence prevention: descriptions and baseline data from 13 evaluation projects. *Am J Prev Med.* 1996;12(5) Suppl.
- DeBruyn L, Wilkins B, Stetter-Burns M, Nelson S. Violence and violence prevention. *IHS Primary Care Provider*. 1997;22:(4):58-60.
- Kuklinski DM, Buchanan CB. Assault injuries on the Hualapai Indian Reservation: a descriptive study. *IHS Primary Care Provider*. 1997;22:(4):60-64.
- Wallace LJD, Calhoun AD, Powell KE, et al. *Homicide and suicide among Native Americans, 1979-1992.* Atlanta, GA: Centers for Disease Control and Prevention, National Center for Injury Prevention and Control, 1996. Violence Surveillance Summary Series, No. 2.
- Perkins R, O'Connor MB. Firearm-related deaths in the Alaska Native population. *IHS Primary Care Provider*. 1996;21(3):33-35.
- 22. Kellerman AL, Rivara FP, Somes G, et al. Suicide in the home in relation to gun ownership. *N Engl J Med.* 1992;327:467-472.
- Sleet DA, Rosenberg ML. Injury control. In: Scutchfield D, Keck W, eds. *Principles of Public Health Practice*. Albany, NY: Delmar Publishers; 1997.
- Public Health Service. *Healthy People 2000: National Health Promotion and Disease Prevention Objectives*. Washington, DC: U.S. Department of Health and Human Services; 1991. Report No: (PHS) 91-50212. □

"We would also like to educate our pre-medical students on the cultural sensitivities and Native American tradition."

Pre-Medical Students Interested in Volunteering With IHS

Duquesne University, a Catholic institution of higher learning, located in the heart of Pittsburgh, PA, stands tall in its mission: to develop the minds and hearts of each generation of youth entering through its corridors. Students from the Schools of Pharmacy, Nursing, Health Sciences, and Pre-Medical programs travel abroad to Tanzania, Kenya, and Nicaragua to volunteer their services in local health care clinics. Some pharmacy and pre-medical students have traveled to Belize to learn about medicinal plants and healing from the Mayan medicine men and women.

As the University prepares for its mission in the 21st century, one of its important goals is the education of future physicians and other health care professionals, particularly in the United States, regarding all segments of its population. For too long, the health and education of the American Indian and Alaska Native population has received less than the attention they deserve. The Indian Health Service (IHS) has served as a step in the right direction. We would like to do our part in the education of future physicians, starting from their undergraduate years as pre-medical students. We would also like to educate our pre-medical students on the cultural sensitivities and Native American tradition. Working with the IHS, they will be able to learn about the traditional and historical knowledge about health, illness, and death from the Native American perspective. As part of their volunteer work, they will have an opportunity to understand the unique nature of Indian health programs, where the IHS, tribal leadership, and community members participate in dealing with the distinctive health care needs of each segment of the population.

To begin with, we would like to have four to five of our pre-medical students work in IHS clinics (one per clinic) during the summer of 1998. Our students would be able to learn under the tutelage of the IHS clinic professionals. This learning and volunteer experience would help them to serve in the future, either with the IHS or similar health service organizations. If anyone from the Indian health program clinics is interested in providing volunteer, preceptor, or pre-medical clerkships, please contact Dr. Tata Subhas, Director, Pre-Health and Post-Baccalaureate Pre-Medical Programs, Duquesne University, B101 Bayer Learning Center, Pittsburgh, PA 15282 (phone: 412-396-6335; e-mail: subhas@duq2.cc.duq.edu).

Tata Subhas, D.V.M., Ph.D. Duquesne University

Secretary's Quality of Work Life Initiative

The Initiative

On December 23, 1996, the Secretary of the Department of Health and Human Services, Donna E. Shalala, distributed a Quality of Work Life Strategy developed in response to concerns about employee commitment and morale during this period of unprecedented change and uncertainty.

The Quality of Work Life Strategy has three objectives: improve employee satisfaction, strengthen workplace learning, and better manage ongoing change and transition. It is believed that when these objectives are met, individual and organizational performance will improve and even better service to our customers and partners will be provided.

To meet these objectives and achieve these outcomes, the Secretary has reconfirmed her commitment to streamline and modernize without involuntary separations, and plans to:

- establish a model work and family center, at headquarters, to coordinate work and family programs and serve as a catalyst for improving work and family programs across the Department.
- establish a Quality of Work Life site on the DHHS Home Page to provide a convenient location for information about this strategy and related efforts to improve the quality of employee work life.
- commission an annual survey of employees, to begin in January 1997, to assess employees' views of organizational effectiveness and the quality of work life.
- ask the DHHS Union-Management Partnership Council to serve as the steering group for implementation of the Quality of Work Life Strategy, and report at least annually on the status of the quality of work life of DHHS employees.
- ask that sign-in sheets (and similar systems for collecting time and attendance information that are incompatible with a work culture based on trust) be eliminated within sixty days of the date of this memorandum (12/23/96).

Agencies of the DHHS were asked to develop their own quality of work life strategy, and submit it to the DHHS Union-Management Partnership Council. As a minimum, the strategy to be designed is to accomplish the following:

improve communication with employees;

- strengthen family-friendly work programs;
- provide all employees Internet access by the end of Calendar Year 1997;
- increase the Agency's investment in workplace learning, to include development of knowledge and skills that employees need to achieve strategic goals and objectives, perform differently, become a learning organization, and improve the effectiveness of supervisors and team leaders;
- evaluate the effectiveness of the Agency's diversity management practices, and make appropriate improvements;
- better plan and manage change and transitions, including briefing all members of your leadership team on change and transition, providing more extensive training when change and transition is imminent, and evaluating and learning from change efforts.

The Secretary's Quality of Work Life Strategy offers a unique opportunity to send a strong signal to DHHS employees that we value and support them.

Quality of Work Life Week

The Department of Health and Human Services celebrated Quality of Work Life Week May 12-16, 1997. During that week the DHHS acknowledged and honored our colleagues and our accomplishments. At the various meetings and events during Quality of Work Life Week, The Secretary reported on what has already been accomplished:

- we renewed our commitment to avoid involuntary separations;
- sequential sign-in sheets are becoming a thing of the past;
- the Quality of Work Life site has been established on the DHHS Internet home page;
- the model Work/Life Center is now open;
- the DHHS Union-Management Partnership Council, which is chaired by the Deputy Secretary, is helping me lead this initiative;
- we completed the first annual employee survey;
- Operating Divisions have developed and are implementing their own plans to improve the quality of work life for their employees, and they can already point to solid achievements.

Results of the Employee Survey

The survey, which establishes a baseline that will be used to measure progress, was distributed in January to a random sample of 12,000 employees; almost 5,000 employees completed it (a 41% response rate). Responses came from employees at all grade levels and throughout the Commissioned Corps.

Seventy-five percent of the survey participants said they were able to balance work and family life by taking advantage of flexible scheduling, flexiplace, leave provisions, and other options. Sixty-nine percent reported that managers and coworkers were supportive of them when they did so. Seventyfive percent felt that they were generally treated fairly. A majority felt that their work was seen as highly effective, that they were encouraged to try new approaches to getting the job done, and that they had the authority they needed to do their jobs.

A majority also said that there was a need for change, with 48% reporting signs of discontent within their work group. Almost two-thirds said there were at least some continuing problems that interfered with efficiency, and just under half said that management practices were generally or very ineffective. Delays in communicating what is happening in the organization were noted by 49% of the respondents. There is room for improvement.

The Secretary, Deputy Secretary, and the Union-Management Partnership Council heard what employees said and are working to identify the appropriate steps to make those improvements. The DHHS Union-Management Partnership Council has been briefed on the survey results. Each of the OPDIVs has received its results, and you [the reader] either already have or should soon hear about them as well. OPDIVs will address issues that were identified through the survey, and will report back to the Partnership Council on follow-up actions. The DHHS will survey a random sample of employees again early next year to help us gauge our progress.

Work/Life Center Opens

A model Work/Life Center opened in the Mary Switzer Building on May 12, 1997. A joint effort of the Program Support Center and the Office of the Assistant Secretary for Management and Budget, the Work/Life Center brings together the employee assistance and career management programs, and creates a new one: the Work and Family Resource Program. The Work and Family Resource Program (temporarily located in Room 1416) has collected and will maintain information on family friendly work policies and programs and updated, accurate referral lists. It will sponsor fairs; support groups; and workshops and seminars on dependent care, parenting, wellness, balancing work and family, and aging/retirement planning. Headquarters staff will be served directly, and telecommunication technology will be used to bring services to regional and field staff. The Work/Life Center will move into its permanent location (Room 1250 in the Switzer Building) in the Fall of this year.

Telephone numbers for the programs that comprise the Work/Life Center are: the Employee Assistance Program: 202-205-5790; the Career Management Program: 202-205-9401; and the Work and Family Program: 202-690-5622.

OPDIV Accomplishments

Operating Divisions (OPDIVs) of the Department of Health and Human Services are taking actions to improve the quality of work life for their employees. A few examples:

- The Centers for Disease Control and Prevention (CDC) published and distributed to all employees a "Career Enhancement Guide." This Guide explains the options CDC offers to help employees improve their careers and current job skills. The Guide contains information on training opportunities; internships; childcare and eldercare; flexiplace; and alternative work schedules.
- The Indian Health Service and the Food and Drug Administration are making organization development consulting services available to managers in order to better manage change and transition.
- The Administration on Aging (AOA) has engaged all employees and key stakeholders in an effort to redesign the way work gets done in AOA.
- The National Institutes of Health is developing an Internet training management and delivery system that will allow employees to receive computer-based training at their workstation (office, home, or other suitably equipped location) at any time that is convenient.
- The Health Resources and Services Administration (HRSA) has established a Career Resource Center to provide employees assistance in planning and managing their career within the context of the HRSA mission and goals.

The Deputy Secretary, on behalf of the DHHS Union-Management Partnership Council, will keep employees informed of some of the best practices to improve the quality of work life for DHHS employees through periodic messages via e-mail.

Suggestions/Comments/Questions

An electronic mail box has been established for the quality of work life initiative so employees have a place to send suggestions, comments, and questions. The Internet address is qwl@os.dhhs.gov. Take advantage of it!

MEETINGS OF INTEREST

American Indian Nursing Summit III October 9-12, 1997 Polson, MT

The American Indian Nursing Summit is a forum for dialogue among American Indian and Alaska Native nurses at grassroots, regional, and national leadership levels. The theme of this year's meeting is "Healing Connections: Empowering Communities Through Nursing Practice."

The conference is sponsored by the Salish Kootenai College Nursing Department. Contributors to conference support include the W.K. Kellogg Foundation, the Health Resources Service Administration (HRSA; an Agency of the federal government) the National AIDS Education and Training Centers Programs, and the Native American Native Alaska - Indian Nurses Association (NANA-INA), among others.

For more information, contact the Salish Kootenai College Nursing Department, Box 117, Pablo, MT 59855 (phone: 406-675-4800, ext. 320 or 343; fax: 406-675-4325).

Safe America: Fourth Annual Injury Control Conference November 19-21, 1997 Washington, DC

The goals of this conference, sponsored by the National Center for Injury Prevention and Control, the Centers for Disease Control and Prevention, are to highlight effective programs linked to the science of injury prevention and control, develop new and stronger partnerships among national, state, and local organizations working to prevent or control injuries, identify resources that will support injury prevention and control research and programs, and communicate injury prevention and control issues and proven methods to the injury control community, policymakers, and the public.

On November 19th, the *Safe America National Injury Control Conference* will merge with the Department of Transportation's *Moving Kids Safely '97 Conference* for joint plenary and breakout sessions involving the participants of both conferences. The purpose of this event is to unite the entire injury control community, thereby stimulating and strengthening partnerships to promote child safety. For information about the *Moving Kids Safely '97* conference, call 1-888-428-KIDS.

For additional information about the *SafeAmerica National Injury Control Conference* and a registration packet, contact Elaine Murray, Prospect Associates, 1801 Rockville Pike, Suite 500, Rockville, MD 20852-1683 (phone: 301-468-6555 ext. 2352; e-mail: emurray@prospectassoc.com).

The Annual NCCD, NCSUD, and NCON Meetings January 5-8, 1998 Phoenix, AZ

The Indian Health Service National Councils of Clinical Directors (NCCD), Service Unit Directors (NCSUD), and Nurses (NCON) will hold their combined meetings at the Hyatt Regency in downtown Phoenix on January 5-8, 1998. Please mark your calendars. An agenda and registration materials will be available later in this calendar year.

International Conference on Emerging Infectious Diseases March 8-11,1998 Atlanta, GA

The International Conference on Emerging Infectious Diseases will be convened to (1) encourage the exchange of scientific and public health information on global emerging infectious disease issues, (2) highlight programs and activities that address emerging infectious disease threats, (3) identify program gaps, (4) increase emerging infectious disease awareness in the public health and scientific communities, and (5) enhance partnerships in addressing emerging infectious diseases.

The meeting will consist of plenary sessions and symposia with invited speakers as well as oral and poster presentations based on submission of an accepted abstract. Major topics will include work on surveillance, epidemiology, research, communications and training, and prevention and control of emerging infectious diseases as well as topics related to emergency preparedness and response.

Abstracts should address new, reemerging, or drug resistant infectious diseases which affect human health including foodborne diseases, tropical diseases, sexually transmitted diseases, infectious respiratory diseases, infectious diseases transmitted by animals and arthropods, infections acquired in health care settings, infectious diseases of infants and children, infectious diseases in immunodeficient persons, infectious diseases in minority and other at-risk populations, blood safety, and xenotransplantation. *The deadline for submission of abstracts is October 31, 1997.*

Registration is limited to 2500 participants.

Additional information on abstract submission, registration, and exhibit opportunities can be obtained by sending an e-mail message to meetinginfo@asmusa.org or by calling 202-942-9248. Proceedings of the conference will be published in the *Emerging Infectious Diseases* journal.

NCME VIDEOTAPES AVAILABLE

Health care professionals employed by Indian health programs may borrow videotapes produced by the Network for Continuing Medical Education (NCME) by contacting the IHS Clinical Support Center, 1616 East Indian School Road, Suite 375, Phoenix, Arizona 85016.

These tapes offer Category 1 or Category 2 credit towards the AMA Physician's Recognition Award. These CME credits can be earned by viewing the tape(s) and submitting the appropriate documentation directly to the NCME.

To increase awareness of this service, new tapes are listed in The IHS Provider on a regular basis.

NCME #713

Cholesterol and Coronary Disease: Clinical Implications of Event Reduction Studies (60 minutes) The benefits of cholesterol-lowering therapy extend to many more patients than have traditionally received treatment, according to the findings of several recent large-scale, international clinical trials. In this program, a noted faculty reviews the expanding indications for cholesterol therapy, including primary prevention in hypercholesterolemic patients with no evidence of coronary disease, as well as secondary prevention in patients with known coronary disease but average cholesterol levels. The selection of specific pharmacotherapy is also discussed in the context of the differential effects of various cholesterollowering agents, particularly with the HMG-coenzyme A reductase inhibitors.

NCME #714

Maximizing Your Practice Within a Managed Care Setting: Part 1 (50 minutes) How do I negotiate a managed care contract? Can a physician have too many contracts? What are the current trends in capitation? How do I control my risk? Should I align with a physician management company? Managed care raises a lot of questions from both new physicians just starting their practices and from veteran physicians who want to position their practices securely. In this program, physicians ask a panel of managed care experts the questions every physician wants to ask on this important topic.

NCME #715


Maximizing Your Practice Within a Managed Care Setting: Part 2 (50 minutes) This is a continuation of NCME #714 (see above).

NCME #716

Lyme Disease and Concomitant Tick-Borne Infections (60 minutes) Lyme Disease is one of the fastest spreading infections in the United States. The deer tick, the tiny arthropod that transmits infection, may carry other (sometimes fatal) diseases. Tick-borne illnesses passed on by the deer tick (Ixodes tick) include Lyme Disease (Lyme Borreliosis), human monocytic ehrlichiosis (HME), human granulocytic ehrlichiosis (HGE), Babesiosis, and Tularemia. There is a great deal that we know about tick vectors, organisms that cause disease, and clinical manifestations of tick-borne infections. Yet, confusion still exists about the most effective methods of diagnosis and treatment. A diagnosis of Lyme Disease should be based on clinical and epidemiologic evidence; serologic findings may be useful in supporting the diagnosis. Prevention, treatment, and cure of Lyme Disease and concomitant tick-borne infections are within our reach. In this program, Dr. Benjamin Luft, an expert on tick-borne diseases, will provide clues to clinical diagnosis, guidelines for effective treatment, and a blueprint for prevention.

NCME #717

Management of Hepatitis C: An Update (60 minutes) An estimated 3.9 million Americans are infected with hepatitis C, most of whom are unaware of their disease. Recent advances have been made in the testing, diagnosis, and prognosis of these patients. This program brings the physician up to date on current progress, including recommendations from the recent NIH Consensus Development Conference.



First Indian Doctors Graduate from the SIHB Family Practice Residency Program

The Seattle Indian Health Board (SIHB), an urban Indian health program located in Seattle, Washington, is pleased to announce the graduation of its first two family practice residents at a ceremony held on June 20, 1997. Dr. Michael H. Trujillo, Director of the Indian Health Service, joined approximately 75 guests to witness the graduation and traditional blanket ceremony to honor these new Indian physicians.

Arne Vainio, MD, and Shannon Weigand, MD, were the first two physicians to enter the three year program conducted in partnership with Providence Hospital Family Practice Residency in Seattle. Both of native heritage, they are returning, in essence, to their homes to practice. Dr. Vainio will be joining the staff at Fond-du-Lac Health Center in Wisconsin, and Dr. Weigund will become a staff physician at the Alaska Native Medical Center in Anchorage.

The Seattle Indian Health Board is the only ACGMEapproved residency targeting Indian health. The program, started in 1994, can accommodate six residents, two for each of the three years of the program. The curriculum includes didactic and practical experience in Indian health including opportunities to work with traditional Indian medicine practitioners from different tribes. The goal of the residency is to prepare family physicians for careers in Indian health.

In addition to Drs. Vainio and Weigand, the SIHB recognized the accomplishments of two Indian physicians who worked at SIHB during their pediatric residency with the Seattle Children's Hospital. Recognized were Dr. Angela Erdich and Dr. Kent Saylor. Dr. Erdich will be going to the Oneida Native Health Center, and Dr. Saylor will be completing a fourth year of training in Montreal, Canada.

For information about the residency program, contact the Seattle Indian Health Board, P.O. Box 3364, Seattle, WA 98114 (phone: 206-324-9360; fax: 206-324-8910; e-mail: www.sihb.org). abstracts and the publishers of full-text articles.

Free **MEDLINE**

On June 26, 1997, the National Library of Medicine (NLM) began to provide all Americans with free access to MEDLINE, the world's most extensive collection of published medical information, over the World Wide Web. The web address for NLM is: www.nlm.nih.gov. Prior to June 26th, only registered users who paid for the service could search MEDLINE, an index of over 8.8 million articles from over 3800 biomedical journals. The opening of MEDLINE at no cost to all who can access it means that individuals can search for medical information on topics of interest to them and their families. As consumers become more informed about diseases and the courses of treatment available, they can use that knowledge in making health care choices.

"PubMed," a free on-line service, will provide direct Web links between MEDLINE abstracts and the publishers of fulltext articles.

Training on MEDLINE is currently being discussed. If you are interested in MEDLINE training, send an e-mail to SHELBY L. ZWICK@OHP.1TUC@IHS so we can assess the amount of interest in this training.

Note: This does not change the prearranged Lonesome Docs program for the IHS *Information Portal Project* participants.

·										
Change of Address or Request for										
New Subscription Form										
Name Job Title										
Address										
City/State/Zip										
Worksite: 🛛 IHS 🗆 Tribal 🗆 Un	rban Indian 🛛 Other									
Service Unit (if applicable) Social Security Number										
Check One: 🗆 New Subscription 🗆 Change of Address										
If change of address, please include old address, below, or attach address label.										
Old Address										
THE IHS PRIMARY CARE PROVIDER										
The Provider is published monthly by the Indian Health Service Clinical Support Center (CSC). Telephone: (602) 640-2140; fax: (602) 640-2138; e-mail: provider@smtp.ihs.gov; timely meeting notices and "Information for Authors" available via the CSC Fax Retrieval Service: 602-640-2140. Previous issues of <i>The Provider</i> (beginning with the December 1994 issue) can be found on the CSC Internet home page (www.csc.ihs.gov).										
Wesley J. Picciotti, MPA Director, CSC preferred. Manuscripts may be received via the IHS Banyon electromail system or the Internet (provider@smtp.ihs.gov).										
Wilma L. Morgan, MSN, FNP Authors should submit one hard copy with each electronic cop E. Y. Hooper, MD, MPH References should be included. All manuscripts are subject to editor John F. Saari, MD Responsibility for obtaining permission from the permission from t										
Thomas J. Ambrose, RPhappropriate tribal authorities and Area Publications Committees to publish manuscripts rests with the author. For those who would like more information, a packet entitled "Information for Authors" is available by contacting the CSC at the address below or through our fax										
Opinions expressed in articles are those of the authors and do not necessarily reflect those of the Indian Health Service or the Editors. retrieval service. Call 602-640-2140, ask for the fax retrieval service, and request document #3005. After business hours, press 8, and follow the instructions.										

Dept. of Health and Human Services Indian Health Service Clinical Support Center 1616 East Indian School Road, Suite 375 Phoenix, Arizona 85016

ADDRESS CORRECTION REQUESTED

OFFICIAL BUSINESS PENALTY FOR PRIVATE USE \$300 BULK RATE Postage and Fees Paid U.S. Dept. of Health & Human Services

Permit No. G-290