

## CORRIGENDUM

BRIAN J. SODEN

*Rosenstiel School for Marine and Atmospheric Science, University of Miami, Miami, Florida*

ISAAC M. HELD

*NOAA/Geophysical Fluid Dynamics Laboratory, Princeton University, Princeton, New Jersey*

There is an error in Fig. 1 of Soden and Held (2006), which displays the values of climate feedbacks from various models. The column of data labeled All in this figure was intended to contain the sum of feedbacks from water vapor, lapse rate, surface albedo, and clouds for each model based on results from both our study and that of Colman (2003). However, the results actually depicted the effective sensitivity of the models, rather than the intended sum of feedbacks. In the revised figure shown below, the All column accurately portrays the sum of water vapor, lapse rate, surface albedo, and cloud feedback calculated from the values listed in Table 1 of our paper and those used in Colman (2003).




FIG. 1

### REFERENCES

- Colman, R., 2003: A comparison of climate feedbacks in GCMs. *Climate Dyn.*, **20**, 865–873.  
 Soden, B. J., and I. M. Held, 2006: An assessment of climate feedbacks in coupled ocean-atmosphere models. *J. Climate*, **19**, 3354–3360.

*Corresponding author address:* Dr. Brian J. Soden, Rosenstiel School for Marine and Atmospheric Science, University of Miami, 4600 Rickenbacker Causeway, Miami, FL 33149.  
 E-mail: bsoden@rsmas.miami.edu