# II. ALTERNATIVES INCLUDING THE PROPOSED ACTION

#### II. ALTERNATIVES INCLUDING THE PROPOSED ACTION

This environmental impact statement (EIS) analyzes five alternatives for the leasing of Federal offshore lands by the U.S. Department of the Interior (USDOI), Minerals Management Service (MMS) for the period from mid-2007 to mid-2012:

- Alternative 1—Proposed Action
- Alternative 2—Exclude North Aleutian Basin
- Alternative 3—Exclude Cook Inlet
- Alternative 4—Exclude Mid-Atlantic
- Alternative 5—Defer Blocks Within 25 Miles of Virginia and Chukchi Sea Coasts
- Alternative 6—Defer Blocks at the Mouth of the Chesapeake Bay
- Alternative 7—Limit Leasing in the North Aleutian Basin Planning Area to Blocks Offered in Lease Sale 92
- Alternative 8—No Action

This chapter describes each alternative and summarizes the potential environmental impacts of the alternatives in comparative form. The summary describes the primary impacts based on the detailed analysis of all potential impacts presented in Chapter IV. The EIS's impact analyses were generated from exploration, development, transportation, and oil-spill scenarios developed specifically for analytical purposes.

# A. Alternative 1—Proposed Action

# 1. Description

The four Outer Continental Shelf (OCS) Regions are divided into 26 OCS planning areas (Fig. II-1). The USDOI is considering leasing in three OCS areas, Gulf of Mexico, Alaska, and mid-Atlantic. Within the Gulf of Mexico OCS Region, leasing is being considered in the Central and Western Gulf of Mexico Planning Areas. In the Mid-Atlantic Planning Area, leasing is being considered offshore Virginia. In addition, the USDOI is considering leasing in 4 of the 14 Alaska Region planning areas: Beaufort Sea, Chukchi Sea, Cook Inlet, and North Aleutian Basin. All other planning areas are not analyzed in this EIS because the USDOI is not considering those areas for leasing in the proposed 5-year program. The proposed action is the USDOI's preferred alternative.

Alternative 1—the Proposed Action calls for 21 sales:

- Central Gulf of Mexico—5 annual areawide lease sales and 1 sale of a portion in 2007 (Fig. II-2)
- Western Gulf of Mexico—5 annual areawide lease sales (Fig. II-2)
- Beaufort Sea—2 lease sales scheduled in 2009 and 2011 (Fig. II-3)
- Chukchi Sea—3 lease sales scheduled in 2007, 2010, and 2012 (Fig. II-3)
- North Aleutian Basin 2 sales scheduled in 2010 and 2012, which are subject to restrictions (Fig. II-4)
- Cook Inlet—2 lease sales scheduled in 2009 and 2011 (Fig. II-5)
- Mid-Atlantic 1 sale in 2011, which is subject to restrictions (Fig. II-6)

Activities that could occur as a result of the 21 lease sales in the proposal may extend over a period of 25-40 years. The impact-causing factors associated with these activities include the placement of offshore infrastructure such as rigs, platforms, and pipelines, and onshore facilities such as support bases and processing plants. Operational impacts include bottom disturbance from platform and pipeline placement; local water quality changes from discharging drilling fluids; and air and noise emissions from platforms, supply boats, and air traffic. The specific estimates of offshore infrastructure required to support exploration and development of the hydrocarbon resources (scenarios) associated with alternative 1 (the proposed action) are provided in Tables IV-1 through IV-3.

Chapter IV presents the basic assumptions about anticipated production, exploration, development, transportation, and oil-spills used to prepare the EIS. Transportation for most oil and gas from the Western and Central Gulf of Mexico Planning Areas would be accomplished by extending and expanding the existing offshore pipeline systems. Some of the oil in deepwater areas and a small amount of the oil from the nearshore areas of the Western and Central Gulf of Mexico Planning Areas would be transported by barge or shuttle tanker.

In the Alaska OCS Region, the lifting of the export ban on Alaskan crude oil has led to infrequent and limited shipments to East Asia. However, the vast majority of oil transported via the Trans-Alaska Pipeline System (TAPS) is still being sent to the U.S. west coast. Oil from the Beaufort Sea and Chukchi Sea Planning Areas would be transported by new subsea and overland pipelines to the TAPS and would eventually be carried to the marine terminal facilities in Valdez where it would be loaded on tankers and shipped primarily to west coast ports. Oil from the Cook Inlet Planning Area would be transported to shore using new subsea pipelines, with new onshore common-carrier pipeline systems delivering the oil to existing refineries in Nikiski and gas to transmission facilities in the Kenai area. In the North Aleutian Basin oil, condensate, and gas will be transported to shore. The gas will be converted to liquefied natural gas and shipped to other markets.

Oil spills are the principal accidental impact-causing event. Table IV-4 presents the number of oil spills assumed to occur as a result of the production and transportation of oil associated with lease sales proposed in alternative 1. The sizes of the assumed large spills are approximately equal to the mean of the historical spills for each spill type (platform, pipeline, tanker, or barge). The assumed spill sizes are: platforms—1,500 barrels (bbl); pipelines—4,600 bbl; and tankers—5,300 bbl for the Gulf of Mexico, and 7,800 bbl for tankers carrying Alaska OCS oil. Assumptions regarding the location of spills are based on the source of the spill, the transportation and market assumptions, the location of existing infrastructure, and the location of the resources being analyzed. Platform spills were assumed to occur in the area proposed for consideration for lease. Pipeline spills were assumed to occur between the proposed area for lease consideration and the existing infrastructure. Tanker and barge spills were assumed to occur along the tanker and barge routes.

Finally, the EIS analyses assume the implementation of all mitigation measures required by statute, regulation, or lease stipulations. The protection afforded by these measures is present in the analysis of the resource being mitigated.

# 2. Summary of Impacts

## a. Impacts on Air Quality

The most commonly emitted air pollutants associated with OCS oil and gas activities include nitrogen dioxide ( $NO_x$ ), sulfur dioxide ( $SO_x$ ), particulate matter less than 10 microns in diameter ( $PM_{10}$ ),

particulate matter less than 2.5 microns in diameter  $(PM_{2.5})$ , carbon monoxide (CO), and volatile organic compounds (VOC). The most common  $NO_x$  sources associated with OCS activities are diesel engines used in construction, drilling and support activities; gas reciprocating engines; turbines; and support vessels. Ozone  $(O_3)$  may form in the atmosphere when  $NO_x$  emissions combine with VOC under the influence of sunlight and warm temperatures. The U.S. Environmental Protection Agency (USEPA) has established national ambient air quality standards (NAAQS) for these pollutants and  $O_3$ . The USEPA standards will not be exceeded as a result of routine OCS activities. Air quality impacts from oil spills and in-situ burning would be localized and of short duration.

Emissions from the 2007-2012 Leasing Program activities on the Gulf of Mexico OCS would cause some slight, localized increases in concentrations of NO<sub>2</sub>, SO<sub>2</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>, and CO in the Gulf coastal areas. Concentrations would be well within the NAAQS and the Prevention of Significant Deterioration (PSD) increments. Portions of the Texas and Louisiana are classified nonattainment for O<sub>3</sub>. Emissions from the proposed Five-Year Program would contribute only very slightly, if at all, toward exceeding the O<sub>3</sub> standard in those areas.

Emissions from 2007-2012 Leasing Program activities on the Alaska OCS would cause some slight, localized increases in concentrations of NO<sub>2</sub>, SO<sub>2</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>, and CO in areas around the Beaufort Sea, Chukchi Sea, Bering Sea, and Cook Inlet. Concentrations would be well within the NAAQS and the PSD increments.

Emissions from the 2007-2012 Leasing Program activities on the Atlantic OCS would cause some slight, localized increases in concentrations of  $NO_2$ ,  $SO_2$ ,  $PM_{10}$ ,  $PM_{2.5}$ , and CO in the southeastern Virginia coastal area. Concentrations would be well within the NAAQS and the PSD increments. Portions of the southeastern Virginia coastal area are classified nonattainment for  $O_3$ . Emissions from the proposed 2007-2012 Program would contribute very slightly, if at all, toward exceeding the  $O_3$  standard in those areas.

#### b. Impacts on Water Quality

In the Gulf of Mexico, routine activities that could potentially affect coastal and marine water quality include structure placement and removal (e.g., platforms, drilling units, and pipeline landfalls) and operational discharges and wastes. Structure placement and removal would increase suspended sediment load in the water column, resulting in temporary impacts on water quality. Operational discharges (e.g., muds, cuttings, and produced water), sanitary and domestic waste, and deck drainage could affect water quality in the immediate vicinity of the release, but such releases are expected to be rapidly diluted and dispersed by currents. However, some areas such as confined portions of some channels may be unable to assimilate bilge water and sanitary wastes, thus resulting in some regional water quality degradation. Trenching operations to bury pipelines and place drilling units and platforms would temporally increase turbidity in the immediate vicinity of the trenching operation. Overall, marine water quality impacts from routine operations would be localized and short-term, as compliance with National Pollutant Discharge Elimination System (NPDES) permits and U.S Coast Guard (USCG) regulations would minimize or avoid most impacts to receiving waters. Also, water quality would recover when discharges cease because of dilution, settling, mixing, and evaporation.

Impacts from accidental releases (including oil spills) on water quality would depend on the type and amount of product spilled, amount of material spilled, dispersion and weathering of the spilled material, and specific location affected. While small oil spills would have measurable impacts on water quality, water quality would rapidly recover without mitigation because of mixing, dilution, and

weathering. A large oil spill may adversely affect water quality and may require cleanup to recover the affected areas. Recovery times could be decreased by cleanup activities.

In Alaska, construction and installation of exploratory and development wells, platforms, pipelines, docks and causeways, and/or artificial islands could impact water quality by disturbing sediments and increasing turbidity in the area of construction. Coastal water quality could also be impacted from construction of onshore pipelines and other infrastructure and pipeline landfalls. Proper facility siting and construction permit requirements would largely mitigate these impacts. Minor water quality contamination could also occur from fluids entrained in ice roads when they breakup in the spring. Most water quality impacts would be localized and short-term, with water quality conditions steadily improving as settling and mixing occurred. Exploration discharges would persist for a few hours within the mixing zone around each rig; however, the NPDES permit limitation on discharge rates would minimize water quality impacts. Most major production facilities would reinject all muds, cuttings, and production waters, thereby eliminating degradation of water quality by these effluents. Where muds, cuttings, and produced water would be discharged directly at the drill sites, nearby water quality would be adversely affected. However, materials entering the water would be quickly diluted and dispersed, and thus would not result in long-term impacts on local water quality.

While small oil spills would have measurable impacts on water quality, water quality would rapidly recover without mitigation because of mixing, dilution, and weathering. A large spill in isolated coastal waters, in shallow water under thick or rapidly freezing ice, or in the open sea when or where access to the spill site is limited could cause sustained degradation of water quality. The decomposition and weathering process for oil are slowed in cold water, thereby increasing the potential for more widespread and long-term water quality impacts. Overall, the impacts from a spill would depend on the spill size and composition, weather conditions and season of year, spill location, and extent and timing of cleanup activities.

In the mid-Atlantic area, the impacts associated with exploration activities on coastal water quality would be localized and short-term, while the impacts associated with development and production activities would be localized, short- to medium-term, and would most likely not result in long-term degradation to the background water quality conditions in the area. The impacts with a potential to affect coastal waters are associated with the discharges from support vessels and helicopters operating nearshore, the contribution of onshore point and nonpoint discharges into mid-Atlantic waters from support facilities, the installation of a natural gas pipeline, and accidental oil spills, including the risk of one large spill from tanker transport of oil.

The overall impacts associated with exploration, development, and production activities on marine water quality would be localized, short- to medium-term, and would most likely not result in long-term degradation to local water quality conditions. These impacts would primarily be generated from drilling activities, including the installation and operation of a drilling rig(s), platform installation and operation, and the discharges from support vessels and helicopters. The impacts would be localized and short-term. Compliance with NPDES permit requirements would minimize or prevent most impacts to receiving waters caused by discharges from routine exploration and development activities. Impacts of accidental releases to water quality would depend on the size of the spill, type of material or product spilled, and environmental factors at the time of the spill. However, there would be no long-term, widespread impairment of marine water quality. Marine water quality could be affected by platform and pipeline installations and operations, discharges, and accidental oil spills. Operational discharges would be in compliance with NPDES permit requirements that would minimize or eliminate most impacts. A large offshore spill could result in short term and localized water quality impacts.

#### c. Impacts on Marine Mammals

In the Gulf of Mexico, routine operations that could affect marine mammals include seismic surveys, construction and operation of offshore facilities, operational discharges and waste generation, vessel and helicopter traffic, and platform removal. Noise generated during exploration and production activities, platform removal, and OCS-related vessels and helicopters may temporarily disturb some individuals. Collisions with OCS-related vessels could also injure or kill some individuals, although the incidence of such collisions is expected to be very low. Many of the effects associated with noise and the presence of OCS-related vessels or structures would likely be short-term and not result in population-level effects. Existing permit requirements, regulatory stipulations, and MMS guidelines targeting many of the routine operations would greatly limit the impact of any potential effects on marine mammals.

An oil spill may expose marine mammals (or their habitats and food resources) to oil or its weathering products. The magnitude of effects from accidental spills would depend on the location, timing, and volume of the spills; the environmental settings of the spills (e.g., restricted coastal waterway, deepwater pelagic location), and the species exposed to the spills. Spill cleanup operations could result in short-term disturbance of marine mammals in the vicinity of the cleanup activity, while a collision with a cleanup vessel could injure or kill the affected individual.

Under the proposed action, some routine operations could affect marine mammals in Alaska. Impacts to marine mammals would be a function of the extent that industrial activities coincided with the species' seasonal occupation of certain regions and the species' tolerance level of the impacts. Noise generated by exploration and operation activities and by OCS-related vessels and helicopters may temporarily disturb some individuals, causing them to leave or avoid the area. Such effects would likely be short-term and not result in population-level effects. However, if the disturbance results in the temporary abandonment of young by adults, survival of young may be reduced. Collisions with OCS-related vessels may injure or kill some individuals, although the incidence of such collisions is expected to be low. Vessels, construction of ice roads, on-ice vehicles, and aircraft have been known to affect polar bear behavior (e.g., they may abandon dens, which could reduce cub survival). However, mitigation should reduce the level of disturbance. Sea otters appear to habituate to regular human activity, and routine operations would have a negligible impact to their populations. Existing permit requirements, regulatory stipulations, and MMS guidelines targeting many of the routine operations would generally limit the likelihood of marine mammals being affected by these operations.

Compliance with the Endangered Species Act and the Marine Mammal Protection Act would limit the likelihood of impacts to both listed and nonlisted marine mammals. Similarly, careful siting of onshore and offshore facilities and OCS-support vessel and aircraft routes to avoid impacts to habitats, particularly during migration, would generally reduce the likelihood and/or magnitude of adverse impacts to marine mammals in Alaska. For example, activity restrictions related to Steller sea lion critical habitat would greatly reduce or avoid the likelihood of routine operations impacting designated rookeries and haulouts, and limit how closely OCS-support vessels may approach critical habitats for this species. Mitigation measures are in place that may subject areas to seasonal drilling restrictions to prevent the endangered bowhead whale from being disrupted during its migration and when it is most likely to be hunted. The endangered fin and humpback whales would also be protected by the same mitigation measures developed to minimize disturbance to bowhead whales.

Accidental oil spills may result in the direct and indirect exposures of marine mammals and their habitats to the oil and subsequent weathering products. Animals could be exposed by the inhalation or

ingestion of oil or contaminated foods, which may result in a variety of lethal and sublethal effects. Fouling of fur of some species (e.g., sea otter and fur seal) could affect thermoregulation and reduce survival. The magnitude of effects from accidental spills would depend on the location, timing, and volume of the spills; the habitats affected by the spills (e.g., coastal habitats); and the species exposed. Spills in open waters may be expected to affect the fewest number of individuals, and not result in population-level impacts to any species. The greatest risk to marine mammals would be associated with large spills reaching rookeries and haulout locations. Spills in these locations have the potential for affecting the greatest number of individuals and could result in population-level impacts to some species (especially the Steller sea lion). Spill cleanup operations could result in short-term disturbance of marine mammals in the vicinity of the cleanup activity, while a collision with a cleanup vessel could injure or kill the affected individual. Also, disturbance of adults with young during cleanup could reduce survival of the young animals.

Under the proposed action, some lease activities could affect marine mammals in the Mid-Atlantic Planning Area offshore Virginia. Underwater noise is expected to be the most prevalent potential impact associated with exploration, development, and production. However, no lethal or physically debilitating impacts are expected from this source. Vessel and aircraft traffic are expected to result in occasional startle reactions and avoidance responses. A limited number of probable lethal collisions between vessels and endangered whales could occur. However, no collisions would be anticipated between vessels and the smaller cetaceans occurring on the Atlantic OCS. Muds and cuttings discharges could release toxic chemicals into the water and possibly contaminate marine mammals directly or through ingestion of contaminated prey species. Ingestion or entanglement with discarded waste could lead to intestinal blocking, reduced mobility, and other lethal or sublethal effects. Installation and operation of offshore facilities could disturb normal behaviors and could lead to avoidance or displacement from the area. The potential for impacts related to oil spills could lead to skin, respiratory, and digestive problems but are expected to be sublethal and non-debilitating. Overall, it is not expected that exploration, development, and production activities would significantly affect populations of marine mammals in the mid-Atlantic. Other than measurable impacts to the extremely endangered right whale population if any individual is killed, such as in the event of a lethal vessel collision, no changes in population size, distribution, or behavior are expected from the proposed action.

#### d. Impacts on Marine and Coastal Birds

In the Gulf of Mexico, routine activities that could affect marine and coastal bird species include infrastructure placement, operational discharges and wastes, and vessel and aircraft traffic. The primary effect would be disturbance of birds in the immediate vicinity of the activity. Because birds tend to habituate to human activities and noise, potential impacts from disturbance would be short-term and not be expected to result in population-level effects. However, construction activities near coastal habitats could disrupt breeding and nesting activities of colonial nesting birds. Some collision mortality with offshore platforms and, to a lesser extent, OCS-related helicopters would be expected. Collisions at offshore platforms may affect several thousand birds each year as they migrate across the Gulf in spring and fall. Because the discharge of production wastes and other materials is regulated, and because permitted production wastes discharged into marine waters would be quickly diluted and dispersed, relatively few birds would be exposed to these discharges and wastes. Loss or alteration of preferred habitat due to pipeline landfalls could result in the displacement and possible decrease in nesting activities. Compliance with regulations and corporate procedures would minimize impacts to birds, including federally-listed species.

Accidental oil spills pose the greatest threat to marine and coastal birds in the Gulf of Mexico. The magnitude and ecological importance of any effects would depend upon the size of the spill, the species and life stages that are exposed, and the size of the local bird population. Exposure to spills in deep water would be largely limited to pelagic birds, while shallow-water spills could affect the greatest variety and number of birds, including shorebirds, waterfowl, wading birds, gulls, and terns. A large near shore oil spill that contacts the endangered whooping crane winter habitat at the Aransas National Wildlife Refuge in Texas could result in population level impacts. Oil washing ashore may contaminate eggs and nest sites as well as foul foraging areas and food resources. Spill cleanup activities could also disturb nesting populations or habitats in nearby areas.

In Alaska, marine and coastal birds may be affected by seismic surveys; construction and operation of onshore and offshore facilities; vessel, aircraft, and on-land vehicle traffic; and noise and human activities during normal operations and maintenance activities. In most cases, disturbed birds would temporarily leave the area, while in other cases, the displacement could be long-term. Construction of onshore facilities and pipelines, offshore pipeline landfalls, and offshore gravel islands (to support drilling platforms) would result in the permanent disturbance of potential habitat within the immediate footprint of the new facilities and gravel excavation areas. Depending on the species present at and in the vicinity of the construction areas, the numbers of birds affected, and the activity (nesting, molting, feeding, staging) that the affected birds were undergoing at the time of disturbance, the displacement could reduce reproductive success, foraging success, and survival, and could result in population-level impacts. New onshore facilities may result in local increases of predator species such as the arctic fox and glaucous gull. Increases in these predators would increase predation pressure of local bird populations and, depending on the birds affected, could result in population-level effects. Compliance with regulations and corporate procedures would minimize impacts to birds, including federally-listed species.

In the event of an accidental oil spill, exposed marine and coastal birds may experience a variety of lethal or sublethal effects, and the magnitude and ecological importance of any such effects would depend upon the size and location of the spill, the species and life stage of the exposed birds, and the size of the local bird population. Although the potential for a large spill is unlikely, it could result in the loss of large numbers of birds, and result in potentially long-term reductions in populations. Spills in offshore locations have the greatest potential for affecting the greatest number of birds, especially if a spill occurs in or reaches an area where birds have congregated and are carrying out important activities (e.g., nesting, molting, and staging). A spill in onshore habitats would affect relatively few birds unless the spill was to reach a surface water body such as a stream, pond, or lake that provides important nesting, brood-rearing, foraging, or staging habitat. Population recovery from high adult mortality during a large oil spill could take many years. Spill cleanup activities may also disturb birds in the vicinity of the cleanup, causing them to leave the vicinity of the cleanup activity.

Marine and coastal bird populations on the mid-Atlantic are not expected to be measurably affected by the routine activities assumed for the proposal. Federally-listed, coastal-oriented endangered species and species of concern are not likely to be adversely affected by exploratory drilling in the proposed lease area. Disturbance from the low level of vessel and helicopter traffic expected in the mid-Atlantic is not likely to cause impacts to marine and coastal birds. Because of the relatively low estimated number of oil spills, there is a low risk of impact resulting in some losses of marine birds, particularly for pelagic birds and sea birds. The long-term effect could be a small reduction in population sizes for a few species. Small spills are expected to result in very few deaths among susceptible species, but it is not expected that these losses would be measurable against natural fluctuations within affected populations. In the unlikely event that a large spill occurred, marine or coastal bird species would not

experience measurable impacts at the population level. Local impacts however could, under certain circumstances, be high depending on the location and time of year in which the spill occurred.

#### e. Impacts on Terrestrial Mammals

In the Gulf of Mexico, routine operations that could affect terrestrial mammals include construction and maintenance of onshore infrastructure and pipelines, and support vessel and aircraft traffic. The terrestrial mammals of primary concern are the federally-listed Alabama, Choctawatchee, St. Andrew, and Perdido Key beach mice, and the Florida salt marsh vole. Because the habitats of these species are located within protected areas, they are buffered from contact with existing OCS industry infrastructure. New pipeline landfalls and onshore OCS-related facilities would not adversely impact these species because the proposed action will not result in activities where the species' habitat occur, and the species occur within protected areas that further preclude the species or their habitats from incurring adverse impacts.

Due to their location on inner dunes, the habitats of the beach mice species are unlikely to be affected by an accidental oil spill. The Florida salt marsh vole is located far from areas where oil leasing and development may occur under the proposed action. Thus, it is unlikely that this species or its habitat would be affected by an accidental oil spill from OCS activities.

In Alaska, the terrestrial mammals that may occur or utilize coastal and/or inland habitats potentially affected by OCS activities (e.g., construction and maintenance of onshore infrastructure and pipelines, and support vehicle and aircraft traffic) include caribou, muskoxen, grizzly and black bears, and foxes. Disturbance of terrestrial mammals during construction would be localized and temporary, affect a relatively small number of individuals, and not be expected to result in long-term impacts to mammal populations. Winter construction could disturb individual bears if their dens are located near construction areas. Overwintering caribou and moose could also be disturbed by construction activities. Disturbance of wildlife could affect overall condition and affect subsequent survival or reproductive success, especially if animals are displaced from calving or wintering habitats to less optimal habitats. Siting onshore facilities to avoid important calving or wintering areas would reduce the likelihood of long-term impacts to the survival or reproduction of affected species.

Facility construction in the Alaskan planning areas would also result in the long-term loss of a relatively small amount of habitat and in the death of a few individuals (primarily small mammals such as mice and voles) unable to flee the construction areas. The amount of habitat loss would be very small compared to that available throughout the area. Operation of onshore facilities could result in the long-term avoidance of adjacent habitats by species sensitive to noise and human activity. The presence of a new pipeline may affect daily or seasonal movements of caribou. The condition of individuals displaced from preferred to less optimal habitats could be adversely affected, thus affecting their subsequent survival or reproductive success. However, some wildlife species, such as bears and foxes, become habituated to human activities and facilities. Grizzly and black bears may incur defense of life or property-related mortality if they become habituated or attracted to new onshore facilities. Increased fox densities caused by human activity and facility presence could result in increased predation on local natural prey species and may also result in an increase in the incidence and transmission of diseases such as rabies.

In the event of an accidental spill in the Alaskan planning areas, terrestrial mammals may be exposed via ingestion of contaminated food, inhalation or airborne oil droplets, and direct ingestion of oil during grooming, which may result in a variety of lethal and sublethal effects. An oil spill may also affect the habitats of terrestrial mammals, resulting in reduced food supply (e.g., oiling of vegetation)

or reduced physical habitat quality (e.g., oiled sands or fouling of nests and burrows). Because most spills would be relatively small (< 50 bbl), relatively few individuals would likely be exposed. A population-level adverse impact would only be expected if the spill was very large or contaminated a crucial habitat area where a large number of individual animals were concentrated (e.g., a riparian area used by moose). However, population-level impacts would not be expected for most species. Cleanup activities could temporarily disturb terrestrial mammals in the vicinity of the cleanup operation, and would not be expected to result in population-level impacts. An exception could be if muskoxen were displaced from preferred overwintering areas in the Arctic. This could adversely affect their overwinter survival and condition, and subsequent calving success, resulting in local population-level impacts.

#### f. Impacts on Fish Resources and Essential Fish Habitats

In the Gulf of Mexico, routine operations that may affect fish resources, including essential fish habitat (EFH), include exploratory surveying and drilling, installation and removal activities, and operational discharges. Seismic survey airgun discharges could result in temporary displacement of fishes. The impact to overall fish population would be negligible since fishes are distributed over wide geographic areas, and seismic operations would be localized. Sediment disturbance during placement of platforms and pipelines will increase turbidity which, in turn, will temporarily impact the water quality of EFH within a small area, causing fish to temporarily disperse. Once put in place, platforms serve as artificial reefs or fish attraction devices, benefiting those species preferring bottom relief. During platform removal, explosives may injure biota and destroy invertebrate and fish communities in the immediate vicinity of the platforms and those that are prey for managed fish species. Considering the small proportion of EFH area that could be affected, potential impacts on EFH due to routine operations under the proposed action would also be limited. Impacts on the Gulf sturgeon are not likely to occur because there is relatively little overlap between the locations that could be affected by the proposed action and the distribution of the threatened Gulf sturgeon.

Effects of oil spills would depend on the location, timing, and volume of the spill, in addition to other environmental factors. Small spills are unlikely to affect a large number of fish or EFH before dilution and weathering would reduce concentrations of toxic fractions to sublethal or nonlethal levels and would, therefore, not have substantial effects on fish populations. It is anticipated that any single large spill in offshore waters would affect only a small proportion of a given fish population (including the Gulf sturgeon) and that fish resources would not be permanently affected. A large spill that reaches submerged seagrass beds or coastal wetlands could have more persistent impacts on EFH and could require remediation. Few impacts would be expected because of the wide dispersal of early life history stages of most fishes in the surface waters of the Gulf of Mexico.

In Alaska, routine operations that may affect fish resources, including EFH, include exploratory surveying and drilling, installation activities, and operational discharges. Seismic survey airgun discharges can affect pelagic fish species with swim bladders. Temporary displacement of fishes is the most probable effect of noise generated by seismic surveys. The impact to overall fish population would be negligible since fishes are distributed over wide geographic areas, and seismic operations would be localized. In Alaska, sediment disturbance, resuspension, and displacement from routine activities would affect EFH in a similar manner as that in the Gulf of Mexico. Assuming compliance with existing Federal, State, and local regulations, policies and consultations, most impacts to fish resources would be minimized, and most impacts to EFH from routine activities under the proposal will be avoided. Although some individual fish are likely to be harmed by activities associated with routine operations, the mitigation measures imposed on leasing activities would limit the scope of these impacts.

Effects of accidental oil spills would depend on the location, timing, and volume of spills; distribution and ecology of affected fish species; and other environmental factors. Small spills would be localized and are unlikely to affect substantial number of fish before dilution and weathering would reduce concentrations of toxic fractions to sublethal or nonlethal levels. Under most circumstances, a large spill would affect only a small proportion of a given fish population, and therefore overall population levels would not be affected. Oiled intertidal areas could lead to considerable mortality of eggs and juvenile stages of some pelagic species in the affected areas. One or more years could be required for population levels to recover. Spilled oil reaching wetland habitat, including salt marshes, could kill vegetation and associated invertebrate species and small fish that are prey species for Pacific salmon. Large spills that reach coastal streams used by Pacific salmon species could have more persistent impacts and require remediation.

Because fishery resources are widespread through the Mid Atlantic Planning Area and the level of exploratory activity is projected to be minimal, impacts in the proposed sale area would only affect a small portion of the population. No impacts to fish population distribution and abundance would occur. Impacts on fish resources may result from the discharge of operational effluents, muds, and cuttings; platform and pipeline emplacement; structure removal; lights on offshore rigs; noise associated with routine drilling operations or geophysical surveys; and discharge of formation or produced waters. Based on the assumptions for the proposal, individual finfish or shellfish are expected to experience sublethal impacts such as reduced biogenic activity, reduced metabolic functions, or disease. Deaths of a few individuals are also expected. However, no measurable decline in whole populations is expected. The anticipated duration of impacts is less than one generation or 1-3 years for most activities and events. Recovery of fish in localized inshore habitats contaminated by an oil spill could require 2-12 years. No measurable impact to species of concern is expected as a result of the proposed activity.

# g. Impacts on Sea Turtles

Five species of sea turtles occur in the Gulf of Mexico: green, hawksbill, Kemp's ridley, leatherback, and loggerhead. All of these species may be found as hatchlings, juveniles, and adults in the Gulf of Mexico program area. All but the hawksbill have been reported to nest on beaches within the Gulf of Mexico planning areas.

Sea turtles could be directly affected by seismic surveys, vessel traffic, construction of offshore and onshore facilities, and removal of platforms. Sea turtles may also be exposed to a variety of waste materials, such as produced water, which have the potential to cause a variety of lethal and sublethal effects. Noise generated during exploration and production activities and platform removal may result in the temporary disturbance of some individuals, while some turtles may be killed during the use of underwater explosives for platform removal. The construction and operation of new onshore facilities may impact nest sites, possibly result in eggs being crushed, and disturb hatchling movement from the nest sites to the water. Sea turtles may also be injured or killed by collisions with OCS vessels. Existing permit requirements, regulatory stipulations, and MMS guidelines and required mitigation measures targeting many of the routine operations could limit the seriousness of any potential effects on sea turtles. Therefore, while routine operations could affect individual sea turtles, population-level impacts are not expected.

Oil spills may result in the exposure of one or more sea turtle life stages to oil or its weathering products. Oil may reduce egg hatching and hatchling survival, and may inhibit hatchling access to water. Hatchlings, juveniles, and adults may inhale or ingest oil and oil vapors, and may incur any of a

variety of physiological impacts. The presence of oil slicks or oiled beaches may alter habitat use and affect nest site access and use. Small spills are unlikely to affect a large number of sea turtles or their habitats and thus are not expected to have substantial or long-term effects. A large spill could affect many more individuals and habitats, including nesting beaches, and potentially incur population-level effects. The magnitude of effects from accidental spills would depend on the location, timing, and volume of the spills; the environmental settings of the spills; and the species and life stages of sea turtle exposed to the spills. The rapid deployment of spill-response teams and implementation of cleanup activities could limit the magnitude of impacts; however, cleanup operations themselves could also impact sea turtles or their habitats due to artificial lighting at night, machine and human activity and related noise, sand removal and cleaning, and the use of dispersants and coagulant chemicals.

No sea turtles occur in the Alaska OCS planning areas.

It is not expected that exploration, development, and production activities would affect populations of marine turtles in the mid-Atlantic. The generally inshore distribution of these animals, as well as their seasonal geographical distribution on the Atlantic OCS, substantially reduces the potential for impacts stemming from routine gas and oil activities offshore. Other than collisions with vessels and accidental oil spills, potential impacts are expected to be sublethal. Impacts from routine activities and small accidental events are not expected to measurably affect the distribution, behavior, or population size of marine turtles on the Atlantic OCS. An oil spill of 1,500 bbl in size would result in more measurable impacts and possibly would affect sea turtle populations in the area.

#### h. Impacts on Coastal Habitats

In the Gulf of Mexico, routine operations could impact coastal habitats as a result of pipeline construction, maintenance dredging of inlets and channels, and vessel traffic. The magnitude of these impacts would depend on the location of new construction, the level of dredging or shipping activity in a specific area, and existing environmental conditions (such as ongoing shoreline degradation). Stabilization of dune margins could be difficult, and establishment of vegetation cover might be slow, possibly resulting in prolonged losses of dune habitat on barrier beaches near pipelines. Direct impacts to barrier islands are frequently avoided during pipeline construction by the use of nonintrusive construction techniques and by pipeline installation using directional boring. Construction could impact wetlands due to habitat fragmentation and isolation and altered hydrology. Additional wetland losses could occur along pipeline routes and navigation channels as a result of the continued erosion of adjacent marsh substrates and subsequent conversion of marsh habitat to open water. Impacts to wetlands from construction could be minimized by maintaining buffers around wetlands and by the use of best management practices for erosion and sedimentation control.

The magnitude of potential oil-spill impacts would depend on a variety of factors, including the location and size of the spill, remediation efforts, existing environmental conditions (such as plant species or substrate type), and natural localized erosion and deposition patterns. If a large oil spill were to reach coastal wetlands, these resources may not fully recover even if remedial actions were undertaken. Cleanup operations might also impact coastal habitats if the removal of contaminated substrates affected beach stability and resulted in accelerated shoreline erosion or if coastal wetlands were damaged (e.g., by trampling or removal of vegetation).

In Alaska, routine operations could have impacts on coastal barrier beaches and dunes and on wetlands, primarily as a result of pipeline construction, shore base construction, and vessel traffic. The magnitude of these impacts would depend on the location of new construction, the level of shipping activity in a specific area, and existing environmental conditions (such as ongoing shoreline

degradation). Small areas of coastal habitat will be lost or degraded from pipeline landfalls and placement of vertical support members for aboveground onshore pipelines, onshore bases, and roads. Also, dredging of intertidal habitats for pipeline burial would disturb benthic communities at the site of the trench. The impacts of buried pipelines and pipeline landfalls and related causeways to benthic communities would be localized.

Potential impacts from oil spills could occur to coastal habitats. The magnitude of these impacts would depend on a variety of factors, including the location and size of the spill, remediation efforts, existing environmental conditions (such as plant species or substrate type), and natural localized erosion and deposition patterns. Cleanup operations might also impact coastal habitats if the removal of contaminated substrates affected beach stability and resulted in accelerated shoreline erosion or if coastal wetlands were damaged (e.g., by trampling or removal of vegetation).

Exploration, development, and production activities could have impacts on the coastal barrier beaches and dunes of the mid-Atlantic area primarily as a result of pipeline construction and vessel traffic. Impacts on wetlands could result from construction activities and indirect impacts as a result of poorer water and air quality and altered hydrology. The magnitude of these impacts would depend on the location of new construction, the level of shipping activity in a specific area, and existing environmental conditions (such as ongoing shoreline degradation). Potential impacts from spills could have direct impacts on wetlands and beach habitats. Impacts to wetlands from small spills are expected to be limited to temporary shifts in flora and fauna ratios in the vicinity of support bases. No measurable impact to shorelines is expected from small spills. The magnitude of impacts from a large spill would depend on a variety of factors, including the location and size of the spill, weather conditions, remediation efforts, beach conditions (e.g., grain size), existing environmental conditions (such as plant species or substrate type), and natural localized erosion and deposition patterns. Cleanup operations themselves might also impact wetlands, estuaries, beaches, and dunes. Adverse impacts on coastal habitats from a large spill can range from insignificant to high degrees of damage, including extensive loss of habitat.

#### i. Impacts on Seafloor Habitats

In the Gulf of Mexico, routine operations that could affect seafloor habitats include placement and removal of structures, vessel traffic, and/or permitted operational discharges. The benthic communities associated with topographic features are sensitive to physical damage (e.g., by placement of structures or anchors) and operational discharges (e.g., drilling muds and cuttings). However, these impacts would be avoided through compliance with the Topographic Features Stipulation and the establishment of No Activity Zones. The installation of mobile drilling units or production platforms and pipeline placement and removal could also impact other seafloor habitats. However, protective lease stipulations exist for seafloor habitats such as live bottom and pinnacle trend areas. Any seafloor habitats that are damaged would eventually recover over a period of years. Most impacts that could occur from drilling muds and cuttings and produced water discharges would be avoided as a result of compliance with the Live Bottom Stipulation that requires the avoidance of live bottom areas and NPDES permit restrictions that limit the types and amounts of discharges allowed near live bottom areas.

Impacts from routine operations to seagrass beds could be avoided by routing the pipelines corridors around seagrass beds. Chemosynthetic (seep) communities would be protected from damage associated with anchoring and placement of structures through siting restriction requirements.

Potential impacts on seafloor habitats and associated benthic communities from accidental oil spills could occur. The magnitude of these impacts would depend upon the location, size, and type of spill; weather conditions; effectiveness of cleanup operations; and other environmental conditions at the time of the spill. For example, given the water depths (e.g., 200 m) of the topographic features, it is unlikely that significant amounts of oil from surface spills would reach the sensitive communities. Because of the Live Bottom Stipulation, the potential for impacts from oil spills is greatly reduced. If a large spill from a pipeline were to occur, there could be lethal effects on the biota in localized areas that receive large quantities of oil. In such cases, the community would recover once the area had been cleared of oil, although full recovery could take many years. In most cases, effects on sensitive biota would be sublethal, with recovery occurring within months to a few years. The benthic community would probably recover more quickly from a shallow-water pipeline spill than from a deepwater pipeline spill, because of the greater potential for wave-induced suspension of sediments in shallow water.

In Alaska, routine operations that could affect seafloor habitats include seismic surveys, drilling of wells, placement of structures, and/or permitted operational discharges. It is anticipated that planning procedures and permitting requirements would avoid or minimize the potential for impacts to the Stefansson Sound Boulder Patch community in the Beaufort Sea Planning Area. Routine activities would not measurably affect local populations of lower trophic-level organisms in other seafloor habitats. At most, platform and pipeline installation would alter species diversity within a small area near the facility. For example, the presence of platforms or artificial islands would favor organisms requiring hard substrates, thereby shifting community composition in some areas. Small areas on intertidal habitat could also be affected by pipeline landfalls. It is anticipated that these development activities could displace some coastal organisms but would have no measurable effect on local populations. The stipulation on protection of biological resources, which requires surveys near, and avoidance of, special benthic habitats, would help to reduce potential impacts to both nearshore and deeper-water biota.

In the event of an oil spill, seafloor benthic communities are unlikely to be heavily oiled since most of the oil would float. Sublethal impacts associated with low concentrations of oil in the water column would be expected in the immediate vicinity of the spill. A large spill associated with a rupture or leakage from buried offshore pipelines could contaminate sediments within the immediate area. Organisms in those sediments could experience high levels of contamination and mortality, but would represent a very small proportion of the overall area available to support similar benthic communities.

In the mid-Atlantic, during exploration there would probably be some unavoidable localized benthic population reductions due to changes in sediment characteristics from the discharge of drilling muds, cuttings, and water, and from the ingestion of oil and other materials in the sediment by benthic organisms. However, it is unlikely that any demersal fish populations would be adversely affected by this low level of activity. During the development and production stages, there would probably continue to be some unavoidable localized benthic population reductions due to changes in sediment characteristics from the discharge of drilling muds and cuttings and from the ingestion of spilled oil in the sediment by benthic organisms. These effects would be most pronounced in areas of high biological productivity and increased ecological sensitivity, such as nearshore areas, hard bottoms (including reef structures and artificial obstructions), and the heads of submarine canyons. However, lease stipulations, the large area over which the activities would occur, and the extensive timeframe during which they would likely happen should result in small, if any, adverse impacts to the environment. Impacts from blowouts, bottom disturbances, and discharges will be mitigated by lease stipulations and regulations to minimize impacts to benthic resources.

#### j. Impacts to Areas of Special Concern

In the Gulf of Mexico, routine activities that could affect areas of special concern such as national marine sanctuaries, parks, reserves, and refuges include placement of structures, pipeline landfalls, operational discharges and wastes, and vessel traffic. However, impacts from these activities are unlikely. Potential impacts on the Flower Garden Banks National Marine Sanctuary due to routine operations under the proposed action would be largely prevented by (1) provisions of the Topographic Features Stipulation that prohibits exploration or development activities in the immediate vicinity of the banks, (2) the subsea location of the features, and (3) ocean currents that circulate around the banks. Among the national parks located in the Gulf of Mexico, only the Padre Island National Seashore and the Gulf Islands National Seashore are located adjacent to regions within which oil and gas activities could occur under the proposed action. No infrastructure (e.g., pipeline landfalls, shore bases) would be sited in national parks, national wildlife refuges (NWR's), or national estuarine research reserves. Some OCS-related trash and debris could wash up on beaches. Over time, vessel wakes can erode shorelines along inlets, channels, and harbors. However, existing mitigation measures limit vessel speeds in inland waterways, so impacts would be reduced. Thus, impacts to areas of special concern would be avoided.

While an oil spill could affect areas of special concern, the magnitude of the potential impact will depend on the location, size, and type of spill; weather conditions; effectiveness of cleanup operations; and other environmental conditions at the time of the spill. However, impacts from oil spills would be unlikely because of the assumption that 75 percent of the hydrocarbons developed as a result of the 2007-2012 Leasing Program would occur in deep water (> 330 m) located far from the shoreline. Also, the Topographic Features Stipulation prohibits exploration or development activities in the immediate vicinity of areas such as the Flower Garden Banks National Marine Sanctuary. Thus, it is unlikely that an oil spill would affect such offshore areas. In the unlikely event of oil contaminating a shore-based area of special concern, impacts could include the destruction of wetland vegetation and associated wildlife, oil saturation and trapping by vegetation and sediments, and mechanical destruction of the wetland area during cleanup.

In Alaska, no OCS-related activities will occur in national park lands in the Arctic Subregion is under the proposed action, thereby minimizing the potential for impacts from routine operations in these areas. Impacts from routine activities in adjacent areas would be minimal under the proposed action. Development may be possible in the Alaska Maritime NWR under the proposed action, but it is anticipated that reviews of individual lease sales would minimize the potential for impacts from routine operations. Routine exploration and development activities in offshore areas distant from NWR areas would have only a limited potential to affect resources on these refuges because they are located relatively far from shoreline areas. However, there could be a potential for effects on wildlife refuges from development and operation of facilities to support offshore oil drilling and production. This could include habitat degradation or loss due to nearshore dredging, construction of access roads, construction and operation of processing and waste facilities, construction and operation of shore bases, and construction of onshore pipelines.

There are no national park or national forest lands that would be affected by exploration, development, or production activities within the North Aleutian Basin Planning Area. Site-specific reviews of individual lease sales would likely minimize the potential for impacts to NWR properties from routine operations due to onshore or nearshore development activities.

Development of onshore facilities within national park lands in the Cook Inlet Planning Area is considered unlikely, thereby minimizing the potential for impacts from routine OCS operations in these areas. However, offshore construction of pipelines and platforms could have temporary effects

on wildlife due to noise and activity levels and on scenic values for park visitors. Development may be allowed in the Gulf of Alaska Unit of the Alaska Maritime NWR. However, it is anticipated that reviews of individual lease sales would minimize the potential for impacts from routine operations due to development activities. No OCS-related development would occur in the Alaska Peninsula Unit of the Alaska Maritime NWR.

Impacts from oil spills that occur adjacent to national park or NWR boundaries would depend on spill location, spill size, type of product spilled, weather conditions, environmental conditions at the time of the spill, and effectiveness of cleanup operations. Large oil spills that contact shoreline areas of NWR's would likely impact coastal habitats and fauna negatively and could also affect subsistence use, commercial or recreational fisheries, and tourism. No affects to national forests would occur because there would be no onshore OCS development within or in the immediate vicinity of national forest lands and because spills would not be expected to come in contact with national forest lands.

Exploratory oil and gas activity in the Atlantic would be prohibited in designated areas of special concern throughout the proposed leasing area. Activities outside these areas may cause impacts within their boundaries that could compromise the integrity of protected marine and coastal environments, thus threatening visitation and use, awareness, and support for current and future protections. The low level of exploratory activity projected and the use of mitigation measures, including lease stipulations, would minimize the risk to these areas. No measurable impacts to these features are expected as a result of exploratory activities. It is unlikely that oil and gas activity would significantly impact areas of special concern within the proposed lease area, although large spills have the potential to threaten protection efforts. Mitigation efforts and thorough contingency planning by multiple parties will minimize the risk to these areas.

#### k. Impacts on Population, Employment, and Regional Income

The main effect of the proposed action on population and employment will be the employment generated by the expected routine OCS oil and gas activity. MMS has developed a new economic impact model (MMS Alaska-GOM Modeling Using IMPLAN). Based on the exploration and development scenarios, potential effects from routine operations would be small in the Gulf of Mexico, Alaska, and the mid-Atlantic. Employment impacts of oil spills reaching landfall can vary considerably given the volume of oil reaching land, land area affected, and sensitivity of local environmental conditions. A large oil spill might have short-term (one season or less) effects.

In the Gulf of Mexico, the proposed action is likely to add 20,250-34,700 jobs in an average year. Most employment effects would concentrate in Louisiana and Texas; however, even in the most affected areas, the added employment demands are not likely to stress the local labor market. Population increases across the region would be small but noticeable in some locations, ranging from 38,500 to 66,000 over the life of the proposed action. Oil spills could affect such activities as beach recreation, diving, commercial diving, recreational fishing, and sightseeing. Studies have shown that there could be a one-time seasonal decline of 5-15 percent in tourist visits associated with a major spill. Employment and regional income impacts from spills would likely be greatest in Texas and Florida. However, the associated loss of business would be localized, with tourist movement to other coastal areas in the region offsetting a reduction in the number of visits to the affected area.

In Alaska, the proposed action is expected to add 12,600 jobs (the largest share on the North Slope) and \$192 million in personal income in an average year. Alaska is projected to grow by 14,000 residents. Workers at North Slope sites stay in enclave housing separate from local communities, and most live in south-central Alaska or the Fairbanks area. Since similar worker-residential patterns are

likely elsewhere, population increases will be concentrated in places such as Anchorage and will have few effects on local communities. A large oil spill could have short-term effects (one season or less) on the tourism and fishing industries. It would also cause some short-term (up to 15 months) local employment, possibly generating moderate local employment effects.

In the mid-Atlantic, during the exploration phase, there would probably be some small increase in local revenue, although given the size of the Hampton Roads economy, it would have limited impact. Some local labor would be needed to support supply boats and helicopters, but at a level so small there would be little impact on overall employment and unemployment. Equipment to support the exploration phase, with the possible exception of crew boats and helicopters, would most likely be brought in from outside the local area. The necessary expertise in development and production of oil and gas does not exist in Hampton Roads, and workers with these skills would have to be imported from other areas where offshore drilling is already being done. However, there is a large labor pool in the Hampton Roads area, including workers skilled in construction and maritime trades. These workers could provide support services in the drilling and pipe-laying phases, as well as in the construction of needed onshore facilities such as the service base, gas processing facility, and pipe coating yard. Any increase in population as a result of development and production is not expected to have a significant impact on the housing market or on the economy. Purchases of food, transportation, and other miscellaneous supplies are not likely to have an impact on the Hampton Roads economy as a whole. Depending on the location, a large spill could affect the recreation, tourism, commercial fishing, and cruise ship economies, and possibly negatively affect the real estate market, resulting in temporary losses of jobs and income.

#### l. Impacts on Sociocultural Systems and Environmental Justice

Impacts on sociocultural systems and environmental justice (EJ) vary across OCS Regions. In the Gulf of Mexico, where sociocultural systems have long experienced offshore operations, and in the Hampton Roads area, where activities would be few and concentrated in industrialized and heterogeneous areas, impacts to sociocultural systems would be few. In Alaska, should a large oil spill occur and impact subsistence or commercial harvesting, effects to sociocultural systems and EJ could be substantial.

In the Gulf of Mexico, the greatest impacts to sociocultural systems are expected to result from the ongoing expansion of deepwater and ultra-deepwater activities. These trends have contributed to the cultural heterogeneity of the area and the offshore workforce; to a wider geographic range from which workers are drawn; and to the creation of jobs that require more specialized skills and longer, unbroken periods of work offshore. Accidents are not expected to alter sociocultural systems.

A large EJ concern is the potential risk to residents from nearby OCS-related infrastructure including helipads, heliports, waste management facilities, pipe coating yards, petrochemical plants, shipyards, platform fabrication yards, supply bases, natural gas storage facilities, repair yards, refineries, port facilities, and terminals. The proposed action is not expected to significantly change the risk exposure of nearby residents. Given current infrastructure distributions, coastal demographic patterns, and the distance from shore of likely deepwater activities, any new infrastructure, air emissions, or oil spills that result from the proposed action are not expected to have disproportionate environmental or health affects to low-income or minority populations.

In Alaska, impacts to sociocultural systems come from two sources. Revenues derived from the petroleum industry provide much of the funding for infrastructure and public services (e.g., education) throughout the State. While these impacts are positive, State revenues from the OCS program are

limited. Impacts can also come from interference with subsistence, an important dietary resource and fundamental expression of Native social organization and culture, and from interference with commercial fishing, an economic mainstay of rural South Alaska and Bering Sea Subregions. Routine operations will not affect fishing and the effects of new onshore infrastructure are expected to have only minor, local effects on terrestrial harvests (by affecting access). Noise disturbance from offshore operations could significantly affect marine mammal harvests and is a particular concern with bowhead whaling, but this is expected to be mitigated through consultation and conflict avoidance measures. The *Exxon Valdez* oil spill has shown that substantial impacts occur from direct effects upon resources and from disruptive cleanup efforts. Potential impacts on sociocultural systems from accidents under the proposal would be determined by the location and timing of the spill.

Alaska's 2000 population is 29 percent minority, including 15.5 percent American Native. About 9.4 percent of its population is below the poverty line. With the exception of the Cook Inlet area in south-central Alaska, all regions addressed by the proposal have significantly larger minority populations than the State as a whole. Only northwest Alaska has a disproportionate percentage of residents below the poverty line. Alaska Native populations are present in many coastal areas and could be near any new onshore infrastructure resulting from the proposal. The importance of marine mammals (such as the bowhead whale) to subsistence raises particular concerns. Should an oil spill occur, its potential environmental and health impacts on Alaska Native populations could be disproportionately high or adverse depending on the geographical location of the spill and its effects on subsistence resources and harvests. Because of these concerns, MMS continues to emphasize consultation and interactions with Native organizations to evaluate potential actions and mitigations.

In the mid-Atlantic, the activities that would occur during exploration would be concentrated in industrial-port areas and would be limited in size and duration. They would be unlikely to have noticeable sociocultural effects. During the exploratory phase of the operation, there would be a limited need for local workers—particularly those with maritime and construction skills—as well as some support businesses such as food service and temporary housing. Availability of workers from the current population in Hampton Roads is assured because of the large, skilled, labor pool that exists in the region, and it is likely that the few jobs available for local workers would be available to all eligible applicants, regardless of race or ethnic origin. A wider range of activities would occur during development than during exploration. While most would occur in industrial-port areas and have limited sociocultural effects, some, such as pipeline landfalls, might occur outside of these areas. Because of the level of population diversity in the Hampton Roads area, opportunities for work would not be constrained by race or ethnic background. Because the Hampton Roads area is ethnically and racially integrated, the limited sociocultural effects of new OCS-related activity and infrastructure would probably not have a disparate impact on minorities or low-income families. Likewise, if a large oil spill occurred in the area, it probably would not have a disparate impact on minorities or lowincome families. The proposal may have a low level of socioeconomic impacts on the Hampton Roads area; however, there would not be disproportionately high and adverse human health or environmental impacts on minority or low-income populations. Thus, the activities likely under the proposal do not raise strong EJ concerns.

#### m. Impacts on Archaeological Resources

Archaeological resources that could be affected by the proposed action include historic shipwrecks and inundated prehistoric sites offshore, and historic and prehistoric sites onshore. Although shipwrecks tend to concentrate in shallow, nearshore waters in all OCS Regions, historic shipwrecks are scattered across the entire continental shelf, and many are found even in deepwater areas. Inundated prehistoric sites may occur on those portions of the continental shelf that were exposed as dry land during the

period of lower sea levels of the last ice age. The extent of the continental shelf that was exposed varies from area to area; however, globally, sea levels were approximately 120 m lower than present approximately 21,000 to 19,000 years ago. Onshore historic properties include sites, structures, and objects such as historic buildings, forts, lighthouses, homesteads, cemeteries, and battlefields. Onshore prehistoric archaeological resources include sites, structures, and objects such as shell middens, earth middens, campsites, kill sites, tool manufacturing areas, ceremonial complexes, and earthworks.

Routine operations associated with the proposed action that may affect archaeological resources in all regions include drilling wells, installing platforms, installing pipelines, anchoring, and constructing onshore infrastructure. Existing Federal, State and local laws and regulations require that archaeological surveys be conducted prior to permitting any activity (onshore or offshore) that might disturb a significant archaeological site. Compliance with existing laws and regulations will protect archaeological resources to the maximum extent possible from most impacts associated with routine activities; however, it is still possible that some impacts could occur.

Should a direct physical contact between a routine activity and a shipwreck site occur, it could destroy fragile ship remains and could disturb the site context, resulting in a loss of data on ship construction, cargo, and the social organization of the vessel's crew, as well as the concomitant loss of information on maritime culture for the time period from which the ship dates. Ferromagnetic debris associated with OCS operations could mask the magnetic signature of historic archaeological resources, making them difficult to detect with magnetometers. Interaction between a routine activity and a prehistoric archaeological site could destroy artifacts or site features and could disturb the stratigraphic context of the site.

Oil spills could affect coastal historic and prehistoric archaeological resources and could result in unavoidable loss of information. The level of this impact would depend on the significance and uniqueness of the information lost. Archaeological resource protection during an oil spill requires specific knowledge of the resource's location, condition, nature, and extent prior to impact; however, the coastal areas of the various OCS Regions have not been systematically surveyed for sites. Existing information indicates that prehistoric sites in all regions occur frequently along the mainland coast and barrier islands, and along the margins of estuaries, bays and lagoons; thus, any spill that contacts these areas could involve a potential impact to a prehistoric site.

#### n. Impacts on Land Use and Existing Infrastructure

In the Gulf of Mexico, routine operations could affect land use and existing infrastructure through construction of pipeline landfalls and onshore support facilities, related petroleum industry support facilities, and new worker in-migration. The impact level would depend upon the amount of new construction and the degree to which the area is already developed. Given the current level of existing infrastructure in the Gulf of Mexico and the region's history with oil and gas operations, impacts to land use and existing infrastructure would be limited. The oil and gas industry has been an integral part of the Gulf of Mexico economy for decades, and the continuation of industry activities is not expected to result in any extensive land-use or infrastructure impacts for the region. Potential impacts resulting from new developments include the conversion of land from other uses, increased demands on roads and utilities, and increased demand for housing and public services (e.g., schools and hospitals) due to the in-migration of workers and their families. The nature and extent of these impacts, especially those associated with the construction of gas-processing facilities, would depend on the specific locations chosen for construction.

Projected impacts to land use and existing infrastructure resulting from an oil spill would likely include stresses of the spill response on existing infrastructure, direct land-use impact (such as impacts of oil contamination to a recreational area or to agricultural land), and restrictions of access to a particular area while the cleanup is being conducted. These impacts are expected to be temporary and localized. The magnitude of the impacts would depend upon the location and size of the spill; the greatest potential would occur from a large spill in shallow water.

In Alaska, routine operations could expand existing land-use infrastructure and transportation systems due to the construction of support bases, terminals, airfields, pipelines, and roads. The proposed action would impact land use in the vicinity of new processing and transport facilities and their associated infrastructure. Routine operations associated with the proposed action could significantly affect land use in the Arctic Subregion by building pipelines (subsea and overland), service roads, and new or expanded marine-support facilities, petroleum processing facilities, and airfields. Impacts associated with platform and pipeline construction would be temporary. Impacts could result from an influx of workers to the region, as housing and expanded community infrastructure could be needed. The impacts of the proposed action on land use and infrastructure would depend on the timing, size, and duration of the activities, and may vary from light but concentrated activities to a large-scale activity in differing drilling locations. The potential for impacts to the road system infrastructure may be high if a new permanent road system is built outside of, and links up with, the existing system (e.g., the Prudhoe Bay/Kuparuk Unit road system in the Arctic Subregion). Depending on the locations chosen for land-based facilities, a change from an existing land use (e.g., undeveloped, commercial, or residential) to an industrial land use would likely occur.

The construction of petroleum industry facilities would expand the area potentially at risk from accidental oil spills, along with the requirement to maintain oil-spill response equipment in those areas. An oil spill could alter land use temporarily but would not likely result in long-term changes. The magnitude of the impacts would depend on the size and location of the spill; the greatest potential would occur from a large spill in shallow water.

In the mid-Atlantic, the existing industrial/maritime infrastructure in the Hampton Roads area can fulfill the requirements of a support base, as well as boat and helicopter traffic. Given the current level of activity in Hampton Roads, there will be minimal impacts on workforce or land use as a result of exploration activity.

#### o. Impacts on Tourism and Recreation

In the Gulf of Mexico, routine operations would have limited effects on recreation and tourism, with potential adverse aesthetic impacts to beach recreation and sightseeing balanced by potential positive impacts to diving and recreational fishing. Additional impacts could be caused by an increase in the amount of trash and debris washing to shore. Onshore facilities would most likely be placed in commercially zoned coastal locations and would not impact recreation or tourism. The proposed program is not expected to result in adverse impacts to recreation and tourism in this region.

Potential impacts to recreation and tourism resulting from an oil spill would likely include direct impacts (e.g., oil contamination of a beach), access restrictions to a particular area (e.g., no diving or fishing while cleanup is being conducted), and aesthetic impacts. These impacts could persist for several months to a year or more pending cleanup completion and any required habitat restoration. The extent of the impacts would depend on the location and size of the spill and the effectiveness of cleanup operations. While oiled beach sediments are usually easily removed via mechanical means, such shoreline activity would effectively close the beach to public use for the duration of cleanup

operations. If beach restoration is required (i.e., to restore the proper beach profile), additional time may be required before public access is allowed. Historical evidence pertinent to the effects of major oil spills has indicated that spills may prompt either a seasonal decline in tourist visits and/or tourist movement to other coastal areas in the region.

In Alaska, routine operations would generally have limited effects on tourism and recreation, with potential adverse aesthetic impacts to water-dependent activities such as fishing (in the Arctic Subregion, fishing is primarily a subsistence activity rather than a recreational activity), boating, sightseeing, and associated land-based activities such as hiking, picnicking, camping, and hunting. However, routine operations in the North Aleutian Basin Planning Area could impact recreation and tourism in concentrated areas, since the North Aleutian Basin has not been previously developed for the oil and gas industry. Impacts would be greatest in the initial construction periods due to the limited infrastructure to accommodate both tourists and construction workers in the region (with regards to boat and air traffic and overnight accommodations) and possible restrictions in touring routes and fishing and boating activities. Impacts on recreational activities would depend on the proximity of the new construction to the recreational-use areas. Areas of new industrialization would be concentrated in a few locations, thus minimizing the impact to the overall region. Any closures of areas to water-oriented recreational activities would be only for short periods of time. Changes in visual quality would be expected to be localized and would be concentrated during periods of high industry activity, such as when drilling and laying of pipelines occur.

Potential impacts to recreation and tourism resulting from an oil spill would likely include direct impacts (e.g., oil contamination of a recreation-use area), access restrictions to a particular area (e.g., no diving or fishing while cleanup is being conducted), and aesthetic impacts (e.g., view of spill and cleanup activities). The magnitude of these impacts would depend on factors such as the size and location of the spill and effectiveness of cleanup operations. Impacts would likely be greatest if the spill occurred during the peak recreational season and in shallow waters.

In the mid-Atlantic area, oil and gas activities during exploration may have visual and audible impacts on coastal visitors, particularly if drilling rig and production platform are within 11 miles of the shoreline. Impacts from routine activities are expected to have no noticeable effects on recreation and tourism activity. Routine activities associated with oil and gas exploration, development and production may result in visual, natural, and branding impacts on tourism and recreation. An oil spill could result in temporary beach closures.

#### p. Impacts on Fisheries

Biological resources that serve as the basis for commercial and recreational fisheries in the Gulf of Mexico are expected to be affected by activities associated with routine operations under the proposed action. These activities include vessel traffic and structure placement, presence, and removal. Turbidity and noise associated with installation and decommission activities (mobile offshore drilling units, pipelines), deposition of cuttings, and drilling activities could temporarily drive fishes away from the area and preclude fishing. Additionally, potential conflicts between exploration activities and fishing gear, bottom trawlers, longliners, and purse netters could also preclude commercial fishing. However, these impacts would be temporary. Once platforms are installed and production activities begin, offshore structures would act as fish attraction devices for both pelagic and reef-associated species; these structures would also be attractive to handline fishers.

If oil spills were to occur, commercial and recreational fisheries could be affected in several ways. The possibility of oil-soaked fishing gear and potentially contaminated fish may reduce fishing efforts.

Individuals of target fish species could be affected directly by exposure to spilled oil, potentially causing fish death or illness. Spills could also indirectly affect fisheries by degrading habitats that are critical for the survival of target species, but would only be serious if they lead to severe declines in target species populations. Highly migratory fish species (tunas, sharks, and billfish) could move away from surface oil spills in deep water, disrupting fishing efforts. The level of effects from accidental oil spills on commercial and recreational fisheries would depend on the location, timing, and volume of spills, in addition to other environmental factors. Small spills are unlikely to have a large effect before dilution and weathering reduces concentrations and, therefore, would not have long-term effects on commercial and recreational fisheries in the Gulf of Mexico. It is anticipated that a large spill would affect only a small proportion of a given fish population within the Gulf of Mexico and that fish resources would not be permanently affected.

In Alaska, routine operations (e.g., vessel traffic, seismic surveys, and structure placement and presence) from the proposed action could have localized and temporary effects on some fishery resources. Commercial and recreational fisheries could be affected by changes in the distribution or abundance of the fishery resource, a reduction in the fish or shellfish catches, preclusion of fishers from access to viable fishing areas, or by causing loss or damage to fishing equipment or vessels. However, overall effects on populations of commercial and recreational fishery resources are not anticipated. Seismic surveys and construction of platforms and pipelines could result in space-use conflicts with commercial fishing activities, although these effects would be localized and, in the case of seismic surveys, temporary. Impacts would be expected to be limited in scope as a result of adherence to mitigation measures and compliance with Federal, State and local requirements.

The level of effects from accidental oil spills on commercial and recreational fisheries would depend on the location, timing, and volume of spills, in addition to other environmental factors. Small spills are unlikely to have a large effect before dilution and weathering reduces concentrations. It is anticipated that any single large spill would affect only a small proportion of a given fish population within Alaska, and that fish resources would not be permanently affected. Substantial temporary effects on populations could occur if important habitat areas were contaminated. Localized effects on commercial and recreational fishing could result as a consequence of reduced catch, contamination of fish tissues, degraded aesthetic values that attract fishers, temporary closure of fishing areas, loss of gear, or loss of fishing opportunities during cleanup and recovery periods.

In the mid-Atlantic, the minimal level of activity projected for the proposed action will result in no discernible impacts for commercial and recreational fisheries from oil and gas exploration. Impacts on commercial fisheries are not expected to produce losses in fish and shellfish stocks that are measurable against natural variation. Economic losses are expected to be limited to a few fishermen and last less than 1 year. No losses distinguishable from natural variation in recreational fish and shellfish stocks are expected. An oil spill in nearshore waters could reduce or prohibit recreational fishing in the affected area for months or longer.

A table comparing the various alternatives to the proposed action (alternative 1) can be found at the end of this chapter.

#### B. Alternative 2—Exclude North Aleutian Basin

# 1. Description

Alternative 2—Exclude North Aleutian Basin would hold 18 sales in 6 OCS planning areas:

- Central Gulf of Mexico—6 annual sales
- Western Gulf of Mexico—5 annual sales
- Beaufort Sea—2 sales
- Chukchi Sea—3 sales
- Cook Inlet—1 sale
- Mid-Atlantic 1 sale

# 2. Comparison of Impacts

As a result of excluding the North Aleutian Basin Planning Area from the 2007-2012 Leasing Program, there would be two fewer sales in the Alaska Region. The OCS oil and gas activities associated with these sales would not occur. The small amount of liquid hydrocarbons assumed to be developed in the North Aleutian Basin under alternative 1 would be replaced with increased domestic production elsewhere or by increased imports. The small amount of liquid hydrocarbons involved would not affect the frequency of import tanker spills versus OCS spills.

Alternative 2 would result in somewhat reduced impacts locally for the resources analyzed in the EIS, but overall, the impacts are expected to be the same as those for alternative 1. Impacts to biological resources would be localized and short-term resulting in no population-level impacts. A possible exception applies to the Steller's eider and the Steller's sea lion. While unlikely under the assumed scenario for the North Aleutian Basin in alternative 1, should a large spill occur and contact large numbers of eiders, which migrate through the area, or important haulout and rookery areas for the Steller's sea lion, populations of the eider in the Arctic Subregion, where they breed, or of the sea lions in Cook Inlet, where they occur during the non-breeding season, could be affected.

In the Gulf of Mexico and mid-Atlantic, the impacts would be the same as for alternative 1. In the Beaufort Sea, Chukchi Sea, and Cook Inlet, the impacts would be the same with the possible exception of the Steller's eider and Steller's sea lion.

#### C. Alternative 3—Exclude Cook Inlet

# 1. Description

Alternative 3 would hold 19 sales in six planning areas:

- Central Gulf of Mexico—6 annual sales
- Western Gulf of Mexico—5 annual sales
- Beaufort Sea—2 sales
- Chukchi Sea—3 sales
- North Aleutian Basin 2 sales
- Mid-Atlantic − 1 sale

Under alternative 3, no sales would be held in Cook Inlet Planning Area. There would be no change from alternative 1 in the remaining areas: annual sales would be held in the Central and Western Gulf of Mexico Planning Areas, 2 sales would be held in the Beaufort Sea, 3 sales would be held in Chukchi, 2 sales in North Aleutian Basin and 1 sale in the mid-Atlantic. The same means of

transporting hydrocarbons to shore from production facilities would be used in all planning areas for alternatives 1 and 3.

## 2. Comparison of Impacts

Under alternative 3, no oil or gas would be produced from the Cook Inlet Planning Area. The estimated oil and gas resources and associated activities would be the same for this alternative as for alternative 1 in the Central and Western Gulf of Mexico, the Beaufort Sea, Chukchi Sea, North Aleutian Basin, and mid-Atlantic. The small amount of liquid hydrocarbons assumed to be developed in Cook Inlet under alternative 1 would be replaced with increased domestic production elsewhere or by increased imports. The small amount of liquid hydrocarbons involved would not affect the frequency of import tanker spills versus OCS spills.

Alternative 3 would result in somewhat reduced impacts locally for the resources analyzed in the EIS, but overall, the impact is expected to be the same as for alternative 1. Impacts to birds, including the Steller's eider, the endangered short-tailed albatross, and the Kittlitz's murrelets could result from oil spills. Population level impacts to these endangered, threatened or candidate species, however, are not expected to occur.

In the Gulf of Mexico, Beaufort Sea, Chukchi Sea, North Aleutian Basin, and mid-Atlantic, the impacts for alternative 3 would be the same as those for alternative 1.

#### D. Alternative 4—Exclude Mid-Atlantic

# 1. Description

Alternative 4 would hold 20 sales in six planning areas:

- Central Gulf of Mexico—6 annual sales
- Western Gulf of Mexico—5 annual sales
- Beaufort Sea—2 sales
- Chukchi Sea—3 sales
- Cook Inlet—2 sales
- North Aleutian Basin 2 sales

Under alternative 4, a lease sale would not be held in the Mid-Atlantic Planning Area. There would be no change from alternative 1 in the remaining areas: annual sales would be held in the Central and Western Gulf of Mexico, 2 sales would be held in the Beaufort Sea, 3 sales would be held in Chukchi, 2 sales in North Aleutian Basin, and 2 sales in Cook Inlet. The same means of transporting hydrocarbons to shore from production facilities would be used in all planning areas for alternatives 1 and 3.

# 2. Comparison of Impacts

Under alternative 4, there would be no difference from alternative 1 for the Gulf of Mexico and Alaska planning areas. There would be a decrease in the expected production from the exclusion of the Atlantic planning area.

Only a small amount of activity and production was estimated to occur in the mid-Atlantic under alternative 1 resulting in some potential small, incremental, short-term and localized impacts in a diversified economic region already supporting a large maritime industry. No population-level impacts to biologic resources were expected under alternative 1. A service vessel or barge/tanker collision with the endangered right whale could affect the population of these whales, which is estimated to number only between 300 and 350 animals. Such an accident is unlikely though given the small amount of activity expected in the area under alternative 1 resulting in an estimated one to five service vessel trips per week. Alternative 4 would preclude this possibility.

In the Gulf of Mexico, Beaufort Sea, Chukchi Sea, North Aleutian Basin, and Cook Inlet, the impacts would be the same as for alternative 1.

# E. Alternative 5—Defer Blocks Within 25 Miles of the Virginia and Chukchi Sea Coasts

# 1. Description

This alternative considers the impacts associated with not leasing within 25 miles of the Virginia and Chukchi Sea coasts.

Alternative 5 would affect the impacts from sales in:

- Chukchi Sea (Alaska) 3 lease sales scheduled in 2007, 2010, and 2012
- Mid-Atlantic (Virginia) 1 sale in 2011, which is subject to restrictions

Sale-related activities identified for the proposed action (alternative 1) would have a reduced impact to shore and nearshore communities under this 25-mile buffer alternative. Offshore infrastructure such as rigs and platforms would be located further from shore. This would make the structures essentially beyond the normal visual field of most individuals. Length of pipelines and length of travel by vessels may increase, but coastal water quality effects from discharge of drilling fluids and air and noise emissions from platforms would be partially reduced.

# 2. Comparison of Impacts

This alternative would reduce potential environmental impacts in resources within the 25-mile buffer zone in the Chukchi Sea (Fig. II-7). Overall impacts on water quality, air quality, marine mammals, marine and coastal birds, benthic communities, and fish resources would be reduced. However, impacts from vessel traffic, aircraft, offshore and onshore pipeline construction, and onshore support facilities would still exist. There would be no gravel island or ice roads constructed in the Chukchi Sea under this alternative, thus eliminating potential impacts arising from these facilities. Possible impacts to polar bears would be reduced. The 25-mile buffer would provide additional protection to the bowhead whales during their spring migration.

The potential for adverse impacts from oil spills would be reduced compared with that for the proposed action. This alternative would also reduce potential effects of a large oil spill on portions of Chukchi Sea Unit of the Alaska Maritime National Wildlife Refuge. The establishment of a 25-mile buffer would substantially reduce potential impacts on Native subsistence. Possible adverse health or environmental impacts from changes in subsistence resources and harvest patterns would be reduced.

A reduction in the likelihood of an oil spill would make adverse effects on Native subsistence resources and harvests less likely.

The impacts to terrestrial animals, coastal habitats, land use and existing infrastructure, population, employment, regional income, tourism, and recreation would be essentially the same as those for the proposed action since the need for onshore support facilities and pipelines would not change.

This alternative would reduce potential environmental impacts in resources within the 25-mile buffer zone off the Virginia coast (Fig. II-8). Overall impacts on water quality, air quality, marine mammals, marine and coastal birds, benthic communities, and fish resources would be reduced. However, impacts from vessel traffic, aircraft, offshore and onshore pipeline construction, and onshore support facilities would still exist. There still would be a risk of collisions between vessels and marine mammals, including the right whale. A reduction in the chance of a nearshore oil spill would also reduce possible longer term adverse effects on tourism. However, a risk of a large spill from tanker transport of oil would still exist. The 25-mile buffer would eliminate the potential visual impacts from beaches and other recreation sites.

The impacts to terrestrial animals, coastal habitats, land use and existing infrastructure, population, employment, and regional income, would be essentially the same as those for alternative 1 since the need for onshore support facilities and pipelines would not change.

#### F. Alternative 6—Defer Blocks at the Mouth of the Chesapeake Bay

# 1. Description

Alternative 6 would hold 21 sales in seven planning areas:

- Central Gulf of Mexico—5 annual sales and 1 sale in a limited area
- Western Gulf of Mexico—5 annual sales
- Beaufort Sea—2 sales
- Chukchi Sea—3 sales
- Cook Inlet—2 sales
- North Aleutian Basin—2 sales
- Mid-Atlantic—1 sale

Under alternative 6, the area offered in the Mid-Atlantic Planning Area would be reduced by eliminating the lease blocks within a wedge-shaped area offshore the entrance to the Chesapeake Bay (Fig. II-9). There would be no change from alternative 1 in the remaining OCS sale areas. It is assumed that the deletion of these lease blocks would not significantly change the resource estimates or the development scenario for the mid-Atlantic sale area.

# 2. Comparison of Impacts

This alternative would affect impact levels only within the mid-Atlantic sale area. In all other OCS areas, there would be no change from the impacts described for alternative 1. Because it is assumed that the deletion of the wedge-shaped area offshore the entrance to the Chesapeake Bay would not significantly change the development scenario for the mid-Atlantic sale area described under alternative 1, the potential impacts described for alternative 1 would also remain unchanged under

alternative 6, with one minor exception. The low potential for direct physical impacts to archaeological resources from exploration and development activities under alternative 1 would be reduced even further under this alternative.

# G. Alternative 7—Limit Leasing in the North Aleutian Basin Planning Area to Blocks Offered in Lease Sale 92

## 1. Description

Alternative 7 would hold 21 sales in 7 planning areas, as described for alternative 1:

- Central Gulf of Mexico—5 annual sales and 1 sale in a limited area
- Western Gulf of Mexico—5 annual sales
- Beaufort Sea—2 sales
- Chukchi Sea—3 sales
- Cook Inlet—2 sales
- North Aleutian Basin 2 sales
- Mid-Atlantic 1 sale

Under alternative 7, the area offered in the North Aleutian Basin lease-sale would be reduced to 990 blocks (approximately 2.27 million hectares) forming a wedge-shaped area that ranges from 18 to about 185 km offshore of the Alaska Peninsula, the same area proposed for the earlier North Aleutian Basin Sale 92 (Fig. II-10). This alternative includes the entire geographic area in which program-related offshore oil and gas activities were projected to occur under the alternative 1 development scenario. Therefore, for the purpose of this analysis, it is assumed that this limitation alternative would not significantly change the resource estimates or the development scenario for the North Aleutian Basin sale area, nor change the general geographic location of sale-related activities. The development scenario under alternative 1 and alternative 7 assume there will be no more than six offshore platforms.

# 2. Comparison of Impacts

This alternative reduces the area offered in the North Aleutian Basin lease sale to 990 blocks. However, it includes the entire geographic area in which program-related offshore oil and gas activities were projected to occur under the alternative 1 development scenario. Therefore, it is not expected to substantially affect the impact levels that would occur under alternative 1. However, by limiting oil-related activities to the southern portion of the planning area, it would reduce slightly the already slight risk of adverse impacts to the Steller sea lion and Steller's eider.

#### H. Alternative 8—No Action

The evaluation of a "no-action" alternative is required by the regulations implementing the National Environmental Policy Act (40 CFR 1502.14(d)). If the Secretary were to adopt this alternative, it would halt OCS presale planning, sales, and new leasing from 2007 to 2012 even in the Central and Western Gulf of Mexico Planning Areas. However, exploration, development, and production stemming from past sales would continue.

This alternative would shut down the OCS leasing program from mid-2007 through mid-2012. The amounts of OCS natural gas (up to 30 trillion cubic feet) and oil (up to 7.5 billion barrels of oil) required to meet national energy needs would be forgone. That amount of energy would have to be replaced by a combination of imports, alternative energy sources, and conservation.

Market forces are expected to be the most important determinant of the substitute mix for OCS oil and gas (see Section IV.F.2). Key market substitutes for forgone OCS oil production would be imported oil (88%), conservation (5%), switching to gas (4%), and onshore production (3%). For OCS natural gas, the principal substitutes would be switching to oil (40%), onshore production (28%), imports (16%), and conservation (16%).

In addition to market-based substitutes, the Nation or individual States might choose to encourage or even impose programs designed to deal with the energy shortfall. To replace oil, these programs might favor alternative vehicle fuels such as ethanol or methanol, vehicles with greater fuel efficiency, or alternate transportation methods such as mass transit.

As a partial replacement for the forgone natural gas, governments might mandate increased reliance on coal, nuclear, hydroelectric, or wind-generated electric power. In addition, governments might give more emphasis to programs encouraging more efficient electricity transmission and more efficient use of gas and electricity in factories, offices, and homes.

A table comparing the alternatives to the proposed action is shown on the following pages.

	ALTERNATIVE 1 ALTERNATIVE 2 ALTERNATIVE 3			
	Proposed	Exclude	Exclude	
RESOURCE	Action	North Aleutian Basin	Cook Inlet	
Air Quality	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	
	Emissions from routine	Same as Alternative 1.	Same as Alternative 1.	
	operations well within national			
	standards. Oil spills and in situ			
	burning effects localized and			
	short-term. Alaska	Alaska	Alaska	
	Emissions from routine	No OCS-related impacts would	No OCS-related impacts would	
	operations well within national	occur in the North Aleutian	occur in Cook Inlet. Impacts in	
	standards. Oil spills and in situ	Basin. Impacts in other Alaska	other program areas in Alaska	
	burning effects localized and	program areas would be the	would be the same as in	
	short-term.	same as those in Alternative 1.	Alternative 1.	
	<u>Atlantic</u>	<u>Atlantic</u>	Atlantic	
	Emissions from routine	Same as Alternative 1.	Same as Alternative 1.	
	operations well within national			
	standards. Oil spills and in situ			
	burning effects localized and			
	short-term.			
Water Quality	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	
	Localized and short-term	Same as Alternative 1.	Same as Alternative 1.	
	impacts from routine			
	operations. Compliance with NPDES permits would			
	minimize impacts to receiving			
	waters. Impacts from a large			
	coastal oil spill could persist for			
	an extended period of time if			
	oil were deposited in wetland			
	and beach sediments or collected in low energy coastal			
	environments.			
	Alaska	<u>Alaska</u>	<u>Alaska</u>	
	Impacts from routine	No OCS-related impacts would	No OCS-related impacts would	
	operations short term and	occur in the North Aleutian	occur in Cook Inlet. Impacts in	
	localized. No drilling	Basin. Impacts in other Alaska	other program areas in Alaska would be the same as in	
	discharges allowed during development. Impacts from oil	program areas would be the same as those in Alternative 1.	Alternative 1.	
	spills would generally be	same as arose in Atternative 1.	Thermative 1.	
	localized. Cleanup operations			
	could be difficult in the Arctic			
	because of cold and ice.			

	ALTERNATIVE 4 Exclude	ALTERNATIVE 5 Defer within 25 Miles	ALTERNATIVE 6 Defer Mouth of	ALTERNATIVE 7 Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Air Quality	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Reduced emissions in nearshore environment.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.
	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.
Water Quality	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Reduced impacts in nearshore areas in Chukchi Sea.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Water Quality (Continued)	Atlantic Localized and short-term impacts from routine operations. Compliance with NPDES permits would	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.
	minimize impacts to receiving waters. Impacts from a large coastal oil spill, though unlikely, could persist for extended period of time if oil were deposited in wetland and beach sediments or collected in low energy coastal environments.		
Marine	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Mammals	Noise from seismic activity and routine operations could disturb individual animals. Collision with service vessels could kill individual sperm whales. Collision impacts to manatees unlikely because they do not occur in program area.	Same as Alternative 1.	Same as Alternative 1.
	Alaska Acoustic impacts sublethal and temporary. Oil-spill impacts could affect survival rates of young animals. Although unlikely, a large spill that contacts Steller sea lion critical habitat in the North Aleutian Basin could have population-level impacts that could also affect the Cook inlet population. Collisions with service vessels could result in mortality, but low activity levels and MMS guidelines and regulation should minimize collisions.	Alaska No potential for OCS-related impacts to Steller sea lions in the North Aleutian Basin. As a result, possible impacts on the seasonal Steller sea lion population in Cook Inlet could not occur. Impacts in the Arctic area would be the same as in Alternative 1.	Alaska No OCS-related impacts would occur in Cook Inlet. Impacts in other program areas in Alaska would be the same as in Alternative 1.

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
	Exclude	<b>Defer within 25 Miles</b>	<b>Defer Mouth of</b>	Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Water Quality (Continued)	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic Reduced impacts in nearshore areas.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.
Marine Mammals	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Reduced impacts in nearshore areas in Chukchi Sea.	Alaska Same as Alternative 1.	Alaska Steller sea lion and Steller eider would face slightly reduced risk from oil spills.

RESOURCE	ALTERNATIVE 1 Proposed Action	ALTERNATIVE 2 Exclude North Aleutian Basin	ALTERNATIVE 3 Exclude Cook Inlet
Marine Mammals (Continued)	Atlantic Acoustic impacts sublethal and temporary. Lethal impacts from collisions with a service vessel and/or tanker is possible although unlikely considering the expected low level of activity. The endangered right whale could experience population-level impacts from collision mortality.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.
Marine and Coastal Birds	Gulf of Mexico Routine operations would disturb birds in immediate vicinity of activity. Impacts would be short-term and would not affect bird populations. Collision mortality with offshore structures could result in the death of several thousand birds per year. Large spills near the coast would impact birds through direct contact as well as from contaminated food and nesting areas. The wild endangered whooping crane population wintering in Texas would be vulnerable to spill impacts.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.

	ALTERNATIVE 4 Exclude	ALTERNATIVE 5 Defer within 25 Miles	ALTERNATIVE 6 Defer Mouth of	ALTERNATIVE 7 Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Marine	Atlantic	<u>Atlantic</u>	<u>Atlantic</u>	<u>Atlantic</u>
Mammals	No OCS-related	Reduced impact in	Slight reduction in	Same as Alternative 1.
(Continued)	impacts would occur	nearshore areas off	risk of collision with	
	offshore Virginia.	Virginia.	marine mammals.	
	Alternative 4 removes			
	the slight possibility of OCS vessel or			
	tanker collision with a			
	right whale, which			
	could affect the right			
	whale population.			
Marine and	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Coastal Birds	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Marine and Coastal Birds (Continued)	Alaska Noise and onshore construction of facilities and pipelines would result in localized and short-term disturbances. Potential for population-level impacts if important breeding and nesting areas and behaviors affected. New facilities could lead to increases in predators, such as arctic fox and glaucous gull, that could lead to population-level impacts on birds. Oil spills could impact birds from direct contact and from tainting of food and nesting material. Although	Alaska Eliminates the possibility of impacts to Steller's eider in North Aleutian Basin, which could affect the breeding population in the Arctic.	Alaska No OCS-related impacts would occur in Cook Inlet. Impacts in other program areas in Alaska would be the same as in Alternative 1.
	unlikely, a large spill contacting Steller's eiders during the winter in North Aleutian Basin could result in population-level impacts and affect populations in the Arctic area.  Atlantic Impacts from routine operations and oil spill would affect birds locally. Because of the low level of activity and the small amount of liquid hydrocarbons estimated to be produced, no population-level impacts are expected.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.
Terrestrial Mammals	Gulf of Mexico No population-level impacts to endangered beach mice and voles from routine operations. Beach mice are unlikely to be contacted by oil spills because of habitat location on inner dunes in the eastern half of the Gulf of Mexico, away from most activities associated with Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
	Exclude	<b>Defer within 25 Miles</b>	<b>Defer Mouth of</b>	Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Marine and Coastal Birds (Continued)	Alaska Same as Alternative 1.	Alaska Reduced impacts in nearshore areas in Chukchi Sea.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.
	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic Reduced impact in nearshore areas off Virginia.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.
Terrestrial Mammals	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Terrestrial Mammals (Continued)	Alaska Construction and operation of onshore facilities and pipelines could result in short- and long-term impacts from disturbances and loss of habitat. While direct impacts would be short-term and localized, stress could affect reproductive success resulting in longer term impacts to populations. Displacement from preferred habitats along pipeline corridors could affect muskoxen calving success.  Development of facilities would result in more contact between humans and bears and possible increased bear mortality in defense of life and property situations. Impacts from oil spills could affect individual mammals but not	Alaska No OCS-related impacts would occur in the North Aleutian Basin. Impacts in other Alaska program areas would be the same as those in Alternative 1.	Alaska No OCS-related impacts would occur in Cook Inlet. Impacts in other program areas in Alaska would be the same as in Alternative 1.
	overall population levels.  Atlantic Impacts to terrestrial mammals were not analyzed in the Atlantic because no endangered, threatened, or listed species occur in the area of potential impact	Atlantic Same as Alternative 1	Atlantic Same as Alternative 1
Fish Resources and EFH	Gulf of Mexico Minimal possibility for impacts to Gulf Sturgeon because of little overlap between where the sturgeon occurs and areas where Alternative 1 activities occur. No population-level impacts to other fish species from routine operations or oil spills. Potential impacts on EFH due to routine operations would be limited because only a small proportion of EFH could be affected under Alternative 1, Large spills that reach coastal wetlands could have persistent impacts on EFH and could require remediation.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
	Exclude	<b>Defer within 25 Miles</b>	<b>Defer Mouth of</b>	Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Terrestrial Mammals (Continued)	Alaska Same as Alternative 1.	Alaska Reduced risk from oil spills in Chukchi Sea.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.
Fish Resources	Atlantic Terrestrial mammals were not analyzed because no ending- ered or threatened species occur in the impact area. Gulf of Mexico	Atlantic Reduced risk from oil spills off Virginia.  Gulf of Mexico	Atlantic Same as Alternative 1.  Gulf of Mexico	Atlantic Same as Alternative 1.  Gulf of Mexico
and EFH	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.

ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
		Exclude
=		
Alaska Some short-term and localized impacts from routine activities, such as seismic surveys, offshore construction, and drilling discharges in the vicinity of the activity. A large spill contacting intertidal areas could lead to considerable mortality of eggs and juvenile stages of some pelagic species, but no population- level impacts are expected. Routine activities could result in localized and short-term impacts to EFH. A large spill that reaches coastal streams and intertidal areas used for spawning by anadromous salmon could have more persistent impacts and require remediation.  Atlantic Localized impacts could occur from routine operations and oil	North Aleutian Basin  Alaska No OCS-related impacts would occur in the North Aleutian Basin. Impacts in other Alaskan program would be the same as those in Alternative 1.  Atlantic Same as Alternative 1.	Alaska No OCS-related impacts would occur in Cook Inlet. Impacts in other program areas in Alaska would be the same as in Alternative 1.  Atlantic Same as Alternative 1.
Gulf of Mexico Impacts to individual turtles could occur from seismic activity, operational discharges, platform removals, or collisions with service vessels, but no population-level impacts are expected. Oil spills could affect turtles at all life stages by direct contact or deterioration of their preferred habitat. Alaska Sea turtles do not occur in	Gulf of Mexico Same as Alternative 1.  Alaska Sea turtles do not occur in	Gulf of Mexico Same as Alternative 1.  Alaska Sea turtles do not occur in
	Action  Alaska Some short-term and localized impacts from routine activities, such as seismic surveys, offshore construction, and drilling discharges in the vicinity of the activity. A large spill contacting intertidal areas could lead to considerable mortality of eggs and juvenile stages of some pelagic species, but no population-level impacts are expected. Routine activities could result in localized and short-term impacts to EFH. A large spill that reaches coastal streams and intertidal areas used for spawning by anadromous salmon could have more persistent impacts and require remediation.  Atlantic Localized impacts could occur from routine operations and oil spills. No population-level impacts expected.  Gulf of Mexico Impacts to individual turtles could occur from seismic activity, operational discharges, platform removals, or collisions with service vessels, but no population-level impacts are expected. Oil spills could affect turtles at all life stages by direct contact or deterioration of their preferred habitat.  Alaska	Action  Alaska Some short-term and localized impacts from routine activities, such as seismic surveys, offshore construction, and drilling discharges in the vicinity of the activity. A large spill contacting intertidal areas could lead to considerable mortality of eggs and juvenile stages of some pelagic species, but no population- level impacts are expected. Routine activities could result in localized and short-term impacts to EFH. A large spill that reaches coastal streams and intertidal areas used for spawning by anadromous salmon could have more persistent impacts and require remediation.  Atlantic  Localized impacts could occur from routine operations and oil spills. No population- level impacts expected.  Gulf of Mexico Impacts to individual turtles could occur from seismic activity, operational discharges, platform removals, or collisions with service vessels, but no population-level impacts are expected. Oil spills could affect turtles at all life stages by direct contact or deterioration of their preferred habitat.  Alaska Sea turtles do not occur in

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
	Exclude	<b>Defer within 25 Miles</b>	<b>Defer Mouth of</b>	Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Fish Resources	Alaska	Alaska	Alaska	Alaska
and EFH	Same as Alternative 1.	Reduced impacts in	Same as Alternative 1.	Same as Alternative 1.
(Continued)		nearshore areas in		
		Chukchi Sea.		
	Atlantic	Atlantic	Atlantic	Atlantic
	No OCS-related	Reduced impact in	Same as Alternative 1.	Same as Alternative 1.
	impacts would occur	nearshore areas off	Sume as internative i.	Sume as internative i.
	offshore Virginia.	Virginia.		
	Impacts in other			
	program areas are			
	same as Alternative 1.			
Sea Turtles	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.
	<u>Alaska</u>	<u>Alaska</u>	<u>Alaska</u>	<u>Alaska</u>
	Sea turtles do not	Sea turtles do not occur	Sea turtles do not	Sea turtles do not
	occur in Alaska.	in Alaska.).	occur in Alaska.	occur in Alaska.

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Sea Turtles	Atlantic	Atlantic	Atlantic
(Continued)	Impacts from routine activities	Same as Alternative 1.	Same as Alternative 1.
,	and small oil spills would not		
	affect sea turtle distribution,		
	behavior, or abundance. Other		
	than collisions with vessels or		
	large oil spills, potential		
	impacts are expected to be		
	sublethal. Although unlikely,		
	a large oil spill contacting turtle		
	habitat could result in		
G . 1	population-level impacts.	0.10.034	G 10 035 :
Coastal	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Habitats	Onshore construction of	Same as Alternative 1.	Same as Alternative 1.
	facilities and pipelines and service vessel traffic could		
	affect coastal wetlands and		
	barrier beaches through direct		
	losses, accelerated erosion, and		
	indirect affects associated with		
	hydrologic changes and water		
	quality impacts. Oil spills and		
	cleanup operations would also		
	affect wetlands.		
	<u>Alaska</u>	<u>Alaska</u>	<u>Alaska</u>
	Onshore construction of	No OCS-related impacts would	No OCS-related impacts would
	facilities and pipelines could	occur in the North Aleutian	occur in Cook Inlet. Impacts in
	affect wetlands and coastal	Basin. Impacts in other	other program areas in Alaska
	barriers through direct loss of	Alaskan program would be the	would be the same as in
	habitat and through indirect	same as those in Alternative 1.	Alternative 1.
	affects of altered hydrology and		
	lowered water quality. Oil-spill		
	impacts could result in habitat	•	
	loss from dieback of vegetation		
	and damage from cleanup		
	operations.		

RESOURCE Sea Turtles (Continued)	ALTERNATIVE 4 Exclude Mid-Atlantic  Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	ALTERNATIVE 5 Defer within 25 Miles Virginia & Chukchi  Atlantic Reduced risk from oil spills off Virginia	ALTERNATIVE 6 Defer Mouth of Chesapeake Bay Atlantic Same as Alternative 1.	ALTERNATIVE 7 Limit Leasing in North Aleutian  Atlantic Same as Alternative 1.
Coastal Habitats (Continued)	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Reduced risk from oil spills in Chukchi Sea.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Coastal	Atlantic	Atlantic	Atlantic
Habitats	Direct and indirect impacts to	Same as Alternative 1.	Same as Alternative 1.
(Continued)	barrier beaches and dunes		
	could occur from pipeline		
	construction and vessel traffic.		
	New construction is expected		
	to occur in industrialized areas,		
	so impacts to sensitive coastal		
	habitats should not occur.		
	Impacts from oil spills could		
	range from an insignificant to a		
	high amount of damage and		
	habitat loss, depending on		
Seafloor	location and size of spill.	Culf of Marian	Culf of Maxico
Habitats	Gulf of Mexico Physical impacts to topographic	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
Habitats	features, pinnacles, and	Same as Atternative 1.	Same as Atternative 1.
	chemosynthetic communities		
	from pipeline and platform		
	emplacements and oil-spill		
	impacts will be minimized by		
	existing MMS lease stipulations		
	that prohibit all or some		
	activities near the features. No		
	long-term population-level		
	impacts are expected. No		
	impacts to seagrasses are		
	expected because large areas of		
	seagrass habitat do not occur in		
	the Gulf of Mexico program		
	areas.	A11.	A11.
	Alaska The Stefansson Boulder Patch	Alaska	Alaska No OCS-related related
	communities in the Arctic	No OCS-related impacts would occur in the North Aleutian	impacts would occur in Cook
	would be vulnerable to impacts	Basin. Impacts in other Alaska	Inlet. Impacts in other program
	from routine operations and oil	program areas would be the	areas in Alaska would be the
	spills, but planning procedures,	same as those in Alternative 1.	same as in Alternative 1.
	lease stipulations and	The state of the s	
	regulations should minimize or		
	eliminate these potential		
	impacts. Possible impacts to		
	other benthic habitats from		
	routine operations, discharges,		
	and oil spills are expected to be		
	localized and short-term.		

RESOURCE Coastal Habitats (Continued)	ALTERNATIVE 4 Exclude Mid-Atlantic  Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	ALTERNATIVE 5 Defer within 25 Miles Virginia & Chukchi  Atlantic Reduced risk from oil spills off Virginia.	ALTERNATIVE 6 Defer Mouth of Chesapeake Bay Atlantic Same as Alternative 1.	ALTERNATIVE 7 Limit Leasing in North Aleutian  Atlantic Same as Alternative 1.
Seafloor Habitats	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Reduced impacts in nearshore areas in Chukchi Sea.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Seafloor	Atlantic	Atlantic	Atlantic
Habitats	Impacts from routine	Same as Alternative 1.	Same as Alternative 1.
(Continued)	operations and oil spills could		
	occur, but lease stipulations and		
	regulations, combined with the		
	low level of activity expected		
	offshore Virginia should result		
	in no impacts to seafloor		
A	habitats.	C 1C · CM·····	C 1C · CM· · · ·
Areas of Special	Gulf of Mexico Routine operations should not	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
Concern	result in impacts to national	Same as Alternative 1.	Same as Alternative 1.
Concern	parks, wildlife refuges, and		
	national estuary program sites		
	because these areas are		
	restricted from development.		
	Impacts from oil spills should		
	be uncommon because most		
	development will occur in		
	deepwater areas far from the coast. Marine sanctuaries are		
	protected by lease stipulations		
	and regulations that prohibit		
	activities within a certain		
	distance of the features.		
	Alaska	<u>Alaska</u>	Alaska
	Impacts to national parks and	No OCS-related impacts would	No OCS-related impacts would
	wildlife refuges from routine	occur in the North Aleutian	occur in Cook Inlet. Impacts in
	operations are unlikely to occur	Basin. Impacts in other Alaska	other program areas in Alaska
	because no OCS related	program areas would be the	would be the same as in
	activities will occur within their boundaries. A large nearshore	same as those in Alternative 1.	Alternative 1.
	oil spill could contact coastal		
	refuges and affect the coastal		
	habitats there.		
	Atlantic	<u>Atlantic</u>	<u>Atlantic</u>
	Routine operations would not	Same as Alternative 1.	Same as Alternative 1.
	affect these areas because of		
	their location relative to		
	activities and the expected		
	small amount of activity. A		
	large coastal spill, although unlikely, could threaten some		
	areas.		
l	uroup.		<u>l</u>

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
DEGOVED CE	Exclude	Defer within 25 Miles	Defer Mouth of	Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Seafloor Habitats (continued)	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other mid- Atlantic program areas would be the same as in Alternative 1.	Atlantic Reduced risk from oil spills off Virginia.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.
Areas of Special Concern	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Reduced impacts in nearshore areas in Chukchi Sea.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.
	Atlantic No potential OCS- related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic Reduced impacts in nearshore areas off Virginia.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Population,	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Employment,	Employment and regional	Same as Alternative 1.	Same as Alternative 1.
and Income	income impacts would be		
	greatest in Texas and		
	Louisiana. Added employment		
	demands are not expected to		
	tax the local labor market. A		
	coastal oil spill could result in a		
	one-time seasonal decline in		
	tourism-related business		
	activity.		
	Alaska	Alaska	Alaska
	No noticeable impacts are	No OCS-related impacts would	Same as Alternative 1.
	anticipated from routine	occur in the North Aleutian	
	operations. A large oil spill could result in noticeable	Basin. Impacts in other Alaska program areas would be the	
	impacts through damage to	same as those in Alternative 1.	
	various industries.	same as mose in Alternative 1.	
	Atlantic	Atlantic	Atlantic
	Some in-migration of labor to	Same as Alternative 1.	Same as Alternative 1.
	fill oil and gas exploration and	24110 45 1 11011141 70 17	
	development jobs because		
	skills don't occur locally.		
	Because of small amount of		
	activity and the robust existing		
	maritime economy of the area,		
	OCS activity is not expected to		
	have an affect on the local		
	housing or labor market.		
Sociocultural	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Systems	The greatest impacts would	Same as Alternative 1.	Same as Alternative 1.
	occur from the ongoing		
	expansion of deepwater		
	activities that result in longer		
	unbroken periods of offshore		
	work and in-migration of part of the workforce.		
	Alaska	Alaska	Alaska
	The largest impacts would	No OCS-related impacts would	No OCS-related impacts would
	occur from oil spills that could	occur in the North Aleutian	occur in Cook Inlet. Impacts in
	affect subsistence hunting and	Basin. Impacts in other Alaska	other program areas in Alaska
	activities.	program areas would be the	would be the same as in
		same as those in Alternative 1.	Alternative 1.

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
	Exclude	<b>Defer within 25 Miles</b>	Defer Mouth of	Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Population,	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Employment, and Income	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.
	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.
	~	G 10 10 1	G 10 03.5	G 10 03 5 1
Sociocultural Systems	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Reduced risk to subsistence hunting in Chukchi.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.

RESOURCE	ALTERNATIVE 1 Proposed Action	ALTERNATIVE 2 Exclude North Aleutian Basin	ALTERNATIVE 3 Exclude Cook Inlet
Sociocultural Systems (Continued)	Atlantic (including Environmental Justice) Limited impacts in these areas are expected because of the small amount of activity anticipated in Alternative 1, the economic and sociocultural	Atlantic (including Environmental Justice) Same as Alternative 1.	Atlantic (including Environmental Justice) Same as Alternative 1.
Environmental Justice	diversity of the population, and the high degree of racial and ethnic integration.  Gulf of Mexico Coastal infrastructure use and	Gulf of Mexico Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.
	construction levels would be similar to previous programs. These activities would not expose residents to notable higher risks than currently occur. No impacts from oil spills are expected because of the movement of OCS activities farther from the coast into deeper water and the demographic pattern of affluent groups living in coastal areas.		
	Alaska Potential for new onshore infrastructure to be located near Native populations and to produce adverse health or environmental impacts if there were impacts on subsistence resources and harvest patterns. Oil-spill environmental and health impacts on Alaska Native populations could be disproportionately high or adverse depending on the geographical location of the spill and the effects on subsistence resources and	Alaska No OCS-related impacts would occur in the North Aleutian Basin. Impacts in other Alaska Program areas would be the same as those in Alternative 1.	Alaska No OCS-related impacts would occur in the North Aleutian Basin. Impacts in other Alaska Program areas would be the same as those in Alternative 1.
	harvests. Atlantic Included in Sociocultural Systems.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
	Exclude	<b>Defer within 25 Miles</b>	<b>Defer Mouth of</b>	Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Sociocultural	Atlantic (including	Atlantic (including	Atlantic (including	Atlantic (including
Systems	Environmental	Environmental Justice)	Environmental Justice)	Environmental Justice)
(Continued)	Justice)	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.
	No OCS-related			
	impacts would occur offshore Virginia.			
	Impacts in other			
	program areas would			
	be the same as in			
	Alternative 1.			
Environmental	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Justice	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.
	Alaska	Alaska	Alaska	Alaska
	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.
	<u>Atlantic</u>	<u>Atlantic</u>	<u>Atlantic</u>	<u>Atlantic</u>
	No OCS-related	No OCS-related	Same as Alternative 1.	Same as Alternative 1.
	impacts would occur	impacts would occur		
	offshore Virginia.	offshore Virginia.		
		1 0		
	Alternative 1.	Alternative 1.		
	Impacts in other program areas would be the same as in	Impacts in other program areas would be the same as in		
	Anternative 1.	Anternative 1.		

	ALTERNATIVE 1 Proposed	ALTERNATIVE 2 Exclude	ALTERNATIVE 3 Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Archaeology	Gulf of Mexico Adherence to existing Federal, State, and local regulations and policies will protect most archaeological resources from impacts from routine opera- tions. Impacts could occur to coastal historic and prehistoric archaeological resources from accidental oil spills. Limited impacts are expected directly from the spill, but some impacts could occur during cleanup operations Alaska Same as Gulf of Mexico.	Alaska No OCS-related impacts would occur in the North Aleutian Basin. Impacts in other Alaska program areas would be the same as those in Alternative 1.  Atlantic Same as Alternative 1.	Alaska No OCS-related impacts would occur in Cook Inlet. Impacts in other program areas in Alaska would be the same as in Alternative 1.  Atlantic Same as Alternative 1.
Land Use and Infrastructure	Gulf of Mexico New facilities are expected to be constructed along the Central and Western Gulf of Mexico in areas already affected by decades of offshore oil and gas development.  Alaska The greatest potential impact would be to expose undeveloped areas to OCS-related onshore construction. Oil spills could result in temporary land-use changes.  Atlantic With the small amount of activity anticipated and the existing maritime economy and associated facilities in the area, no impacts to infrastructure are expected.	Gulf of Mexico Same as Alternative 1.  Alaska No OCS-related impacts would occur in the North Aleutian Basin. Impacts in other Alaskan program would be the same as those in Alternative 1.  Atlantic Same as Alternative 1.	Gulf of Mexico Same as Alternative 1.  Alaska No OCS-related impacts would occur in Cook Inlet. Impacts in other program areas in Alaska would be the same as in Alternative 1.  Atlantic Same as Alternative 1.

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
RESOURCE	Exclude Mid-Atlantic	Defer within 25 Miles Virginia & Chukchi	Defer Mouth of Chesapeake Bay	Limit Leasing in North Aleutian
Archaeology	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.
	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic Reduced risk of direct physical contact with archaeological resources.	Atlantic Reduced risk because of smaller area of possible activity.
Land Use and	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Infrastructure	Same as Alternative 1.  Alaska Same as Alternative 1.	Same as Alternative 1.  Alaska Same as Alternative 1.	Same as Alternative 1.  Alaska Same as Alternative 1.	Same as Alternative 1.  Alaska Same as Alternative 1.
	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	<b>North Aleutian Basin</b>	Cook Inlet
Tourism and	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Recreation	Routine operations would have	Same as Alternative 1.	Same as Alternative 1.
	limited effects, with potential		
	adverse aesthetic impacts to		
	beach recreation and		
	sightseeing and potential		
	positive impacts for diving and		
	recreational fishing.		
	<u>Alaska</u>	<u>Alaska</u>	<u>Alaska</u>
	Routine operations could have	No OCS-related impacts would	No OCS-related impacts would
	limited effects, particularly in	occur in the North Aleutian	occur in Cook Inlet. Impacts in
	previously undeveloped oil and	Basin. Impacts in other Alaska	other program areas in Alaska
	gas areas where there is limited	program areas would be the	would be the same as in
	infrastructure to accommodate	same as those in Alternative 1.	Alternative 1.
	both tourists and construction		
	workers. New onshore and		
	offshore infrastructure could		
	limit access to recreational		
	users. Oil spills would affect		
	tourism and recreation		
	temporarily.		
	<u>Atlantic</u>	<u>Atlantic</u>	<u>Atlantic</u>
	Although routine activities could	Same as Alternative 1	Same as Alternative 1
	result in some visual and		
	branding impacts on tourism		
	An oil spill could result in		
	temporary beach closings. Any		
	impacts that do occur are		
	expected to be short-term and/or		
	localized.		
Fisheries	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
	Some short-term and localized	Same as Alternative 1.	Same as Alternative 1.
	impacts to recreational and		
	commercial fisheries from		
	routine operations. An oil spill		
	would not affect the population		
	of these fish but could		
	temporarily affect recreational		
	and commercial fishing in areas		
	contacted by the spill.		

RESOURCE	ALTERNATIVE 4 Exclude Mid-Atlantic	ALTERNATIVE 5 Defer within 25 Miles Virginia & Chukchi	ALTERNATIVE 6 Defer Mouth of Chesapeake Bay	ALTERNATIVE 7 Limit Leasing in North Aleutian
Tourism and	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico	Gulf of Mexico
Recreation	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.	Same as Alternative 1.
	Alaska Same as Alternative 1.	Alaska Same as Alternative 1 in Chukchi Sea.	Alaska Same as Alternative 1.	Alaska Same as Alternative 1.
	Atlantic No OCS-related impacts would occur offshore Virginia. Impacts in other program areas would be the same as in Alternative 1	Atlantic Reduced risk from oil spills off Virginia.	Atlantic Reduced risk from oil spills off Virginia and elimination of visual impacts from shore	Atlantic Same as Alternative 1.
Fisheries	Gulf of Mexico Same as Alternative 1	Gulf of Mexico Same as Alternative 1	Gulf of Mexico Same as Alternative 1	Gulf of Mexico Same as Alternative 1

	ALTERNATIVE 1	ALTERNATIVE 2	ALTERNATIVE 3
	Proposed	Exclude	Exclude
RESOURCE	Action	North Aleutian Basin	Cook Inlet
Fisheries	<u>Alaska</u>	Alaska	<u>Alaska</u>
(Continued)	Commercial and recreational	No OCS-related impacts would	No OCS-related impacts would
,	fisheries in the Arctic are	occur in the North Aleutian	occur in Cook Inlet. Impacts in
	relatively small and localized.	Basin. Impacts in other Alaska	other program areas in Alaska
	Impacts from routine	program areas would be the	would be the same as in
	operations and accidents would	same as those in Alternative 1.	Alternative 1.
	not occur unless located in the		
	direct vicinity of the fishery.		
	Local and short-term effects		
	from routine operations and oil		
	spills could occur in the Cook		
	Inlet and North Aleutian Basin		
	areas, but no population-level		
	impacts are expected.		
	<u>Atlantic</u>	<u>Atlantic</u>	<u>Atlantic</u>
	Because of the small amount of	Same as Alternative 1.	Same as Alternative 1.
	activity expected offshore		
	Virginia, no long-term impacts		
	to fisheries from discharges,		
	seismic activity, noise, or lights		
	on offshore rigs are expected to		
	occur. Death or reduced		
	biological functioning could		
	occur to some economically		
	important fish, but the impacts		
	to fish populations are not		
	expected to be distinguishable		
	from natural variations.		

	ALTERNATIVE 4	ALTERNATIVE 5	ALTERNATIVE 6	ALTERNATIVE 7
	Exclude	Defer within 25 Miles	<b>Defer Mouth of</b>	Limit Leasing in
RESOURCE	Mid-Atlantic	Virginia & Chukchi	Chesapeake Bay	North Aleutian
Fisheries	<u>Alaska</u>	<u>Alaska</u>	<u>Alaska</u>	<u>Alaska</u>
(Continued)	Same as Alternative 1.	Reduced risk from oil	Same as Alternative 1.	Same as Alternative 1.
		spills in Chukchi Sea.		
	Atlantic No OCS-related impacts would occur offshore Virginia.	Atlantic Reduced risk fro oil spills off Virginia.	Atlantic Same as Alternative 1.	Atlantic Same as Alternative 1.
	Impacts in other program areas would be the same as in Alternative 1.			

77	A Ita	rnatives
"	AIII	T HULLVES