

SCALE DEPENDENCE IN THE EFFECTS OF FOREST COVERAGE ON PARASITIZATION BY BROWN-HEADED COWBIRDS

W. M. HOCHACHKA*, T. E. MARTIN*, V. ARTMAN, C. R. SMITH, S. J. HEJL, D. E. ANDERSEN, D. CURSON, L. PETIT, N. MATHEWS, T. DONOVAN, E. E. KLAAS, P. B. WOOD, J. C. MANOLIS, K. P. MCFARLAND, J. V. NICHOLS, J. C. BEDNARZ, D. M. EVANS, J. P. DUGUAY, S. GARNER, J. TEWKSBURY, K. L. PURCELL, J. FAABORG, C. B. GOGUEN, C. RIMMER, R. DETTMERS, M. KNUTSON, J. A. COLLAZO, L. GARNER, D. WHITEHEAD, AND G. GEUPEL

Abstract. Previous work has shown that the rate at which Brown-headed Cowbirds (*Molothrus ater*) parasitize forest nesting birds is affected by the proportion of a local landscape that is forested. However, much of the previous work has been restricted to a relatively small part of the cowbird's range, and has looked at forest coverage in very restricted areas around study plots. We used data from a wider geographical area, the entire width of the United States, and examined forest coverage in relatively large areas (10-km and 50-km radii) around study plots to determine if forest coverage is a generally useful statistic for predicting rates of brood parasitization. As was found in previous studies, we showed that increased amounts of forest coverage within 10 km of an area resulted in lower rates of parasitization by cowbirds. This pattern held not only among widely separated sites, but also within local clusters of study plots. However, we found that increased amounts of forest within 50 km of a study site resulted in slightly increased rates of parasitization in sites west of the Great Plains, contrary to previous research findings. Forest structure, as indicated by the relationship between forest coverage and other measures of forest distribution and abundance, differed across the United States. However, differences in forest structure were not obviously related to differences in the manner that parasitization and forest coverage covaried from east to west across the continent. Even given the variable patterns found, management for higher proportions of forest within 10-km radius areas should result in decreased rates of parasitization of host species; however, the impact of such a management strategy will vary across the continent.

Key Words: Brown-headed Cowbird, forest coverage, geographical variation, landscape structure, *Molothrus ater*, parasitization rate, scale.

Areas containing a greater proportion of forest have a lower abundance of Brown-headed Cowbirds (*Molothrus ater*) (Donovan et al. 1997, Donovan et al. in press, Tewksbury et al. 1998), and show a lower rate of parasitization of the nests of host species (Robinson et al. 1995b, Thompson et al. in press). The conclusion of this previous research is that larger proportions of forest, relative to all terrestrial habitats (the landscape), will result in a lower impact of Brown-headed Cowbirds on their hosts.

However, the majority of work relating forest coverage to rates of parasitization is from the eastern edge of the Great Plains (e.g., Robinson et al. 1995b, Donovan et al. 1997, Donovan et al. in press, Thompson et al. in press; but see Coker and Capen 1995, Tewksbury et al. 1998 for exceptions). We might expect the relationship between forest coverage and parasitization to differ away from the Midwest for a number of reasons. Variation in cowbird abundance may not only affect absolute rates of parasitization (Thompson et al. in press), but also the pattern

of variation in parasitization rate with varying forest coverage. Cowbirds in different parts of the continent encounter communities of hosts with different lengths of exposure (e.g., Mayfield 1965) and responses (e.g., Briskie et al. 1992) to parasitization, and host species with longer exposure to cowbirds may be resistant to parasitization regardless of the proportion of forest in a landscape.

Geographical variation in the relationship between forest coverage and parasitization rate also may result because of geographical differences in the pattern of forest in a landscape. Cowbirds may respond to the amount of edge (Gates and Gysel 1978, Brittingham and Temple 1983, Thompson et al. in press), distance from foraging sites (Donovan et al. in press), or other features correlated with forest coverage. Within a region, the proportion of forest in a landscape may correlate well with measures such as the amount of edge (Robinson et al. 1995b). However, land-use practices and topography vary across the continent, such that the relationship between forest coverage and features such as edge may vary across the continent.

The relationship between cowbird parasitiza-

* Ordering of names of authors subsequent to T. E. Martin determined using a random number generator.

tion and forest coverage may vary as a function of the local area over which forests were measured, in addition to varying among widely separate regions of the continent. Research relating forest coverage to rates of cowbird parasitization initially examined effects of variation in the size of individual forest patches and distance from forest edges (e.g., Paton 1994), and only recently has looked at local landscapes around individual forest patches (e.g., Robinson et al. 1995b, Donovan et al. in press, Tewksbury et al. 1998). Within these local areas, forest coverage varied in its power to predict parasitization, depending on the size of the area over which forest coverage was measured (Donovan et al. in press, Tewksbury et al. 1998). However, it is still not clear whether the range of areas measured (up to 10-km radius) encompass those that give the best predictions of the rate of parasitization. Local variation in forest coverage may only affect the movements of individual cowbirds (functional responses). Better predictions of the rate of cowbird parasitization may be provided by measuring forest coverage over larger areas than previously considered, if forest coverage over larger regions predict the abundance of cowbirds (a numeric response) and rates of parasitization are better predicted by cowbird abundance than the behavior of individual cowbirds. Knowledge of the most appropriate scale on which to manage forest coverage is essential for informed decisions about land management.

Differences in forest coverage may not predict the same change in the rate of parasitization depending on whether the sites being compared are widely separated. To date, studies have looked at variation in cowbird abundance or parasitization in relation to either local (e.g., Tewksbury et al. 1998) or regional (e.g., Robinson et al. 1995b, Thompson et al. in press) variation in forest coverage, but not both simultaneously. It is still unclear whether parasitization rates vary with local differences in forest coverage in the same manner as they respond to differences in forest coverage among more widely spaced sites, because the proportion of forest in a local landscape may be highly correlated with the proportion of forest within a far wider region.

This paper examines four questions: (1) does the relationship between forest coverage and other measures of landscape structure (e.g., amount of edge, size of forest patches) vary across the continent? (2) do changes in forest coverage over small distances predict the same variation in parasitization rates as changes in forest coverage among sites more widely separated? (3) does the relationship between forest coverage and cowbird parasitization vary with the size of the region over which forest coverage

is measured? and (4) does the relationship between forest coverage and parasitization differ among the eastern, central, and western United States? In conducting our analyses, we had no prior expectations of the patterns that would emerge. Our goal was to document patterns that could affect the way land managers use the previously described pattern of lower cowbird parasitization in areas containing a higher proportion of forested land.

METHODS

The data on parasitization rates of forest birds come from the Breeding Biology Research and Monitoring Database (BBIRD), with data from 23,448 individual nests being represented in our analyses. BBIRD is a collaborative project in which researchers across the United States have monitored nests and recorded data following a standardized protocol (Martin et al. 1997). There were 26 study sites (Fig. 1) on which the nesting success of forest-nesting birds was monitored. Data from five sites were previously used in the analyses of Robinson et al. (1995b). Each study site included 2 to 31 separate study plots (median = 9), with a total of 366 study plots in the data set. The spatial arrangement of study plots into local groups (termed "study sites") allowed us to contrast the effects of local (within tens of kilometers), and large-scale (across hundreds of kilometers) variation in forest coverage. This comparison was made by examining the relationship between forest coverage and the rate of parasitization both within study sites and among study sites.

The data obtained from each study plot were the proportion of nests containing cowbird eggs or young; potential hosts were only included when at least one nest of a species was recorded as having been parasitized in our database. Proportions were calculated across all species of hosts combined. Roughly 75% of all variance in the rate of parasitization occurred among plots within individual study sites (calculated following Sokal and Rohlf [1981:216]; we excluded data from sites on which cowbirds were not present). Given the high proportion of variance in parasitization rate that occurred within individual sites, we treated each of the study plots as an independent data point; i.e., we treated the data from each study plot as independent estimates of the rate of parasitization within the area that encompassed the separate study plots that compose a site.

The data on landscape structure came from an ARC/INFO GIS layer that was produced for the USDA Forest Service's Forest and Rangeland Renewable Resources Planning Act (RPA) 1993 Assessment Update (Anonymous no date). Data




FIGURE 1. Locations of study sites. Diamond-shaped points indicate sites designated as "eastern", triangles as "Midwestern", and squares as "western". Each site plotted on this map is composed of several independent study plots.

were derived from NOAA satellite images (AVHRR data), with the Forest Service project being completed at the end of 1992. The finest resolution of the GIS layer is a 1 km square that is classified as either water, non-forest, or forest; within forested areas the type of forest was specified as one of 22 types (e.g., oak-hickory, pinon-juniper). The relatively coarse resolution of the GIS layer placed constraints on our use and interpretation of the data on forest coverage. Each one of the 1-km squares could easily represent multiple patches of forest, detail that would be lost from our analyses. Additionally, our circles were approximate, with edge pixels from the GIS layer being included within a "circle" if >50% of that pixel was included within the circle. Because of the coarse resolution of the GIS layer, we used circles of 10-km radius (over 300 km²) as the minimum area in which forest coverage was measured. We made this decision in order to average measurement errors caused by individual pixels in the GIS layer containing fractions of both forested and non-forested land. However, in interpreting our results, we do not know what fraction of the unexplained variance in parasitization rates was caused by variation in the spatial arrangement of forest at a resolution finer than was provided by our GIS layer.

Statistics describing landscape structure were obtained using FRAGSTATS (McGarigal and Marks 1995). The areas in which landscape structure was described were circles of 10-km and 50-km radius surrounding each study plot. The 10-km radius, chosen to allow comparison with Robinson et al. (1995b), was based on observations of distances that female cowbirds fly between feeding and nesting areas in the Midwest (Thompson 1994). Although female cowbirds have also been found flying distances of under 10 km in California (Rothstein et al. 1984), work in New Mexico (C. B. Goguen and D. R. Curson, unpubl. data) has found female cowbirds flying in excess of 10 km between foraging and nesting sites. Thus landscape structure further than 10 km from study plots can potentially affect cowbirds' presence and abundance. Fifty km was arbitrarily chosen to represent larger spatial scales. The circles of 50-km radius contain 25 times the surface area as the 10-km circles and roughly 9 times greater area than was used in any previous study examining effects of forest coverage on cowbird abundance (Donovan et al. 1997). We did not use data from 50-km circles in comparisons of the rate of parasitization within study sites, because within individual study sites the study plots were often so closely spaced that 50-km forest coverage were

essentially identical among the plots within a single study site. In analyses examining presence and absence of parasitization among study sites, forest coverage for each site was calculated as the weighted average forest coverage around each study plot. Forest coverages were weighted by the proportion of a site's potential hosts that were found on each plot.

The proportion of a landscape in forest was used as the primary measure of landscape structure in this paper following the conventions of previous studies (e.g., Robinson et al. 1995b). However, other metrics generated by FRAGSTATS were also collated for each study plot: size of largest patch (as a proportion of the landscape), number of forest patches, mean size of forest patches, standard deviation in patch size, edge density (m/ha of edge), and the number of types of forest. Some of these metrics require further explanation because our FRAGSTATS calculations were done separately for each of the types of forest recognized in the original data set. As a result, we calculated edge density as the amount of non-forest edge, assuming that most non-forest edges were with forest. Additionally, the largest patch of forest in a landscape may be contiguous with other areas of forest of a different type, and the number of patches may not represent the actual number of discrete units of forest because patches of one type of forest may be nested within another type of forest. Still, these metrics represent some aspects of the spatial complexity of a landscape. Mean and standard deviation of patch size were calculated by decomposing the mean and SD for each forest type into sums and sums of squares and then calculating an overall mean and SD by combining this information across forest types.

Analyses relating parasitization rates to forest coverage were of two types: those examining whether variation in forest coverage affected whether any nests were parasitized, and those examining variation in the rate of parasitization given that at least some nests were parasitized. The former analyses concerned the presence or absence of parasitization, and we tested for patterns using logistic regression. For the latter analyses we used generalized linear models, and excluded sites on which no parasitization was found. Plots varied in the number of nests monitored, and thus the accuracy of our estimates of parasitization rates also varied. This varying accuracy was taken into account in our analyses by weighting each data point by $1/SE$ of the estimated rate of parasitization, which resulted in greater importance being placed on those data that were estimated with the greatest accuracy. In all analyses, continent-wide geographical variation in patterns were examined by dividing

study sites into three regions (Fig. 1): west of the Great Plains, Midwest (eastern edge of the Great Plains), and east. Data were also divided into two categories, east or west of the Great Plains, to test if better predictions were made when two or three regions were used in analyses.

Data from all sites were used simultaneously in analyses that tested for variation in parasitization rate within individual sites. To use data from all sites in a single analysis, we standardized forest coverages and rates of parasitization to have a mean value of zero within each group of study plots. This standardizing eliminated overall differences in forest coverage and rate of parasitization among these sites, and thus analyses of within-site variation exclusively examine variation relative to the average parasitization rate and forest coverage for a site. Forest coverages used in this analysis were within a 10-km radius of each study.

All statistical analyses were conducted using SPSS 7 (SPSS 1996). We refer to results from statistical tests as being "statistically significant" when $P \leq 0.05$. However, because statistical significance is not necessarily an indication of biological reality or importance (e.g., Thomas 1997), we have also noted instances in which the results of statistical tests approached but did not meet the arbitrary criterion of $P = 0.05$. In these instances, we have presented confidence limits (e.g., Greenland 1988, Steidl et al 1997, Thomas 1997) around parameters estimated in the analyses as a more refined indication of the potential biological significance of results.

RESULTS

Our results are divided into three sections. First, we examined landscape structure to show that landscape structure differed across the continent. These differences could provide a biological explanation for differences in the relationship between forest coverage and rates of cowbird parasitization across the continent. The second set of analyses examined whether variation in forest coverage was associated with the presence or absence of cowbird parasitization in a study area. Finally, where cowbirds were present, we show how the rate at which nests were parasitized was associated with forest coverage. These last two sets of analyses tested for geographical variation in parasitization rates, as well as for differences in the predicted effects of forest coverage that resulted from varying the area over which forest coverage was measured.

We examined the relationship between forest coverage and parasitization rates, both within local clusters of study plots and among widely separated study areas. The within-site analyses

TABLE 1. VARIATION IN LANDSCAPE STRUCTURE ALONG A GRADIENT OF FOREST COVER (10 KM SCALE). RESULTS ARE FROM GENERALIZED LINEAR MODELS

Forest metric	% Forest (10 km)			Region			Interaction ^c			R ²
	β	SE	P	β	SE	P	β	SE	P	
% Forest (50 km)	0.630	0.046	<0.001	6.736 ^a	5.860	<0.001	-0.096 ^a	0.075	<0.001	0.74
# Forest Types	0.028	0.004	<0.001	-17.068 ^b	4.358	<0.001	0.217 ^b	0.059	<0.001	0.35
# Forest Patches	0.196	0.038	<0.001	-1.703 ^a	0.451	<0.001	0.00915 ^a	0.006	<0.001	0.18
Mean Patch Size (ha)	12.131	13.713	0.001	1.185 ^b	0.335	<0.001	-0.0230 ^b	0.061	<0.001	0.12
SD Patch Size (ha)	30.978	7.721	<0.001	-4.775 ^a	4.732	<0.001	-0.00219 ^a	0.048	0.354	0.38
Edge Density (m/ha)	0.121	0.012	<0.001	19.971 ^b	3.513	0.899	-0.262 ^b	0.006	<0.001	0.73
Max. Patch Size (%)	-0.001 ^d	0.000	<0.001	-36.895 ^a	1687.8	<0.001	31.379 ^a	21.795	<0.001	0.71
	0.450	0.055	<0.001	-521.68 ^b	1253.1	0.014	11.016 ^b	17.212	<0.001	
				640.56 ^a	950.3		8.117 ^a	12.272		
				-1547.6 ^b	705.6		30.363 ^b	9.691		
				4.496 ^a	0.447		-0.0467 ^a	0.006		
				3.588 ^b	0.335		-0.0473 ^b	0.005		
				-8.296 ^a	6.553	0.003	0.330 ^a	0.085		
				-16.447 ^b	4.866		0.328 ^b	0.067		

^a Regression coefficients for eastern sites (see Fig. 1).

^b Regression coefficients for sites from the Midwest. Coefficients for western sites were set to zero in the analysis.

^c Statistical interaction between % Forest and Region.

^d Coefficient for quadratic term of the regression.




FIGURE 2. Relationship of forest coverage measured on different scales for the same study plots. Regression coefficients are given in Table 1. Regions noted in the legend correspond to those shown in Fig. 1.

were used to determine whether parasitization varied with local variation in landscape structure, whereas the among-site analyses show whether parasitization rates varied with differences in average forest coverage among widely separated regions.

RELATIONSHIPS AMONG FOREST METRICS

Measuring forest coverage at one scale predicts forest coverage at other scales, but the statistical relationships differed among geographical regions across the continent (Table 1). Low forest coverages, measured within 10-km radii of study plots, indicated even lower proportions of forest within 50 km in the Midwest than in either eastern or western landscapes (Fig. 2).

The relationship between forest coverage and most of the other measures of landscape structure that we compiled also differed across the United States. The only exception was mean size of forest patches; as the proportion of forest in the landscape increased, the mean size of forest patches increased consistently across the United States. Edge density was always highest at intermediate levels of forest coverage, and for a given amount of forest cover the amount of edge was highest in eastern forests and lowest in western forests (Fig. 3).

All other forest metrics varied linearly with increasing forest coverage, and the patterns were typically that landscapes with greater forest coverage also contained a larger number of forest types, larger size for the biggest forest patch, greater variation in patch size, and greater number of forest patches (Table 1). The one exception was for numbers of forest patches; in eastern and western sites greater forest coverage meant a larger number of patches, but in the Midwest greater forest coverage meant fewer




FIGURE 3. Relationship between forest coverage and edge density in different regions. Regions noted in the legend correspond to those shown in Fig. 1.

patches. This relationship at least partially resulted from different types of forest being treated as separate patches, in combination with the number of forest types remaining relatively unchanged with increased forest coverage in the Midwest (Table 1).

FOREST COVERAGE AND PRESENCE OF COWBIRD PARASITIZATION

We found no indication that local variation in forest coverage affected the presence or absence of cowbird parasitization on a given study plot. Twelve of 26 sites had plots both with and without detected cowbird parasitization. For each of these 12 sites, we determined whether increased forest coverage (measured within a 10-km radius of each study plot) resulted in a change in the probability of finding cowbird eggs or nestlings. No single regression was statistically significant (range $P = 0.11$ to $P = 0.99$), which may reflect the low statistical power resulting from the small number ($N = 5$ – 31) of data points in each analysis.

Further, we also found no indication of an effect even when results from individual analyses were combined in a meta-analysis. The meta-analysis used the regression coefficients from the individual logistic regressions as data points. Each regression coefficient was weighted by $1/$

SE of the coefficient, meaning that the coefficients that were estimated more precisely were given greater importance in the analysis. These weighted regression coefficients were used in a 1-sample t-test to determine if on average greater forest coverage lead to a greater or lower probability of detection of parasitization on study plots. The results of the meta-analysis were not significant ($P = 0.64$, $df = 11$, weighted mean regression coefficient = -0.0121 ± 0.025 SE), again indicating that when cowbirds were present in a region (i.e., at least one nest was parasitized on a study plot within a site) they did not avoid parasitizing nests on specific study plots in relation to local variation in forest coverage.

Sites with greater forest coverage tended to have a lower chance of cowbird parasitization, although the pattern only approached statistical significance (Table 2). For this analysis each of the separate study sites was treated as a single data point. The probability of detecting cowbird parasitization was not significantly affected by forest coverage on either scale of measurement (10-km or 50-km radii; Table 2). However, confidence limits around the regression coefficients showed a 95.3% probability that increased forest coverage within 10 km of study plots resulted in a decreased likelihood of cowbird parasitization at that site. Confidence limits also indicated a 92.7% probability that sites east of the Great Plains were less likely to have any cowbird parasitization.

FOREST COVERAGE AND THE RATE OF PARASITIZATION

Although we found some evidence that forest coverage affected the presence or absence of cowbird parasitization (above), we found more consistent evidence that the proportion of nests that were parasitized was related to forest coverage. Hosts were parasitized at lower rates when there was greater forest coverage, in comparisons both among study plots within the same study site and among widely separate study sites.

We examined the effects of local variation in forest coverage on the rate of brood parasitiza-

TABLE 2. VARIATION IN FOREST COVERAGE, AND PRESENCE OR ABSENCE OF COWBIRD PARASITIZATION. RESULTS ARE FROM LOGISTIC REGRESSIONS

Scale, forest coverage	% Forest			Region ^a		
	β	SE	P	β	SE	P
10-km radius	-0.055	0.033	0.09	1.55	1.07	0.15
50-km radius	-0.0088	0.019	0.64	1.33	0.93	0.64

^a Denotes whether sites were east or west of the Great Plains; results were similarly non-significant when data were divided into east, Midwest, and west. Regression coefficient is for data east of Great Plains; regression coefficient for west of Great Plains is zero.

TABLE 3. VARIATION IN FOREST COVERAGE AND THE PROPORTION OF NESTS PARASITIZED. RESULTS ARE FROM GENERALIZED LINEAR MODELS

Test	% Forest			Region			Interaction ^c			R ²
	β	SE	P	β	SE	P	β	SE	P	
Within Site (10 km)	-0.00099	0.0003	<0.001	-0.011 ^a -0.004 ^b	0.013 0.010	0.71				0.08
Among Site (10 km)	-0.00054	0.0003	0.001	0.235 ^a -0.004 ^b	0.070 0.038	0.003	-0.0020 ^a 0.0004 ^b	0.001 0.001	0.031	0.13
Among Site (50 km)	0.0014	0.0005	0.082	0.317 ^a 0.103 ^b	0.079 0.036	<0.001	-0.0046 ^a -0.0019 ^b	0.001 0.0007	<0.001	0.16

^a Regression coefficients for eastern sites (see Fig. 1).

^b Regression coefficients for sites from the Midwest. Coefficients for western sites were set to zero in the analysis.

^c Statistical interaction between % Forest and Region.

tion by comparing forest coverage and the rate of parasitization among study plots within the same study site. A 10% increase in forest coverage was predicted to result in a roughly 1% decrease in the proportion of nests that were parasitized (Table 3). This effect did not vary across the continent, either when sites were divided as east or west of the Great Plains, or east, Midwest, and west. We added forest coverage as a quadratic term to the statistical model to test for non-linear relationships between forest coverage and parasitization rate. No quadratic effect approached statistical significance, and we con-

clude that non-linearity in the relationship was minimal.

Both forest coverage and geographical location affected the rate of parasitization in comparisons among widely separate regions; additionally, the effect of forest coverage varied with the scale at which forest coverage was measured (Table 3). The typical pattern was as expected: the rate of parasitization was lower with increased forest coverage. However, an increase in parasitization with increased forest coverage was found from sites west of the Great Plains, but only when forest coverages were measured within 50-km radii of study plots (Table 3, Fig. 4). Confidence intervals around this regression coefficient indicate that there was only a 0.4% chance that the true pattern was for parasitization to be lower in areas of higher forest coverage. Regression models better fit the data when study sites were divided into 3 regions than when only categorized as being either east or west of the Great Plains. When forest coverage was added as a quadratic term to the models, the goodness of fit of regressions was identical or improved over the relationships given in Table 3. However, the qualitative patterns shown in Fig. 4 remained unchanged.

The magnitude of the effect of forest coverage on parasitization rate (i.e., slope of the regression) was greater when differences in forest coverage were measured among widely separated sites; however, this result was not robust. Within a given geographical region, the slopes of the within- and among-site regressions were within 2 SE (a roughly 95% confidence interval) of each other, with confidence intervals calculated assuming that the main and interaction effects in the among-site analyses were independent. To further test for differences within and among sites, we calculated separate regressions for each geographic region, both within and among sites; in this case, regression coefficients within a re-




FIGURE 4. Variation in the rate of parasitization of nests as a function of forest coverage. Different point and line styles correspond to the legends in Figs. 2, 3.

gion all overlapped in confidence limits of 1 SE (roughly 68% confidence limits).

DISCUSSION

Generally, we found that rates of parasitization were lower in areas of greater forest coverage (Fig. 4), as previously described (Robinson et al. 1995b, Donovan et al. in press, Tewksbury et al. 1998). This pattern was perhaps minimally due to increased forest coverage tending to result in a lower probability of any cowbird parasitization (Table 2). However, the clearer effect was a statistically significant decrease in the proportion of nests parasitized with increasing forest coverage (Table 3, Fig. 4). The relationship between greater forest coverage and lower rates of parasitization held regardless of whether we examined variation in forest coverage among plots within a local area or among widely separated study sites (Table 3). The presence of a relationship between forest coverage and parasitization rate, even within single study sites, suggests that behavioral decisions of individual cowbirds were at least partially responsible for the larger-scale variation in parasitization rate previously found (e.g., Robinson et al. 1995b).

However, the generalization that lower rates of parasitization are associated with a greater proportion of forest is not universal; greater rates of parasitization were found in areas of greater forest coverage in sites west of the Great Plains (Fig. 4, bottom panel) when forest coverage was measured within a 50-km radius of study plots. We suspect that traits other than landscape structure, such as human land-use practices (e.g., Tewksbury et al. 1998) may be responsible for our findings (Fig. 4, bottom panel). This result was not an artifact of a narrower range of forest coverages from the western sites (Fig. 4), nor did data from a single site create the pattern. Although landscape structure varied with changes in forest coverage across the continent (Table 1; Figs. 2, 3), we found no traits for which western forests differed qualitatively from both eastern and mid-western forests. Hence, we do not think that our results (Fig. 4) were due to differences in landscape structure east and west of the Great Plains. Neither are we aware of any substantial differences in the behavior and habitat requirements among the races of Brown-headed Cowbird (Lowther 1993). We also do not think that our results (Fig. 4) were an artifact of combining data from all host species into a single measure of parasitization, because an artifact of differing species composition would be manifested at both scales of measurement of forest coverage (top and bottom panels of Fig. 4). Finally, although cowbird abundance declined westward, away from the

center of the cowbird's range (Thompson et al. in press), the lower abundance of cowbirds in the west should simply lower the rate of parasitization but not cause a completely opposite response of parasitization rate to variation in forest coverage.

Our results indicate that the predicted rate of parasitization can be affected by the area over which forest coverage is measured (Table 3; Fig. 4, compare top and bottom panels). Previous work (Donovan et al. in press, Tewksbury et al. 1998) has shown that some scales of measuring forest coverage provide better predictions of the rate of parasitization than other scales. Our results indicate that not only the goodness of fit (measured as a correlation), but the actual predicted rates of parasitization (regression intercept and slope) were dependent on the scale at which forest coverage was measured (Table 3). However, we were not able to estimate the effects of variation in forest coverage on parasitization with great accuracy. The 95% confidence limits around the effect of forest coverage (10-km radius) in the eastern U.S. (Table 3) showed that the estimated effect could be somewhere within a 35-fold range of values! If this variation is due to insufficient sampling, the variation is probably sufficiently large to make the current estimates unsuitable for attempts to model (i.e., Hilborn and Mangel 1997, Starfield 1997) the demographic consequences to host species of modifying forest coverage. If the variation is biologically real, then our results indicate that relying on measurement of forest coverage to accurately predict rates of parasitization is probably not a fruitful endeavor.

The low accuracy of estimates is an indication that forest coverage explains only a small fraction of variation in the rate of parasitization (Table 3). As noted above, roughly 75% of all variance in the rate of parasitization was within local clusters of study plots, even though less than 23% of all variance in forest coverage was found among study plots within these same local clusters. While some of the within-site variance in the rate of parasitization was due to sampling error, variation in species composition of hosts among plots, and other random effects, we feel that the importance of non-forest landscape features (e.g., Tewksbury et al. 1998) should not be underestimated. One known reason is the need by female cowbirds to have both feeding sites and breeding areas in close proximity (Rothstein et al. 1984, Thompson 1994, Donovan et al. in press), and feeding sites are often human-related features of landscapes (Verner and Ritter 1983, Airola 1986).

The one consistent finding of this study was that lower rates of parasitization of host species

occurred with greater forest coverage within 10 km of a location, a result that held in spite of the different communities of hosts and their histories of exposure to cowbirds (Mayfield 1965) from east to west across the continent. This consistent result suggests that management for greater forest coverage even over relatively small spatial extents can decrease rates of brood parasitization. However, managers should realize that variation in forest coverage may show qualitatively different relationships with the rate of parasitization across the continent (Table 3, Fig. 4). The most extreme case was the sites from west of the Great Plains (Fig. 4), but we feel that data from additional sites are needed to substantiate the relationship between larger scale (50-km radius) forest coverage and rates of parasitization that we have found.

Nevertheless, it is clear that patterns found in one part of the continent should not be blindly extrapolated to other regions. Managers should also be aware that non-forest features such as feeding sites can play an important role in determining the rate of parasitization by cowbirds in a region (e.g., Airola 1986, Tewksbury et al. 1998, Thompson et al. in press). The effects of non-forest features should be carefully examined

if demographic modeling is to be a useful part of a research and management strategy (e.g., Starfield 1997), because the effects of forest coverage alone on rates of parasitization are variable enough that accurate predictions of parasitization rate were not possible, even with a data set as large as was available for this study.

ACKNOWLEDGMENTS

The long list of authors reflects the highly collaborative effort among principal investigators associated with the BBIRD project; each has contributed significantly to the preparation of this manuscript, including the provision of previously unpublished data. The results presented here are the product of the labor of well in excess of 100 field workers; thank-you all, anonymous though you must remain. The data on landscape structure were calculated by M. Thornton and W. Williams of the Wildlife Spatial Analysis Lab, at the University of Montana. Comments by D. Reinking, P. Vickery, T. Rich, members of the Montana Cooperative Wildlife Research Unit, and one anonymous reviewer improved this manuscript. Funding and logistical support for this work was provided by at least 27 federal, state, local, and private sources, with primary funding sources being: Biological Resources Division of the US Geological Survey, USDA Forest Service, US National Fish and Wildlife Foundation, US Fish & Wildlife Service, and the US National Science Foundation.

LITERATURE CITED

- AGENBROAD, L. D. 1978. Buffalo jump complexes in Owyhee County, Idaho. *Plains Anthropologist* 23: 213–221.
- AHLERS, D., AND L. WHITE. 1995. 1995 Southwestern Willow Flycatcher survey results: selected sites along the Rio Grande from Velarde, New Mexico, to the headwaters of Elephant Butte Reservoir. Technical Service Center, Bureau of Reclamation, Denver, CO.
- AIROLA, D. A. 1986. Brown-headed Cowbird parasitism and habitat disturbance in the Sierra Nevada. *Journal of Wildlife Management* 50:571–575.
- ALLEN, J. A. 1877. History of the American bison, *Bison americanus*. Annual Report of the U.S. Geological and Geographical Surveys of the Territories 9:443–587.
- ALVAREZ, F. 1993. Proximity of trees facilitates parasitism by Cuckoos *Cuculus canorus* on Rufous Warblers *Cercotrichas galactotes*. *Ibis* 135:331.
- ALVERSON, W. S., W. KUHLMANN, AND D. M. WALLER. 1994. Wild forests: conservation biology and public policy. Island Press, Washington, D.C.
- ANDERSON, B. W., AND R. D. OHMART. 1986. Vegetation. Pp. 639–659 in A. Y. Cooperrider, R. J. Boyd, and H. R. Stuart (editors). Inventory and monitoring of wildlife habitat. U.S. Bureau of Land Management Service Center, Denver, CO.
- ANDERSON, W. L., AND R. W. STORER. 1976. Factors influencing Kirtland's Warbler nesting success. *Jack-Pine Warbler* 54:105–115.
- ANKNEY, C. D., AND D. M. SCOTT. 1980. Changes in nutrient reserves and diet of breeding Brown-headed Cowbirds. *Auk* 97:684–696.
- ANONYMOUS. No date. Forest land distribution data for the United States [online]. Available: <http://www.epa.gov/docs/grd/forestinventory/> [1998, September 13].
- ANTHONY, A. W. 1893. Birds of San Pedro Martir, Lower California. *Zoe* 4:228–247.
- ANTHONY, A. W. 1895. Birds of San Fernando, Lower California. *Auk* 12:134–143.
- ARCESE, P., AND J. N. M. SMITH. 1988. Effects of population density and supplemental food on reproduction in Song Sparrows. *Journal of Animal Ecology* 57:119–136.
- ARCESE, P., AND J. N. M. SMITH. In press. Impacts of nest depredation and brood parasitism on the productivity of North American passerines. In N. Adams and R. Slotow (editors). Proceedings of the 22nd International Ornithological Congress. University of Natal, Durban, South Africa.
- ARCESE, P., J. N. M. SMITH, AND M. I. HATCH. 1996. Nest predation by cowbirds and its consequences for passerine demography. Proceedings of the National Academy Science (USA) 93:4608–4611.
- ARCESE, P., J. N. M. SMITH, W. M. HOCHACHKA, C. M. ROGERS, AND D. LUDWIG. 1992. Stability, regulation and the determination of abundance in an insular Song Sparrow population. *Ecology* 73:805–822.
- ARMSTRONG, D. M. 1972. Distribution of mammals in Colorado. Monograph of the University of Kansas Museum of Natural History 3:1–415.
- ARNOLD, K. A. 1983. Annual adult survival rates for Brown-headed Cowbirds wintering in southeast Texas. *Wilson Bulletin* 95:150–153.
- ARNQVIST, G., AND D. WOOSTER. 1995. Meta-analysis: synthesizing research findings in ecology and evolution. *Trends in Evolution and Ecology* 10:236–240.
- ASKINS, R. A. 1995. Hostile landscapes and the decline of migratory birds. *Science* 67:1956–1957.
- ASKINS, R. A., J. F. LYNCH, AND R. GREENBERG. 1990. Population declines in migratory birds in eastern North America. *Current Ornithology* 7:1–57.
- AVERILL, A. 1996. Brown-headed cowbird parasitism of neotropical migrant songbirds in riparian areas along the lower Colorado River. M.S. thesis. University of Arizona, Tucson, AZ.
- BAICICH, P. J., AND C. J. O. HARRISON. 1997. A guide to the nests, eggs, and nestlings of North American birds, 2nd ed. Academic Press, San Diego, CA.
- BAILEY, F. M. 1923. Birds recorded from the Santa Rita Mountains in southern Arizona. *Pacific Coast Avifauna* No. 15. Cooper Ornithological Club, Berkeley, CA.
- BAIRD, S. F. 1858. Explorations and surveys for a railroad route from the Mississippi River to the Pacific Ocean. Volume IX: Birds. Part II—General report upon the zoology of the several Pacific railroad routes. Beverley Tucker (printer), Washington, D.C.
- BALAZS, I. J., M. BAIRD, M. CLYNE, AND E. MEADE. 1989. Human population genetic studies of five hypervariable DNA loci. *American Journal Human Genetics* 44:182–190.
- BALAZS, I. J., A. CHIMERA, D. EISENBERG, D. ENDEAN, R. GILES, J. LATELY, S. MAGUIRE, B. SCHALL, A. TURCK, AND R. MCKEE. 1990. Accuracy, precision, and site-to-site reproducibility in analysis of DNA polymorphisms for identity testing. Pp. 54–56 in H. F. Polesky and W. R. Mayr (editors). *Advances in forensic haemogenetics* 3. Springer-Verlag, Berlin, Germany.
- BANKS, A. J. 1997. Variation among host species in probability of parasitism by Brown-headed Cowbirds: the role of everyday host activity. M. S. thesis. University of Montana, Missoula, MT.
- BARBER D. R., AND T. E. MARTIN. 1997. Influence of alternate host densities on Brown-headed Cowbird parasitism rates in Black-capped Vireos. *Condor* 99: 595–604.
- BARLOW, J. C. 1962. Natural history of the Bell's Vireo, *Vireo bellii* Audubon. University of Kansas Publications of the Museum of Natural History 12:241–296.
- BEAL, F. E. L. 1900. Food of the bobolink, blackbirds, and grackles. U.S. Department of Agriculture, Biological Survey Bulletin 13.
- BEEDEY, E. C., AND S. L. GRANHOLM. 1985. *Discovering Sierra Birds*. Yosemite Natural History Association.
- BEEZLEY, J. A., AND J. P. RIEGER. 1987. Least Bell's Vireo management by cowbird trapping. *Western Birds*:55–61.
- BEHLE, W. H., E. D. SORENSON, AND C. M. WHITE. 1985. *Utah Birds: a revised checklist*. Utah Museum of Natural History, University of Utah, Salt Lake City, UT.
- BEIDLEMAN, R. G. 1955. An altitudinal record for bison

- in northern Colorado. *Journal of Mammalogy* 36: 470-471.
- BENEDICT, J. B. 1993. Excavations at Bode's Draw, a women's work area in the mountains near Estes Park, Colorado. Research Report, Center for Mountain Archeology 6:1-42.
- BENSON R. H., AND K. L. P. BENSON. 1990. Estimated size of Black-capped Vireo populations in Coahuila, Mexico. *Condor* 92: 777-779.
- BENT, A. C. 1958. Life histories of North American blackbirds, orioles, tanagers, and allies. U.S. National Museum Bulletin No. 211, Washington, D.C.
- BENT, A. C. 1968. Life histories of North American cardinals, grosbeaks, buntings, towhees, finches, sparrows, and allies. U.S. National Museum Bulletin No. 237 (parts 1 and 2), Washington, D.C.
- BERGER, A. J. 1951. The cowbird and certain host species in Michigan. *Wilson Bulletin* 63:26-34.
- BERGIN, T. M., L. B. BEST, AND K. E. FREEMARK. 1997. An experimental study of predation on artificial nests in roadsides adjacent to agricultural habitats in Iowa. *Wilson Bulletin* 109:437-448.
- BERGTOLD, W. H. 1929. Bison in Colorado. *Journal of Mammalogy* 10:107.
- BEST, L. B. 1978. Field Sparrow reproductive success and nesting ecology. *Auk* 95:9-22.
- BEST, L. B. 1979. Effects of fire on a Field Sparrow population. *American Midland Naturalist* 101:434-442.
- BEST, L. B., AND D. F. STAUFFER. 1980. Factors affecting nesting success in riparian bird communities. *Condor* 82:149-158.
- BEYER, D. E. 1987. Population and habitat management of elk in Michigan. Ph.D. dissertation. Michigan State University, East Lansing, MI.
- BIERMANN, G. C., W. B. MCGILLIVRAY, AND K. E. NORDIN. 1987. The effect of cowbird parasitism on Brewer's Sparrow productivity in Alberta. *Journal of Field Ornithology* 58: 350-354.
- BLINCOE, B. J. 1935. A cowbird removes a robin's egg. *Wilson Bulletin* 47:158.
- BLONDEL, J., C. FERY, AND B. FROCHOT. 1981. Point counts with unlimited distance. *Studies in Avian Biology* 6:414-420.
- BOCK, C. E., V. A. SAAB, T. D. RICH, AND D. S. DOBKIN. 1993. Effects of livestock grazing on neotropical migratory landbirds in Western North America. Pp. 296-309 in D. M. Finch and P. W. Stangel (editors). Status and management of neotropical migratory birds. USDA Forest Service Gen. Tech. Rep. RM-229. USDA Forest Service, Rocky Mountain Forest and Range Experimental Station, Ft. Collins, CO.
- BÖHNING-GAESE, K., M. L. TAPER, AND J. H. BROWN. 1993. Are declines in North American insectivorous songbirds due to causes on the breeding range? *Conservation Biology* 7:76-86.
- BONHAM, C. D. 1989. Measurements for terrestrial vegetation. John Wiley and Sons, New York, NY.
- BRADEN, G. T., R. L. MCKERNAN, AND S. M. POWELL. 1997a. Association of within-territory vegetation characteristics and fitness components of California Gnatcatchers. *Auk* 114:601-609.
- BRADEN, G. T., R. L. MCKERNAN, AND S. M. POWELL. 1997b. Effects of nest parasitism by the Brown-headed Cowbird on nesting success of the California Gnatcatcher. *Condor* 99:858-865.
- BRAGG, T. B. 1995. The physical environment of Great Plains grasslands. Pp. 49-81 in A. Joern and K. H. Keeler (editors). The changing prairie: North American grasslands. Oxford University Press, New York, NY.
- BRAULT, S., AND H. CASWELL. 1993. Pod-specific demography of killer whales (*Orcinus orca*). *Ecology* 74:1444-1454.
- BRAWN, J. D., AND S. K. ROBINSON. 1996. Source-sink dynamics may complicate the interpretation of long-term census data. *Ecology* 77:3-11.
- BRAY, O. E., J. W. DEGRAZIO, J. L. GUARINO, AND R. G. STREETER. 1974. Recoveries of Brown-headed Cowbirds banded at Sand Lake, South Dakota. *Inland Bird Banding News* 46:204-209.
- BRISKIE, J. V., AND S. G. SEALY. 1990. Evolution of short incubation periods in the parasitic cowbirds, *Molothrus* spp. *Auk* 107:789-794.
- BRISKIE, J. V., S. G. SEALY, AND K. A. HOBSON. 1990. Differential parasitism of Least Flycatchers and Yellow Warblers by the Brown-headed Cowbird. *Behavioral Ecology and Sociobiology* 27:403-410.
- BRISKIE, J. V., S. G. SEALY, AND K. A. HOBSON. 1992. Behavioral defenses against brood parasitism in sympatric and allopatric host populations. *Evolution* 46:334-340.
- BRITTINGHAM, M. C., AND S. A. TEMPLE. 1983. Have cowbirds caused forest songbirds to decline? *BioScience* 33:31-35.
- BRITTINGHAM, M. C., AND S. A. TEMPLE. 1996. Vegetation around parasitized and non-parasitized nests within a deciduous forest. *Journal of Field Ornithology* 67:406-413.
- BROCKNER, W. W. 1984. Brown-headed Cowbird parasitizing Mountain Chickadee nest. C. F. O. [Colorado Field Ornithologists] *Journal* 18:109-110.
- BROOKER, M. G., AND L. C. BROOKER. 1989a. The comparative breeding behaviour of two sympatric cuckoos, Horsfield's Bronze-Cuckoo *Chrysococcyx basalis* and the Shining Bronze-Cuckoo *C. lucidus*, in Western Australia: a new model for the evolution of egg morphology and host specificity in avian brood parasites. *Ibis* 131:528-547.
- BROOKER, M. G., AND L. C. BROOKER. 1989b. Cuckoo hosts in Australia. *Australian Zoological Reviews* 2: 1-67.
- BROOKER, M. G., AND L. C. BROOKER. 1992. Evidence for individual female host specificity in two Australian Bronze-Cuckoos (*Chrysococcyx* spp.). *Australian Journal of Zoology* 40:485-493.
- BROWN, B. T. 1988. Breeding ecology of a Willow Flycatcher population in Grand Canyon, Arizona. *Western Birds* 19: 25-33.
- BROWN, B. T. 1994. Rates of brood parasitism by Brown-headed Cowbirds on riparian passerines in Arizona. *Journal of Field Ornithology* 65: 160-168.
- BROWN, H. 1903. Arizona bird notes. *Auk* 20:43-50.
- BROWN, J. L. 1975. The evolution of behavior. W.W. Norton and Co., New York, NY.
- BROWNING, M. R. 1993. Comments on the taxonomy of *Empidonax traillii* (Willow Flycatcher). *Western Birds* 24:241-257.
- BUCKLAND, S. T., D. R. ANDERSON, K. P. BURNHAM,

- AND J. L. LAAKE. 1996. Distance sampling. Chapman and Hall, New York, NY.
- BUECH, R. R. 1982. Nesting ecology and cowbird parasitism of Clay-colored, Chipping, and Field sparrows in a Christmas tree plantation. *Journal of Field Ornithology* 53:363–369.
- BURGER, L. D., L. W. BURGER, JR., AND J. FAABORG. 1994. Effects of prairie fragmentation on predation on artificial nests. *Journal of Wildlife Management* 58:249–254.
- BURGHAM, M. C. J., AND J. PICMAN. 1989. Effect of Brown-headed Cowbirds on the evolution of Yellow Warbler anti-parasite strategies. *Animal Behavior* 38:298–308.
- BURGMAN, M. A., S. FERSON, AND H. R. AKÇAKAYA. 1993. Risk assessment in conservation biology. Chapman and Hall, New York, NY.
- BURGMAN, M. A., AND V. A. GERARD. 1990. A stage-structured, stochastic population model for the giant kelp *Macrocystis pyrifera*. *Marine Biology* 105:15–23.
- BURHANS, D. E. 1997. Habitat and microhabitat features associated with cowbird parasitism in two forest edge cowbird hosts. *Condor* 99:866–872.
- BURHANS, D. E. In press. Dawn nest arrivals in cowbird hosts: their role in aggression, cowbird recognition, and response to parasitization. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- BURKE, T., N. B. DAVIES, M. W. BRUFORD, AND B. J. HATCHWELL. 1989. Parental care and mating behavior of polyandrous dunnocks (*Prunella modularis*) related to paternity by DNA fingerprinting. *Nature* 338:249–251.
- BURNHAM, K. P., D. R. ANDERSON, AND J. L. LAAKE. 1980. Estimation of density from line transect sampling of biological populations. *Wildlife Monographs* 72:1–202.
- BUTCHER, G. S. (CHAIR), AND CORRESPONDENTS. 1992. Needs assessment: monitoring neotropical migratory birds. *Partners in Flight*. Cornell University, Ithaca, NY.
- BUTLER, R. B. 1978. Bison hunting in the desert west before 1800: the paleo-ecological potential and the archaeological reality. *Plains Anthropologist* 23:107–112.
- CAMPBELL, L. 1995. Endangered and threatened animals of Texas—their life history and management. Texas Parks and Wildlife Press, Austin, TX.
- CANNON, R. W. 1979. Lesser Prairie Chicken responses to range fires at the booming ground. *Wildlife Society Bulletin* 7:44–46.
- CAROTHERS, S. W. 1974. Population structure and organization of southwestern riparian birds. *American Zoologist* 14:97–108.
- CARTER, M. D. 1986. The parasitic behavior of the Bronzed Cowbird in south Texas. *Condor* 88:11–25.
- CASWELL, H. 1989. Matrix population models. Sinauer Associates, Inc., Sunderland, MA.
- CAVALCANTI, R. B. 1981. Nest desertion: theory and tests of its adaptive significance in birds. Ph.D. dissertation. McGill University, Montreal, PQ.
- CHACE, J. F. 1995. The factors affecting the reproductive success of the Solitary Vireo (*Vireo solitarius plumbeus*) in Colorado. M.A. thesis. University of Colorado, Boulder, CO.
- CHACE, J. F. AND A. CRUZ. 1996. Knowledge of the Colorado host relations of the parasitic Brown-headed Cowbird. C. F. O. [Colorado Field Ornithologists] *Journal* 30:67–81.
- CHACE, J. F. AND A. CRUZ. 1998. Range of the Brown-headed Cowbird in Colorado—past and present. *Great Basin Naturalist* 58:245–249.
- CHACE, J. F., A. CRUZ, AND R. E. MARVIL. In press. Reproductive interactions of the Brown-headed Cowbird and Solitary Vireo in Colorado. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- CHASKO, G. G., AND E. J. GATES. 1982. Avian habitat suitability along a transmission-line corridor in an oak-hickory forest region. *Wildlife Monographs* 82:1–41.
- CHRISTMAN, G. M. 1971. The mountain bison. *American West* 8:44–47.
- CHRISTY, B. H. 1925. Summer birds of Huron Mountain, Michigan. *Wilson Bulletin* 37:208–215.
- CLARK, K. L., AND R. J. ROBERTSON. 1979. Spatial and temporal multi-species nesting aggregations in birds as anti-parasite and anti-predator defenses. *Behavioral Ecology and Sociobiology* 5:359–371.
- CLARK, K. L., AND R. J. ROBERTSON. 1981. Cowbird parasitism and evolution of anti-parasite strategies in the Yellow Warbler. *Wilson Bulletin* 93:249–258.
- CLOTFELTER, E. D. 1998a. Impact of Brown-headed Cowbird brood parasitism on Red-winged Blackbirds and factors influencing patterns of parasitism. Ph.D. dissertation. University of Wisconsin, Madison, WI.
- CLOTFELTER, E. D. 1998b. What cues do Brown-headed Cowbirds use to locate Red-winged Blackbird host nests? *Animal Behaviour* 55:1181–1189.
- CLOTFELTER, E. D., AND T. BRUSH. 1995. Unusual parasitism by the Bronzed Cowbird. *Condor* 97:814–815.
- CODY, M. L. 1985. Habitat selection in grassland and open-country birds. Pp. 191–226 in M. L. Cody (editor). *Habitat selection in birds*. Academic Press, Orlando, FL.
- COHEN, J. 1988. Statistical power analysis for the behavioral sciences, 2nd ed. Academic Press, New York, NY.
- COKER, D. R., AND D. E. CAPEN. 1995. Landscape-level habitat use by Brown-headed Cowbirds in Vermont. *Journal of Wildlife Management* 59:631–637.
- COKER, D. R., AND D. E. CAPEN. In press. Distribution and habitat associations of Brown-headed Cowbirds in the Green Mountains of Vermont. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- COLLAR, N. J., M. J. CROSBY, AND A. J. STATTERSFIELD. 1994. *Birds to watch 2: the world list of threatened birds*. BirdLife International, Cambridge, U.K.
- COLLAR, N. J., L. P. GONZAGA, N. KRABBE, A. MADRONO NIETO, L. G. NARANJO, T. A. PARKER, III, AND

- D. C. WEGE. 1992. Threatened birds of the Americas: the ICBP/IUCN Red Data Book. International Council for Bird Preservation, Cambridge, U.K.
- COLLINS, C. T., L. R. HAYS, A. DAVENPORT, AND D. WILLICK. 1988. Status and management of Least Bell's Vireo within the Prado Basin, California. Report submitted to State of California, Dept. of Transportation, District 8, San Bernardino, CA.
- COOK, T. L., J. A. KOLOSZAR, M. D. GOERING, AND L. L. SANCHEZ. 1998. The spatial and temporal response of Brown-headed Cowbirds (*Molothrus ater*) to cattle removal. Pp. 76–96 in *The Nature Conservancy, Summary of 1997 Research Activities*. Texas Conservation Data Center, The Nature Conservancy, Fort Hood, Texas.
- COOKE, W. W. 1897. The birds of Colorado. Bulletin of the State Agriculture Collection 37 (Technical Series) 2:1–224.
- COOPER, C. A. 1996. Summary of 1995 Surveys for Willow Flycatchers in New Mexico. Contract #96–516.51. New Mexico Department of Game and Fish, Santa Fe, NM.
- COOPER, C. A. 1997. Summary of 1996 Surveys for Willow Flycatchers in New Mexico. Contract #96–516.81. New Mexico Department of Game and Fish, Santa Fe, NM.
- COOPER, J. G. 1861. New California animals. Proceedings of the California Academy of Science 2:118–123.
- COX, G. W. 1990. Laboratory manual of general ecology, 6th ed. Wm. C. Brown Publishers, Dubuque, IA.
- CROUSE, D. T., L. B. CROWDER, AND H. CASWELL. 1987. A stage-based population model for loggerhead sea turtles and implication for conservation. *Ecology* 68: 1412–1423.
- CRUZ, A., W. POST, J. W. WILEY, C. P. ORTEGA, T. K. NAKAMURA, AND J. W. PRATHER. 1998. Potential impacts of cowbird range expansion in Florida. Pp. 313–336 in S. I. Rothstein and S. K. Robinson (editors). *Parasitic birds and their hosts*. Oxford University Press, Oxford, U.K.
- CUNNINGHAM, R. 1993. The cowbird peril: a resources management problem and an interpretive story. Internal report of the National Park Service, Western Regional Office, San Francisco, CA.
- CURSON, D. R. 1996. Nest predation and brood parasitism of passerine birds in pinyon-juniper woodland in northeast New Mexico. M.S. thesis. University of Wisconsin, Madison, WI.
- DARLEY, J. A. 1971. Sex ratio and mortality in the Brown-headed Cowbird. *Auk* 88:560–566.
- DARLEY, J. A. 1983. Territorial behavior of the female Brown-headed Cowbird (*Molothrus ater*). *Canadian Journal of Zoology* 61:65–69.
- DAUBENMIRE, R. F. 1968. Ecology of fire in grasslands. *Advances in Ecological Research* 5:209–266.
- DAUBENMIRE, R. F. 1988. Steppe vegetation of Washington. Washington State University, Pullman, WA.
- DAVIES, N. B., AND M. DE L. BROOKE. 1988. Cuckoos versus reed warblers: adaptations and counteradaptations. *Animal Behaviour* 36: 262–284.
- DAVIS, S. K., AND S. G. SEALY. In press. Cowbird parasitization and predation in grassland fragments of southwestern Manitoba. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- DAVIS, W. B. 1935. Mammals of the Ross Expedition (1824) in Idaho. *Murrelet* 16:7–10.
- DAVISON, W. 1998. Starvation and nestling ejection as sources of mortality in parasitized Lazuli Bunting nests. *Great Basin Naturalist* 58:285–288.
- DECAPITA, M. E. In press. Brown-headed Cowbird control on Kirtland's Warbler nesting areas in Michigan, 1972–1995. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- DELLASALA, D. A. 1985. The Yellow Warbler in southeastern Michigan: factors affecting its productivity. *Jack-Pine Warbler* 63:52–60.
- DELLASALA, D. A., D. M. OLSON, S. E. BARTH, S. L. CRANE, AND S. A. PRIMM. 1995. Forest health: moving beyond rhetoric to restore healthy landscapes in the inland Northwest. *Wildlife Society Bulletin* 23: 346–356.
- DESANTE, D. F., AND T. L. GEORGE. 1994. Population trends in the landbirds of western North America. *Studies in Avian Biology* 15:173–190.
- DHONDT, A. A. 1979. Summer dispersal and survival of juvenile Great Tits in southern Sweden. *Oecologia* 42:139–157.
- DOAK, D. F., AND L. S. MILLS. 1994. A useful role for theory in conservation. *Ecology* 75:615–626.
- DOBKIN, D. S. 1994. Conservation and management of neotropical migrant landbirds in northern Rockies and Great Plains. University of Idaho Press, Moscow, ID.
- DOBKIN, D. S., AND B. A. WILCOX. 1986. Analysis of natural forest fragments: riparian birds in the Toyabe Mountains, Nevada. Pp. 293–299 in J. Verner, M. L. Morrison, and C. J. Ralph (editors). *Wildlife 2000: modeling habitat relationships of terrestrial vertebrates*. University of Wisconsin Press, Madison, WI.
- DOBLER, F. C., J. EBY, C. PERRY, S. RICHARDSON, AND M. VANDER HAEGEN. 1996. Status of Washington's shrubsteppe ecosystem: extent, ownership, and wildlife/vegetation relationships. Research Report. Washington Department of Fish and Wildlife, Olympia, WA.
- DOLBEER, R. A. 1982. Migratory patterns for age and sex classes of blackbirds and starlings. *Journal of Field Ornithology* 53:28–46.
- DONOVAN, T. M., P. W. JONES, E. M. ANNAND, AND F. R. THOMPSON, III. 1997. Variation in local-scale edge effects: mechanisms and landscape context. *Ecology* 78:2064–2075.
- DONOVAN, T. M., R. H. LAMBERSON, F. R. THOMPSON, III, AND J. FAABORG. 1995a. Modeling the effects of habitat fragmentation on source and sink demography of neotropical migrant birds. *Conservation Biology* 9:1396–1407.
- DONOVAN, T. M., F. R. THOMPSON, III, AND J. FAABORG. In press. Ecological trade-offs and the influence of scale on Brown-headed Cowbird distribution. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and*

- management of cowbirds and their hosts. University of Texas Press, Austin, TX.
- DONOVAN, T. M., F. R. THOMPSON, III, J. FAABORG, AND J. R. PROBST. 1995b. Reproductive success of migratory birds in habitat sources and sinks. *Conservation Biology* 9:1380-1395.
- DRAPER, N. R., AND H. SMITH. 1981. Applied regression analysis. John Wiley & Sons, New York, NY.
- DRAWE, D. L., A. D. CHAMRAD, AND T. W. BOX. 1978. Plant communities of the Welder Wildlife Refuge, 2nd ed. Welder Wildlife Foundation, Sinton, TX.
- DRENT, R. H., AND S. DAAN. 1980. The prudent parent: energetic adjustments in avian breeding. *Ardea* 68: 225-252.
- DREW, F. M. 1885. On the vertical range of birds in Colorado. *Auk* 2:16.
- DRILLING, N. E., AND C. F. THOMPSON. 1988. Natal and breeding dispersal in House Wrens (*Troglodytes aedon*). *Auk* 105:480-491.
- DUFTY, A. M., JR. 1982a. Movements and activities of radio-tracked Brown-headed Cowbirds. *Auk* 99: 316-327.
- DUFTY, A. M., JR. 1982b. Responses of Brown-headed Cowbirds to simulated conspecific intruders. *Animal Behavior* 30:1043-1052.
- DUFTY, A. M., JR. 1983. Variation in the egg markings of the Brown-headed Cowbird. *Condor* 85:109-111.
- DUNN, J. D., AND K. L. GARRETT. 1997. A field guide to the warblers of North America. Houghton Mifflin, Boston, MA.
- DUNNING, J. B., JR. 1993. CRC handbook of avian body masses. CRC Press, Boca Raton, FL.
- EATON, S. W. 1958. A life history study of the Louisiana Waterthrush. *Wilson Bulletin* 70:211-236.
- EDWARDS, T. C., JR., E. T. DESHLER, D. FOSTER, AND G. G. MOISEN. 1996. Adequacy of wildlife habitat relation models for estimating spatial distribution of terrestrial vertebrates. *Conservation Biology* 10: 263-270.
- EHRlich, P. R., D. S. DOBKIN, AND D. WHEYE. 1988. The birder's handbook: a field guide to the natural history of North American Birds. Simon and Shuster, New York, NY.
- EHRlich, P. R., D. S. DOBKIN, AND D. WHEYE. 1992. Birds in jeopardy. Stanford University Press, Stanford, CA.
- ELLIOT, P. F. 1978. Cowbird parasitism in the Kansas tallgrass prairie. *Auk* 95:161-167.
- EVANS, D. R., AND J. E. GATES. 1997. Cowbird selection of breeding areas: the role of habitat and bird species abundance. *Wilson Bulletin* 109:470-480.
- EVANS, H. E. 1997. The natural history of the Long expedition to the Rocky Mountains 1819-1820. Oxford University Press, New York, NY.
- FANKHAUSER, D. P. 1971. Survival rates of blackbirds and starlings. *Bird-Banding* 42:36-42.
- FERNANDEZ-DUQUE, E., AND C. VALEGGIA. 1994. Meta-analysis: a valuable tool in conservation research. *Conservation Biology* 8:555-561.
- FERSON, S. 1994. RAMAS/stage: generalized stage-based modeling for population dynamics (version 1.4). Applied Biomathematics, Setauket, NY.
- FERSON, S., AND H. R. AKÇAKAYA. 1991. RAMAS/age: modeling fluctuations in age-structured populations. Exeter Software, Setauket, NY.
- FIGGINS, J. D. 1933. The bison of the western area of the Mississippi Basin. *Proceedings of the Denver Museum of Natural History* 12:16-33.
- FINCH, D. M. 1983. Brood parasitism of the Abert's Towhee: timing, frequency, and effects. *Condor* 85: 355-359.
- FINCH, D. M. 1991. Population ecology, habitat requirements, and conservation of neotropical migratory birds. USDA Forest Service Gen. Tech. Rep. RM-205. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- FISHER, A. K. 1893a. Report on the ornithology of the Death Valley Expedition of 1891. *North American Fauna* 7:7-158.
- FISHER, A. K. 1893b. The Death Valley expedition: a biological survey of parts of California, Nevada, Arizona, and Utah. U.S. Government Printing Office, Washington, D.C.
- FLEISCHER, R. C. 1985. A new technique to identify and assess the dispersion of eggs of individual brood parasites. *Behavioral Ecology and Sociobiology* 17: 91-99.
- FLEISCHER, R. C., AND S. I. ROTHSTEIN. 1988. Known secondary contact and rapid gene flow among subspecies and dialects in the Brown-headed Cowbird. *Evolution* 42:1146-1158.
- FLEISCHER, R. C., S. I. ROTHSTEIN, AND L. S. MILLER. 1991. Mitochondrial DNA variation indicates gene flow across a zone of known secondary contact between two subspecies of the Brown-headed Cowbird. *Condor* 93:185-189.
- FLEISCHER, R. C., A. P. SMYTH, AND S. I. ROTHSTEIN. 1987. Temporal and age-related variation in the laying rate of the parasitic Brown-headed Cowbird in the eastern Sierra Nevada, California. *Canadian Journal of Zoology* 65: 2724-2730.
- FLESHMAN, C., AND D. S. KAUFMAN. 1984. The South Fork (Kern River) Wildlife Area: will the commitment be forgotten? Pp. 482-494 in R. E. Warner and K. M. Hendrix (editors). *California riparian systems: ecology, conservation and productive management*. University of California Press, Berkeley, CA.
- FONDELL, T. F. 1997. Nest density and nest success of ground-nesting grassland birds relative to grazing in western Montana. M. S. thesis. University of Montana, Missoula, MT.
- FORD, R., Z. MA, S. BARSNESS, AND R. REDMOND. 1997. Rule-based aggregation of classified imagery. *Proceedings of the 1997 ACSM/ASPRS Convention and Exposition* 3:115-123.
- FRAGA, R. M. 1985. Host-parasite interactions between Chalk-browed Mockingbirds and Shiny Cowbirds. *Ornithological Monographs* 36:829-844.
- FRANZREB, K. E. 1989a. Ecology and conservation of the endangered least Bell's vireo. U.S. Fish and Wildlife Service, Biological Report 89(1). U.S. Fish and Wildlife Service Publications Unit, Washington, D.C.
- FRANZREB, K. E. 1989b. Ecology and conservation of the Least Bell's Vireo (*Vireo bellii pusillus*) in California. *Western Birds* 18:43-49.
- FRANZREB, K. E. 1990. An analysis of options for reintroducing a migratory, native passerine, the endangered Least Bell's Vireo *Vireo bellii pusillus* in the

- Central Valley, California. *Biological Conservation* 53:105-123.
- FREEMAN, S. D., F. FORI, AND S. ROHWER. 1990. Red-winged Blackbirds and Brown-headed Cowbirds: some aspects of a host-parasite relationship. *Condor* 92:336-340.
- FRIEDMANN, H. 1928. The origin of host specificity in the parasitic habit in the Cuculidae. *Auk* 45:33-38.
- FRIEDMANN, H. 1929. The cowbirds: a study in the biology of social parasitism. Charles C. Thomas, Springfield, IL.
- FRIEDMANN, H. 1963. Host relations of the parasitic cowbirds. *U.S. National Museum Bulletin* 233:1-276.
- FRIEDMANN, H. 1967a. Alloxyenia in three sympatric African species of *Cuculus*. *Proceedings of the United States National Museum* 124:1-14.
- FRIEDMANN, H. 1967b. Evolutionary terms for parasitic species. *Systematic Zoology* 16:175.
- FRIEDMANN, H. 1971. Further information on the host relations of the parasitic cowbirds. *Auk* 88:239-255.
- FRIEDMANN, H., AND L. F. KIFF. 1985. The parasitic cowbirds and their hosts. *Proceedings of the Western Foundation of Vertebrate Zoology* 2: 226-304.
- FRIEDMANN, H., L. F. KIFF, AND S. I. ROTHSTEIN. 1977. A further contribution to the knowledge of the host relations of the parasitic cowbirds. *Smithsonian Contributions in Zoology* 235:1-75.
- FRYXELL, F. M. 1926. A new high altitude limit for the American bison. *Journal of Mammalogy* 7:102-109.
- FRYXELL, F. M. 1928. The former range of the American bison in the Rocky Mountains. *Journal of Mammalogy* 9:129-139.
- FULLER, W. A. 1962. The biology and management of the bison of Wood Buffalo National Park. *Wildlife Management Bulletin, Canadian Wildlife Service Series 1*, 16:1-52.
- GAILLARD, J., M. FESTA-BIANCHET, AND N. G. YOCOZ. 1998. Population dynamics of large herbivores: variable recruitment with constant adult survival. *Trends in Ecology and Evolution* 13:58-63.
- GAINES, D. 1974. A new look at the nesting riparian avifauna of the Sacramento Valley, California. *Western Birds* 5:61-80.
- GAINES, D. 1977. Birds of the Yosemite Sierra: a distributional survey. *California Syllabus*, Oakland, CA.
- GAINES, D. 1988. Birds of Yosemite and the East Slope. Artemisia Press, Lee Vining, CA.
- GALBRAITH, D. A., P. T. BOAG, H. L. GIBBS, AND B. N. WHITE. 1991. Sizing bands on autoradiograms: a study of precision for scoring DNA fingerprints. *Electrophoresis* 12: 210-220.
- GARDALI, T., A. M. KING, AND G. R. GEUPEL. 1998. Cowbird parasitism and nest success of the Lazuli Bunting in the Sacramento Valley. *Western Birds* 29: 174-179.
- GARRETT, K. L., AND J. D. DUNN. 1981. Birds of southern California: status and distribution. The Artesian Press, Los Angeles, CA.
- GATES, J. E., AND N. R. GIFFEN. 1991. Neotropical migrant birds and edge effects at a forest-stream ecotone. *Wilson Bulletin* 103:204-217.
- GATES, J. E., AND L. W. GYSEL. 1978. Avian nest dispersion and fledging success in field-forest ecotones. *Ecology* 59:871-883.
- GAVIN, T. A. 1991. Why ask "why": the importance of evolutionary biology in wildlife science. *Journal of Wildlife Management* 55:760-766.
- GEORGES, M., A. S. LEQUARRE, M. CASTILLI, R. HANSET, AND G. LESSART. 1988. DNA fingerprinting in domestic animals using four different microsatellite probes. *Cytogenetic and Cellular Genetics* 47: 127-131.
- GILL, F. B. 1990. *Ornithology*. W. H. Freeman and Co., New York, NY.
- GILL, S. A., P. M. GRIEF, L. M. STAIB, AND S. G. SEALY. 1997. Does nest defense deter or facilitate cowbird parasitism? A test of the nesting-cue hypothesis. *Ethology* 103: 56-71.
- GILPIN, M. E., AND I. HANSKI (EDITORS). 1991. *Metapopulation dynamics: empirical and theoretical investigations*. Academic Press, London, U.K.
- GLITZENSTEIN, J. S., C. D. CANHAM, M. J. McDONNELL, AND D. R. STRENG. 1990. Effects of environment and land-use history on upland forests of the Cary Arboretum, Hudson Valley, New York. *Bulletin of Torrey Botanical Club* 117:106-122.
- GOCHFELD, M. 1979. Brood parasite and host coevolution: interactions between Shiny Cowbirds and two species of meadowlarks. *American Naturalist* 113:855-870.
- GOGUEN, C. B., AND N. E. MATHEWS. 1996. Nest desertion by Blue-gray Gnatcatchers in association with Brown-headed Cowbird parasitism. *Animal Behaviour* 52:613-619.
- GOGUEN, C. B., AND N. E. MATHEWS. 1998. Songbird community composition and nesting success in grazed and ungrazed pinyon-juniper woodlands. *Journal of Wildlife Management* 62:474-484.
- GOLDWASSER, S., D. A. GAINES, AND S. R. WILBUR. 1980. The Least Bell's Vireo in California: a de facto endangered race. *American Birds* 34:742-745.
- GOOSSEN, J. P., AND S. G. SEALY. 1982. Production of young in a dense nesting population of Yellow Warblers, *Dendroica petechia*, in Manitoba. *Canadian Field-Naturalist* 96:189-199.
- GOTMARK, F. 1992. The effects of investigator disturbance on nesting birds. *Current Ornithology* 9:63-104.
- GOWATY, P. A. 1996. Field studies of parental care in birds: new data focus questions on variation among females. *Advances in the Study of Behavior* 25:477-531.
- GRABER, J. W. 1961. Distribution, habitat requirements, and life history of the Black-capped Vireo (*Vireo atricapilla*). *Ecological Monographs* 31:313-336.
- GRAHAM, D. S. 1988. Responses of five host species to cowbird parasitism. *Condor* 90:588-591.
- GRANT, P. R., AND P. T. BOAG. 1980. Rainfall on the Galápagos and the demography of Darwin's finches. *Auk* 97:227-244.
- GRAY, M. V., AND J. GREAVES. 1984. Riparian forest as habitat for the Least Bell's Vireo. Pp. 605-611 in R. E. Warner and K. M. Hendrix (editors). *California riparian systems: ecology, conservation and productive management*. University of California Press, Berkeley, CA.
- GREENE, E., V. MUEHTER, AND W. DAVISON. 1996. Laz-

- uli Bunting (*Passerina amoena*). In A. Poole and F. Gill (editors). The birds of North America, no. 232. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, D.C.
- GREENLAND, S. 1988. On sample-size and power calculations for studies using confidence intervals. *American Journal of Epidemiology* 128:231-237.
- GREENLAW, J. S. 1996. Spotted Towhee (*Pipilo maculatus*). In A. Poole and F. Gill (editors). The birds of North America, no. 263. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, D.C.
- GREENWOOD, P. J. AND P. H. HARVEY. 1982. The natal and breeding dispersal of birds. *Annual Review of Ecology and Systematics* 13:1-21.
- GRIFFITH, J. T., AND J. C. GRIFFITH. In press. Cowbird control and the endangered Least Bell's Vireo: a management success story. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). The biology and management of cowbirds and their hosts. University of Texas Press, Austin, TX.
- GRINNELL, J. 1914. An account of the mammals and birds of the lower Colorado River Valley with especial reference to the distributional problems presented. University of California Publications in Zoology 12:51-294.
- GRINNELL, J., AND A. H. MILLER. 1944. The distribution of the birds of California. Pacific Coast Avifauna No. 27. Cooper Ornithological Club, Berkeley, CA.
- GRINNELL, J., AND T. I. STORER. 1924. Animal life in Yosemite. University of California Press, Berkeley, CA.
- GRINNELL, J., AND H. S. SWARTH. 1913. An account of the birds and mammals of the San Jacinto area of southern California. University of California Publications in Zoology 10:197-406.
- GROENEVELD, D. P., AND T. E. GRIEPENTROG. 1985. Interdependence of groundwater, riparian vegetation, and streambank stability: a case study. Pp. 44-48 in R. R. Johnson, C. D. Ziebell, D. R. Patton, and others (technical coordinators). Riparian ecosystems and their management: reconciling conflicting uses. USDA Forest Service Gen. Tech. Rep. RM-120. USDA Forest Service, Rocky Mountain Forest and Range Experimental Station, Fort Collins, CO.
- GRZYBOWSKI, J. A. 1991. Survivorship, dispersal, and population structure of the Black-capped Vireo at the Kerr Wildlife Management Area, Texas. Resource Protection Division, Texas Parks and Wildlife Department, Austin, TX.
- GRZYBOWSKI, J. A. 1995. Black-capped Vireo (*Vireo atricapillus*). In A. Poole and F. Gill (editors). The birds of North America, no. 181. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, D.C.
- GRZYBOWSKI, J. A., R. B. CLAPP, AND J. T. MARSHALL, JR. 1986. History and current population status of the Black-headed Vireo in Oklahoma. *American Birds* 40:1151-1161.
- GRZYBOWSKI, J. A., AND C. M. PEASE. In press. Comparing the relative effects of brood parasitism and nest predation on seasonal fecundity in passerine birds. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). The biology and management of cowbirds and their hosts. University of Texas Press, Austin, TX.
- GUREVITCH, J., AND L. V. HEDGES. 1993. Meta-analysis: combining the results of independent experiments. Pp. 378-398 in S. M. Scheiner and J. Gurevitch (editors). Design and analysis of ecological experiments. Chapman and Hall, New York, NY.
- GUSTAFSSON, L., AND W. J. SUTHERLAND. 1988. The costs of reproduction in the Collared Flycatcher *Ficedula albicollis*. *Nature* 335:813-815.
- GUTZWILLER, K. J., AND S. H. ANDERSON. 1992. Interception of moving organisms: influences of patch shape, size, and orientation on community structure. *Landscape Ecology* 6:293-303.
- GYSEL, L. W., AND L. J. LYON. 1980. Habitat analysis and evaluation. Pp. 305-327 in S. D. Schemnitz (editor). Wildlife management techniques manual. The Wildlife Society, Washington, D.C.
- HAHN, D. C., AND R. C. FLEISCHER. 1995. DNA fingerprint similarity between female and juvenile Brown-headed Cowbirds trapped together. *Animal Behavior* 49:1577-1580.
- HAHN, D. C., AND J. S. HATFIELD. 1995. Parasitism at the landscape scale: cowbirds prefer forests. *Conservation Biology* 6:1415-1424.
- HAHN, D. C., AND J. S. HATFIELD. In press. Host selection in the forest interior: cowbirds target ground-nesting species. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). Ecology and management of cowbirds and their hosts. University of Texas Press, Austin, TX.
- HAIK, S. M., J. D. BALLOU, AND N. J. CASNA. 1994. Identification of kin structure in Guam Rails: analysis of pedigrees and DNA profiles. *Molecular Ecology* 3:109-119.
- HAIK, S. M., J. D. BALLOU, AND N. J. CASNA. 1995. Genetic identification of kin in Micronesian Kingfishers. *Journal of Heredity* 86: 423-431.
- HALL, L. S., M. L. MORRISON, AND W. M. BLOCK. 1997. Songbird status and roles. Pp. 69-88 in W. M. Block and D. M. Finch (editors). Songbird ecology in southwestern ponderosa pine forests: a literature review. USDA Forest Service Gen. Tech. Rep. RM-292. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- HAMILTON, W. J., III, AND G. H. ORIANS. 1965. Evolution of brood parasitism in altricial birds. *Condor* 67:361-382.
- HANEY, J. C., D. S. LEE, AND M. WALSH-MCGEHEE. 1998. A quantitative analysis of winter distribution and habitats of Kirtland's Warblers in the Bahamas. *Condor* 100:201-217.
- HANKA, L. R. 1985. Recent altitudinal range expansion by the Brown-headed Cowbird in Colorado. *Western Birds* 16:183-184.
- HANN, H. W. 1937. Life history of the Oven-bird in southern Michigan. *Wilson Bulletin* 49:145-237.
- HANN, W. J., J. L. JONES, M. G. KARL, P. F. HESSBURG, R. E. KEANE, D. G. LONG, J. P. MENAKIS, C. H. MCNICOLL, S. G. LEONARD, R. A. GRAVENMIER, AND B. G. SMITH. 1997. Landscape dynamics of the basin. Volume II. Pp. 337-1055 in T. M. Quigley and S. J. Arbelbide (technical editors). An assessment of ecosystem components in the Interior Columbia Basin and portions of the Klamath and Great Basins.

- USDA Forest Service Gen. Tech. Rep. PNW-405. USDA Forest Service Pacific Northwest Research Station, Portland, OR.
- HANNA, W. C. 1928. Notes on the Dwarf Cowbird in southern California. *Condor* 30:161-162.
- HANSEN, P. L., R. D. PFISTER, K. BOGGS, B. J. COOK, J. JOY, AND D. K. HINCKLEY. 1995. Classification and management of Montana's riparian and wetland sites. Montana Forest and Conservation Experiment Station, Miscellaneous Publication No. 54. University of Montana, Missoula, MT.
- HANSKI, I. 1982. Dynamics of regional distribution: the core and satellite species hypothesis. *Oikos* 38:210-221.
- HARRIS, J. H. 1991. Effects of brood parasitism by Brown-headed Cowbirds on Willow Flycatcher nesting success along the Kern River, California. *Western Birds* 22:13-26.
- HARRIS, J. H., S. D. SANDERS, AND M. A. FLETT. 1987. Willow Flycatcher surveys in the Sierra Nevada. *Western Birds* 18:27-36.
- HARRISON, S. 1991. Local extinction in a metapopulation context: an empirical evaluation. *Biological Journal of the Linnean Society* 42:73-88.
- HARRISON, S., AND J. F. QUINN. 1989. Correlated environments and the persistence of metapopulations. *Oikos* 56:293-298.
- HAUFLER, J. B. 1998. A strategy for bird research in forested ecosystems of the western United States. Pp. 219-229 in J. M. Marzluff and R. Sallabanks (editors). *Avian conservation: research and management*. Island Press, Washington, D.C.
- HAYDEN, T. J., AND D. J. TAZIK. 1991. Project status report: 1991 field studies of two endangered species (the Black-capped Vireo and the Golden-cheeked Warbler) and the cowbird control program on Fort Hood, Texas. Report submitted to HQ III Corps and Fort Hood, DEH, Fort Hood, Texas.
- HAYDEN, T. J., D. J. TAZIK, R. H. MELTON, AND J. D. CORNELIUS. In press. Cowbird control program at Fort Hood, Texas: lessons for mitigation of cowbird parasitism on a landscape scale. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- HAYWARD, C. L. 1941. Notes on the nesting habits some mountain-dwelling birds in Utah. *Great Basin Naturalist* 2:1-8.
- HAYWARD, C. L. 1945. Biotic communities of the southern Wasatch and Unita Mountains, Utah. *Great Basin Naturalist* 6:1-124.
- HAYWARD, C. L., C. COTTAM, A. M. WOODBURY, AND H. H. FROST. 1976. Birds of Utah. *Great Basin Naturalist Memoirs* 1:1-229.
- HEINSOHN, R. G. 1987. Age-dependent vigilance in winter aggregations of cooperatively breeding White-winged Choughs (*Corcorax melanoramphos*). *Behavioral Ecology and Sociobiology* 20:303-306.
- HEINSOHN, R. G. 1991. Slow learning of foraging skills and extended parental care in cooperatively breeding White-winged Choughs. *American Naturalist* 137:864-881.
- HEIL, S. J. 1992. The importance of landscape patterns to bird diversity: a perspective from the Northern Rocky Mountains. *Northwest Environmental Journal* 8:119-137.
- HEIL, S. J. 1994. Human-induced changes in bird populations in coniferous forests in western North America during the past 100 years. *Studies in Avian Biology* 15:232-246.
- HEIL, S. J., R. L. HUTTO, C. R. PRESTON, AND D. M. FINCH. 1995. Effects of silvicultural treatments in the Rocky Mountains. Pp. 220-244 in T. E. Martin and D. M. Finch (editors). *Ecology and management of neotropical migratory birds*. Oxford University Press, New York, NY.
- HEIL, S. J., AND L. C. PAIGE. 1994. A preliminary assessment of birds in continuous and fragmented forests of western red-cedar/western hemlock in northern Idaho. Pp. 189-197 in D. M. Baumgartner and J. E. Lotan (editors). *Proceedings of a symposium on interior cedar-hemlock-white pine forests: ecology and management*. Washington State University Cooperative Extension, Pullman, WA.
- HEIL, S. J., AND R. E. WOODS. 1991. Bird assemblages in old-growth and rotation-aged Douglas-fir/ponderosa pine stands in the northern Rocky Mountains: a preliminary assessment. Pp. 93-100 in D. M. Baumgartner and J. E. Lotan (editors). *Proceedings of a symposium on interior Douglas-fir: the species and its management*. Washington State University Cooperative Extension, Pullman, WA.
- HENDERSON, A. B. 1870. Narrative of a prospecting expedition to the east fork and Clark's fork of the Yellowstone. In M. Meagher, *The bison of Yellowstone National Park*. 1973 National Park Service Science Monograph 1:1-161.
- HENSHAW, H. W. 1875. Chapter III. Report upon the ornithological collections made in portions of Nevada, Utah, California, Colorado, New Mexico, and Arizona during the years 1871, 1872, 1873, and 1874. U.S. Government Printing Office, Washington, D.C.
- HENSLER, G. L. 1985. Estimation and comparison of functions of daily nest survival probabilities using the Mayfield method. Pp. 289-301 in B. J. T. Morgan and P. M. North (editors). *Statistics in ornithology*. Springer-Verlag, New York, NY.
- HENSLER, G. L., AND J. D. NICHOLS. 1981. The Mayfield method of estimating nesting success: a model, estimators and simulation results. *Wilson Bulletin* 93:42-53.
- HERGENRADER, G. L. 1962. The incidence of nest parasitism by the Brown-headed Cowbird (*Molothrus ater*) on roadside nesting birds in Nebraska. *Auk* 79:85-88.
- HERKERT, J. R. 1994a. Breeding bird communities of midwestern prairie fragments: the effects of prescribed burning and habitat area. *Natural Areas Journal* 14:128-135.
- HERKERT, J. R. 1994b. The effects of habitat fragmentation on midwestern grassland bird communities. *Ecological Applications* 4:461-471.
- HIGGINS, K. F. 1986. A comparison of burn season effects on nesting birds in North Dakota mixed-grass prairie. *Prairie Naturalist* 18:219-228.
- HILBORN, R., AND M. MANGEL. 1997. *The ecological detective: confronting models with data*. Princeton University Press, Princeton, NJ.

- HILL, D. P., AND S. G. SEALY. 1994. Desertion of nests parasitized by cowbirds: have Clay-coloured Sparrows evolved an anti-parasite defense? *Animal Behaviour* 48:1063–1070.
- HILL, R. A. 1976. Host-parasite relationships of the Brown-headed Cowbird in a prairie habitat of west-central Kansas. *Wilson Bulletin* 88:555–565.
- HOBSON, K. A., P. PERRINE, E. B. ROBERTS, M. L. FOSTER, AND P. WOODIN. 1985. A breeding bird survey of Salt Marsh Yellowthroats, *Geothlypis trichas sinuosa*, in the San Francisco Bay Region. San Francisco Bay Bird Observatory. U.S. Fish and Wildlife Service Contract No. 84–57. U.S. Fish and Wildlife Service, Sacramento, CA.
- HOBSON, K. A., AND S. G. SEALY. 1989. Responses of Yellow Warblers to the threat of cowbird parasitism. *Animal Behavior* 38:510–519.
- HOCHACHAKA, W. 1990. Seasonal decline in reproductive performance of Song Sparrows. *Ecology* 71:1279–1288.
- HOFSLUND, P. B. 1957. Cowbird parasitism of the northern Yellow-Throat. *Auk* 74:42–48.
- HOFSLUND, P. B. 1959. A life history study of the Yellowthroat, *Geothlypis trichas*. *Proceedings of the Minnesota Academy of Sciences* 27:144–174.
- HOLFORD, K. C., AND D. D. ROBY. 1993. Factors limiting fecundity of captive Brown-headed Cowbirds. *Condor* 95:536–545.
- HOLMES, B. 1993. An avian arch-villain gets off easy. *Science* 262:1514–1515.
- HOLMES, N. D., D. S. SMITH, AND A. JOHNSTON. 1979. Effect of grazing by cattle on the abundance of grasshoppers on fescue grasslands. *Journal of Range Management* 32:310–311.
- HOLMES, R. T., AND T. W. SHERRY. 1992. Site fidelity of migratory warblers in temperate breeding and neotropical wintering areas: implications for population dynamics, habitat selection, and conservation. Pp 563–575 in J. M. Hagan, III, and D. W. Johnston (editors). *Ecology and conservation of neotropical migrant landbirds*. Smithsonian Institution Press, Washington, D.C.
- HOOVER, J. P., AND M. C. BRITTINGHAM. 1993. Regional variation in brood parasitization of Wood Thrushes. *Wilson Bulletin* 105:228–238.
- HOOVER, J. P., M. C. BRITTINGHAM, AND L. J. GOODRICH. 1996. Effects of forest patch size on nesting success of Wood Thrushes. *Auk* 112:146–155.
- HORN, H. S., AND D. I. RUBENSTEIN. 1984. Behavioral adaptations and life history. Pp. 279–300 in J. Krebs and N. B. Davies (editors). *Behavioral ecology: an evolutionary approach*. Blackwell, Oxford, U.K.
- HOSMER, D. W., JR., AND S. LEMESHOW. 1989. *Applied logistic regression*. Wiley Series in Probability and Statistics. John Wiley and Sons, New York, NY.
- HOWE, H. F. 1994. Managing species diversity in tall-grass prairie: assumptions and implications. *Conservation Biology* 8:691–704.
- HOWE, W. H., AND F. L. KNOPF. In press. The role of vegetation on cowbird parasitism on Yellow Warblers. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- HOY, G., AND J. OTTOW. 1964. Biological and oological studies of the molothrine cowbirds (Icteridae) of Argentina. *Auk* 81:186–203.
- HUBER, G. E., AND A. A. STEUTER. 1984. Vegetation profile and grassland bird response to spring burning. *Prairie Naturalist* 16:55–61.
- HUNT, J. 1991. Status of black-capped vireos on Fort Hood, Texas. Preliminary report submitted to U.S. Army Construction Engineering Laboratories, Champaign, IL.
- HUNTER, W. C., R. D. OHMART, AND B. W. ANDERSON. 1988. Use of exotic saltcedar (*Tamarix chinensis*) by birds in arid riparian systems. *Condor* 90:113–123.
- HURLBERT, L. C. 1988. Causes of fire effects in tall-grass prairie. *Ecology* 69:46–58.
- HUTTO, R. L. 1995a. Composition of bird communities following stand-replacement fires in northern Rocky Mountain (U.S.A.) conifer forests. *Conservation Biology* 9:1041–1058.
- HUTTO, R. L. 1995b. USFS Northern Region Songbird Monitoring Program: Distribution and Habitat Relationships. USDA Forest Service Report R1–95–05. (Also available at www.umt.edu/biology/dbs/land-bird.html).
- HUTTO, R. L., S. M. PLETSCHE, AND P. HENDRICKS. 1986. A fixed radius point count method for non-breeding and breeding season use. *Auk* 103:593–602.
- IMMELMANN, K. 1971. Ecological aspects of periodic reproduction. Pp. 342–489 in D. S. Farner and J. R. King (editors). *Avian biology*. Vol 1. Academic Press, New York, NY.
- INGOLD, D. J. 1996. Delayed nesting decreases reproductive success in Northern Flickers: implications for competition with European Starlings. *Journal of Field Ornithology* 67:321–326.
- INTERNATIONAL BIRD CENSUS COMMITTEE. 1970. An international standard for a mapping method in bird census work recommended by the International Bird Census Committee. *Audubon Field Notes* 24:722–726.
- JACKSON, N. H., AND D. D. ROBY. 1992. Fecundity and egg-laying patterns of captive yearling Brown-headed Cowbirds. *Condor* 94:585–589.
- JAMES, F. R., AND C. E. MCCULLOCH. 1990. Multivariate analysis in ecology and systematics: panacea or Pandora's box. *Annual Review of Ecology and Systematics* 21:129–166.
- JAMES, F. C., AND H. H. SHUGART, JR. 1970. A quantitative method of habitat description. *Audubon Field Notes* 24:727–736.
- JEPSON-INNES, K., AND C. E. BOCK. 1989. Response of grasshoppers (Orthoptera:Acrididae) to livestock grazing in southeastern Arizona: differences between seasons and subfamilies. *Oecologia* 78:430–431.
- JETTE, L. A., T. J. HAYDEN, AND J. D. CORNELIUS. 1998. Demographics of the Golden-cheeked Warbler (*Dendroica chrysoparia*) on Fort Hood, Texas. USA-CERL Technical Report 98/52. U.S. Army Construction Engineering Laboratories, Champaign, IL.
- JOHNSON, M., AND M. SOGGE. 1995. Cowbird concentrations at livestock corrals in Grand Canyon National Park. Pp. 275–284 in C. Van Riper (editor). *Proceedings of the second biennial conference on research in Colorado Plateau National Parks*. Na-

- tional Park Service Proceedings NPS/NRNAU/NRTP-95/11. Flagstaff, AZ.
- JOHNSON, R. G., AND S. A. TEMPLE. 1986. Assessing habitat quality for birds nesting in fragmented tallgrass prairies. Pp. 245–250 in J. Verner, M. L. Morrison, and C. J. Ralph (editors). *Wildlife 2000: modeling habitat relationships of terrestrial vertebrates*. University of Wisconsin Press, Madison, WI.
- JOHNSON, R. G., AND S. A. TEMPLE. 1990. Nest predation and brood parasitism of tallgrass prairie birds. *Journal of Wildlife Management* 54:106–111.
- JOHNSON, R. R., AND L. T. HAIGHT. 1984. Riparian problems and initiatives in the American southwest: a regional perspective. Pp. 404–411 in R. E. Warner and K. M. Hendrix (editors). *California riparian systems: ecology, conservation and productive management*. University of California Press, Berkeley, CA.
- JOHNSON, R. R., L. T. HAIGHT, AND J. M. SIMPSON. 1977. Endangered species vs. endangered habitats: a concept. Pp. 68–79 in R. R. Johnson and D. A. Jones (technical coordinators). *Importance, preservation and management of riparian habitat: a symposium*. USDA Forest Service Gen. Tech. Rep. RM-166. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- KANTRUD, H. A. 1981. Grazing intensity effects on the breeding avifauna of North Dakota native grasslands. *Canadian Field-Naturalist* 95:404–417.
- KANTRUD, H. A., AND R. L. KOLOGOWSKI. 1982. Effects of soils and grazing on breeding birds of uncultivated upland grasslands of the Northern Great Plains. U.S. Fish and Wildlife Service Research Report No. 15, Washington, D.C.
- KEELER-WOLF, T., V. KEELER-WOLF, AND W. A. CALDER. 1972. Bird fauna of the vicinity of the Rocky Mountain Biological Laboratory. *Colorado Field Ornithology* 15: 22–25.
- KELLY, S. T., AND M. E. DECAPITA. 1982. Cowbird control and its effect on Kirtland's Warbler reproductive success. *Wilson Bulletin* 94:63–365.
- KEPLER, C. B., W. G. IRVINE, M. E. DECAPITA, AND J. WEINRICH. 1996. The conservation and management of Kirtland's Warbler *Dendroica kirtlandii*. *Bird Conservation International* 6:11–22.
- KING, A. P. 1979. Variables affecting parasitism in the North American cowbird (*Molothrus ater*). Ph.D. dissertation. Cornell University, Ithaca, NY.
- KLAAS, E. E. 1975. Cowbird parasitism and nesting success in the Eastern Phoebe. *Occasional Papers of the Kansas Museum of Natural History* 41:1–18.
- KLEINBAUM, D. G., L. L. KUPPER, AND K. E. MULLER. 1988. Applied regression analysis and other multivariable methods. Duxbury Press, Belmont, CA.
- KNAPTON, R. W. 1978. Breeding ecology of the Clay-colored Sparrow. *Living Bird* 17:137–158.
- KNOPF, F. L. 1985. Significance of riparian vegetation to breeding birds across an altitudinal cline. Pp. 105–111 in R. R. Johnson, C. D. Ziebell, D. R. Patton, P. F. Ffolliot and R. H. Hamre (editors). *Riparian ecosystems and their management: reconciling conflicting uses*. USDA Forest Service Gen. Tech. Rep. RM-120. USDA Forest Service, Rocky Mountain Forest and Range Experimental Station, Fort Collins, CO.
- KNOPF, F. L. 1994. Avian assemblages on altered grasslands. *Studies in Avian Biology* 15:247–257.
- KNOPF, F. L., R. R. JOHNSON, T. RICH, F. B. SAMSON, AND R. C. SZARO. 1988. Conservation of riparian ecosystems in the United States. *Wilson Bulletin* 100:272–282.
- KNOPF, F. L., AND F. B. SAMSON. 1994. Scale perspectives on avian diversity in western riparian ecosystems. *Conservation Biology* 8:669–676.
- KNOPF, F. L., J. A. SEDGWICK, AND R. W. CANNON. 1988. Guild structure of a riparian avifauna relative to seasonal cattle grazing. *Journal of Wildlife Management* 52:280–290.
- KNUTSON, H., AND J. B. CAMPBELL. 1976. Relationships of grasshoppers (Acrididae) to burning, grazing, and range sites of native tallgrass prairie in Kansas. *Proceedings of the Tall Timbers Conference on Ecology of Animal Control by Habitat Management* 6:107–120.
- KOFORD, R. R., B. S. BOWEN, J. T. LOKEMOEN, AND A. D. KRUSE. In press. Cowbird parasitism in grassland and cropland in the northern Great Plains. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- KOLOSZAR, J. A. 1998. Field studies of the Black-capped Vireo (*Vireo atricapillus*) on Fort Hood, Texas. Pp. 7–27 in *The Nature Conservancy, Summary of 1997 Research Activities*. Texas Conservation Data Center, The Nature Conservancy of Texas, Fort Hood, Texas.
- KOZLOVIC, D. R. In press. Cowbird parasitism and productivity of House Finch hosts. *Canadian Journal of Zoology*.
- KOZLOVIC, D. R., R. W. KNAPTON, AND J. C. BARLOW. 1996. Unsuitability of the House Finch as a host of the Brown-headed Cowbird. *Condor* 98:253–258.
- KREBS, J., AND N. B. DAVIES. 1993. *An introduction to behavioral ecology*. Blackwell Scientific Publications, Cambridge, MA.
- KRUSE, A. D., AND J. L. PIEHL. 1986. The impact of prescribed burning on ground-nesting birds. Pp. 153–156 in G. K. Clambey and R. H. Pemble (editors). *The prairie—past, present, and future*. North Dakota State University, Fargo, ND.
- KUEHL, R. O. 1994. *Statistical principles of research design and analysis*. Duxbury Press, Belmont, CA.
- LAAKE, J. L., S. T. BUCKLAND, D. R. ANDERSON, AND K. P. BURNHAM. 1993. *DISTANCE User's Guide*. Colorado Cooperative Fish and Wildlife Research Unit, Colorado State University, Fort Collins, CO.
- LANGEN, T. A., D. T. BOLGER, AND T. J. CASE. 1991. Predation on artificial bird nests in chaparral fragments. *Oecologia* 86:395–401.
- LANTZ, G. 1976. *Compartment prescription and environmental analysis report*. USDA Forest Service, 2410–1, Harrisville, MI.
- LARISON, B. 1996. *Avian responses to riparian restoration*. M.S. thesis. San Francisco State University, San Francisco, CA.
- LARISON, B., S. A. LAYMON, P. L. WILLIAMS, AND T. B. SMITH. 1998. Song Sparrows vs. cowbird brood parasites: impacts of forest structure and nest-site selection. *Condor* 100:93–101.

- LAWTON, J. H., J. R. BEDDINGTON, AND R. BONSER. 1974. Switching in invertebrate predators. Pp. 141–158 in M. B. Usher and M. H. Williamson (editors). Ecological stability. Chapman and Hall, London, U.K.
- LAYMON, S. A. 1987. Brown-headed cowbirds in California: historical perspectives and management opportunities in riparian habitats. *Western Birds* 18:63–70.
- LAYMON, S. A. 1988. Ecology of the Spotted Owl in the central Sierra Nevada, California. PhD dissertation. University of California, Berkeley, CA.
- LAYMON, S. A., P. L. WILLIAMS, M. D. HALTERMAN, AND S. D. ROWE. 1996. Monitoring of riparian habitat restoration sites at the South Fork Kern River, California: breeding birds and habitat characteristics, 1988 to 1995. Nature Conservancy Report. Kern River Research Center, Weldon, CA.
- LAYMON, S. A., P. L. WILLIAMS, M. D. HALTERMAN, AND S. D. ROWE. 1997. Riparian restoration at the Kern River Preserve: 1996 bird monitoring results. Nature Conservancy Report. Kern River Research Center, Weldon, CA.
- LEVIN, S. 1974. Dispersion and population interactions. *American Naturalist* 108:207–228.
- LINDSDALE, J. M. 1936. The birds of Nevada. Pacific Coast Avifauna No. 23. Cooper Ornithological Club, Berkeley, CA.
- LITTELL, R. C., G. A. MILLIKEN, W. W. STROUP, AND R. D. WOLFINGER. 1996. SAS System for Mixed Models. SAS Institute, Inc., Cary, NC.
- LOSENSKY, B. J. 1993. Historical vegetation in Region One by climatic section. USDA Forest Service, Draft Report for the Northern Region, P. O. Box 7669, Missoula, MT 59807.
- LOWTHER, P. E. 1979. Nest selection by Brown-headed Cowbirds. *Wilson Bulletin* 91:118–122.
- LOWTHER, P. E. 1993. Brown-headed Cowbird (*Molothrus ater*). In A. Poole and F. Gill (editors). The birds of North America, no. 47. Academy of Natural Sciences, Philadelphia, and American Ornithologists' Union, Washington, D.C.
- LOWTHER, P. E. 1995. Bronzed Cowbird (*Molothrus aeneus*). In A. Poole and F. Gill (editors). The birds of North America, no. 144. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, D.C.
- LYMAN, R. L., AND S. D. LIVINGSTON. 1983. Late Quaternary mammalian zoogeography of eastern Washington. *Quaternary Research* 20: 360–373.
- LYNCH, M. 1988. Estimation of relatedness by DNA fingerprinting. *Molecular Biology and Evolution* 5: 584–599.
- LYNCH, M. 1990. The similarity index and DNA fingerprinting. *Molecular Biology and Evolution* 7: 478–484.
- MA, Z. 1995. Using a rule-based merging algorithm to eliminate "salt-pepper" and small regions of classified images. Proceedings of the Ninth Annual Symposium on Geographic Information Systems 11: 834–837.
- MAJOR, R. E. 1990. The effect of human observers on the intensity of nest predation. *Ibis* 132:608–612.
- MANN, C. 1990. Meta-analysis in the breach. *Science* 249:476–480.
- MARTIN, T. E. 1987. Artificial nest experiments: effects of nest appearance and type of predator. *Condor* 89: 925–928.
- MARTIN, T. E. 1988. On the advantage of being different: nest predation and the coexistence of bird species. *Proceedings of the National Academy of Science (USA)* 85:2196–2199.
- MARTIN, T. E. 1992. Breeding productivity considerations: what are the appropriate habitat features for management? Pp. 455–473 in J. M. Hagan, III, and D. W. Johnston (editors). Ecology and conservation of neotropical migrant landbirds. Smithsonian Institution Press, Washington, D.C.
- MARTIN, T. E. 1993. Nest predation among vegetative layers and habitat types: revising the dogmas. *American Naturalist* 141: 897–913.
- MARTIN, T. E., AND G. R. GEUPEL. 1993. Nest-monitoring plots: methods for locating nests and monitoring success. *Journal of Field Ornithology* 64:507–519.
- MARTIN, T. E., W. M. HOCHACHKA, C. J. CONWAY, AND J. W. JENKINS. 1996. BBIRD field protocol. Montana Cooperative Wildlife Research Unit. University of Montana, Missoula, MT.
- MARTIN, T. E., C. R. PAINE, C. J. CONWAY, W. M. HOCHACHKA, P. ALLEN, AND J. W. JENKINS. 1997. BBIRD field protocols [online]. Available: <http://pica.wru.umt.edu/bbird/protocol/protocol.htm> [1998, September 13].
- MARTIN, T. E., AND J. J. ROPER. 1988. Nest predation and nest-site selection of a western population of the Hermit Thrush. *Condor* 90:51–57.
- MARVIL, R. E., AND A. CRUZ. 1989. Impact of Brown-headed Cowbird parasitism on the reproductive success of the Solitary Vireo. *Auk* 106:276–480.
- MASON, P. 1986. Brood parasitism in a host generalist, the Shiny Cowbird: I. The quality of different species as hosts. *Auk* 103:52–60.
- MATSUOKA, S. M., C. M. HANDEL, AND D. D. ROBY. 1997. Nesting ecology of Townsend's Warblers in relation to habitat characteristics in a mature boreal forest. *Condor* 99:271–281.
- MAY, R. M., AND S. K. ROBINSON. 1985. Population dynamics of avian brood parasitism. *American Naturalist* 126:475–494.
- MAYFIELD, H. F. 1960. The Kirtland's Warbler. Cranbrook Institute of Science, Bloomfield Hills, MI.
- MAYFIELD, H. F. 1961. Nesting success calculated from exposure. *Wilson Bulletin* 73:255–261.
- MAYFIELD, H. F. 1965. The Brown-headed Cowbird, with old and new hosts. *Living Bird* 4:13–28.
- MAYFIELD, H. F. 1975. Suggestions for calculating nest success. *Wilson Bulletin* 87:456–466.
- MAYFIELD, H. F. 1977. Brown-headed cowbird: agent of extinction? *American Birds* 31:107–113.
- MAYNARD, W. R. 1994. Summary of 1994 survey efforts in New Mexico for Southwestern Willow Flycatchers (*Empidonax traillii extimus*). Contract #94–516–69. New Mexico Department of Game and Fish, Albuquerque, NM.
- MCCULLAGH, P., AND J. A. NELDER. 1983. Generalized linear models. Chapman and Hall, New York, NY.
- MCCUNE, B. 1983. Fire frequency reduced two orders of magnitude in the Bitterroot Canyons, Montana. *Canadian Journal of Forest Research* 13:212–218.

- MCDONALD, D. B., AND H. CASWELL. 1993. Matrix methods for avian demography. *Current Ornithology* 10:139–185.
- MCDONALD, J. N. 1981. North American bison: their classification and evolution. University of California Press, Berkeley, CA.
- MCGARIGAL, K., AND B. J. MARKS. 1995. FRAGS-TATS: spatial pattern analysis program for quantifying landscape structure. USDA Forest Service Gen. Tech. Rep. PNW-351. USDA Forest Service Pacific Northwest Research Station, Portland, OR.
- MCGEEN, D. S. 1972. Cowbird-host relationships. *Auk* 89:360–380.
- MCMASTER, D. G., AND S. G. SEALY. 1997. Host-egg removal by Brown-headed Cowbirds: a test of the host incubation limit hypothesis. *Auk* 114:212–220.
- MEAGHER, M. 1976. Winter weather as a population-regulating influence on free-ranging bison in Yellowstone National Park. *Research in the Parks: transactions of the National Centennial Symposium* 1:29–38.
- MEAGHER, M. 1986. *Bison bison*. *Mammalian Species* 266:1–8.
- MEANEY, C. A., AND D. VAN VUREN. 1993. Recent distribution of bison in Colorado west of the Great Plains. *Proceedings of the Denver Museum Natural History, Series 3*, 4:1–10.
- MEHTA, C., AND N. PATEL. 1995. StatXact 3, User manual. Cytel Software Corporation, Cambridge, MA.
- MENG, A., R. E. CARTER, AND D. T. PARKIN. 1989. The variability of DNA fingerprints in three species of swan. *Heredity* 64:73–80.
- MERRILL, J. C. 1876. Notes on Texan birds. *Bulletin of the Nuttall Ornithological Club* 1:88–89.
- MICHIGAN WEATHER SERVICE. 1974. Climate of Michigan bystations. Michigan Department of Agriculture Cooperating with NOAA—National Weather Service. U.S. Department of Commerce, East Lansing, MI.
- MIDDLETON, A. L. A. 1977. Effect of cowbird parasitism on American Goldfinch nesting. *Auk* 94:304–307.
- MIDDLETON, A. L. A. 1991. Failure of Brown-headed Cowbird parasitism in nests of the American Goldfinch. *Journal of Field Ornithology* 62:200–203.
- MILLS, G. S., J. B. DUNNING, JR., AND J. M. BATES. 1991. The relationship between breeding bird density and vegetation volume. *Wilson Bulletin* 103:468–479.
- MILLS, L. S., D. F. DOAK, AND M. J. WISDOM. In press. The reliability of conservation actions based on elasticity analysis of matrix models. *Conservation Biology*.
- MINNICH, R. A. 1985. Evolutionary convergence or phenotypic plasticity? Responses to summer rain by California chaparral. *Physical Geography* 6:272–287.
- MORRIS, D. L., AND F. R. THOMPSON, III. 1998. Effects of habitat and invertebrate density on abundance and foraging behavior of Brown-headed Cowbirds. *Auk* 115:376–385.
- MORRISON, M. L., AND B. G. MARCOT. 1995. An evaluation of resource inventory and monitoring program used in national forest planning. *Environmental Management* 19:147–156.
- MORSE, D. H. 1988. American warblers: an ecological and behavioral perspective. Harvard University Press, Cambridge, MA.
- MORSE, S., AND S. K. ROBINSON. In press. Nesting success of a neotropical migrant in a multiple-use forested landscape. *Conservation Biology*.
- MORTON, E. S., L. FORMAN, AND M. BRAUN. 1990. Extra-pair fertilizations and the evolution of colonial breeding in Purple Martins. *Auk* 107:275–283.
- MORTON, M. L. 1992. Effects of sex and birth date on premigration biology, migration schedules, return rates and natal dispersal in the Mountain White-crowned Sparrow. *Condor* 94:117–133.
- MOSCONI, S. L., AND R. L. HUTTO. 1982. The effects of grazing on land birds of a western Montana riparian habitat. Pp. 221–223 in J. M. Peek and P. D. Dalke (editors). *Wildlife-livestock relationships symposium*. University of Idaho, Forest, Wildlife, and Range Experiment Station, Moscow, ID.
- MUIZNIEKS, B. D., T. E. CORMAN, S. J. SFERRA, M. K. SOGGE, AND T. J. TIBBITTS. 1994. Arizona Partners in Flight 1993 Southwestern Willow Flycatcher survey. Nongame and Endangered Wildlife Program Tech. Rpt. 52. Arizona Game and Fish Department, Phoenix, AZ.
- MUMFORD, R. E. 1952. Bell's Vireo in Indiana. *Wilson Bulletin* 64:224–233.
- NAKAMURA, T. K., AND A. CRUZ. In press. The ecology of egg puncture by the Shiny Cowbird (*Molothrus bonariensis*) in southwestern Puerto Rico. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION. 1994. Climatological data annual summary, Arizona. Vol. 97, No. 13. US Department of Commerce, Asheville, NC.
- NERNEY, N. J. 1958. Grasshopper infestation in relation to range conditions. *Journal of Range Management* 3:308–315.
- NEUDORF, D. L., AND S. G. SEALY. 1992. Reactions of four passerine species to threats of predation and cowbird parasitization: enemy recognition or generalized response? *Behaviour* 123:84–105.
- NEUDORF, D. L., AND S. G. SEALY. 1994. Sunrise nest attentiveness in cowbird hosts. *Condor* 96:162–169.
- NEWMAN, G. A. 1970. Cowbird parasitism and nesting success of Lark Sparrows in southern Oklahoma. *Wilson Bulletin* 82:304–309.
- NEWTON, I. (EDITOR). 1989. Lifetime reproduction in birds. Academic Press, London, U.K.
- NEWTON, S. F., AND A. V. NEWTON. 1997. The effect of rainfall and habitat on abundance and diversity of birds in a fenced protected area in the central Saudi Arabian desert. *Journal of Arid Environments* 35:715–735.
- NICE, M. M. 1937. Studies in the life history of the song sparrow. I. *Transactions of the Linnean Society of New York* 4:1–247.
- NICE, M. M. 1954. Problems of incubation periods in North American birds. *Condor* 56:173–197.
- NOLAN, V., JR. 1978. The ecology and behavior of the Prairie Warbler, *Dendroica discolor*. *Ornithological Monographs* 26:1–595.

- NOON, B. R. 1981. Techniques for sampling avian habitats. Pp. 42–52 in D. E. Capen (editor). The use of multivariate statistics in the studies of wildlife habitat. USDA Forest Service Gen. Tech. Rep. RM-87. USDA Forest Service, Rocky Mountain Forest and Range Experimental Station, Fort Collins, CO.
- NORMAN, R. F., AND R. J. ROBERTSON. 1975. Nest-searching behavior in the Brown-headed Cowbird. *Auk* 92:610–611.
- NORRIS, R. T. 1947. The cowbirds of Preston Frith. *Wilson Bulletin* 59:83–103.
- NUR, N. 1988. The cost of reproduction in birds: an examination of the evidence. *Ardea* 76: 155–168.
- O'LEARY, J. F. 1990. Coastal sage scrub: general characteristics and consideration for biological conservation. Pp. 24–41 in A. A. Schoenherr (editor). Endangered plant communities of southern California. Southern California Botanists Special Publication No. 3.
- OBERHOLSER, H. C. 1974. The birdlife of Texas. Vol. 2. University of Texas Press, Austin, TX.
- O'CONNOR, R. J., AND J. FAABORG. 1992. The relative abundance of the Brown-headed Cowbird (*Molothrus ater*) in relation to exterior and interior edges in forests of Missouri. *Transactions of the Missouri Academy of Sciences* 26:1–9.
- OHMART, R. D. 1994. The effects of human-induced changes on the avifauna of western riparian habitats. *Studies in Avian Biology* 15:273–285.
- ORIAN, G. H., E. RØSKAFT, AND L. D. BELETSKY. 1989. Do Brown-headed Cowbirds lay their eggs at random in the nests of Red-winged Blackbirds? *Wilson Bulletin* 101:599–605.
- ORING, L. W., K. P. ABLE, D. W. ANDERSON, L. F. BAPTISTA, A. S. GAUNT, F. B. GILL, AND J. C. WINGFIELD. 1988. Guidelines for the use of wild birds in research. *Auk* (supplement) 105:1A–14A.
- ORING, L. W., R. C. FLEISCHER, J. M. REED, AND K. MARSDEN. 1992. Cuckoldry via sperm storage in the polyandrous Spotted Sandpiper. *Nature* 359:631–633.
- ORR, R. T., AND J. MOFFITT. 1971. Birds of the Lake Tahoe region. California Academy of Sciences, San Francisco, CA.
- ORTEGA, C. P., AND A. CRUZ. 1988. Mechanisms of egg acceptance by marsh-dwelling blackbirds. *Condor* 90: 349–358.
- ORTEGA, C. P., AND A. CRUZ. 1991. A comparative study of cowbird parasitism in Yellow-headed Blackbirds and Red-winged Blackbirds. *Auk* 108: 16–24.
- ORTEGA, C. P., J. C. ORTEGA, AND A. CRUZ. 1994. Use of artificial cowbird eggs as a potential management tool in deterring parasitism. *Journal of Wildlife Management* 58:488–492.
- OTT, L. 1988. An introduction to statistical methods and data analysis. PWS-Kent Publishing Co., Boston, MA.
- PATON, P. W. C. 1994. The effect of edge on avian nest success: how strong is the evidence? *Conservation Biology* 8:17–26.
- PATTIE, D. L., AND N. A. M. VERBEEK. 1966. Alpine birds of the Beartooth Mountains. *Condor* 68:167–176.
- PATTIE, D. L., AND N. A. M. VERBEEK. 1967. Alpine mammals of the Beartooth Mountains. *Northwest Science* 41:110–117.
- PAYNE, R. B. 1973. The breeding season of a parasitic bird, the Brown-headed Cowbird. *Condor* 75: 80–89.
- PAYNE, R. B. 1976. The clutch size and numbers of eggs of Brown-headed Cowbirds: effects of latitude and breeding season. *Condor* 78:337–342.
- PAYNE, R. B. 1977. The ecology of brood parasitism in birds. *Annual Review of Ecology and Systematics* 8:1–28.
- PAYNE, R. B. 1989. Indigo Bunting. Pp. 153–172 in I. Newton (editor). Lifetime reproduction in birds. Academic Press, London, U.K.
- PAYNE, R. B. 1992. Indigo Bunting. In A. Poole and F. Gill (editors). The birds of North America, no. 4. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, D.C.
- PAYNE, R. B. 1997. Avian brood parasitism. Pp. 338–369 in D. H. Clayton and J. Moore (editors). Host-parasite evolution: general principles and avian models. Oxford University Press, Oxford, U.K.
- PAYNE, R. B. 1998. Brood parasitism in birds: strangers in the nest. *BioScience* 48:377–386.
- PAYNE, R. B., AND K. PAYNE. 1967. Cuckoo hosts in southern Africa. *Ostrich* 38:135–143.
- PAYNE, R. B., AND L. L. PAYNE. 1998. Brood parasitism by cowbirds: risks and effects on reproductive success and survival in Indigo Buntings. *Behavioral Ecology* 9:64–73.
- PEASE, C. M., AND J. A. GRYBOWSKI. 1995. Assessing the consequences of brood parasitism and nest predation on seasonal fecundity in passerine birds. *Auk* 112:343–363.
- PECK, G. K., AND R. D. JAMES. 1987. Breeding birds of Ontario. Nidology and distribution, Vol. 2. Passerines. Life Sciences Miscellaneous Publications, Royal Ontario Museum, Toronto, ON.
- PEER, B. D. 1998. An experimental investigation of egg rejection behavior in the grackles (*Quiscalus*). Ph.D. dissertation. University of Manitoba, Winnipeg, MB.
- PEER, B. D., AND E. K. BOLLINGER. 1997. Explanations for the infrequent cowbird parasitism on Common Grackles. *Condor* 99:151–161.
- PEER, B. D., AND E. K. BOLLINGER. In press. Why do female Brown-headed Cowbirds remove host eggs? A test of the incubation efficiency hypothesis. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). The biology and management of cowbirds and their hosts. University of Texas Press, Austin, TX.
- PEER, B. D., AND S. G. SEALY. 1999. Laying time of the Bronzed Cowbird. *Wilson Bulletin*. 111:138–140.
- PERRINS, C. M. 1965. Population fluctuations and clutch-size in the Great Tit, *Parus major*. *Journal of Animal Ecology* 34:601–647.
- PERRINS, C. M. 1970. The timing of birds' breeding seasons. *Ibis* 112:242–255.
- PERRINS, C. M., AND R. H. MCCLEERY. 1989. Laying dates and clutch size in the Great Tit. *Wilson Bulletin* 101:236–253.
- PETERJOHN, B. G., J. R. SAUER, AND S. SCHWARTZ. In press. Temporal and geographic patterns in popula-

- tion trends of Brown-headed Cowbirds. *In* J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- PETIT, L. J. 1991. Adaptive tolerance of cowbird parasitism by Prothonotary Warblers: a consequence of nest-site limitation? *Animal Behaviour* 41:425-432.
- PETIT, L. J., AND D. R. PETIT. *In press*. Brown-headed Cowbird parasitism of migratory birds: effects of forest area and surrounding landscape. *In* J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- PETIT, L. J., D. R. PETIT, AND T. E. MARTIN. 1995. Landscape-level management of migratory birds: looking past the trees to see the forest. *Wildlife Society Bulletin* 23:420-429.
- PETRINOVICH, L., AND T. L. PATTERSON. 1978. Cowbird parasitism on the White-crowned Sparrow. *Auk* 95: 415-417.
- PETRINOVICH, L., AND T. L. PATTERSON. 1983. The White-crowned Sparrow: reproductive success (1975-1980). *Auk* 100:811-825.
- PHILLIPS, R. S. 1951. Nest location, cowbird parasitism, and nesting success of the Indigo Bunting. *Wilson Bulletin* 63:206-207.
- PICMAN, J. 1992. Egg destruction by Eastern Meadowlarks. *Wilson Bulletin* 104:520-525.
- PLISSNER, J. H., AND P. A. GOWATY. 1996. Patterns of natal dispersal, turnover, and dispersal costs in Eastern Bluebirds. *Animal Behavior* 51:1307-1322.
- PLUMB, G. E., AND J. L. DODD. 1993. Foraging ecology of bison and cattle on a mixed prairie: implications for natural area management. *Ecological Applications* 3:631-643.
- POST, W., A. CRUZ, AND D. B. MCNAIR. 1993. The North American invasion pattern of the Shiny Cowbird. *Journal of Field Ornithologists* 64:32-41.
- POST, W., AND J. W. WILEY. 1977. Reproductive interactions of the Shiny Cowbird and the Yellow-shouldered Blackbird. *Condor* 79:176-184.
- PRICE, J., S. DROEGE, AND A. PRICE. 1995. *The summer atlas of North American birds*. Academic Press, New York, NY.
- PULICH, W. M. 1976. *The Golden-cheeked Warbler—a bioecological study*. Texas Parks and Wildlife Press, Austin, TX.
- PULLIAM, H. R. 1988. Sources, sinks, and population regulation. *American Naturalist* 137:550-566.
- PURCELL, K. L. 1997. Use of a fiberscope for examining cavity nests. *Journal of Field Ornithology* 68: 283-286.
- PYLE, P., S. N. G. HOWELL, R. P. YUNICK, AND D. DESANTE. 1987. *Identification guide to North American passerines*. Slate Creek Press, Bolinas, CA.
- PYLYPEC, B. 1991. Impacts of fire on bird populations in a fescue prairie. *Canadian Field-Naturalist* 105: 346-349.
- QUIGLEY, T. M., AND S. J. ARBELBIDE (TECHNICAL EDITORS). 1997. An assessment of ecosystem components in the interior Columbia basin and portions of the Klamath and Great Basins. Vol. 2. USDA Forest Service Gen. Tech. Rep. PNW-405. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- RAIM, A. *In press*. Spatial patterns of breeding female cowbirds on an Illinois site. *In* J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- RALPH, C. J., S. DROEGE, AND J. R. SAUER. 1995. Managing and monitoring birds using point counts: standards and applications. Pp. 161-175 *in* C. J. Ralph, J. R. Sauer, and S. Droegge (editors). *Monitoring bird populations by point counts*. USDA Forest Service Gen. Tech. Rep. PSW-149. USDA Forest Service Pacific Southwest Research Station, Albany, CA.
- RALPH, C. J., G. R. GEUPEL, P. PYLE, T. E. MARTIN, AND D. F. DESANTE. 1993. *Handbook of field methods for monitoring landbirds*. USDA Forest Service Gen. Tech. Rep. PSW-144. USDA Forest Service Redwood Sciences Laboratory, Arcata, CA.
- REDMOND, R. L., AND M. L. PRATHER. 1996. Mapping existing vegetation and land cover across western Montana and northern Idaho. Executive summary. Wildlife Spatial Analysis Lab, Montana Cooperative Wildlife Research Unit. University of Montana, Missoula, MT.
- REDMOND, R. L., AND WILDLIFE SPATIAL ANALYSIS LAB. 1996. Mapping existing vegetation and land cover across western Montana and northern Idaho. Final Report. USDA Forest Service, Northern Regional Office, Missoula, MT.
- REEVES, B. O. K. 1978. Bison killing in the southwestern Alberta Rockies. *Plains Anthropologist* 23: 63-78.
- REGOSIN, J. V. 1994. Scissor-tailed Flycatchers eject Brown-headed Cowbird eggs. *Journal of Field Ornithology* 65:508-511.
- REYNOLDS, R. T., J. M. SCOTT, AND R. A. NUSSBAUM. 1980. A variable circular-plot method for estimating bird numbers. *Condor* 82:309-313.
- REYNOLDS, T. D. 1981. Nesting of the Sage Thrasher, Sage Sparrow and Brewers Sparrow in southeast Idaho. *Condor* 83: 61-64.
- REYNOLDS, T. D., AND C. H. TROST. 1981. Grazing, crested wheatgrass, and bird populations in southeastern Idaho. *Northwest Science* 55:225-234.
- RICE, L. 1932. The effect of fire on prairie animal communities. *Ecology* 13:392-401.
- RICH, A. C., D. S. DOBKIN, AND L. J. NILES. 1994. Defining forest fragmentation by corridor width: the influence of narrow forest-dividing corridors on forest-nesting birds in southern New Jersey. *Conservation Biology* 8:1109-1121.
- RICH, T. D. 1978. Cowbird parasitism of Sage and Brewers sparrows. *Condor* 80:438.
- RICH, T. D., AND S. I. ROTHSTEIN. 1985. Sage Thrashers reject cowbird eggs. *Condor* 87:561-562.
- RICKLEFS, R. E. 1969. An analysis of nesting mortality in birds. *Smithsonian Contributions in Zoology* 9:1-48.
- RICKLEFS, R. E. 1973. Fecundity, mortality, and avian demography. Pp. 366-435 *in* D. S. Farner (editor). *Breeding biology of birds*. National Academy of Sciences, Washington, D.C.
- RIDGWAY, R. 1880. Report of the geological exploration

- tion of the fortieth parallel made by order of the Secretary of War according to acts of Congress of March 2, 1867, and March 3, 1869, under the direction of A. A. Humphreys, Chief of Engineers. Part III: Ornithology. U.S. Government Printing Office, Washington, D.C.
- RISCH, N. J., AND B. DEVLIN. 1992. On the probability of matching DNA fingerprints. *Science* 255:717-720.
- ROBBINS, C. S., J. R. SAUER, R. S. GREENBERG, AND S. DROEGE. 1989. Population declines in North American birds that migrate to the neo-tropics. *Proceedings of the National Academy of Sciences (USA)* 86:7658-7662.
- ROBERTSON, R. J., AND R. F. NORMAN. 1976. Behavioral defenses to brood parasitism by potential hosts of the Brown-headed Cowbird. *Condor* 78: 166-173.
- ROBERTSON, R. J., AND R. F. NORMAN. 1977. The function and evolution of aggressive host behavior towards the Brown-headed Cowbird (*Molothrus ater*). *Canadian Journal of Zoology* 55:508-518.
- ROBINSON, S. K. 1990. Effects of forest fragmentation on nesting songbirds. *Illinois Natural History Reports* 296:1-2
- ROBINSON, S. K. 1992. Population dynamics of breeding neotropical migrants in a fragmented Illinois landscape. Pp. 408-418 in J. M. Hagan, III, and D. W. Johnston (editors). *Ecology and conservation of neotropical migrant landbirds*. Smithsonian Institution Press, Washington, D.C.
- ROBINSON, S. K., J. A. GRZYBOWSKI, S. I. ROTHSTEIN, M. C. BRITTINGHAM, L. J. PETIT, AND F. R. THOMPSON, III. 1993. Management implications of cowbird parasitization on neotropical migrant songbirds. Pp. 93-102 in D. M. Finch and P. W. Stangel (editors). *Status and management of neotropical migratory birds*. USDA Forest Service Gen. Tech. Rep. RM-229. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- ROBINSON, S. K., J. HOOVER, J. R. HERKERT, AND R. JACK. In press. Cowbird parasitism in a fragmented landscape: effects of tract size, habitat, and abundance of cowbirds and hosts. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- ROBINSON, S. K., S. I. ROTHSTEIN, M. C. BRITTINGHAM, L. J. PETIT, AND J. A. GRZYBOWSKI. 1995a. Ecology of cowbirds and their impact on host populations. Pp. 428-460 in T. E. Martin and D. M. Finch (editors). *Ecology and management of neotropical migratory birds*. Oxford University Press, New York, NY.
- ROBINSON, S. K., AND J. N. M. SMITH. In press. Cowbird parasitization at multiple spatial scales. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- ROBINSON, S. K., F. R. THOMPSON III, T. M. DONOVAN, D. R. WHITEHEAD AND J. FAABORG. 1995b. Forest fragmentation and the regional population dynamics of songbirds. *Science* 267:1987-1990.
- ROBINSON, S. K., AND D. S. WILCOVE. 1994. Forest fragmentation in the temperate zone and its effects on migratory songbirds. *Bird Conservation International* 4:233-249.
- RODENHOUSE, N. L., T. W. SHERRY, AND R. T. HOLMES. 1997. Site-dependent regulation of population size: a new synthesis. *Ecology* 78:2025-2042.
- ROE, F. G. 1970. *The North American buffalo*, 2nd ed. University of Toronto Press, Toronto, ON.
- ROGERS, C. M., M. J. TAITT, J. N. M. SMITH, AND G. J. JONGEJAN. 1997. Nest predation and cowbird parasitism create a population sink in a wetland breeding population of Song Sparrows. *Condor* 99:622-633.
- ROHWER, S., AND C. D. SPAW. 1988. Evolutionary lag versus bill-size constraints: a comparative study of the acceptance of cowbird eggs by old hosts. *Evolutionary Ecology* 2:27-36.
- ROMIG, G. P., AND R. D. CRAWFORD. 1995. Clay-colored Sparrows in North Dakota parasitized by Brown-headed Cowbirds. *Prairie Naturalist* 27:193-203.
- ROSA, S. M., AND M. T. MURPHY. 1994. Trade-offs and constraints on Eastern Kingbird parental care. *Wilson Bulletin* 106:668-678.
- ROSENBERG, K. V., A. A. DHONDT, AND J. D. LOWE. 1996. Lessons from the landscape. *Birdscope* 10:1-3.
- ROSENBERG, K. V., R. D. OHMART, W. C. HUNTER, AND B. W. ANDERSON. 1991. *Birds of the lower Colorado River Valley*. University of Arizona Press, Tucson, AZ.
- ROSENBERG, K. V., AND M. G. RAPHAEL. 1986. Effects of forest fragmentation on vertebrates in Douglas-fir forests. Pp. 263-272 in J. Verner, M. L. Morrison, and C. J. Ralph (editors). *Wildlife 2000: modeling habitat relationships of terrestrial vertebrates*. University of Wisconsin Press, Madison, WI.
- RØSKAFT, E., G. H. ORIANS, AND L. D. BELETSKY. 1990. Why do Red-winged Blackbirds accept eggs of Brown-headed Cowbirds? *Evolutionary Ecology* 4: 35-42.
- RØSKAFT, E., S. ROHWER, AND C. D. SPAW. 1993. Cost of puncture ejection compared with costs of rearing cowbird chicks for Northern Orioles. *Ornis Scandinavica* 24:28-32.
- ROTEBERRY, J. T., AND J. A. WIENS. 1980. Temporal variation in habitat structure and shrubsteppe bird dynamics. *Oecologia* 47:1-9.
- ROTH, R. R., M. S. JOHNSON, AND T. J. UNDERWOOD. 1996. Wood Thrush (*Hylocichla mustelina*). In A. Poole and F. Gill (editors). *The birds of North America*, no. 246. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, D.C.
- ROTH, R. R., AND R. K. JOHNSON. 1993. Long-term dynamics of a Wood Thrush population breeding in a forest fragment. *Auk* 110:37-48.
- ROTHSTEIN, S. I. 1975a. An experimental and teleonomic investigation of avian brood parasitism. *Condor* 77:250-271.
- ROTHSTEIN, S. I. 1975b. Evolutionary rates and host defenses against avian brood parasitism. *American Naturalist* 109:161-176.
- ROTHSTEIN, S. I. 1976. Cowbird parasitism of the Cedar Waxwing and its evolutionary implications. *Auk* 93:498-509.

- ROTHSTEIN, S. I. 1976. Experiments on defenses Cedar Waxwings use against cowbird parasitism. *Auk* 93: 675-691.
- ROTHSTEIN, S. I. 1977. Cowbird parasitism and egg recognition of the Northern Oriole. *Wilson Bulletin* 89:21-32.
- ROTHSTEIN, S. I. 1978. Geographical variation in the nestling coloration of parasitic cowbirds. *Auk* 95: 152-160.
- ROTHSTEIN, S. I. 1982. Successes and failures in avian egg and nestling recognition with comments on the utility of optimality reasoning. *American Zoologist* 22:547-560.
- ROTHSTEIN, S. I. 1990. A model system for coevolution: avian brood parasitism. *Annual Review of Ecology and Systematics* 21:481-508.
- ROTHSTEIN, S. I. 1994. The cowbird's invasion of the Far West: history, causes and consequences experienced by host species. *Studies in Avian Biology* 15: 301-315.
- ROTHSTEIN, S. I. AND T. L. COOK. In press. Cowbird management, host population regulation and efforts to save endangered species. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- ROTHSTEIN, S. I. AND S. K. ROBINSON. 1994. Conservation and coevolutionary implications of brood parasitism by cowbirds. *Trends in Ecology and Evolution* 9:162-164.
- ROTHSTEIN, S. I., AND S. K. ROBINSON (EDITORS). 1998. *Parasitic birds and their hosts*. Oxford University Press, Oxford, U.K.
- ROTHSTEIN, S. I., J. VERNER, AND R. C. FLEISCHER. 1986a. Social dominance, mating and spacing systems, female fecundity, and vocal dialects in captive and free-ranging Brown-headed Cowbirds. *Current Ornithology* 3:127-185.
- ROTHSTEIN, S. I., J. VERNER, AND E. STEVENS. 1980. Range expansion and diurnal changes in dispersion of the Brown-headed Cowbird in the Sierra Nevada. *Auk* 97:253-267.
- ROTHSTEIN, S. I., J. VERNER, AND E. STEVENS. 1984. Radio-tracking confirms a unique diurnal pattern of spatial occurrence in the parasitic Brown-headed Cowbird. *Ecology* 65:77-88.
- ROTHSTEIN, S. I., J. VERNER, E. STEVENS, AND L. V. RITTER. 1987. Behavioral differences among sex and age classes of the Brown-headed Cowbird and their relation to the efficacy of a control program. *Wilson Bulletin* 99:322-337.
- ROTHSTEIN, S. I., D. A. YOKEL, AND R. C. FLEISCHER. 1986b. Social dominance, mating and spacing systems, female fecundity, and vocal dialects in captive and free-ranging Brown-headed Cowbirds. *Current Ornithology* 3:127-185.
- SAAB, V. A., C. E. BOCK, T. D. RICH, AND D. S. DOBKIN. 1995. Livestock grazing effects in western North America. Pp 311-353 in T. Martin and D. Finch (editors). *Ecology and management of neotropical migrant birds*. Oxford University Press, New York, NY.
- SAAB, V. A., AND C. R. GROVES. 1992. Idaho's migratory landbirds: description, habitats and conservation. Idaho Department of Fish and Game, Nongame Wildlife Leaflet No. 10.
- SAAB, V. A., AND T. D. RICH. 1997. Large-scale conservation assessment for neotropical migratory land birds in the interior Columbia River Basin. USDA Forest Service Gen. Tech. Rep. PNW-399. USDA Forest Service, Pacific Northwest Research Station, Portland, OR.
- SABADELL, J. E. 1982. Desertification in the United States. U.S. Bureau of Land Management, Washington, D.C.
- SAETHER, B.-E. 1990. Age-specific variation in reproductive performance of birds. *Current Ornithology* 7:251-283.
- SAS INSTITUTE. 1985. SAS/STAT user's guide, Release 6.03. SAS Institute, Inc., Cary, NC.
- SAS INSTITUTE. 1997. SAS/STAT software: changes and enhancements through Release 6.12. SAS Institute, Inc., Cary, NC.
- SAUER, J. R., AND S. DROEGE. 1992. Geographic patterns in population trends of neotropical migrants in North America. Pp. 26-42 in J. M. Hagan, III, and D. W. Johnston (editors). *Ecology and conservation of neotropical migrant landbirds*. Smithsonian Institution Press, Washington, D.C.
- SAUER, J. R., J. E. HINES, G. GOUGH, I. THOMAS, AND B. G. PETERJOHN. 1997. The North American Breeding Bird Survey: results and analysis. Version 96.4 (<http://www.mbr.nbs.gov/bbs/>). Patuxent Wildlife Research Center, Laurel, MD.
- SAUNDERS, A. A. 1921. A distributional list of the birds of Montana. *Pacific Coast Avifauna* 14. Cooper Ornithological Club, Berkeley, CA.
- SCHMIEGELOW, F. K. A., C. S. MACHTANS, AND S. J. HANNON. 1997. Are boreal birds resilient to forest fragmentation? An experimental study of short-term community responses. *Ecology* 78:1914-1932.
- SCHROEDL, G. F. 1973. The archaeological occurrence of bison in the southern plateau. Washington State University Laboratory of Anthropology, Reports of Investigations 51. Washington State University, Pullman, WA.
- SCHULZ, T. T., AND W. C. LEININGER. 1991. Nongame wildlife communities in grazed and ungrazed montane riparian sites. *Great Basin Naturalist* 51:286-292.
- SCLATER, W. L. 1912. *A history of the birds of Colorado*. Witherby and Co., London, U.K.
- SCOTT, D. M. 1963. Changes in the reproductive activity of the Brown-headed Cowbird within the breeding season. *Wilson Bulletin* 75:123-129.
- SCOTT, D. M. 1977. Cowbird parasitism on the Gray Catbird at London, Ontario. *Auk* 94:18-27.
- SCOTT, D. M., AND C. D. ANKNEY. 1979. Evaluation of a method for estimating the laying rate of Brown-headed Cowbirds. *Auk* 96:483-488.
- SCOTT, D. M., AND C. D. ANKNEY. 1980. Fecundity of the Brown-headed Cowbird in southern Ontario. *Auk* 97:677-683.
- SCOTT, D. M., AND C. D. ANKNEY. 1983. The laying cycle of Brown-headed Cowbirds: passerine chickens? *Auk* 100:583-592.
- SCOTT, D. M., AND R. E. LEMON. 1996. Differential reproductive success of Brown-headed Cowbirds

- with Northern Cardinals and three other hosts. *Condor* 98:259-271.
- SCOTT, P. E., AND B. R. MCKINNEY. 1994. Brown-headed cowbird removes Blue-grey Gnatcatcher nestlings. *Journal of Field Ornithology* 65:363-364.
- SCOTT, T. W. 1979. Growth and age determination of nestling Brown-headed Cowbirds. *Wilson Bulletin* 91:464-466.
- SCURLOCK, D., AND D. M. FINCH. 1997. A historical review. Pp. 43-68 in W. M. Block and D. M. Finch (editors). *Songbird ecology in southwestern ponderosa pine forests: a literature review*. USDA Forest Service Gen. Tech. Rep. RM-292. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- SEALY, S. G. 1992. Removal of Yellow Warbler eggs in association with cowbird parasitism. *Condor* 94:40-54.
- SEALY, S. G. 1994. Observed acts of egg destruction, egg removal, and predation on nests of passerine birds at Delta Marsh, Manitoba. *Canadian Field-Naturalist* 108:41-51.
- SEALY, S. G. 1996. Evolution of host defenses against brood parasitization: implications of puncture-ejection by a small passerine. *Auk* 113:346-355.
- SEALY, S. G., AND R. C. BAZIN. 1995. Low frequency of observed cowbird parasitism on eastern kingbirds: host rejection, effective nest defense, or parasite avoidance? *Behavioral Ecology* 6:140-145.
- SEALY, S.G., J. E. SÁNCHEZ, R. G. CAMPOS, AND M. MARIN. 1997. Bronzed Cowbird hosts: new records, trends in host use, and cost of parasitism. *Ornitologia Neotropical* 8:175-184.
- SEARCY, W. A., AND K. YASUKAWA. 1995. Polygyny and sexual selection in Red-winged Blackbirds. Princeton University Press, Princeton, NJ.
- SEASTEDT, T. R. 1984. Belowground macroarthropods of annually burned and unburned tallgrass prairie. *American Midland Naturalist* 11:405-407.
- SEDGWICK, J. A., AND F. L. KNOFF. 1988. A high incidence of Brown-headed Cowbird parasitism of Willow Flycatchers. *Condor* 90:253-256.
- SERENA, M. 1982. The status and distribution of the Willow Flycatcher (*Empidonax traillii*) in selected portions of the Sierra Nevada, 1982. California Department of Fish and Game, Wildlife Management Branch Administrative Report 82-5.
- SFERRA, S. J., T. E. CORMAN, C. E. PARADZICK, J. W. ROURKE, J. A. SPENCER, AND M. W. SUMNER. 1997. Arizona Partners in Flight Southwestern Willow Flycatcher survey: 1993-1994. Nongame and Endangered Wildlife Program Tech. Rep. 113. Arizona Game and Fish Department, Phoenix, AZ.
- SFERRA, S. J., R. A. MEYER, AND T. E. CORMAN. 1995. Arizona Partners in Flight 1994 Southwestern Willow Flycatcher survey. Nongame and Endangered Wildlife Program Tech. Rep. 69. Arizona Game and Fish Department, Phoenix, AZ.
- SHAKE, W. F. AND J. P. MATTON. 1975. Three years of cowbird control: an effort to save the Kirtland's Warbler. *Jack-Pine Warbler* 53:48-53.
- SHARP, B. E. 1995. Brown-headed Cowbirds and grazing on National Forests in the Pacific Northwest. *Northwestern Naturalist* 76:121-126.
- SHAW, J., AND T. S. CARTER. 1990. Bison movements in relation to fire and seasonality. *Wildlife Society Bulletin* 18:426-430.
- SHEPPARD, J. M. 1996. Nestling Kentucky Warblers and cowbird attacked by Brown-headed Cowbird. *Journal of Field Ornithology* 67:384-386.
- SHERRY, T. W., AND R. T. HOLMES. 1992. Population fluctuations in a long-distance neotropical migrant: demographic evidence for the importance of breeding season events in the American Redstart. Pp. 431-442 in J. M. Hagan, III, and D. W. Johnston (editors). *Ecology and conservation of neotropical migrant landbirds*. Smithsonian Institution Press, Washington, D.C.
- SIMPSON, T. B., P. E. STUART, AND B. V. BARNES. 1990. Landscape ecosystems and cover types of the reserve area and adjacent lands of the Huron Mountain Club. Occasional Papers of the Huron Mountain Wildlife Foundation. No. 4.
- SKAGGS, R. W. 1996. Population size, breeding biology, and habitat of Willow Flycatchers in the Cliff Gila Valley, New Mexico. New Mexico Dept. Game and Fish, Santa Fe, NM.
- SLACK, R. D. 1976. Nest guarding behavior by male gray catbirds. *Auk* 93:292-300.
- SMITH, C. C. 1940. The effect of overgrazing and erosion upon the biota of the mixed grass prairie of Oklahoma. *Ecology* 21:381-397.
- SMITH, J. N. M., T. L. COOK, S. I. ROTHSTEIN, S. K. ROBINSON, AND S. G. SEALY (EDITORS). In press. The biology and management of cowbirds and their hosts. University of Texas Press, Austin, TX.
- SMITH, J. N. M. 1981. Cowbird parasitism, host fitness, and age of the host female in an island Song Sparrow population. *Condor* 83:152-161.
- SMITH, J. N. M., AND P. ARCESE. 1994. Brown-headed Cowbird and an island population of Song Sparrows: a 16-year study. *Condor* 96:916-934.
- SMITH, J. N. M., AND J. R. MERKT. 1980. Development and stability of single-parent family units in the Song Sparrow. *Canadian Journal of Zoology* 58:1869-1875.
- SMITH, J. N. M., AND I. H. MYERS-SMITH. 1998. Spatial variation in parasitization of song sparrows by Brown-headed Cowbirds. Pp. 296-312 in S. I. Rothstein and S. K. Robinson (editors). *Parasitic birds and their hosts*. Oxford University Press, Oxford, U.K.
- SMITH, J. N. M., M. J. TAITT, C. M. ROGERS, P. ARCESE, L. F. KELLER, A. L. E. V. CASSIDY, AND W. M. HOCHACHKA. 1996. A metapopulation approach to the population biology of the Song Sparrow. *Ibis* 138:120-128.
- SOGGE, M. K., T. J. TIBBITTS, AND J. R. PETERSON. 1997. Status and breeding ecology of the Southwestern Willow Flycatcher in the Grand Canyon. *Western Birds* 28:142-157.
- SOKAL, R. R., AND F. J. ROHLF. 1981. *Biometry*, 2nd ed. W. H. Freeman and Co., New York, NY.
- SOKAL, R. A., AND F. J. ROHLF. 1995. *Biometry*, 3rd ed. W. H. Freeman and Co., New York, NY.
- SOLER, M. 1990. Relationships between the Great Spotted Cuckoo *Clamator glandarius* and its corvid hosts in a recently colonized area. *Ornis Scandinavica* 21:212-223.
- SOLER, M., J. J. SOLER, AND J. G. MARTINEZ. 1997.

- Great Spotted Cuckoos improve their reproductive success by damaging magpie host eggs. *Animal Behaviour* 54:1227-1233.
- SOLER, M., J. J. SOLER, J. G. MARTINEZ, AND A. P. MØLLER. 1995. Magpie host manipulation by Great-Spotted Cuckoos: evidence for an avian *mafia*? *Evolution* 49:770-775.
- SOULÉ, M. E., A. C. ALBERTS, AND D. T. BOLGER. 1992. The effects of fragmentation on chaparral plants and vertebrates. *Oikos* 63:39-47.
- SOULÉ, M. E., D. T. BOLGER, A. C. ALBERTS, J. WRIGHT, M. SORICE, AND S. HILL. 1988. Reconstructed dynamics of rapid extinctions of chaparral-requiring birds in urban habitat islands. *Conservation Biology* 2:75-92.
- SOUTHERN, W. E. 1958. Nesting of the Red-eyed Vireo in the Douglas Lake region, Michigan. *Jack-Pine Warbler* 36:105-130, 185-207.
- SPENCER, J. A., S. J. SFERRA, T. E. CORMAN, J. W. ROURKE AND M. W. SUMNER. 1996. Arizona Partners in Flight 1995 Southwestern Willow Flycatcher survey. Nongame and endangered Wildlife Program Tech. Rep. 97. Arizona Game and Fish Department, Phoenix, AZ.
- SPENCER, R. A. 1985. Brown-headed Cowbird feeding incidents. C. F. O. [Colorado Field Ornithologists] *Journal* 19:39.
- SPSS. 1992. SPSS/PC+ advanced statistics, version 5.0. SPSS Inc., Chicago, IL.
- SPSS. 1996. SPSS Base 7.0 for Windows users guide. SPSS Inc., Chicago, IL.
- SPSS. 1997. Sigmplot 4.0 for Windows Users Manual. SPSS Inc., Chicago, IL.
- STAAB, C. A. 1995. Host and nest selection by Brown-Headed cowbirds within a riparian area in central Arizona. M.S. thesis. University of Arizona, Tucson, AZ.
- STAMPS, J. 1994. Territorial behavior: testing the assumptions. *Advances in the Study of Behavior* 23: 173-232.
- STANSBURY, H. 1852. Exploration and survey of the valley of Great Salt Lake of Utah. Lippincott, Grambo and Co., Philadelphia, PA.
- STARFIELD, A. M. 1997. A pragmatic approach to modeling for wildlife management. *Journal of Wildlife Management* 61:261-270.
- STEELE, B. M., J. C. WINNE, AND R. L. REDMOND. 1998. Estimation and mapping of misclassification probabilities for thematic land cover maps. *Remote Sensing of Environment*. 66:192-202.
- STEIDL, R. J., J. P. HAYES, AND E. SCHAUBER. 1997. Statistical power analysis in wildlife research. *Journal of Wildlife Management* 61:270-279.
- STEVENS, D. R. 1980. The deer and elk of Rocky Mountain National Park: a 10 year study. Report to Rocky Mountain National Park, ROMO-N-13.
- STEWART, R. E. 1953. A life study history of the Yellowthroat. *Wilson Bulletin* 65:99-115.
- STEWART, R. M., R. P. HENDERSON, AND K. DARLING. 1977. Breeding ecology of the Wilson's Warbler in the high Sierra Nevada, California. *Living Bird* 16: 83-102.
- STOLESON, S. H., M. J. WHITFIELD, AND M. K. SOGGE. In press. Demography of the Southwestern Willow Flycatcher. Chapter 7 in D. M. Finch, R. Perriman, S. H. Stoleson, S. J. Sferra, and M. J. Whitfield (editors). Southwestern Willow Flycatcher conservation assessment. Final report. USDA Forest Service, Rocky Mountain Research Station, Albuquerque, NM.
- STRAUSBERGER, B. M., AND M. V. ASHLEY. 1997. Community-wide patterns of parasitization of a host "generalist" brood-parasitic cowbird. *Oecologia* 112:254-262.
- STRIBLEY, J. M. 1993. Factors influencing cowbird distributions in forested landscapes of northern Michigan. M.S. thesis. Michigan State University, East Lansing, MI.
- STUTCHBURY, B. 1997. Effects of female cowbird removal on reproductive success of Hooded Warblers. *Wilson Bulletin* 109:74-81.
- STUTCHBURY, B., J. M. RHYMER, AND E. S. MORTON. 1994. Extra-pair paternity in the Hooded Warbler. *Behavioral Ecology* 5:384-339.
- SULLIVAN, K. A. 1988. Ontogeny of time budgets in Yellow-eyed Juncos: adaptations to ecological constraints. *Ecology* 69:118-124.
- SWARTH, H. S. 1914. A distributional list of the birds of Arizona. Pacific Coast Avifauna No. 10. Cooper Ornithological Club, Hollywood, CA.
- SWENGEL, S. R. 1996. Management responses of three species of declining sparrows in tallgrass prairie. *Bird Conservation International* 6:241-253.
- SYKES, P. W., JR., AND M. H. CLENCH. 1998. Winter habitat of Kirtland's Warbler: an endangered nearctic-neotropical migrant. *Wilson Bulletin* 110:244-261.
- SZARO, R. C. 1989. Riparian forest and scrubland community types of Arizona and New Mexico. *Desert Plants* 9:70-138.
- TATE, J., JR. 1967. Cowbird removes nestling warbler from nest. *Auk* 84:822.
- TAYLOR, D. M. 1986. Effects of cattle grazing on passerine birds nesting in riparian habitat. *Journal of Range Management* 39:254-258.
- TAYLOR, D. M., AND C. D. LITTLEFIELD. 1986. Willow Flycatcher and Yellow Warbler response to cattle grazing. *American Birds* 40:1169-1173.
- TAZIK, D. J., AND J. D. CORNELIUS. 1993. Status of the Black-capped Vireo population at Fort Hood, Texas. Volume III: Population and nesting ecology. USA-CERL Technical Report EN-94/01, Vol III. U.S. Army Construction Engineering Laboratories, Champaign, IL.
- TAZIK, D. J., J. D. CORNELIUS, AND C. A. ABRAHAMSON. 1993. Status of the Black-capped Vireo at Fort Hood, Texas. Volume I: Distribution and abundance. USACERL Technical Report EN-94/01, Vol I. U.S. Army Construction Engineering Laboratories, Champaign, IL.
- TEATHER, K. L., AND R. J. ROBERTSON. 1985. Female spacing patterns in Brown-headed Cowbirds. *Canadian Journal of Zoology* 63:218-222.
- TELFAIR, R. C., II. 1994. Cattle Egret (*Bubulcus ibis*). In A. Poole and F. Gill (editors). *The birds of North America*, no. 113. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, D.C.
- TELFER, E. S. AND J. P. KELSALL. 1984. Adaptations of

- some large North American mammals for survival in snow. *Ecology* 65:1828–1834.
- TEMPLE, S. A., AND J. R. CARY. 1988. Modeling dynamics of habitat-interior bird populations in fragmented landscapes. *Conservation Biology* 2:340–347.
- TERBORGH, J. 1989. *Where have all the birds gone?* Princeton University Press, Princeton, NJ.
- TESTER, J. R., AND W. H. MARSHALL. 1961. A study of certain plant and animal interrelationships on a prairie in northwestern Minnesota. University of Minnesota, Minneapolis, MN.
- TEWKSBURY, J. J., S. J. HEJL, AND T. E. MARTIN. 1998. Breeding productivity does not decline with increasing fragmentation in a western landscape. *Ecology* 79:2890–2903.
- THOMAS, J. W., C. MASER, AND J. E. REDOIK. 1979. Riparian zones. Pp. 40–47 in J. W. Thomas (editor). *Wildlife habitats in managed forests: the Blue Mountains of Oregon and Washington*. Agricultural Handbook No. 553. USDA Forest Service, Washington, D.C.
- THOMAS, L. 1997. Retrospective power analysis. *Conservation Biology* 11:276–280.
- THOMPSON, C. F., AND V. NOLAN, JR. 1973. Population biology of the Yellow-breasted Chat (*Icteria virens* L.) in southern Indiana. *Ecological Monographs* 43:145–171.
- THOMPSON, F. R., III. 1993. Simulated response of a forest-interior bird population to forest management options in central hardwoods forests of the United States. *Conservation Biology* 7:325–333.
- THOMPSON, F. R., III. 1994. Temporal and spatial pattern of breeding in Brown-headed Cowbirds in the midwestern United States. *Auk* 111:979–990.
- THOMPSON, F. R., III, AND W. D. DIJAK. In press. Differences in movements, home range, and habitat preferences of female Brown-headed Cowbirds in three midwestern landscapes. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- THOMPSON, F. R., III, W. D. DIJAK, T. G. KULOWIEC, AND D. A. HAMILTON. 1992. Breeding bird populations in Missouri Ozark forests with and without clearcutting. *Journal of Wildlife Management* 56:23–30.
- THOMPSON, F. R., III, S. K. ROBINSON, T. M. DONOVAN, J. FAABORG, AND D. R. WHITEHEAD. In press. Biogeographic, landscape, and local factors affecting cowbird abundance and host parasitism levels. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- TONHASCA, A., JR., AND D. N. BYRNE. 1994. The effects of crop diversification on herbivorous insects: a meta-analysis approach. *Ecological Entomology* 19:239–244.
- TRAIL, P. W. 1992. Nest invaders. *Pacific Discovery Summer 1992*: 32–37.
- TRAIL, P. W., AND L. F. BAPTISTA. 1993. The impact of Brown-headed Cowbird parasitism on populations of the Nuttall's White-crowned Sparrow. *Conservation Biology* 7:309–315.
- TRAUTMAN, M. B. 1940. The birds of Buckeye Lake, Ohio. *Miscellaneous Publications of University of Michigan Museum of Zoology* 44:1–466.
- TRINE, C. L. 1998. Wood Thrush population sinks and implications for the scale of regional conservation strategies. *Conservation Biology* 12:576–585.
- TRINE, C. L. In press. Effects of multiple parasitism on cowbird and Wood Thrush nesting success. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- TRINE, C. L., W. D. ROBINSON, AND S. K. ROBINSON. 1998. Consequences of Brown-headed Cowbird brood parasitization for host population dynamics. Pp. 273–295 in S. I. Rothstein and S. K. Robinson (editors). *Parasitic birds and their hosts*. Oxford University Press, Oxford, U.K.
- TUCKER, K., S. P. RUSHTON, R. A. SANDERSON, E. B. MARTIN, AND J. BLAIKLOCK. 1997. Modelling bird distributions—a combined GIS and Bayesian rule-based approach. *Landscape Ecology* 12:77–93.
- TULJAPURKAR, S. D., AND S. H. ORZACK. 1980. Population dynamics in variable environments I. Long-run growth rates and extinction. *Theoretical Population Biology* 18:314–342.
- TWOMEY, A. C. 1945. The bird population of an elm-maple forest with special reference to aspektion, territorialism, and coactions. *Ecological Monographs* 15:173–205.
- U.S. DEPARTMENT OF COMMERCE. 1979. *Climate atlas of the United States*. National Oceanic and Atmospheric Administration, Washington, D.C.
- U.S. FISH AND WILDLIFE SERVICE. 1973. Building and operating a decoy trap for live-capturing cowbirds and other birds. Report AC-211. U.S. Fish and Wildlife Service, Twin Cities, MN.
- U.S. FISH AND WILDLIFE SERVICE. 1984. Starling control with live traps. Report GPO 794–791. U.S. Fish and Wildlife Service, Olympia, WA.
- U.S. FISH AND WILDLIFE SERVICE. 1986. Final rule determining endangered status for the Least Bell's Vireo. *Federal Register* 51(85):16474–16482.
- U.S. FISH AND WILDLIFE SERVICE. 1991. Black-capped Vireo (*Vireo atricapillus*) recovery plan. U.S. Fish and Wildlife Service, Austin, TX.
- U.S. FISH AND WILDLIFE SERVICE. 1992. Golden-cheeked Warbler recovery plan. U.S. Fish and Wildlife Service, Albuquerque, NM.
- U.S. FISH AND WILDLIFE SERVICE. 1993a. Biological opinion. U.S. Fish and Wildlife Service, Austin, TX.
- U.S. FISH AND WILDLIFE SERVICE. 1993b. Proposed Rule to list the Southwestern Willow Flycatcher as endangered with critical habitat. *Federal Register* 58:39495–39522.
- U.S. FISH AND WILDLIFE SERVICE. 1995. Final rule determining endangered status for the Southwestern Willow Flycatcher. *Federal Register* 60(38):10694–10715.
- U.S. FISH AND WILDLIFE SERVICE. 1998. Draft recovery plan for the Least Bell's Vireo. U.S. Fish and Wildlife Service, Portland, OR.

- UNITT, P. 1987. *Empidonax traillii extimus*: an endangered subspecies. *Western Birds* 18:137-162.
- UYEHARA, J. C., AND P. M. NARINS. 1995. Nest defense by Willow Flycatchers to brood-parasitic intruders. *Condor* 97: 361-368.
- UYEHARA, J. C., AND M. J. WHITFIELD. In press. Cowbird parasitism and vegetative cover in territories of Southwestern Willow Flycatchers. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.
- UYEHARA, J. C., M. J. WHITFIELD, AND L. GOLDWASSER. In press. The ecology of Brown-headed Cowbirds and their effects on Southwestern Willow Flycatchers. Chapter 8 in D. M. Finch, R. Perriman, S. H. Stoleson, S. J. Sferra, and M. J. Whitfield (editors). *Southwestern Willow Flycatcher conservation assessment*. Final report. USDA Forest Service Rocky Mountain Research Station, Albuquerque, NM.
- VAN TIENDEREN, P. H. 1995. Life cycle trade-offs in matrix population models. *Ecology* 76:2482-2489.
- VAN VUREN, D. 1983. Group dynamics and summer home range of bison in southern Utah. *Journal of Mammalogy* 64:329-332.
- VAN VUREN, D., AND M. P. BRAY. 1985. The recent geographic distribution of *Bison bison* in Oregon. *Murrelet* 66:56-58.
- VAN VUREN, D., AND M. P. BRAY. 1986. Population dynamics of bison in the Henry Mountains, Utah. *Journal of Mammalogy* 67:503-511.
- VERHULST, S., J. H. VAN BALEN, AND J. M. TINBERGEN. 1995. Seasonal decline in reproductive success of the Great Tit: variation in time or quality. *Ecology* 76: 2392.
- VERNER, J. 1985. Assessment of counting techniques. *Current Ornithology* 2:247-302.
- VERNER, J. 1988. Optimizing the duration of point counts for monitoring trends in bird populations. USDA Forest Service Res. Note PSW-395. USDA Forest Service, Pacific Southwest Research Station, Berkeley, CA.
- VERNER, J., AND A. S. BOSS. 1980. California wildlife and their habitats: western Sierra Nevada. USDA Forest Service Gen. Tech. Rep. PSW-37. USDA Forest Service Pacific Southwest Research Station, Berkeley, CA.
- VERNER, J., AND L. V. RITTER. 1983. Current status of the Brown-headed Cowbird in the Sierra National Forest. *Auk* 100:355-368.
- VERNER, J., AND S. I. ROTHSTEIN. 1988. Implications of range expansion into the Sierra Nevada by the parasitic Brown-headed Cowbird. Pp. 92-98 in D. Bradley (editor). *Proceedings, State of the Sierra Symposium 1985-1986*. Pacific Publications, San Francisco, CA.
- VICKERY, P. D., M. L. HUNTER, JR., AND J. V. WELLS. 1992. Evidence of incidental nest predation and its effects on nests of threatened grassland birds. *Oikos* 63:281-288.
- VILLARD, M.-A., K. FREEMARK, AND G. MERRIAM. 1989. Metapopulation theory and neotropical migrant birds in temperate forests: an empirical investigation. Pp. 474-482 in J. M. Hagan, III, and D. M. Johnston (editors). *Ecology and conservation of neotropical migrant birds*. Smithsonian Institution Press, Washington D.C.
- WALKINSHAW, L. H. 1972. Kirtland's Warbler—endangered. *American Birds* 26:3-9.
- WALKINSHAW, L. H. 1983. Kirtland's Warbler: the natural history of an endangered species. Cranbrook Institute of Science, Bloomfield Hills, MI.
- WALTERS, C. 1986. Adaptive management of renewable resources. Macmillan, New York, NY.
- WARD, D., AND J. N. M. SMITH. 1998. Morphological differentiation of Brown-headed Cowbirds in the Okanagan Valley, British Columbia. *Condor* 100: 1-7.
- WARREN, E. R. 1906. Mammals of Colorado. Colorado College Publications, General Series, 19 (Science Series, 46):225-274.
- WARREN, E. R. 1927. Altitude limit of bison. *Journal of Mammalogy* 8:60-61.
- WEATHERHEAD, P. J. 1989. Sex ratios, host-specific reproductive success, and impacts of Brown-headed Cowbirds. *Auk* 106:358-366.
- WEBSTER, M. S., AND D. F. WESTNEAT. 1998. The use of molecular markers to study kinship in birds: techniques and questions. Pp. 7-35 in R. DeSalle and B. Schierwater (editors). *Molecular approaches to ecology and evolution*. Birkhauser Verlag, Boston, MA.
- WEINBERG, H. J., T. J. HAYDEN, AND J. D. CORNELIUS. 1998. Local and installation-wide Black-capped Vireo dynamics on the Fort Hood, Texas, Military Reservation. USACERL Technical Report 98/54. U.S. Army Construction Engineering Laboratories, Champaign, IL.
- WELLS, J., AND J. TURNBULL. 1998. Sensitive species survey results for Pine Creek and Hauser Canyon Wilderness Areas. Cleveland National Forest, San Diego, CA.
- WESTEMEIER, R. L., AND J. E. BUHNERKEMPE. 1983. Responses of nesting wildlife to prairie grass management on prairie chicken sanctuaries in Illinois. Pp. 36-46 in R. Brewer (editor). *Proceedings of the Eighth North American Prairie Conference*. Western Michigan University, Kalamazoo, MI.
- WESTMAN, W. E. 1981. Diversity relationships and succession in California coastal sage scrub. *Ecology* 62: 170-184.
- WESTMAN, W. E. 1983. Xeric Mediterranean-type shrubland associations of Alta and Baja California and the community/continuum debate. *Vegetatio* 52: 3-19.
- WESTNEAT, D. F. 1990. Genetic parentage in Indigo Buntings: a study using DNA fingerprinting. *Behavior Ecology and Sociobiology* 27:67-76.
- WHITE, J. M. 1973. Breeding biology and feeding patterns of the Oregon Junco in two Sierra Nevada habitats. Ph.D. dissertation. University of California, Berkeley, CA.
- WHITFIELD, M. J. 1990. Willow Flycatcher reproductive response to Brown-headed Cowbird parasitism. M.S. thesis. California State University, Chico, CA.
- WHITFIELD, M. J. In press. Results of a Brown-headed Cowbird program for the Southwestern Willow Flycatcher. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). *The biology and management of cowbirds and their hosts*. University of Texas Press, Austin, TX.

- WHITFIELD, M. J., AND K. M. ENOS. 1996. A Brown-headed Cowbird control program and monitoring for the Southwestern Willow Flycatcher, South Fork Kern River, California, 1996. Report: U.S. Army Corps of Engineers, Sacramento District, Environmental Resources Division. Purchase order DACW05-96-P-0900.
- WHITFIELD, M. J., AND C. M. STRONG. 1995. A Brown-headed Cowbird control program and monitoring for the Southwestern Willow Flycatcher, South Fork Kern River, CA, 1995. California Department of Fish and Game Bird and Mammal Conservation Program, Report 95-4.
- WIEDENFELD, D. A. In press. Cowbird population changes and their relationship to changes in some host species. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). The biology and management of cowbirds and their hosts. University of Texas Press, Austin, TX.
- WIENS, J. A. 1963. Aspects of cowbird parasitism in southern Oklahoma. *Wilson Bulletin* 75:130-138.
- WIENS, J. A. 1974. Climatic instability and the "ecological saturation" of bird communities in North American grasslands. *Condor* 76:385-400.
- WIENS, J. A. 1992. The ecology of bird communities. Vol. 2: processes and variations. Cambridge University Press, Cambridge, U.K.
- WILCOVE, D. 1985. Nest predation in forest tracts and the decline of migratory songbirds. *Ecology* 66:1211-1214.
- WILKINSON, L. 1997. SYSTAT: the system for statistics. SYSTAT Inc., Evanston, IL.
- WINTERNITZ, B. L., AND D. W. CRUMPACKER (EDITORS). 1985. Species of special concern. Unpubl. Rep., Colorado Wildlife Workshop, Denver, CO.
- WISDOM, M. J., AND L. S. MILLS. 1997. Using sensitivity analysis to guide population recovery: prairie chickens as an example. *Journal of Wildlife Management* 61:302-312.
- WITTENBERGER, J. F. 1981. Animal social behavior. Duxbury Press, Boston, MA.
- WOLF, L. 1987. Host-parasite interactions of Brown-headed Cowbirds and Dark-eyed Juncos in Virginia. *Wilson Bulletin* 99:338-350.
- WOODBURY, A. M., AND N. H. RUSSELL, JR. 1945. Birds of the Navajo Country. University of Utah, Salt Lake City, UT.
- WOODWARD, P. W. 1983. Behavioral ecology of fledgling Brown-headed Cowbirds and their hosts. *Condor* 85:151-163.
- WOODWARD, P. W., AND J. C. WOODWARD. 1979. Survival of fledgling Brown-headed Cowbirds. *Bird Banding* 50:66-68.
- WOODWORTH, B. L. 1997. Brood parasitism, nest predation, and season-long reproductive success of a tropical island endemic. *Condor* 99:605-621.
- WOOTTON, J. T. 1996. Purple Finch (*Carpodacus purpureus*). In A. Poole and F. Gill (editors). The birds of North America, no. 208. Academy of Natural Sciences, Philadelphia, PA, and American Ornithologists' Union, Washington, D.C.
- WOOTTON, J. T., AND D. A. BELL. 1992. A metapopulation model of the Peregrine Falcon in California: viability and management strategies. *Ecological Applications* 2:307-321.
- YAHNER, R. H., AND C. A. DELONG. 1992. Avian predation and parasitism on artificial nests and eggs in two fragmented landscapes. *Wilson Bull.* 104: 162-168.
- YAMASAKI, M., T. M. MCLELLAN, R. M. DEGRAAF, AND C. A. COSTELLO. In press. Effects of land-use and management practices on the presence of Brown-headed Cowbirds in the White Mountains of New Hampshire and Maine. In J. N. M. Smith, T. L. Cook, S. I. Rothstein, S. K. Robinson, and S. G. Sealy (editors). The biology and management of cowbirds and their hosts. University of Texas Press, Austin, TX.
- YASUKAWA, K., J. L. MCCLURE, R. A. BOLEY, AND J. ZANOCCO. 1990. Provisioning of nestlings by male and female Red-winged Blackbirds, *Agelaius phoeniceus*. *Animal Behaviour* 40:153-166.
- YOKEL, D. A. 1989. Intrasexual aggression and mating behavior of Brown-headed Cowbird. *Condor* 91:43-51.
- YOUNG, B. E. 1991. Annual molts and interruption of the fall migration for molting in Lazuli Buntings. *Condor* 93: 236-250.
- ZAHAVI, A. 1979. Parasitism and nest predation in parasitic cuckoos. *American Naturalist* 113:157-159.
- ZAR, J. H. 1984. Biostatistical analysis, 2nd ed. Prentice Hall, Englewood Cliffs, NJ.
- ZIMMERMAN, J. L. 1983. Cowbird parasitism of Dickcissels in different habitats and at different nest densities. *Wilson Bulletin* 95:7-22.
- ZIMMERMAN, J. L. 1992. Density-independent factors affecting the avian diversity of the tallgrass prairie community. *Wilson Bulletin* 104:85-94.