Resource Evaluation Program Report

Geological & Geophysical Data Acquisition

Outer Continental Shelf Through 1999

By George Dellagiarino, Patricia Fulton, Keith Meekins, and David Zinzer

Contents

| Acronyms and Abbreviations | V |
|--|----|
| Introduction | 1 |
| Permits, Data Acquisition, and Reimbursement | 3 |
| Geophysical Data Surveys | 3 |
| Geological Data Collection | 4 |
| Bottom Sampling and Shallow Coring | |
| Deep Stratigraphic Tests | |
| Bibliography | 12 |
| Appendix | 13 |
| Alaska Tables for Data Acquisitions, Permits, Expenditures | |
| Atlantic Tables for Data Acquisitions, Permits, Expenditures | 19 |
| Gulf of Mexico Tables for Data Acquisitions, Permits, Expenditures | 23 |
| Pacific Tables for Data Acquisitions, Permits, Expenditures | |
| Glossary | 31 |

Illustration

Tables

| 1. | Summary of estimates of CDP (2-D) seismic miles in MMS inventory through FY 1999 by planning area | 5 |
|----|--|----|
| 2. | Summary of 2-D seismic data acquisition for FY 1968-1999 | 6 |
| 3. | Summary of 3-D seismic data acquisition for FY 1968-1999 | 6 |
| 4. | Summary of geological and geophysical data acquisition by data type and region, FY 1968-1999 | 7 |
| 5. | Total number of permits issued for geological and geophysical exploration | 8 |
| 6. | Summary of total annual expenditures by MMS for geological and geophysical data acquisition by region, FY 1968-1999 (in dollars) | 9 |
| 7. | Summary of geological and geophysical data acquisition expenditures by data type and region, FY 1968-1999 | 10 |
| 8. | Summary of average cost per mile by MMS for 2-D seismic data, FY 1968-1999 | 11 |

Acronyms and Abbreviations

CDP Common depth point seismic data

CFR Code of Federal Regulations

COST Continental Offshore Stratigraphic Test

DOI Department of the Interior

DST Deep stratigraphic test (well)

FY Fiscal Year

G&G Geological and geophysical

GOM Gulf of Mexico

GRAV Gravity data

HRD High-resolution seismic data

MAG Magnetics data

MMS Minerals Management Service

OCS Outer Continental Shelf

OCSLA Outer Continental Shelf Lands Act

RE Resource Evaluation

REP Resource Evaluation Program

SEG Society of Exploration Geophysicists

2-D Two-dimensional seismic data

3-D Three-dimensional seismic data

4-D Four-dimensional seismic data

Introduction

This report addresses the general role of the MMS Resource Evaluation Program (REP) in geological and geophysical (G&G) data acquisition and permitting activities.

The Minerals Management Service (MMS) administers the provisions of the Outer Continental Shelf Lands Act (OCSLA) through regulations found at Title 30 of the Code of Federal Regulations (CFR). The regulations govern permitting, data acquisitions and release, leasing, and postlease operations on the Outer Continental Shelf (OCS). The OCS is divided into planning areas for administrative purposes as shown in figure 1.

With regard to the REP, authority has been vested in the Secretary of the Interior under 30 CFR Part 251 to regulate prelease G&G exploration for oil, gas, and sulphur resources on the OCS. Part 251 applies not only to G&G exploration but to scientific research as well. The purpose of these regulations is to prescribe (1) when a permit or the filing of a notice is required to conduct G&G activities on the OCS and (2) operating procedures for conducting exploration, as well as requirements for disclosing data and information, conditions for reimbursing permittees for certain costs, and other conditions under which exploration shall be conducted. Similar regulations addressing prelease prospecting activities for minerals other then oil, gas, or sulphur can be found in 30 CFR Part 280.

In this report, the totals for permits issued, mileage acquired, and expenditures reflect the overall trends of oil and gas pricing, limitations of areas due to offshore moratoria, and the shift of industry emphasis to foreign theatres. Also reflected is the trend among the MMS Regions with diminished leasing activity to obtain digital tapes of in-house analog data in preparation for data release commencing with the expiration of proprietary terms in 2001 as discussed by Fulton (1998).




Figure 1. Federal Outer Continental Shelf Planning Areas

Permits, Data Acquisition, and Reimbursement

The permits, issued by the Resource Evaluation (RE) Regional Supervisors, set forth the specific details for each data-gathering activity, which include the area where the data are collected, the timing of the data-gathering activity, approved equipment and methods, and other similar detailed information relevant to each specific permit. The primary source of the G&G data used by the REP is the oil and gas industry, which conducts exploration, development, and production activities on OCS lands. While the MMS does not perform any direct datacollection activities, it does issue permits to industry for collecting prelease G&G data. Lessees and operators are also required by regulations to provide data from their leases to MMS. The MMS has access to the permitted data and information as a condition set forth in the permit. The MMS selectively obtains copies from these activities. Data from prelease permits constitute approximately 90 percent of the MMS database. Reimbursement for data collected by permittees and lessees in their normal conduct of business is for only the cost of data reproduction. However, if industry has collected data in areas not under MMS jurisdiction, e.g., State waters or adjacent foreign waters, and MMS selects such data, MMS pays the significantly higher "market price" for obtaining such data.

Geophysical Data Surveys

Common Depth Point, Gravity, Magnetic, 3-D, and 4-D Surveys

A large percentage of the geophysical data in the MMS inventory is two-dimensional (2-D) common depth point (CDP) seismic information. Also known as common midpoint or common reflection point data, it is derived from a common location in the ocean subbottom where sound waves originating from various positions of the seismic (sound) source near the ocean surface are reflected back toward the surface. Estimates of the amount of these data in the MMS inventory by planning area are shown in table 1.

Magnetic surveys measure the magnetic field or a component (such as the vertical component) at a series of different locations over an area of interest usually to locate concentrations of magnetic anomalies or to determine depth to basement.

Gravity surveys produce measurements of the gravitational field at a series of different locations over an area of interest. The objective in exploration work is to map density differences that may indicate different rock types. Gravity data usually are displayed as anomaly maps.

While the main use of three-dimensional (3-D) seismic data is still in reservoir development, the evolution of 3-D seismic data and information in conjunction

with interactive computer workstations has made it possible to more closely define and assess the potential for oil and gas occurrence on the OCS, especially with regard to subsalt prospects. The 3-D information is used to delineate, in greater detail than that of traditional 2-D information, subsurface geologic conditions associated with the occurrence of natural gas and oil.

As 3-D seismic technology evolved, the use of 3-D reflection techniques not only portrayed subsurface structure and stratigraphy but revealed information about fluids within the subsurface. Three dimensional seismic surveys that are shot over the same area at different times can detect, where present, changes from one fluid to another, e.g., oil to water, fluid flows, and pressure changes.

Thus timelapse 3-D seismic surveys, known more commonly as 4-D seismic surveys, have been used to monitor fluid movement in producing reservoirs where changes in fluid content are imaged with seismic techniques over a period of time. Its chief use to date has been reservoir management, e.g., determining where and how long to drain hydrocarbon-bearing areas and to monitor gas injection or steam or water flooding during enhanced recovery operations.

The proprietary term for geophysical information (i.e., seismic sections, etc.) purchased by MMS is 25 years; for raw geophysical data it is 50 years.

Geological Data Collection

Bottom Sampling and Shallow Coring

Bottom samples are obtained by dropping a weighted tube to the ocean floor and recovering it with an attached wire line; they can also be obtained from dredging. Shallow coring is performed by conventional rotary drilling equipment to obtain a near-surface sample of the rocks of the seabed. The proprietary term for these kinds of geological information is 10 years.

Deep Stratigraphic Tests

A deep stratigraphic test, as defined in 30 CFR 251, means "drilling which involves the penetration into the sea bottom of more than 50 feet (15.2 meters) of consolidated rock or a total of more than 300 feet (91.4 meters)." These wells are also known as Continental Offshore Stratigraphic Test (COST) wells and are drilled primarily to gather geological information. Conversely, shallow test drilling, as defined in the same regulations, means "drilling into the sea bottom to depths less than those specified in the definition of a deep stratigraphic test." Three COST wells have encountered hydrocarbons: the COST B-3 (Atlantic), Point Conception No.1 (California), and the Norton COST No. 2 (Alaska). The proprietary term for a COST well is 25 years or, if a lease sale is held in the area, 60 days after the issuance of a lease within 50 geographic miles of the test. A discussion of the cost well program is described by Dellagiarino (1991).

Table 1. Summary of Estimates of CDP (2-D) Seismic Miles in MMS Inventory Through FY 1999 by Planning Area (Rounded off to Nearest 1,000 Miles)

| Planning Area | | Estimated Mileage |
|--|----------------|---|
| | Alaska | |
| Gulf of Alaska Cook Inlet Kodiak Shumagin North Aleutian St. George Basin Aleutian Arc Bowers Basin Aleutian Basin St. Matthew-Hall Norton Basin Navarin Basin Hope Basin Chukchi Sea Beaufort Sea Total | | 36,000 21,000 23,000 10,000 43,000 50,000 < 500 <1,000 10,000 25,000 55,000 9,000 95,000 61,000 440,000 |
| | Atlantic | |
| North Atlantic Mid-Atlantic South Atlantic Straits of Florida Total | | 69,000 60,000 54,000 <u>7,000</u> 190,000 |
| | Gulf of Mexico | |
| Eastern GOM Central GOM Western GOM Total | | 137,000 454,000 <u>309,000</u> 900,000 |
| | Pacific | |
| Southern Calif. Central California Northern Calif. Wash./Oregon Total | | 85,000 21,000 19,000 <u>8,000</u> 133,000 |

Table 2. Summary of 2-D Seismic Data Acquisition for FY 1968-1999

| FY | Total Miles |
|-----------|---------------|
| 1968-1975 | 269,814 |
| 1976 | 108,922 |
| 1977 | 42,808 |
| 1978 | 54,426 |
| 1979 | 31,489 |
| 1980 | 19,400 |
| 1981 | 69,904 |
| 1982 | 79,961 |
| 1983 | 120,743 |
| 1984 | 89,853 |
| 1985 | 71,521 |
| 1986 | 47,287 |
| 1987 | 113,680 |
| 1988 | 78,920 |
| 1989 | 53,494 |
| 1990 | 85,280 |
| 1991 | 40,513 |
| 1992 | 49,191 |
| 1993 | 25,482 |
| 1994 | 7,138 |
| 1995 | 8,930 |
| 1996 | 33,296 |
| 1997 | 39,682 |
| 1998 | 90,981 |
| 1999 | <u>30,135</u> |
| Total | 1,662,850 |

Figures may vary by 1-2%.

Table 3. Summary of 3-D Seismic Data Acquisition for FY 1968-1999

| FY | Total Blocks |
|-----------|--------------|
| 1968-1992 | 0 |
| 1993 | 1,563 |
| 1994 | 1,420 |
| 1995 | 1,826 |
| 1996 | 1,470 |
| 1997 | 3,129 |
| 1998 | 3,460 |
| 1999 | <u>3,226</u> |
| Total | 16,094 |

Table 4. Summary of Geological and Geophysical Data Acquisition by Data Type and Region, FY 1968-1999

| Data Type | Region | Mileage * | |
|------------------------|----------------|--------------------------|--|
| 2-D Seismic | Alaska | 440,179 | |
| | Atlantic | 189,858 | |
| | Gulf of Mexico | 899,972 | |
| | Pacific | <u>132,841</u> | |
| | Total | 1,662,850 | |
| High Resolution | Alaska | 59,855 | |
| _ | Atlantic | 49,509 | |
| | Gulf of Mexico | 145,768 | |
| | Pacific | <u>30,582</u> | |
| | Total | 285,714 | |
| CDP | Alaska | 84,683 | |
| Interpretations | Atlantic | 44,801 | |
| | Gulf of Mexico | 139,418 | |
| | Pacific | <u>42,365</u> | |
| | Total | 311,267 | |
| Gravity and | Alaska | 370,849 | |
| Magnetics | Atlantic | 15,783 | |
| | Gulf of Mexico | 306,247 | |
| | Pacific | <u>110,150</u> | |
| | Total | 803,029 | |
| 3-D Seismic | Alaska | 10 | |
| | Atlantic | 0 | |
| | Gulf of Mexico | 16,043 | |
| | Pacific | <u>41</u> | |
| | Total | 16,094 | |
| Deep | Alaska | 14 | |
| Stratigraphic Atlantic | | 5 | |
| Tests | Gulf of Mexico | 3 | |
| | Pacific | 5 3 <u>2</u> 24 | |
| | Total | 24 | |

 $^{^{\}star}$ 3-D seismic is measured in blocks and Deep Stratigraphic Test units are wells drilled. bl = Blocks

Table 5. Total Number of Permits Issued for Geological and Geophysical Exploration

| | | | | | E | F | G |
|-------|--------------------|-------------------|-----------|----------|-----------|------------|---------------|
| 1960- | 2,353 | | | | | | |
| 1968 | 0=0 | 0.40 | | | | | • |
| 1969 | 258 | 249 | 9 | 0 | 0 | 0 | 0 |
| 1970 | 213 | 203 | 10 | 0 | 0 | 0 | 0 |
| 1971 | 210 | 205 | 5 | 0 | 0 | 0 | 0 |
| 1972 | 220 | 210 | 10 | 0 | 0 | 0 | 0 |
| 1973 | 339 | 321 | 18 | 0 | 0 | 0 | 0 |
| 1974 | 357 | 345 | 12 | 2 | 0 | 0 | 0 |
| 1975 | 510 | 487 | 23 | 3 | 0 | 0 | 0 |
| 1976 | 420 | 400 | 20 | 7 | 0 | 0 | 0 |
| 1977 | 452 | 436 | 16 | 4 | 0 | 0 | 0 |
| 1978 | 342 | 329 | 13 | 2 | 0 | 0 | 0 |
| 1979 | 276 | 265 | 11 | 0 | 0 | 0 | 0 |
| 1980 | 318 | 302 | 16 | 1 | 0 | 0 | 0 |
| 1981 | 394 | 383 | 11 | 0 | 0 | 0 | 0 |
| 1982 | 502 | 490 | 12 | 3 | 0 | 0 | 0 |
| 1983 | 574 | 542 | 32 | 1 | 16 | 0 | 0 |
| 1984 | 543 | 518 | 25 | 0 | 18 | 0 | 0 |
| 1985 | 398 | 382 | 16 | 0 | 38 | 0 | 0 |
| 1986 | 211 | 207 | 4 | 0 | 32 | 0 | 0 |
| 1987 | 298 | 282 | 16 | 0 | 42 | 0 | 0 |
| 1988 | 313 | 289 | 24 | 0 | 45 | 0 | 0 |
| 1989 | 249 | 237 | 12 | 1 | 47 | 0 | 0 |
| 1990 | 251 | 241 | 9 | 0 | 57 | 1 | 0 |
| 1991 | 170 | 156 | 12 | 0 | 45 | 2 | 0 |
| 1992 | 141 | 137 | 3 | 0 | 53 | 1 | 0 |
| 1993 | 147 | 135 | 11 | 0 | 70 | 1 | 0 |
| 1994 | 133 | 117 | 16 | 0 | 53 | 0 | 0 |
| 1995 | 104 | 92 | 11 | 0 | 50 | 1 | 0 |
| 1996 | 136 | 120 | 16 | 0 | 59 | 0 | 0 |
| 1997 | 159 | 139 | 20 | 0 | 69 | 0 | 1 |
| 1998 | 157 | 143 | 14 | 0 | 59 | 0 | 1 |
| 1999 | <u>111</u> | <u>98</u> | <u>13</u> | <u>0</u> | <u>44</u> | <u>0</u> | |
| Total | 11, 259 | 8,4 60 | 440 | 24 | 797 | <u>-</u> 6 | <u>0</u> 2 |

A=Total Number of Geological, Geophysical, and Strategic Minerals Permits

B=Number of Geophysical Permits

C=Number of Geological Permits

D=Number of Geological Permits Issued for Deep Stratigraphic Tests

E=Number of Geophysical Permits Issued for 3-D Seismic Data

F=Number of Permits Issued for Strategic (Nonenergy) Minerals

G=Number of Permits Issued for 4-D Seismic Data

Dashed lines = Individual breakouts not established

Figures may vary 1-2%.

Table 6. Summary of Total Annual Expenditures by MMS for Geological and Geophysical Data Acquisition by Region, FY 1968-1999 (in dollars)

| | | | Gulf of | | |
|-------|----------------|------------|------------------|---------------|------------------|
| FY | Alaska | Atlantic | Mexico | Pacific | Total |
| 1968- | 3,162,548 | 361,686 | 9,414,042 | 1,443,987 | 14,891,780* |
| 1975 | | • | | | |
| 1976 | 3,496,607 | 2,504,710 | 3,281,698 | 581,670 | 9,914,882* |
| 1977 | 450,161 | 2,287,390 | 3,764,678 | 1,147,968 | 7,719,974* |
| 1978 | 3,421,269 | 906,989 | 1,842,701 | 416,463 | 6,587,422 |
| 1979 | 6,240,687 | 232,085 | 1,573,094 | 2,272,407 | 11,020,298* |
| 1980 | 6,972,885 | 4,469,762 | 4,388,508 | 1,412,062 | 17,243,217 |
| 1981 | 6,842,045 | 1,530,898 | 1,168,618 | 866,656 | 10,408,217 |
| 1982 | 1,864,661 | 1,945,270 | 2,943,602 | 1,996,271 | 8,749,804 |
| 1983 | 5,673,514 | 1,738,427 | 3,802,409 | 1,312,596 | 12,526,946 |
| 1984 | 4,751,354 | 1,580,008 | 4,246,742 | 1,286,598 | 11,864,702 |
| 1985 | 3,676,375 | 318,261 | 2,959,989 | 861,687 | 7,816,312 |
| 1986 | 2,904,246 | 87,307 | 1,834,553 | 363,564 | 5,189,670 |
| 1987 | 2,579,190 | 438,792 | 1,840,609 | 939,558 | 5,798,149 |
| 1988 | 1,382,560 | 71,510 | 1,078,713 | 114,168 | 2,646,951 |
| 1989 | 389,960 | 259,629 | 913,481 | 96,354 | 1,659,424 |
| 1990 | 886,402 | 150 | 865,083 | 0 | 1,751,635 |
| 1991 | 539,986 | 2,790 | 1,003,066 | 31,000 | 1,576,842 |
| 1992 | 99,797 | 1,932 | 794,104 | 0 | 1,490,798** |
| 1993 | 322,410 | 0 | 1,014,853 | 26,700 | 1,363,963 |
| 1994 | 582,132 | 0 | 760,245 | 11,806 | 1,454,183** |
| 1995 | 379,395 | 0 | 628,752 | 21,125 | 1,142,817** |
| 1996 | 283,764 | 0 | 1,697,494 | 40,867 | 2,022,125 |
| 1997 | 204,655 | 0 | 1,180,893 | 19,594 | 1,471,967** |
| 1998 | 278,606 | 0 | 1,804,694 | 10,264 | 2,094,400** |
| 1999 | <u>543,775</u> | <u>0</u> | <u>1,400,781</u> | <u>13,350</u> | <u>1,957,906</u> |
| Total | 57,928,984 | 18,737,596 | 56,203,402 | 15,286,715 | 150,364,384 |
| | | | | | |

^{*} Included in the budget for these years were General Account funds that were transferred to the U.S. Geological Survey, Branch of Marine Geology, for G&G data acquisition. These accounts included \$509,517 in the interval between FY 1968-1975, \$50,197 in FY 1976, \$69,777 in FY 1977, and \$702,025 in FY 1979.

^{**}Included in the budget for these years were funds that were used for special projects related to G&G activities. In FY 1992, \$494,965 was obligated toward the purchase of geologic interpretive workstations and \$100,000 was obligated toward the initiation of the Offshore Northern Gulf of Mexico Oil and Gas Atlas Series. In FY 1994, \$100,000 was again obligated toward the preparation of the Offshore Northern Gulf of Mexico Oil and Gas Atlas and in FY 1995, \$100,000 was obligated toward the completion of the Atlas, and \$13,545 was obligated toward finalization of a well log data conversion contract in the Gulf of Mexico. In FY 1997, \$5,000 was obligated towards the curation of Atlantic well samples. In FY 1998, \$836 was obligated toward updating the MMS AAPG CD-ROM investment. All figures have been rounded.

Table 7. Summary of Geological and Geophysical Data Acquisition Expenditures by Data Type and Region, FY 1968-1999

| Data Type Expenditures (\$)* | |
|---|--|
| Alasi | ка |
| 2-D Seismic High Resolution CDP Interpretations Gravity and Magnetics 3-D Seismic Total | 40,944,258 11,125,798 439,793 1,026,956 <u>358,155</u> 53,894,960 |
| Atlan | tic |
| 2-D Seismic High Resolution CDP Interpretations Gravity and Magnetics 3-D Seismic Total | 7,791,438 9,751,232 55,274 2,902 <u>0</u> 17,600,846 |
| Gulf of N | lexico |
| 2-D Seismic High Resolution CDP Interpretations Gravity and Magnetics 3-D Seismic Total | 31,198,570 12,729,139 856,526 529,734 <u>5,898,594</u> 51,212,563 |
| Pacif | ic |
| 2-D Seismic High Resolution CDP Interpretations Gravity and Magnetics 3-D Seismic Total | 9,553,194 3,696,394 72,175 534,363 <u>27,275</u> 13,883,401 |

^{*}MMS has had additional expenditures through its G&G data acquisition budget for other general purchases such as field tapes, special processing, navigation tapes, interpretive hardware and software for evaluation purposes, geological studies, and acquisition of digital tapes of in-house analog data.

Table 8. Summary of Average Cost Per Mile by MMS for 2-D Seismic Data, FY 1968-1999

| FY | Average Cost (\$/Mile) |
|-----------|------------------------|
| 1968-1975 | 33.60 |
| 1976 | 34.90 |
| 1977 | 30.00 |
| 1978 | 73.60 |
| 1979 | 99.70 |
| 1980 | 91.50 |
| 1981 | 100.70 |
| 1982 | 107.00 |
| 1983 | 102.50 |
| 1984 | 121.10 |
| 1985 | 105.90 |
| 1986 | 102.00 |
| 1987 | 48.30 |
| 1988 | 32.70 |
| 1989 | 26.10 |
| 1990 | 18.00 |
| 1991 | 19.86 |
| 1992 | 7.49 |
| 1993 | 13.33 |
| 1994 | 75.84 |
| 1995 | 22.02 |
| 1996 | 39.04 |
| 1997 | 5.45 |
| 1998 | 3.18 |
| 1999 | 1.40 |

Note: Totals reflect average cost per mile for all CDP information acquired, both State and Federal. Average costs reflect only those dollars assigned to the bureauwide G&G budget and do not reflect monies allocated from Regional funds.

Bibliography

Dellagiarino, George, 1991, The Offshore Deep Stratigraphic Test Well Program, U.S. Department of the Interior, Minerals Management Service, OCS Report MMS 90-0028, 36 p.

Dellagiarino, George, Fulton, Patricia, Meekins, Keith, and Zinzer, David, 1999, Geological and Geophysical Data Acquisition, Outer Continental Shelf Through 1998, U.S. Department of the Interior, Minerals Management Service, OCS Report MMS 99-0058, 32 p.

Dellagiarino, George, Fulton, Patricia, and Zinzer, David, 1997, Geological and Geophysical Data Acquisition, A Twenty-Year Retrospective, 1976-1996, U.S. Department of the Interior, Minerals Management Service, OCS Report MMS 97-0035, 39 p.

Dellagiarino, George, and Meekins, Keith, 1998, The Resource Evaluation Program: Structure and Mission on the Outer Continental Shelf, U.S. Department of the Interior, Minerals Management Service, OCS Report MMS 98-0028, 43 p.

Fulton, Patricia, 1998, Prelease Permits for Oil, Gas, and Sulphur on the Outer Continental Shelf, U.S. Department of the Interior, Minerals Management Service, OCS Report MMS 98-0009, 119 p.

Geophysics, Society of Exploration Geophysicists, Tulsa, Oklahoma.

Geophysics: The Leading Edge of Exploration, Society of Exploration Geophysicists, Tulsa, Oklahoma.

Robinson, E.S., and Coruh, C., 1988, Basic Exploration Geophysics, John Wiley and Sons, 562 p.

Wood, Larry C., 1982, "Imaging the Subsurface" *in* Concepts and Techniques in Oil and Gas Exploration, Society of Exploration Geophysicists, Tulsa, Oklahoma, pgs. 45-90.

Appendix

| <u>Table</u> | | |
|--------------|---|----|
| | Alaska | |
| A-1 | Summary of geological and geophysical data acquisition by fiscal year for Alaska | 15 |
| A-2 | Number of permits issued for geological and geophysical exploration in Alaska | 16 |
| A-3 | Summary of expenditures by MMS for geological and geophysical data acquisition by fiscal year for Alaska (in dollars) | 17 |
| A-4 | Summary of average cost per mile by MMS for 2-D seismic data by fiscal year for Alaska (in dollars) | 18 |
| ۸ ۶ | | |
| A-5 | Summary of geological and geophysical data acquisition by fiscal year for the Atlantic | 19 |
| A-6 | Number of permits issued for geological and geophysical exploration in the Atlantic | 20 |
| A-7 | Summary of expenditures by MMS for geological and geophysical data acquisition by fiscal year for the Atlantic (in dollars) | 21 |
| A-8 | Summary of average cost per mile by MMS for 2-D seismic data by fiscal year for the Atlantic (in dollars) | 22 |
| A-9 | Summary of geological and geophysical data acquisition by fiscal year for the Gulf of Mexico | 23 |
| A-10 | Number of permits issued for geological and geophysical exploration in the Gulf of Mexico | 24 |
| A-11 | Summary of expenditures by MMS for geological and geophysical data acquisition by fiscal year for the Gulf of Mexico (in dollars) | 25 |
| A-12 | Summary of average cost per mile by MMS for 2-D seismic data by fiscal year for the Gulf of Mexico (in dollars) | 26 |

| | Pacific | |
|------|--|----|
| A-13 | Summary of geological and geophysical data acquisition by fiscal year for the Pacific | 27 |
| A-14 | Number of permits issued for geological and geophysical exploration in the Pacific | 28 |
| A-15 | Summary of expenditures by MMS for geological and geophysical data acquisition by fiscal year for the Pacific (in dollars) | 29 |
| A-16 | Summary of average cost per mile by MMS for 2-D seismic data by fiscal year for the Pacific (in dollars) | 30 |

Alaska Tables

Table A-1. Summary of Geological and Geophysical Data Acquisition by Fiscal Year for Alaska

| Year | 2-D | HRD | Interpretations | Grav/Mag | 3-D | DST |
|-----------|----------|----------|-----------------|----------|-----------|----------|
| 1968-1975 | 70,306 | 5,500 | 32,819 | 55,710 | 0 | 1 |
| 1976 | 37,785 | 19,163 | 30,164 | 0 | 0 | 4 |
| 1977 | 11,952 | 5,606 | 21,700 | 23,470 | 0 | 4 |
| 1978 | 28,524 | 0 | 0 | 36,625 | 0 | 0 |
| 1979 | 8,538 | 5,412 | 0 | 25,465 | 0 | 0 |
| 1980 | 10,109 | 7,703 | 0 | 0 | 0 | 1 |
| 1981 | 35,430 | 4,590 | 0 | 14,969 | 0 | 0 |
| 1982 | 16,624 | 0 | 0 | 0 | 0 | 2 |
| 1983 | 51,903 | 0 | 0 | 0 | 0 | 2 |
| 1984 | 30,961 | 7,904 | 0 | 5,850 | 0 | 0 |
| 1985 | 30,270 | 0 | 0 | 0 | 0 | 0 |
| 1986 | 21,603 | 1,600 | 0 | 0 | 0 | 0 |
| 1987 | 49,532 | 470 | 0 | 80,826 | 0 | 0 |
| 1988 | 14,963 | 1,741 | 0 | 0 | 0 | 0 |
| 1989 | 3,136 | 166 | 0 | 9,543 | 0 | 0 |
| 1990 | 8,557 | 0 | 0 | 11,046 | 0 | 0 |
| 1991 | 3,964 | 0 | 0 | 1,500 | 0 | 0 |
| 1992 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 1,893 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 2,422 | 0 | 0 | 102,845 | 0 | 0 |
| 1995 | 737 | 0 | 0 | 3,000 | 0 | 0 |
| 1996 | 315 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 382 | 0 | 0 | 0 | 3* | 0 |
| 1998 | 273 | 0 | 0 | 0 | 0 | 0 |
| 1999 | <u>0</u> | <u>0</u> | <u>0</u> | <u>0</u> | <u>7*</u> | <u>0</u> |
| Total | 440,179 | 59,855 | 84,683 | 370,849 | 10* | 14 |

Note: * Acquisition for 3-D seismic data are measured in blocks; all other acquisitions in this table are measured in miles. DST dates are assigned based upon completion dates and are measured in terms of wells completed. All other data are measured in terms of miles.

Table A-2. Number of Permits Issued for Geological and Geophysical Exploration in Alaska

| Year | Α | В | С | D | Е | F |
|-----------|----------|----------|-----------------|----------------|---------------------|---------------|
| 1960-1968 | 75 | | | | | |
| 1969 | 31 | 28 | 3 | 0 | 0 | 0 |
| 1970 | 40 | 36 | 4 | 0 | 0 | 0 |
| 1971 | 27 | 26 | 1 | 0 | 0 | 0 |
| 1972 | 17 | 17 | 0 | 0 | 0 | 0 |
| 1973 | 33 | 32 | 1 | 0 | 0 | 0 |
| 1974 | 47 | 44 | 3 | 0 | 0 | 0 |
| 1975 | 82 | 74 | 8 | 1 | 0 | 0 |
| 1976 | 69 | 61 | 8 | 4 | 0 | 0 |
| 1977 | 33 | 29 | 4 | 4 | 0 | 0 |
| 1978 | 9 | 8 | 1 | 0 | 0 | 0 |
| 1979 | 32 | 30 | 2 | 0 | 0 | 0 |
| 1980 | 41 | 36 | 5 | 1 | 0 | 0 |
| 1981 | 54 | 49 | 5 | 0 | 0 | 0 |
| 1982 | 85 | 79 | 6 | 3 | 0 | 0 |
| 1983 | 103 | 80 | 23 | 1 | 0 | 0 |
| 1984 | 70 | 62 | 8 | 0 | 0 | 0 |
| 1985 | 63 | 56 | 7 | 0 | 0 | 0 |
| 1986 | 18 | 17 | 1 | 0 | 0 | 0 |
| 1987 | 18 | 14 | 4 | 0 | 0 | 0 |
| 1988 | 13 | 9 | 4 | 0 | 0 | 0 |
| 1989 | 17 | 14 | 3 | 0 | 0 | 0 |
| 1990 | 19 | 15 | 3 | 0 | 0 | 1 |
| 1991 | 7 | 4 | 1 | 0 | 0 | 2 |
| 1992 | 7 | 6 | 0 | 0 | 0 | 1 |
| 1993 | 11 | 10 | 0 | 0 | 2 | 1 |
| 1994 | 3 | 3 | 0 | 0 | 1 | 0 |
| 1995 | 1 | 1 | 0 | 0 | 1 | 0 |
| 1996 | 6 | 6 | 0 | 0 | 5 | 0 |
| 1997 | 5 | 4 | 1 | 0 | 0 | 0 |
| 1998 | 2 | 2 | 0 | Ō | 2 | 0 |
| 1999 | <u>2</u> | <u>2</u> | $\frac{0}{0}$ | <u>0</u> | 2 <u>2</u> 13 | |
| Total | 1,040 | 854 | $10\frac{3}{6}$ | $1\frac{3}{4}$ | 13 | <u>0</u> 5 |

A=Total Number of Geological, Geophysical, and Strategic Minerals Permits

B=Number of Geophysical Permits

C=Number of Geological Permits

D=Number of Geological Permits Issued for Deep Stratigraphic Tests E=Number of Geophysical Permits Issued for 3-D Seismic Data

F=Number of Permits Issued for Strategic (Nonenergy) Minerals

Dashed lines = Individual breakouts not established

Table A-3. Summary of Expenditures by MMS for Geological and Geophysical Data Acquisition by Fiscal Year for Alaska (in dollars)

| Year | 2-D | HRD | Interpretations | Grav/Mag | 3-D | DST |
|-----------|------------|------------|-----------------|-----------|---------|-----|
| 1968-1975 | 2,803,939 | 119,700 | 160,832 | 7,515 | 0 | NA |
| 1976 | 1,628,153 | 1,598,789 | 268,961 | 0 | 0 | NA |
| 1977 | 271,035 | 36,473 | 10,000 | 49,450 | 0 | NA |
| 1978 | 2,956,280 | 0 | 0 | 408,679 | 0 | 0 |
| 1979 | 2,180,700 | 2,019,512 | 0 | 125,148 | 0 | 0 |
| 1980 | 1,086,423 | 5,789,936 | 0 | 0 | 0 | NA |
| 1981 | 5,231,130 | 1,531,458 | 0 | 69,286 | 0 | 0 |
| 1982 | 1,817,736 | 0 | 0 | 0 | 0 | NA |
| 1983 | 5,673,514 | 0 | 0 | 0 | 0 | NA |
| 1984 | 4,118,626 | 19,238 | 0 | 27,072 | 0 | 0 |
| 1985 | 3,669,129 | 0 | 0 | 0 | 0 | 0 |
| 1986 | 2,780,556 | 950 | 0 | 0 | 0 | 0 |
| 1987 | 2,301,780 | 400 | 0 | 249,951 | 0 | 0 |
| 1988 | 1,339,007 | 3,425 | 0 | 0 | 0 | 0 |
| 1989 | 347,872 | 5,917 | 0 | 21,851 | 0 | 0 |
| 1990 | 832,476 | 0 | 0 | 51,681 | 0 | 0 |
| 1991 | 518,613 | 0 | 0 | 15,573 | 0 | 0 |
| 1992* | 0 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 139,117 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 579,129 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 167,170 | 0 | 0 | 750 | 0 | 0 |
| 1996 | 113,071 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 195,855 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 192,947 | 0 | 0 | 0 | 0 | 0 |
| 1999 | <u>0</u> | <u>0</u> | <u>0</u> | <u>0</u> | 358,155 | _0 |
| Total | 40,944,258 | 11,125,798 | 439,793 | 1,026,956 | 358,155 | |

^{*}In FY 1992, the Alaska Region spent funds from the G&G budget and Regional funds to acquire digital copies of data already in their inventory and did not acquire any new or additional data.

Note: NA represents "not applicable" as no G&G funds are used to acquire information from a DST. Where no DST was completed, a zero is entered into the expenditure column.

Table A-4. Summary of Average Cost Per Mile by MMS for 2-D Seismic Data by Fiscal Year for Alaska (in dollars)

| Year | Average Cost (\$/Mile) |
|-----------|------------------------|
| 1968-1975 | 39.88 |
| 1976 | 43.09 |
| 1977 | 22.68 |
| 1978 | 103.64 |
| 1979 | 255.41 |
| 1980 | 107.47 |
| 1981 | 147.65 |
| 1982 | 109.34 |
| 1983 | 109.31 |
| 1984 | 133.03 |
| 1985 | 121.21 |
| 1986 | 128.71 |
| 1987 | 46.47 |
| 1988 | 89.49 |
| 1989 | 110.93 |
| 1990 | 97.29 |
| 1991 | 130.85 |
| 1992 | 0 |
| 1993 | 73.48 |
| 1994 | 239.18 |
| 1995 | 475.85 |
| 1996 | 358.96 |
| 1997 | 512.71 |
| 1998 | 706.77 |
| 1999 | 0 |

Note: Totals reflect average cost per mile for all CDP information acquired, both State and Federal. Average costs reflect only those dollars assigned to the bureauwide G&G budget and do not reflect monies allocated from Regional funds.

Zero indicates G&G dollars were not spent on CDP information.

Atlantic Tables

Table A-5. Summary of Geological and Geophysical Data Acquisition by Fiscal Year for the Atlantic

| Year | 2-D | HRD | Interpretations | Grav/Mag | 3-D | DST |
|-----------|---------|--------|-----------------|----------|---------------|-----|
| 1968-1975 | 41,958 | 1,740 | 11,802 | 14,267 | 0 | 0 |
| 1976 | 25,211 | 23,867 | 29,822 | 1,076 | 0 | 2 |
| 1977 | 21,032 | 6,100 | 3,177 | 440 | 0 | 2 |
| 1978 | 14,281 | 0 | 0 | 0 | 0 | 0 |
| 1979 | 6,877 | 0 | 0 | 0 | 0 | 1 |
| 1980 | 585 | 10,660 | 0 | 0 | 0 | 0 |
| 1981 | 9,950 | 7,142 | 0 | 0 | 0 | 0 |
| 1982 | 19,074 | 0 | 0 | 0 | 0 | 0 |
| 1983 | 30,077 | 0 | 0 | 0 | 0 | 0 |
| 1984 | 9,386 | 0 | 0 | 0 | 0 | 0 |
| 1985 | 1,640 | 0 | 0 | 0 | 0 | 0 |
| 1986 | 424 | 0 | 0 | 0 | 0 | 0 |
| 1987 | 2,356 | 0 | 0 | 0 | 0 | 0 |
| 1988 | 827 | 0 | 0 | 0 | 0 | 0 |
| 1989 | 2,730 | 0 | 0 | 0 | 0 | 0 |
| 1990 | 31 | 0 | 0 | 0 | 0 | 0 |
| 1991 | 1,042 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 2,377 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | $\frac{0}{0}$ | _0 |
| Total | 189,858 | 49,509 | 44,801 | 15,783 | 0 | 5 |

Note: DST dates are assigned based upon completion dates and are measured in terms of wells completed. All other data are measured in terms of miles.

Table A-6. Number of Permits Issued for Geological and Geophysical Exploration in the Atlantic

| Year | Α | В | С | D | E | F |
|-----------|-----|-----|----|---|---|---|
| 1960-1968 | 45 | | | | | |
| 1969 | 7 | 7 | 0 | 0 | 0 | 0 |
| 1970 | 4 | 3 | 1 | 0 | 0 | 0 |
| 1971 | 4 | 4 | 0 | 0 | 0 | 0 |
| 1972 | 4 | 4 | 0 | 0 | 0 | 0 |
| 1973 | 4 | 4 | 0 | 0 | 0 | 0 |
| 1974 | 2 | 2 | 0 | 0 | 0 | 0 |
| 1975 | 29 | 23 | 6 | 1 | 0 | 0 |
| 1976 | 35 | 28 | 7 | 3 | 0 | 0 |
| 1977 | 20 | 20 | 0 | 0 | 0 | 0 |
| 1978 | 17 | 13 | 4 | 1 | 0 | 0 |
| 1979 | 9 | 9 | 0 | 0 | 0 | 0 |
| 1980 | 15 | 15 | 0 | 0 | 0 | 0 |
| 1981 | 17 | 16 | 1 | 0 | 0 | 0 |
| 1982 | 11 | 11 | 0 | 0 | 0 | 0 |
| 1983 | 10 | 10 | 0 | 0 | 0 | 0 |
| 1984 | 6 | 6 | 0 | 0 | 0 | 0 |
| 1985 | 2 | 1 | 1 | 0 | 0 | 0 |
| 1986 | 3 | 2 | 1 | 0 | 0 | 0 |
| 1987 | 2 | 0 | 2 | 0 | 0 | 0 |
| 1988 | 4 | 4 | 0 | 0 | 0 | 0 |
| 1989 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1990 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1991 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 1 | 0 | 0 | 0 | 0 | 1 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 2 | 1 | 1 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 254 | 184 | 24 | 5 | 0 | 1 |

A=Total Number of Geological, Geophysical, and Strategic Minerals Permits

Dashed lines = Individual breakouts not established

B=Number of Geophysical Permits

C=Number of Geological Permits

D=Number of Geological Permits Issued for Deep Stratigraphic Tests
E=Number of Geophysical Permits Issued for 3-D Seismic Data
F=Number of Permits Issued for Strategic (Nonenergy) Minerals

Table A-7. Summary of Expenditures by MMS for Geological and Geophysical Data Acquisition by Fiscal Year for the Atlantic (in dollars)

| Year | 2-D | HRD | Interpretations | Grav/Mag | 3-D | DST |
|-----------|-----------|-----------|-----------------|----------|-----|-----|
| 1968-1975 | 309,029 | 4,900 | | | 0 | 0 |
| 1976 | 196,687 | 2,256,167 | 45,282 | 2,902 | 0 | NA |
| 1977 | 242,868 | 1,968,513 | 9,992 | 0 | 0 | NA |
| 1978 | 581,562 | 0 | 0 | 0 | 0 | 0 |
| 1979 | 119,250 | 0 | 0 | 0 | 0 | NA |
| 1980 | 51,096 | 4,278,448 | 0 | 0 | 0 | 0 |
| 1981 | 179,682 | 1,243,204 | 0 | 0 | 0 | 0 |
| 1982 | 1,882,723 | 0 | 0 | 0 | 0 | 0 |
| 1983 | 1,718,584 | 0 | 0 | 0 | 0 | 0 |
| 1984 | 1,500,298 | 0 | 0 | 0 | 0 | 0 |
| 1985 | 287,135 | 0 | 0 | 0 | 0 | 0 |
| 1986 | 87,307 | 0 | 0 | 0 | 0 | 0 |
| 1987 | 438,792 | 0 | 0 | 0 | 0 | 0 |
| 1988 | 71,510 | 0 | 0 | 0 | 0 | 0 |
| 1989 | 120,042 | 0 | 0 | 0 | 0 | 0 |
| 1990 | 150 | 0 | 0 | 0 | 0 | 0 |
| 1991 | 2,790 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 1,933 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | _0 | _0 |
| Total | 7,791,438 | 9,751,232 | 55,274 | 2,902 | 0 | |

Note: NA represents "not applicable" as no G&G funds are used to acquire information from a DST. Where no DST was completed, a zero is entered into the expenditure column.

No expenditures are available for CDP interpretations or gravity and magnetic data for 1968-1975.

Table A-8. Summary of Average Cost Per Mile by MMS for 2-D Seismic Data by Fiscal Year for the Atlantic (in dollars)

| Year | Average Cost (\$/Mile) |
|-----------|------------------------|
| 1968-1975 | 7.37 |
| 1976 | 7.80 |
| 1977 | 11.55 |
| 1978 | 40.72 |
| 1979 | 17.34 |
| 1980 | 87.34 |
| 1981 | 18.06 |
| 1982 | 98.70 |
| 1983 | 57.14 |
| 1984 | 159.85 |
| 1985 | 175.08 |
| 1986 | 205.91 |
| 1987 | 186.24 |
| 1988 | 86.47 |
| 1989 | 43.97 |
| 1990 | 4.84 |
| 1991 | 2.68 |
| 1992 | 0.81 |
| 1993 | |
| 1994 | |
| 1995 | |
| 1996 | |
| 1997 | |
| 1998 | |
| 1999 | |
| | |

Note: Totals reflect average cost per mile for all CDP information acquired, both State and Federal. Average costs reflect only those dollars assigned to the bureauwide G&G budget and do not reflect monies allocated from Regional funds.

Dashed lines indicate no funding for that year.

Gulf of Mexico Tables

Table A-9. Summary of Geological and Geophysical Data Acquisition by Fiscal Year for the Gulf of Mexico

| Year | 2-D | HRD | Interpretations | Grav/Mag | 3-D | DST |
|-----------|---------|---------|-----------------|----------|---------|-----|
| 1968-1975 | 143,458 | 88,549 | 120,038 | 19,670 | 0 | 2 |
| 1976 | 31,474 | 9,367 | 19,380 | 56,272 | 0 | 0 |
| 1977 | 4,485 | 18,119 | 0 | 0 | 0 | 0 |
| 1978 | 7,188 | 8,275 | 0 | 0 | 0 | 0 |
| 1979 | 11,681 | 5,018 | 0 | 0 | 0 | 0 |
| 1980 | 4,758 | 15,940 | 0 | 0 | 0 | 0 |
| 1981 | 16,454 | 500 | 0 | 0 | 0 | 0 |
| 1982 | 28,700 | 0 | 0 | 0 | 0 | 0 |
| 1983 | 26,290 | 0 | 0 | 0 | 0 | 0 |
| 1984 | 40,828 | 0 | 0 | 0 | 0 | 0 |
| 1985 | 31,430 | 0 | 0 | 0 | 0 | 0 |
| 1986 | 22,616 | 0 | 0 | 0 | 0 | 0 |
| 1987 | 43,073 | 0 | 0 | 0 | 0 | 0 |
| 1988 | 56,265 | 0 | 0 | 0 | 0 | 0 |
| 1989 | 43,121 | 0 | 0 | 0 | 0 | 1 |
| 1990 | 76,692 | 0 | 0 | 0 | 0 | 0 |
| 1991 | 35,507 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 46,814 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 23,589 | 0 | 0 | 0 | 1,563* | 0 |
| 1994 | 4,416 | 0 | 0 | 0 | 1,420* | 0 |
| 1995 | 8,193 | 0 | 0 | 0 | 1,826* | 0 |
| 1996 | 32,797 | 0 | 0 | 0 | 1,458* | 0 |
| 1997 | 39,300 | 0 | 0 | 0 | 3,105* | 0 |
| 1998 | 90,708 | 0 | 0 | 178,305 | 3,452* | 0 |
| 1999 | 30,135 | 0 | 0 | 52,000 | 3,219* | _0 |
| Total | 899,972 | 145,768 | 139,418 | 306,247 | 16,043* | 3 |

Note: *Acquisitions for 3-D seismic data are measured in blocks; all other acquisitions in this table are measured in miles.

DST dates are assigned based upon completion dates and are measured in terms of wells completed.

Table A-10. Number of Permits Issued for Geological and Geophysical Exploration in the Gulf of Mexico

| Year | Α | В | С | D | E | F | G |
|-----------|------------|--------------------|------------------|---------------|-----------|---------------|---------------|
| 1960-1968 | 2,071 | | | | | | |
| 1969 | 207 | 204 | 3 | 0 | 0 | 0 | 0 |
| 1970 | 166 | 162 | 4 | 0 | 0 | 0 | 0 |
| 1971 | 179 | 175 | 4 | 0 | 0 | 0 | 0 |
| 1972 | 198 | 188 | 10 | 0 | 0 | 0 | 0 |
| 1973 | 272 | 264 | 8 | 0 | 0 | 0 | 0 |
| 1974 | 284 | 275 | 9 | 2 | 0 | 0 | 0 |
| 1975 | 353 | 348 | 5 | 0 | 0 | 0 | 0 |
| 1976 | 292 | 289 | 3 | 0 | 0 | 0 | 0 |
| 1977 | 368 | 361 | 7 | 0 | 0 | 0 | 0 |
| 1978 | 278 | 278 | 0 | 0 | 0 | 0 | 0 |
| 1979 | 211 | 204 | 7 | 0 | 0 | 0 | 0 |
| 1980 | 231 | 225 | 6 | 0 | 0 | 0 | 0 |
| 1981 | 283 | 280 | 3 | 0 | 0 | 0 | 0 |
| 1982 | 344 | 341 | 3 | 0 | 0 | 0 | 0 |
| 1983 | 416 | 416 | 0 | 0 | 16 | 0 | 0 |
| 1984 | 411 | 408 | 3 | 0 | 18 | 0 | 0 |
| 1985 | 300 | 295 | 5 | 0 | 38 | 0 | 0 |
| 1986 | 170 | 169 | 1 | 0 | 32 | 0 | 0 |
| 1987 | 258 | 252 | 6 | 0 | 42 | 0 | 0 |
| 1988 | 263 | 251 | 12 | 0 | 45 | 0 | 0 |
| 1989 | 232 | 223 | 9 | 1 | 47 | 0 | 0 |
| 1990 | 227 | 222 | 5 | 0 | 57 | 0 | 0 |
| 1991 | 163 | 152 | 11 | 0 | 45 | 0 | 0 |
| 1992 | 134 | 131 | 3 | 0 | 53 | 0 | 0 |
| 1993 | 136 | 125 | 11 | 0 | 68 | 0 | 0 |
| 1994 | 130 | 114 | 16 | 0 | 52 | 0 | 0 |
| 1995 | 102 | 91 | 11 | 0 | 49 | 0 | 0 |
| 1996 | 130 | 114 | 16 | 0 | 54 | 0 | 0 |
| 1997 | 152 | 134 | 18 | 0 | 69 | 0 | 1 |
| 1998 | 155 | 141 | 14 | 0 | 57 | 0 | 1 |
| 1999 | <u>109</u> | <u>96</u> | <u>13</u> | 0 | <u>42</u> | <u>0</u> | <u>0</u> |
| Total | 9,225 | $6,9\overline{28}$ | $2\overline{26}$ | <u>0</u> 3 | 784 | $\frac{0}{0}$ | <u>0</u> 2 |

A=Total Number of Geological, Geophysical, and Strategic Minerals Permits

B=Number of Geophysical Permits

C=Number of Geological Permits

D=Number of Geological Permits Issued for Deep Stratigraphic Tests

E=Number of Geophysical Permits Issued for 3-D Seismic Data F=Number of Permits Issued for Strategic (Nonenergy) Minerals

G=Number of Permits Issued for 4-D Seismic Data

Dashed lines = Individual breakouts not established

Table A-11. Summary of Expenditures by MMS for Geological and Geophysical Data Acquisition by Fiscal Year for the Gulf of Mexico (in dollars)

| Year | 2-D | HRD | Interpretations | Grav/Mag | 3-D | DST |
|-----------|------------|------------|-----------------|--------------|-----------|-----|
| 1968-1975 | 5,255,068 | 2,795,562 | 722,442 | 129,500 | 0 | NA |
| 1976 | 1,489,665 | 514,141 | 134,084 | 385,234 | 0 | 0 |
| 1977 | 579,583 | 3,072,088 | 0 | 0 | 0 | 0 |
| 1978 | 330,183 | 1,438,856 | 0 | 0 | 0 | 0 |
| 1979 | 492,299 | 949,697 | 0 | 0 | 0 | 0 |
| 1980 | 388,329 | 3,926,990 | 0 | 0 | 0 | 0 |
| 1981 | 939,506 | 31,805 | 0 | 0 | 0 | 0 |
| 1982 | 2,936,727 | 0 | 0 | 0 | 0 | 0 |
| 1983 | 3,678,684 | 0 | 0 | 0 | 0 | 0 |
| 1984 | 3,999,326 | 0 | 0 | 0 | 0 | 0 |
| 1985 | 2,768,574 | 0 | 0 | 0 | 0 | 0 |
| 1986 | 1,600,031 | 0 | 0 | 0 | 0 | 0 |
| 1987 | 1,824,927 | 0 | 0 | 0 | 0 | 0 |
| 1988 | 1,075,515 | 0 | 0 | 0 | 0 | 0 |
| 1989 | 885,748 | 0 | 0 | 0 | 0 | NA |
| 1990 | 704,670 | 0 | 0 | 0 | 0 | 0 |
| 1991 | 289,266 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 376,893 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 200,407 | 0 | 0 | 0 | 537,908 | 0 |
| 1994 | 26,946 | 0 | 0 | 0 | 647,592 | 0 |
| 1995 | 21,535 | 0 | 0 | 0 | 592,223 | 0 |
| 1996 | 1,151,587 | 0 | 0 | 0 | 526,471 | 0 |
| 1997 | 44,103 | 0 | 0 | 0 | 1,150,050 | 0 |
| 1998 | 96,771 | 0 | 0 | 12,000 | 1,289,773 | 0 |
| 1999* | 42,227 | 0 | 0 | <u>3,000</u> | 1,154,577 | 0 |
| Total | 31,198,570 | 12,729,139 | 856,526 | 529,734 | 5,898,594 | |

Note: NA represents "not applicable" as no G&G funds are used to acquire information from a DST. Where no DST was completed, a zero is entered into the expenditure column.

^{*} In FY 1999, the Gulf of Mexico Region also spent funds to acquire digital copies of data and information that were already in their inventory or purchased as two differing displays.

Table A-12. Summary of Average Cost Per Mile by MMS for 2-D Seismic Data by Fiscal Year for the Gulf of Mexico (in dollars)

| Year | Average Cost (\$/Mile) |
|-----------|------------------------|
| 1968-1975 | 36.63 |
| 1976 | 47.33 |
| 1977 | 129.23 |
| 1978 | 45.94 |
| 1979 | 42.15 |
| 1980 | 81.62 |
| 1981 | 57.10 |
| 1982 | 102.33 |
| 1983 | 139.93 |
| 1984 | 97.96 |
| 1985 | 88.09 |
| 1986 | 70.75 |
| 1987 | 42.37 |
| 1988 | 19.12 |
| 1989 | 20.54 |
| 1990 | 9.19 |
| 1991 | 8.14 |
| 1992 | 8.05 |
| 1993 | 8.49 |
| 1994 | 6.10 |
| 1995 | 2.63 |
| 1996 | 35.11 |
| 1997 | 1.01 |
| 1998 | 1.07 |
| 1999 | 1.40 |

Note: Totals reflect average cost per mile for all CDP information acquired, both State and Federal. Average costs reflect only those dollars assigned to the bureauwide G&G budget and do not reflect monies allocated from Regional funds.

Pacific Tables

Table A-13. Summary of Geological and Geophysical Data Acquisition by Fiscal Year for the Pacific

| Year | 2-D | HRD | Interpretations | Grav/Mag | 3-D | DST |
|-----------|---------|--------|-----------------|----------|----------|-----|
| 1968-1975 | 14,092 | 9,971 | 15,552 | 87,637 | 0 | 1 |
| 1976 | 14,452 | 2,429 | 2,288 | 1,851 | 0 | 0 |
| 1977 | 5,339 | 5,979 | 24,525 | 3,950 | 0 | 0 |
| 1978 | 4,433 | 1,155 | 0 | 0 | 0 | 1 |
| 1979 | 4,393 | 6,578 | 0 | 0 | 0 | 0 |
| 1980 | 3,948 | 4,470 | 0 | 0 | 0 | 0 |
| 1981 | 8,070 | 0 | 0 | 3,662 | 0 | 0 |
| 1982 | 15,563 | 0 | 0 | 13,050 | 0 | 0 |
| 1983 | 12,473 | 0 | 0 | 0 | 0 | 0 |
| 1984 | 8,678 | 0 | 0 | 0 | 0 | 0 |
| 1985 | 8,181 | 0 | 0 | 0 | 0 | 0 |
| 1986 | 2,644 | 0 | 0 | 0 | 0 | 0 |
| 1987 | 18,719 | 0 | 0 | 0 | 0 | 0 |
| 1988 | 6,865 | 0 | 0 | 0 | 0 | 0 |
| 1989 | 4,507 | 0 | 0 | 0 | 0 | 0 |
| 1990 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1991 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 300 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 184 | 0 | 0 | 0 | 12* | 0 |
| 1997 | 0 | 0 | 0 | 0 | 21* | 0 |
| 1998 | 0 | 0 | 0 | 0 | 8* | 0 |
| 1999 | 0 | 0 | 0 | 0 | <u>0</u> | _0 |
| Total | 132,841 | 30,582 | 42,365 | 110,150 | 41* | 2 |

Note: *Acquisitions for 3-D seismic data are measured in blocks; all other acquisitions in this table are measured in miles.

DST dates are assigned based upon completion dates and are measured in terms of wells completed.

Table A-14. Number of Permits Issued for Geological and Geophysical Exploration in the Pacific

| Year | Α | В | С | D | Е | F |
|-----------|-----|-----|----|---|---|---|
| 1960-1968 | 162 | | | | | |
| 1969 | 13 | 10 | 3 | 0 | 0 | 0 |
| 1970 | 3 | 2 | 1 | 0 | 0 | 0 |
| 1971 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1972 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1973 | 30 | 21 | 9 | 0 | 0 | 0 |
| 1974 | 24 | 24 | 0 | 0 | 0 | 0 |
| 1975 | 46 | 42 | 4 | 1 | 0 | 0 |
| 1976 | 24 | 22 | 2 | 0 | 0 | 0 |
| 1977 | 31 | 26 | 5 | 0 | 0 | 0 |
| 1978 | 38 | 30 | 8 | 1 | 0 | 0 |
| 1979 | 24 | 22 | 2 | 0 | 0 | 0 |
| 1980 | 31 | 26 | 5 | 0 | 0 | 0 |
| 1981 | 40 | 38 | 2 | 0 | 0 | 0 |
| 1982 | 62 | 59 | 3 | 0 | 0 | 0 |
| 1983 | 45 | 36 | 9 | 0 | 0 | 0 |
| 1984 | 56 | 42 | 14 | 0 | 0 | 0 |
| 1985 | 33 | 30 | 3 | 0 | 0 | 0 |
| 1986 | 20 | 19 | 1 | 0 | 0 | 0 |
| 1987 | 20 | 16 | 4 | 0 | 0 | 0 |
| 1988 | 33 | 25 | 8 | 0 | 0 | 0 |
| 1989 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1990 | 4 | 3 | 1 | 0 | 0 | 0 |
| 1991 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 740 | 494 | 84 | | 0 | 0 |

A=Total Number of Geological, Geophysical, and Strategic Minerals Permits

B=Number of Geophysical Permits

C=Number of Geological Permits

D=Number of Geological Permits Issued for Deep Stratigraphic Tests

E=Number of Geophysical Permits Issued for 3-D Seismic Data F=Number of Permits Issued for Strategic (Nonenergy) Minerals

Dashed lines = Individual breakouts not established

Table A-15. Summary of Expenditures by MMS for Geological and Geophysical Data Acquisition by Fiscal Year for the Pacific (in dollars)

| Year | 2-D | HRD | Interpretations | Grav/Mag | 3-D | DST |
|-----------|-----------|-----------|-----------------|----------|----------|-----|
| 1968-1975 | 697,733 | 175,000 | 49,617 | 415,913 | 0 | NA |
| 1976 | 486,139 | 57,660 | 20,596 | 17,275 | 0 | 0 |
| 1977 | 188,930 | 752,400 | 1,962 | 11,796 | 0 | 0 |
| 1978 | 137,754 | 23,685 | 0 | 0 | 0 | NA |
| 1979 | 346,612 | 1,588,695 | 0 | 0 | 0 | 0 |
| 1980 | 249,048 | 1,098,954 | 0 | 0 | 0 | 0 |
| 1981 | 689,372 | 0 | 0 | 20,029 | 0 | 0 |
| 1982 | 1,918,891 | 0 | 0 | 69,350 | 0 | 0 |
| 1983 | 1,309,608 | 0 | 0 | 0 | 0 | 0 |
| 1984 | 1,262,030 | 0 | 0 | 0 | 0 | 0 |
| 1985 | 848,777 | 0 | 0 | 0 | 0 | 0 |
| 1986 | 356,700 | 0 | 0 | 0 | 0 | 0 |
| 1987 | 921,422 | 0 | 0 | 0 | 0 | 0 |
| 1988 | 93,748 | 0 | 0 | 0 | 0 | 0 |
| 1989 | 44,273 | 0 | 0 | 0 | 0 | 0 |
| 1990 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1991 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1992 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1993 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1994 | 443 | 0 | 0 | 0 | 0 | 0 |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 1,714 | 0 | 0 | 0 | 10,452 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 13,479 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 3,344 | 0 |
| 1999* | 0 | 0 | 0 | 0 | <u>0</u> | 0 |
| Total | 9,553,194 | 3,696,394 | 72,175 | 534,363 | 27,275 | |

Note: NA represents "not applicable" as no G&G funds are used to acquire information from a DST. Where no DST was completed, a zero is entered into the expenditure column.

^{*} In FY 1999, the Pacific Region spent funds to acquire digital copies of data already in their inventory and did not acquire any new or additional data.

Table A-16. Summary of Average Cost Per Mile by MMS for 2-D Seismic Data by Fiscal Year for the Pacific (in dollars)

| Year | Average Cost (\$/Mile) | |
|-----------|------------------------|--|
| 1968-1975 | 49.51 | |
| 1976 | 33.64 | |
| 1977 | 35.39 | |
| 1978 | 31.08 | |
| 1979 | 78.90 | |
| 1980 | 63.08 | |
| 1981 | 85.42 | |
| 1982 | 123.30 | |
| 1983 | 105.00 | |
| 1984 | 145.43 | |
| 1985 | 103.75 | |
| 1986 | 134.91 | |
| 1987 | 49.22 | |
| 1988 | 13.66 | |
| 1989 | 9.82 | |
| 1990 | 0 | |
| 1991 | 0 | |
| 1992 | | |
| 1993 | 0 | |
| 1994 | 1.48 | |
| 1995 | 0 | |
| 1996 | 9.32 | |
| 1997 | 0 | |
| 1998 | 0 | |
| 1999 | 0 | |

Note: Totals reflect average cost per mile for all CDP information acquired, both State and Federal. Average costs reflect only those dollars assigned to the bureauwide G&G budget and do not reflect monies allocated from Regional funds.

Dashed line indicates no funding for that year. Zero indicates G&G dollars were not spent on CDP information.

Glossary

Block - a geographically defined section of the Outer Continental Shelf (OCS) designated by a number on an Official Protraction Diagram or Leasing Map prepared by the Minerals Management Service (MMS). A block normally is a 9-square-mile area (3 miles x 3 miles) consisting of 5,760 acres. A single block is the smallest unit that can be leased for oil and gas exploration on the OCS.

Common Depth Point - a common location in the ocean subbottom where sound waves originating from various positions of the seismic (sound) source near the ocean surface are reflected back toward the surface. The traces from different seismic profiles corresponding to the same reflection point are mathematically summed (stacked) for reflection points beneath the survey line. Also known as common midpoint or common reflection point.

COST Wells - Continental Offshore Stratigraphic Test Wells - deep stratigraphic wells drilled to determine the geological character or stratigraphy of rock strata. These wells, which may be more than 20,000 feet deep, provide information that can be used by Government and industry to evaluate tracts to be offered in a lease sale.

Fair Market Value - the amount in cash, or on terms reasonably equivalent to cash, for which in all probability the property would be sold by a knowledgeable purchaser who desired, but is not obligated, to buy. This market value that is sought is not merely theoretical or hypothetical but represents, insofar as it is possible to estimate, the actual selling price.

High-Resolution - a range of seismic frequencies above the normal range of frequencies used in exploration, with an improvement in resolution in the shallow portions of the subbottom but with less total penetration into the subbottom.

Lease - any form of authorization that is used under section 8 or maintained under section 6 of the Outer Continental Shelf Lands Act and that authorizes exploration for and development and production of minerals or the area covered by that authorization, whichever is required of the context.

Lease Sale - an MMS proceeding by which leases for certain OCS tracts are offered for sale by competitive bidding and during which bids are received, publicly announced, and recorded.

Outer Continental Shelf - all submerged lands lying seaward and outside of the area of lands beneath navigable waters as defined in section 2 of the Submerged Lands Act and of which the subsoil and seabed appertain to the United States and are subject to its jurisdiction and control.

Outer Continental Shelf Lands Act - law passed by Congress on August 7, 1953, and amended in 1975, 1978, and 1985.

Permit - the contract or agreement, other than a lease, approved for a specified period of not more than 1 year under which a person acquires the right to conduct (1) geological exploration for mineral resources, (2) geophysical exploration for mineral resources, (3) geological scientific research, or (4) geophysical scientific research.

Planning Area - a subdivision of an offshore area used as the initial basis for considering blocks to be offered for lease in the DOI's offshore oil and gas leasing program.

Shallow Hazards - potential geological and manmade hazards to exploration on the OCS that are in the shallow portion of the subbottom. Examples include seismicity, active faults, shallow gas deposits, steep slopes, unstable soil conditions, pipelines, anchors, and sunken ships. Shallow hazards may occur in shallow or deep waters.