

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76CH03073

PPPL-3813
UC-70


PPPL-3813

**Observations of Anisotropic Ion Temperature
during RF Heating in the NSTX Edge**

by

T.M. Biewer, R.E. Bell, D.S. Darrow, C.K. Phillips, and J.R. Wilson

May 2003



**PRINCETON PLASMA PHYSICS LABORATORY
PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY**

PPPL Reports Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Availability

This report is posted on the U.S. Department of Energy's Princeton Plasma Physics Laboratory Publications and Reports web site in Fiscal Year 2003. The home page for PPPL Reports and Publications is: http://www.pppl.gov/pub_report/

DOE and DOE Contractors can obtain copies of this report from:

U.S. Department of Energy
Office of Scientific and Technical Information
DOE Technical Information Services (DTIS)
P.O. Box 62
Oak Ridge, TN 37831

Telephone: (865) 576-8401

Fax: (865) 576-5728

Email: reports@adonis.osti.gov

This report is available to the general public from:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

Telephone: 1-800-553-6847 or
(703) 605-6000

Fax: (703) 321-8547

Internet: <http://www.ntis.gov/ordering.htm>

Observations of Anisotropic Ion Temperature during RF Heating in the NSTX Edge

T.M. Biewer*, R.E. Bell*, D.S. Darrow*, C.K. Phillips* and J.R. Wilson*

**Princeton Plasma Physics Laboratory, Princeton, NJ 08543*

Abstract. A new spectroscopic diagnostic on the National Spherical Torus Experiment (NSTX) measures the velocity distribution of ions in the plasma edge with both poloidal and toroidal views. An anisotropic ion temperature is measured during the presence of high power HHFW RF heating in He plasmas, with the poloidal T_i roughly twice the toroidal T_i . Moreover, the measured spectral distribution suggests that two populations have temperatures of 500 eV and 50 eV with rotation velocities of -50 km/s and -10 km/s, respectively. This bi-modal distribution is observed in both the toroidal and poloidal views (in both He II and C III ions), and is well correlated with the period of RF power application to the plasma. The temperature of the edge ions is observed to increase with the applied RF power, which was scanned between 0 and 4.3 MW. The ion heating mechanism from HHFW RF power has not yet been identified.

INTRODUCTION

A new spectroscopic diagnostic with both toroidal and poloidal views has been implemented in the edge of NSTX plasmas. This edge rotation diagnostic (ERD) is designed to measure the velocity and temperature of ions in the edge of NSTX. Correlated with HHFW, a strong heating of He II and C III ions is observed. The apparent temperature is anisotropic; the T_i viewing poloidally is roughly twice the T_i measured with the toroidal view. Moreover, a bi-modal distribution of these ions is observed. The presence of two distributions of ions is consistent with calculated ionization and thermalization times for He II. No direct method of edge RF ion heating has yet been identified.

OBSERVATIONS

The edge rotation diagnostic (ERD)[1] measures the intrinsic emission of light from the plasma edge. There are 7 toroidally directed views and 6 poloidally directed views of the outboard plasma edge. The poloidal views are ~ 20 cm (toroidally) from the RF antenna, and the toroidal views are ~ 2 m away. The intersection of the diagnostic sightline with the intrinsic emission shell provides the localization of the measurement. The sightlines are nearly tangent to the flux surfaces. The C III triplet near 4651 Å and the He II line at 4685 Å are measured. In the results presented here, He is the bulk, "working" ion of the discharge.

The National Spherical Torus Experiment (NSTX)[2] is a large spherical tokamak[3] with a major radius of 0.85 m and a minor radius of 0.65 m. Pulse lengths for these




FIGURE 1. The He II spectrum from Shot 110144 during adjacent 10 ms time slices (centered on 455 and 465 ms), showing the difference in the poloidal view and toroidal view when HHFW RF heating is applied to the plasma. RF heating ends at 460 ms. Both views are tangent to the flux surface at ~ 146 cm. Error bars indicate statistical uncertainty.




FIGURE 2. The He II spectrum under the influence of RF heating, fit with 2 Gaussians.

NSTX discharges are $\simeq 600$ ms, with an on-axis toroidal magnetic field of ~ 0.3 T. The plasma current was 500 kA. The on-axis electron temperature and density are ≤ 2 keV and $\sim 2 \times 10^{19} \text{ m}^{-3}$, respectively with ≤ 4.3 MW of HHFW RF auxiliary heating[4].

During the application of 30 MHz RF heating, phased to drive current, a distortion to the spectrum of both He II and C III is observed, as shown in Fig. 1. The distortion is more pronounced in the poloidal view, but it is also present in the toroidal view. Under the influence of RF power, the spectrum is non-Maxwellian. However, fitting the spectrum with 2 Gaussians yields a very accurate representation of the measured data, as shown in Fig. 2, suggesting "hot" and "cold" components are present.

Fig. 3 shows the time evolution of NSTX Shot 110144. HHFW heating is applied from 60 to 460 ms. The effect of the RF heating is apparent in the edge measured brightness, velocity, and rotation. After the RF is applied, ~ 100 ms are needed to establish a new equilibrium rotation, as indicated by the velocity trace. Once an equilibrium is established, however, the plasma is very robust, as indicated by the near constant level of velocity and temperature. When the RF heating is terminated, the hot component




FIGURE 3. Time evolution of NSTX Shot 110144, showing (a) I_p , T_e , n_e , and P_{RF} and the hot and cold components of poloidally measured He II (b) brightness, (c) velocity, and (d) temperature at a tangency radius of 146 cm.




FIGURE 4. The temperature of the "hot" component scales as $P_{RF}^{0.47}$, whereas the temperature of the "cold" component is not effected by the amount of RF power applied to the plasma. He II data from Shots 110133-110145 are shown.

disappears promptly from the plasma. The amount of RF power was increased over a number of plasma discharges. Fig. 4 shows how T_i and v of the hot and cold components vary with the amount of RF power that is applied to the plasma.

DISCUSSION

One interpretation of the observed bi-modal distribution, is that there are two populations of He ions in the plasma. In this scenario, the RF excites (by some unidentified mechanism) a high-temperature population of He⁺ ions. The emission time scale is ~ 1 ns, implying that light from both populations (hot and cold) would be readily observed, since the time scale for thermalization between two populations of He ions is ~ 1 ms. However, the time scale for ionization is ~ 100 μ s. Hence, thermalization between the hot and cold populations would not be observed before ionization occurs. Presumably, thermalization occurs among He²⁺ ions, which are not observed by the ERD.

Key to this interpretation of the observed bi-modal distribution is the heating mechanism of the RF which excites the hot component of He ions. The HHFW launched by the NSTX antenna is not expected to heat edge ions[5], but should damp on electrons in the core. Resonant heating at the ion cyclotron frequency (27^{th} sub-harmonic for He, and 41^{st} for C) is unlikely. One possibility for an ion heating mechanism is stochastic heating by the RF wave. In the near-field of the antenna, a compressional wave is generated which stochastically heats He (and C) ions[6]. Another possibility for ion heating is parametric decay of the launched HHFW into a daughter HHFW and an ion quasi-mode at the fundamental ion cyclotron resonance. In both of these scenarios, since these heated ions are "born" at the antenna, they would be more prevalent in the poloidal view than in the toroidal view, due to the proximity of the poloidal view to the RF antenna. Single particle simulations suggest that particles born near the midplane, precess poloidally while drifting toroidally, and could eventually make it into the line of sight of the toroidal ERD view.

The authors wish to recognize the many contributions of the NSTX group and collaborators at Oak Ridge National Lab. This research was supported by the U.S. D.O.E. under contract: DE-AC02-76CH03073.

REFERENCES

1. Biewer, T. M., Bell, R. E., Feder, R., Johnson, D. J., and Palladino, R. W., in preparation (2003).
2. Spitzer, J., Ono, M., Peng, M., Bashore, et al., *Fusion Technology*, **30**, 1337 (1996).
3. Peng, Y.-K., and Strickler, D., *Nuclear Fusion*, **26**, 769 (1986).
4. Ono, M., *Phys. Plasmas*, **2**, 4075 (1995).
5. Majeski, R., Menard, J., Batchelor, et al., in *Radio Frequency Power in Plasmas—Thirteenth Topical Conference, Annapolis, MD*, AIP Press, New York, 1999, p. 296.
6. Ono, M., and Peng, M., personal communications (2003).

External Distribution

Plasma Research Laboratory, Australian National University, Australia
Professor I.R. Jones, Flinders University, Australia
Professor João Canalle, Instituto de Fisica DEQ/IF - UERJ, Brazil
Mr. Gerson O. Ludwig, Instituto Nacional de Pesquisas, Brazil
Dr. P.H. Sakanaka, Instituto Fisica, Brazil
The Librarian, Culham Laboratory, England
Mrs. S.A. Hutchinson, JET Library, England
Professor M.N. Bussac, Ecole Polytechnique, France
Librarian, Max-Planck-Institut für Plasmaphysik, Germany
Jolan Moldvai, Reports Library, MTA KFKI-ATKI, Hungary
Dr. P. Kaw, Institute for Plasma Research, India
Ms. P.J. Pathak, Librarian, Institute for Plasma Research, India
Ms. Clelia De Palo, Associazione EURATOM-ENEA, Italy
Dr. G. Grosso, Instituto di Fisica del Plasma, Italy
Librarian, Naka Fusion Research Establishment, JAERI, Japan
Library, Plasma Physics Laboratory, Kyoto University, Japan
Research Information Center, National Institute for Fusion Science, Japan
Dr. O. Mitarai, Kyushu Tokai University, Japan
Dr. Jiangang Li, Institute of Plasma Physics, Chinese Academy of Sciences, People's Republic of China
Professor Yuping Huo, School of Physical Science and Technology, People's Republic of China
Library, Academia Sinica, Institute of Plasma Physics, People's Republic of China
Librarian, Institute of Physics, Chinese Academy of Sciences, People's Republic of China
Dr. S. Mirnov, TRINITI, Troitsk, Russian Federation, Russia
Dr. V.S. Strelkov, Kurchatov Institute, Russian Federation, Russia
Professor Peter Lukac, Katedra Fyziky Plazmy MFF UK, Mlynska dolina F-2, Komenskeho Univerzita, SK-842 15 Bratislava, Slovakia
Dr. G.S. Lee, Korea Basic Science Institute, South Korea
Institute for Plasma Research, University of Maryland, USA
Librarian, Fusion Energy Division, Oak Ridge National Laboratory, USA
Librarian, Institute of Fusion Studies, University of Texas, USA
Librarian, Magnetic Fusion Program, Lawrence Livermore National Laboratory, USA
Library, General Atomics, USA
Plasma Physics Group, Fusion Energy Research Program, University of California at San Diego, USA
Plasma Physics Library, Columbia University, USA
Alkesh Punjabi, Center for Fusion Research and Training, Hampton University, USA
Dr. W.M. Stacey, Fusion Research Center, Georgia Institute of Technology, USA
Dr. John Willis, U.S. Department of Energy, Office of Fusion Energy Sciences, USA
Mr. Paul H. Wright, Indianapolis, Indiana, USA

The Princeton Plasma Physics Laboratory is operated
by Princeton University under contract
with the U.S. Department of Energy.

Information Services
Princeton Plasma Physics Laboratory
P.O. Box 451
Princeton, NJ 08543

Phone: 609-243-2750
Fax: 609-243-2751
e-mail: pppl_info@pppl.gov
Internet Address: <http://www.pppl.gov>