


United States
Department of
Agriculture

Animal and
Plant Health
Inspection
Service

Veterinary
Services

PART II: Beef Cow/Calf Reproductive & Nutritional Management Practices



Beef

CHAPA

Cow/Calf Health & Productivity Audit

January 1994

Acknowledgements

This report has been prepared from material received and analyzed by the U.S. Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS), Veterinary Services (VS).

The Beef Cow/Calf Health and Productivity Audit was a cooperative effort between State and Federal animal health officials, university researchers, and extension personnel. We want to thank the State and Federal Veterinary Medical Officers (VMO's) who visited the farms and collected the data for their hard work and dedication to the National Animal Health Monitoring System (NAHMS).

The roles of the producer, Area Veterinarian in Charge (AVIC), NAHMS Coordinator, Veterinary Medical Officer (VMO), Animal Health Technician (AHT), and enumerators from the National Agricultural Statistics Service (NASS) were critical in providing quality data for this report. All participants are to be commended for their efforts, particularly the producers whose voluntary efforts made the study possible.

Table of Contents

Introduction	1
PART II: Beef cow/calf reproduction & nutritional management practices	
A. Participant profile	3
1. Descriptive statistics of responding operations	3
B. Population estimates based on data collected	4
1. Calving management	4
2. Replacement management during 1992	9
3. Operation records and marketing	14
4. Nutrition	20
5. Animal health	25
6. Sources of animal health or beef production information	28
PART III: Beef cow/calf health & health management	
A. Participant profile	31
1. Descriptive statistics of responding operations	31
B. Population estimates based on data collected	32
1. Bull management	32
2. Replacement heifer and cow management	34
3. Calves weaned in 1992 - average weight and value at weaning	35
4. Breeding-age animals sold in 1992	35
5. Death loss and health concerns	36
6. Producer opinions on beef health	40
7. Vaccine usage	42

List of Illustrations

States participating in data collection for this report and percent of U.S. beef cow/calf operations represented, January 1, 1992	2
Examples of 95% confidence intervals	2
Percent of responding operations by beef cow herd size	3
Percent of operations by age of replacement heifers at first calving	4
Percent of calves born by location	5
Percent of operations by number of hours females were allowed to labor before given assistance	6
Percent of females requiring various levels of assistance during calving	7

List of Illustrations (continued)

Selected management practices routinely performed on calves within 24 hours after birth	8
Factors in purchasing or selecting a bull by level of importance	10
Factors in culling a bull by level of importance	11
Percent of operations by the majority of breed makeup of cows and calves	12
Most important factor for determining when to wean calves	13
Number of identification methods used per operation	14
Individual ID status for cows & calves in beef cow/calf industry	15
Percent of operations recording various kinds of information	16
Percent of operations by method of record keeping	17
Use of marketing methods for weaned steers/bulls	18
Percent of operations and calves by percent of calves forward priced	20
Laboratory nutritional analysis completed on purchased or raised feed	21
Cow herd access to pasture and crop residue by month	22
Implant practices	24
Identification of minerals as deficient or cause of health problems in herd in last 5 years	25
Percent of operations by deworming practice	26
Percent of operations treating cattle	27
Sources of animal health information	28
Sources of beef production information	29
Sources of nutritional information	30
Percent of responding operations by beef cow herd size	31
Percent of operations and cows on these operations performing selected procedures on bulls	33
Percent of operations using selected management procedures on cows and replacement heifers in the last 12 months	34
Average weight and value per head of breeding age animals sold in 1992	35
Percent of calves affected by selected conditions during the past 12 months	37
Percent of cows and replacement heifers affected by selected conditions during the past 12 months	38
Percent of total animals that died or were lost due to perceived causes	39
Top conditions that producers agreed had a significant economic impact on the cow/calf operation during the past 12 months	41
Percent of producers that agreed the following conditions are significant problems for the U.S. beef cattle industry	41
Percent of operations using general vaccines by animal class	42
Percent of operations using the following vaccines by animal class:	
—Reproductive system	43
—Clostridial	44
—Digestive system	45

Introduction

As part of the National Animal Health Monitoring System (NAHMS), USDA:APHIS: Veterinary Services conducted a National study of beef production designed to provide both participants and the industry with information on animal health, productivity, and management practices of cow/calf producers. The National Agricultural Statistics Service (NASS) collaborated with USDA:APHIS:VS to select a producer sample that was statistically designed to provide inferences about the nation's cow/calf population.

NASS enumerators contacted producers in the 48 continental States by computer-assisted telephone interview and asked them a series of questions about management practices and the health of their animals. The 3,397 cow/calf producers participating represented all U.S. cow/calf operations. Results of NASS telephone contacts for the Beef Cow/Calf Health and Productivity Audit (CHAPA) were released in August 1993 as *Part I: Beef Cow/Calf Herd Management Practices in the United States*.

NASS enumerators collected data for Part II of this report, *Nutritional & Reproductive Management Practices*, from November 9 through December 4, 1992, by personal interview from a subset of producers responding to the first NASS contact. Producers participating in this portion of the study were required to have five or more beef cows (or beef replacement heifers) and 50 percent or more of their 1992 calf crop born between January 1 and June 30, 1992. Data collection was limited to 18 of the largest cow/calf-producing States (shown on the next page). The 18 States with producers participating represented 70 percent of the U.S. beef cow inventory.

The target population represents:

- 49 percent of beef cows in the U.S.
- 42 percent of beef operations in the U.S.
- 71 percent of beef cows on predominantly spring calving beef operations in the U.S. with 5 or more beef cows (or replacement heifers).
- 75 percent of predominantly spring calving beef operations in the U.S. with 5 or more beef cows (or replacement heifers).

Data for *Part III: Beef Cow/Calf Health & Health Management* were collected from 540 producers from the subset described above. Federal and State Veterinary

Part I: Beef Cow/Calf Herd Management Practices in the U.S.

- States surveyed: 48
- Target population: all U.S. beef cow/calf producers
- Participating producers: 3,397
- Data collection period: 9/29-10/9/92

Part II: Beef Cow/Calf Reproductive & Nutritional Management Practices

- States surveyed: 18
- Target population: beef cow/calf producers with 5 or more beef cows and with 50 percent or more of 1992 calves born from January through June
- Participating producers: 799
- Data collection period: 11/9-12/4/92

Part III: Beef Cow/Calf Health & Health Management


- States surveyed: 18
 - Target population: beef cow/calf producers with 5 or more beef cows and with 50 percent or more of 1992 calves born from January through June
 - Participating producers: 540
-

Officers (VMO's) conducted personal interviews with the producers between January 4 and February 28, 1993.


Descriptive tables in this report are divided into two parts:

- **The Participant Profile** contains descriptive results from only the subset of operations that completed the respective personal interviews for Parts II and III.
- **Population Estimates Based on Data Collected** are population estimates, such as averages and proportions which have been weighted to represent the cow/calf population. Most of the estimates are provided with a measure of variability called the standard error and denoted by (\pm). Chances are 95 out of 100 that the interval created by the estimate plus or minus two standard errors will contain the true population value. In the example at right, an estimate of 7.5 with a standard error of ± 1.0 results in a range of 5.5 to 9.5 (two times the standard error above and below the estimate).

States Participating in Data Collection for this Report and Percent of U.S. Cow Inventory, January 1, 1992



Examples of 95% Confidence Intervals



Subsequent Beef CHAPA activities collected additional data from 540 producers in the 18 States. Additional Beef Cow/Calf Health and Productivity Audit (CHAPA) results will be released as they are completed. If you have questions about this report contact NAHMS at:

Centers for Epidemiology and Animal Health
 USDA:APHIS:VS, Attn. NAHMS
 555 South Howes, Suite 200
 Fort Collins, Colorado 80521
 (303) 490-7800

Part II: A. Participant Profile¹

1. Descriptive statistics of responding operations

a. Beef cow herd size:	<u>Number of Operations</u>	
1-4		15
5-9		41
10-49		255
50-99		148
100-299		187
300+		<u>153</u>
Total		799
b. Breed make-up - majority of cows:	<u>Number of Operations</u>	
Purebred or straightbred (only one breed)		142
Crossbred (two breeds)		359
Crossbred (three or more breeds)		<u>298</u>
Total		799
c. Number of head reported:	<u>Number of Head</u>	
Cows		245,273
Calf crop		224,315
Cows and replacement heifers	287,184	

¹ Actual sample values; not population estimates.

Part II: B. Population Estimates Based on Data Collected

1. Calving management

a. First calving


i. Age of replacement heifers at first calving: Number of
Months Standard Error

Operation average	25.2	(±0.2)
Replacement heifer average ¹	24.8	(±0.2)

ii. Percent of operations (and percent of replacement heifers on these operations) by age of replacement heifers at first calving:

<u>Age in Months</u>	<u>Percent of Operations</u>	<u>Standard Error</u>	<u>Percent of Replacement Heifers¹</u>	<u>Standard Error</u>
Less than 20	3.8	(±1.3)	5.2	(±2.1)
20-26	74.7	(±2.8)	80.3	(±2.9)
27-32	17.1	(±2.5)	11.3	(±1.9)
32-38	4.4	(±1.3)	3.2	(±0.9)
39 or more	<u>0.0</u>	(±0.0)	<u>0.0</u>	(±0.0)
Total	100.0		100.0	

Percent of Operations by Age of Replacement Heifers at First Calving



*Calculated as if all replacement heifers on an operation would calve at the same month of age.

iii. Percent of operations separating replacement heifers from cows, at least:

	<u>Percent of Operations</u>	<u>Standard Error</u>
30 days before calving	35.9	(±2.8)
30 days after calving	21.4	(±2.4)

¹ Calculated if all replacement heifers on an operation would calve at the same month of age.

1. Calving management (continued)

b. Calving location

i. Percent of operations that separate cow/calf pairs from pregnant cows within a week after calving:

	<u>Percent</u>	<u>Standard Error</u>
Percent of operations	14.9	(±2.1)

ii. Percent of cows on operations that separate cow/calf pairs from pregnant cows within a week after calving:

25.7	(±2.9)
------	--------

iii. Percent of operations where one or more calves were born in each location in the last 12 months:

<u>Location</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
Special calving pastures that allow increased observation and/or shelter	32.8	(±2.7)
Calving lots	8.5	(±1.5)
Individual calving pens	1.6	(±0.6)
Covered sheds or barns	5.9	(±1.4)
Other locations	76.7	(±2.3)

iv. Percent of calves born by location:

<u>Location</u>	<u>Percent of Calves</u>	<u>Standard Error</u>	<u>Operation Average Percent</u>	<u>Standard Error</u>
Special calving pastures that allow increased observation and/or shelter	33.5	(±2.6)	21.8	(±2.1)
Calving lots	8.1	(±1.4)	4.3	(±0.7)
Individual calving pens	0.8	(±0.3)	0.6	(±0.3)
Covered sheds or barns	2.9	(±1.2)	1.8	(±0.6)
Other locations	<u>54.7</u>	(±2.7)	<u>71.5</u>	(±2.3)
Total	100.0		100.0	

1. Calving management (continued)

c. Observing females during calving season

i. Operation average number of times females were observed over a 24-hour period during the calving season:

<u>Female Group</u>	<u>Number of Times</u>	<u>Standard Error</u>
Replacement heifers	2.9 ¹	(±0.2)
Mature cows	1.9	(±0.1)

ii. Percent of operations by number of times females were observed over a 24-hour period:

<u>Number of Times Observed</u>	<u>Replacement Heifers¹</u>		<u>Mature Cows</u>	
	<u>Percent of Operations</u>	<u>Standard Error</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
0	4.7	(±1.7)	7.8	(±1.5)
1-2	57.2	(±3.5)	72.9	(±2.5)
3-4	21.7	(±2.8)	13.5	(±1.9)
5 or more	<u>16.4</u>	(±2.3)	<u>5.8</u>	(±1.0)
Total	100.0		100.0	

d. Calving assistance


i. Operation average number of hours females were allowed to labor before given assistance:

	<u>Number of Hours</u>	<u>Standard Error</u>
Replacement heifers	2.9 ¹	(±0.1)
Mature cows	2.6	(±0.1)

ii. Percent of operations by number of hours females were allowed to labor before given assistance:

<u>Number of Hours Observed</u>	<u>Replacement Heifers¹</u>		<u>Mature Cows</u>	
	<u>Percent of Operations</u>	<u>Standard Error</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
0	0.0	(±0.0)	12.7	(±2.2)
1-2	54.8	(±3.3)	40.1	(±2.9)
3-4	32.4	(±3.3)	36.3	(±3.0)
5 or more	<u>12.8</u>	(±2.3)	<u>10.9</u>	(±1.9)
Total	100.0		100.0	

% Operations by Number of Hours Females Were Allowed to Labor Before Given Assistance




¹ Only included operations with replacement heifers.

1. Calving management (continued)

iii. Percent of females requiring various levels of assistance during calving:

	<u>Replacement Heifers</u> ¹		<u>Mature Cows</u>	
	<u>Percent</u>	<u>Standard Error</u>	<u>Percent</u>	<u>Standard Error</u>
No assistance	82.8	(±1.6)	97.8	(±0.2)
Easy pull	9.4	(±1.4)	1.4	(±0.2)
Hard pull	7.4	(±0.8)	0.8	(±0.1)
Cesarean section	<u>0.4</u>	(±0.1)	<u>0.0</u>	(±0.0)
Total	100.0		100.0	

Percent of Females Requiring Various Levels of Assistance During Calving



iv. Operation average percent of assisted births attended by a veterinarian:

<u>Percent of Assisted Births</u>	<u>Standard Error</u>
12.8	(±1.7)

v. Percent of assisted births attended by a veterinarian:

<u>Percent of Assisted Births</u>	<u>Standard Error</u>
15.3	(±2.0)

1 Only included operations with replacement heifers.


1. Calving management (continued)

e. Management practices at birth

i. Percent of operations where selected management practices were routinely performed on calves within 24 hours after birth:

<u>Management Practices</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
Treat navel with disinfectant	16.0	(±2.2)
Vitamin A injection	6.8	(±1.4)
Selenium injection	1.7	(±0.6)
Weigh calf	7.6	(±1.5)
Individual identification	28.5	(±2.5)
Provide a colostrum supplement	9.1	(±1.7)

Selected Management Practices Routinely Performed on Calves Within 24 Hours After Birth



f. Factors determining calving season

i. Percent of operations by the most used factor in determining the timing of the 1992 calving season:

<u>Determining Factor</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
No set calving season	52.7	(±2.9)
Market cycle	4.9	(±1.4)
Maximize age/weight at weaning	5.2	(±1.3)
Forage availability	5.6	(±1.1)
Tradition	11.9	(±1.7)
Labor availability	2.9	(±0.9)
Time of cattle movement	0.8	(±0.4)
Weather during calving	14.2	(±1.9)
Other	<u>1.8</u>	(±0.7)
Total	100.0	

2. Replacement management during 1992

a. Source of replacement females

i. Operation average percent of replacement females:

<u>Source</u>	<u>Percent of Females</u>	<u>Standard Error</u>
Purchased	21.1	(±2.2)
Raised	<u>78.9</u>	(±2.2)
Total	100.0	

ii. Percent of replacement females:

<u>Source</u>	<u>Percent of Females</u>	<u>Standard Error</u>
Purchased	11.6	(±1.9)
Raised	<u>88.4</u>	(±1.9)
Total	100.0	

b. Importance of factors in purchasing or selecting females:

<u>Factors</u>	<u>Percent of Operations by Level of Importance</u>			
	<u>Not</u>	<u>Moderate</u>	<u>Very</u>	<u>Extreme</u>
Breed	13.5	30.8	33.1	22.6
Standard Error	(±2.1)	(±3.0)	(±3.0)	(±2.4)
Reputation of breeder	38.1	20.8	28.1	13.0
Standard Error	(±3.1)	(±2.3)	(±2.9)	(±2.1)
Birth weight	34.9	27.9	28.1	9.1
Standard Error	(±3.1)	(±2.9)	(±2.8)	(±1.5)
Weaning weight/yearling weight	25.7	21.7	37.6	15.0
Standard Error	(±2.8)	(±2.6)	(±3.0)	(±2.1)
Hip height/frame score	27.0	29.9	34.3	8.8
Standard Error	(±2.9)	(±2.8)	(±3.0)	(±1.4)
Price	19.9	18.4	35.2	26.5
Standard Error	(±2.2)	(±2.2)	(±3.2)	(±2.8)
Pelvic area	23.9	30.4	33.3	12.4
Standard Error	(±2.6)	(±3.0)	(±2.9)	(±1.8)
Reproductive tract score	32.4	26.5	30.9	10.2
Standard Error	(±3.0)	(±2.7)	(±3.2)	(±1.7)
Appearance	4.2	17.5	53.6	24.7
Standard Error	(±1.1)	(±2.2)	(±3.2)	(±2.6)
Temperament	7.9	14.5	44.4	33.2
Standard Error	(±1.8)	(±2.1)	(±3.1)	(±2.8)
Sire information	20.5	23.6	36.1	19.8
Standard Error	(±2.4)	(±2.8)	(±3.1)	(±2.3)
Longevity of reproductive life	19.8	21.4	42.0	16.8
Standard Error	(±2.5)	(±2.6)	(±3.2)	(±2.2)

2. Replacement management during 1992 (continued)

c. Importance of factors in purchasing or selecting a bull:


<u>Factors</u>	<u>Percent of Operations by Level of Importance</u>			
	<u>Not</u>	<u>Moderate</u>	<u>Very</u>	<u>Extreme</u>
Breed	3.1	8.9	36.3	51.7
Standard Error	(±1.1)	(±1.7)	(±3.0)	(±3.1)
Reputation of breeder	15.8	22.3	41.7	20.2
Standard Error	(±2.2)	(±2.7)	(±3.2)	(±2.1)
Birth weight	20.3	20.0	38.0	21.7
Standard Error	(±2.5)	(±2.5)	(±3.0)	(±2.3)
Weaning weight/yearling weight	20.2	15.7	42.9	21.2
Standard Error	(±2.5)	(±2.2)	(±3.0)	(±2.5)
Hip height/frame score	14.2	27.0	42.6	16.2
Standard Error	(±2.1)	(±2.7)	(±3.1)	(±2.1)
Expected progeny difference (EPD)	30.5	25.3	31.5	12.7
Standard Error	(±3.0)	(±2.8)	(±3.0)	(±1.7)
Temperament	2.8	10.9	37.0	49.3
Standard Error	(±1.0)	(±1.9)	(±3.0)	(±3.0)
Scrotal circumference	19.2	23.6	42.2	15.0
Standard Error	(±2.4)	(±2.5)	(±3.0)	(±2.0)
Price	8.1	23.7	37.9	30.3
Standard Error	(±1.6)	(±2.6)	(±2.9)	(±2.9)
Structural soundness/appearance	2.5	3.0	43.3	51.2
Standard Error	(±0.9)	(±1.0)	(±3.2)	(±3.2)

2. Replacement management during 1992 (continued)

d. Importance of factors in culling bulls:

<u>Factors</u>	Percent of Operations by Level of Importance			
	<u>None</u>	<u>Moderate</u>	<u>Very</u>	<u>Extreme</u>
Age/bad teeth	15.2	18.9	38.4	27.5
Standard Error	(±2.1)	(±2.4)	(±3.0)	(±2.7)
Infertility	1.6	3.1	18.2	77.1
Standard Error	(±0.5)	(±1.3)	(±2.6)	(±2.8)
Physical unsoundness (injury/lameness)	1.5	3.4	38.4	56.7
Standard Error	(±0.5)	(±1.2)	(±3.1)	(±3.1)
Performance of offspring	4.7	10.6	39.1	45.6
Standard Error	(±1.3)	(±2.0)	(±3.1)	(±3.1)
Too many offspring in herd	15.5	14.3	28.8	41.4
Standard Error	(±2.2)	(±2.1)	(±2.9)	(±3.0)
Temperament	6.1	12.4	31.0	50.5
Standard Error	(±1.6)	(±2.0)	(±2.9)	(±3.1)
Size	9.3	23.4	46.3	21.0
Standard Error	(±2.0)	(±2.6)	(±3.1)	(±2.4)
Disease	3.2	7.1	28.2	61.5
Standard Error	(±0.7)	(±1.7)	(±2.9)	(±3.1)
Bad eye(s)	8.8	16.0	32.2	43.0
Standard Error	(±1.8)	(±2.3)	(±2.9)	(±3.1)

**Cow/Calf Health and Productivity Audit
Factors in Culling a Bull
by Level of Importance**




2. Replacement management during 1992 (continued)

e. Breed makeup

i. Percent of operations (and percent of cows on these operations) by breed makeup of the majority of the cows:

<u>Breed Makeup of Cows</u>	<u>Percent of Operations</u>	<u>Standard Error</u>	<u>Percent of Cows</u>	<u>Standard Error</u>
Purebred or straightbred (only one breed)	16.0	(±2.1)	16.8	(±2.0)
Crossbred (two breeds)	47.8	(±3.1)	45.1	(±3.0)
Crossbred (three or more breeds)	<u>36.2</u>	(±3.0)	<u>38.1</u>	(±2.9)
Total	100.0		100.0	

Percent of Operations by the Majority of Breed Makeup of Cows & Calves



ii. Percent of operations (and percent of calves on these operations) by breed makeup of the majority of the 1992 calf crop:

<u>Breed Makeup of Calf Crop</u>	<u>Percent of Operations</u>	<u>Standard Error</u>	<u>Percent of Calves</u>	<u>Standard Error</u>
Purebred or straightbred (only one breed)	12.4	(±1.8)	13.1	(±1.7)
Crossbred (two breeds)	42.6	(±3.2)	37.9	(±3.0)
Crossbred (three or more breeds)	<u>45.0</u>	(±3.0)	<u>49.0</u>	(±3.1)
Total	100.0		100.0	

2. Replacement management during 1992 (continued)

f. Factors determining weaning time

i. Percent of operations by the most important factor for determining when to wean calves:

<u>Determining Factor</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
Calf age/weight	52.9	(±3.1)
End of grazing lease or permit	2.3	(±0.9)
Forage availability	7.0	(±1.4)
Body condition of dam	6.8	(±1.7)
Market price or contract	9.0	(±2.2)
Cash flow	7.2	(±1.9)
Tradition	<u>14.8</u>	(±2.0)
Total	100.0	

g. Fall weight of mature cows (producer estimates)

i. Average weight of mature cows in the fall:

	<u>Pounds per Cow</u>	<u>Standard Error</u>
Operation average mature cow weight	1,022.9	(±9.7)
Mature cow average weight	1,047.2	(±7.5)


3. Operation records and marketing

a. Methods of identification

i. Number of identification methods used per operation:

Number Method	Individual Calf		Individual Cow		Herd ¹	
	Percent of Operations	Stand. Error	Percent of Operations	Stand. Error	Percent of Operations	Stand. Error
0	46.9	(±2.9)	39.7	(±3.1)	54.3	(±3.0)
One	29.2	(±2.6)	29.2	(±2.7)	30.0	(±2.7)
Two	16.0	(±1.9)	16.9	(±2.0)	13.4	(±1.8)
Three or more	7.9	(±1.4)	14.2	(±1.8)	2.3	(±0.7)
Total	100.0		100.0		100.0	

Number of Identification Methods Used per Operation



*Herd ID: all animals have same mark or number, such as a brand.


1 A herd level identification is one where all animals have the same mark or number, such as a brand.

3. Operation records and marketing (continued)

ii. Percent of operations using the following methods of identification:

Method	Individual Calf		Individual Cow		Herd ¹	
	Percent of Operations	Stand. Error	Percent of Operations	Stand. Error	Percent of Operations	Stand. Error
Brucellosis ear tag	21.7	(±2.2)	28.3	(±2.4)	NA ²	NA ²
Other metal ear tag	1.4	(±0.5)	1.5	(±0.5)	1.0	(±0.4)
Plastic ear tag	40.8	(±2.8)	45.3	(±3.0)	27.1	(±2.7)
Ear tattoo	10.2	(±1.5)	12.9	(±1.7)	6.8	(±1.2)
Hot iron brand	12.1	(±1.7)	17.9	(±2.1)	21.4	(±2.2)
Freeze brand	0.5	(±0.3)	2.1	(±0.8)	1.2	(±0.5)
Microchip transponder	0.4	(±0.4)	0.7	(±0.5)	0.4	(±0.4)
Neck chain	0.0	(±0.0)	0.4	(±0.4)	0.0	(±0.0)
Horn brand	0.0	(±0.0)	0.4	(±0.4)	0.1	(±0.1)
Ear notch	NA ²	NA ²	NA ²	NA ²	6.5	(±1.1)
Brisket tag	0.0	(±0.0)	0.4	(±0.4)	0.0	(±0.0)
None	46.9	(±2.9)	39.7	(±3.1)	54.3	(±3.0)

Individual ID Status for Cows & Calves in Beef Cow/Calf Industry



1 A herd level identification is one where all animals have the same mark or number, such as a brand.

2 NA: Not applicable.

3. Operation records and marketing (continued)

iii. Percent of animals (calves, cows, and total cattle) on operations using the following methods of identification:


Method	Individual Calf		Individual Cow		Herd	
	Percent of Calves	Stand. Error	Percent of Cows	Stand. Error	Percent of Total Cattle	Stand. Error
Brucellosis ear tag	30.4	(±2.8)	40.3	(±3.1)	NA ¹	NA ¹
Other metal ear tag	2.1	(±0.6)	2.4	(±0.8)	1.6	(±0.6)
Plastic ear tag	55.9	(±2.9)	61.3	(±2.8)	32.8	(±2.9)
Ear tattoo	13.6	(±1.7)	20.2	(±2.5)	11.3	(±2.2)
Hot iron brand	22.0	(±2.8)	31.3	(±3.1)	40.2	(±2.8)
Freeze brand	0.9	(±0.5)	2.6	(±1.1)	1.5	(±0.6)
Microchip transponder	0.5	(±0.4)	0.4	(±0.3)	0.2	(±0.2)
Neck chain	0.2	(±0.2)	0.4	(±0.3)	0.1	(±0.1)
Horn brand	0.2	(±0.2)	0.2	(±0.2)	0.3	(±0.2)
Ear notch	NA ¹	NA ¹	NA ¹	NA ¹	17.5	(±2.1)
Brisket tag	0.2	(±0.2)	0.7	(±0.3)	0.3	(±0.2)
None	29.8	(±2.6)	21.5	(±2.2)	34.1	(±2.9)

b. Record keeping

i. Percent of operations recording the following information for individual animals or on a herd basis:

Information Type	Percent of Operations			
	Individual	Stand. Error	Basis	Stand. Error
Birthdates or age	48.7	(±3.2)	NA ¹	NA ¹
Birth weight	6.2	(±1.1)	NA ¹	NA ¹
Weaning weight/yearling weight	13.3	(±1.9)	8.3	(±1.3)
Vaccinations and health records	36.2	(±2.9)	32.5	(±2.9)
Culling/death cause and date	30.1	(±2.7)	NA ¹	NA ¹

Percent of Operations Recording Various Kinds of Information




1 NA: Not applicable.

3. Operation records and marketing (continued)

ii. Percent of operations using the following methods to keep records:

<u>Record Type</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
Hand-written records	65.2	(±3.1)
Computer located on-farm	4.7	(±1.1)
Computer located off-farm	3.8	(±1.1)
Other	2.0	(±0.6)
No records are kept	27.6	(±3.2)

Percent of Operations by Method of Record Keeping




3. Operation records and marketing (continued)

c. Marketing methods in 1992

i. Percent of operations using the following methods by class of animal:

<u>Marketing Methods</u>	<u>Percent of Operations</u>			
	<u>Weaned Steers/Bulls</u>	<u>Weaned Heifers</u>	<u>Cull Cows</u>	<u>Cull Bulls</u>
Auction	85.2	84.0	94.3	90.6
Standard Error	(±1.9)	(±2.1)	(±1.4)	(±1.7)
Direct video	0.3	0.2	0.0	0.0
Standard Error	(±0.1)	(±0.1)	(±0.0)	(±0.0)
Direct private treaty	8.0	7.8	0.6	2.8
Standard Error	(±1.5)	(±1.5)	(±0.2)	(±0.9)
Consignment	1.6	1.1	0.2	0.0
Standard Error	(±0.6)	(±0.5)	(±0.1)	(±0.0)
Forward contract	0.5	0.2	0.0	0.0
Standard Error	(±0.3)	(±0.1)	(±0.0)	(±0.0)
Carcass basis	0.7	0.6	0.3	0.3
Standard Error	(±0.5)	(±0.5)	(±0.1)	(±0.1)
Another method	1.5	1.4	0.6	0.8
Standard Error	(±0.5)	(±0.5)	(±0.3)	(±0.5)
None marketed	2.2	4.7	4.0	5.5
Standard Error	(±0.8)	(±1.2)	(±1.4)	(±1.4)
Total	100.0	100.0	100.0	100.0

Use of Marketing Methods for Weaned Steers/Bulls



3. Operation records and marketing (continued)

ii. Percent of animals on operations (as a percent of all animals) using the following methods by class of animal:

<u>Marketing Methods</u>	<u>Percent of Operations</u>			
	<u>Weaned Steers/Bulls</u>	<u>Weaned Heifers</u>	<u>Cull Cows</u>	<u>Cull Bulls</u>
Auction	53.3	67.8	93.4	89.4
Standard Error	(±7.1)	(±3.6)	(±1.2)	(±1.9)
Direct video	1.5	1.4	0.1	0.1
Standard Error	(±0.6)	(±0.5)	(±0.0)	(±0.0)
Direct private treaty	20.1	16.8	2.7	4.3
Standard Error	(±5.7)	(±2.8)	(±0.6)	(±1.4)
Consignment	1.8	0.7	0.1	0.2
Standard Error	(±1.2)	(±0.3)	(±0.1)	(±0.2)
Forward contract	7.7	3.6	0.0	0.0
Standard Error	(±3.8)	(±2.0)	(±0.0)	(±0.0)
Carcass basis	5.9	1.3	1.3	2.2
Standard Error	(±2.8)	(±0.6)	(±0.3)	(±1.0)
Another method	3.6	2.4	0.9	0.5
Standard Error	(±2.3)	(±0.9)	(±0.4)	(±0.2)
None marketed	6.1	6.0	1.5	3.3
Standard Error	(±2.5)	(±1.5)	(±0.9)	(±0.9)
Total	100.0	100.0	100.0	100.0

d. Forward pricing of 1992 calf crop


	<u>Percent</u>	<u>Standard Error</u>		
i. Percent of operations forward pricing:	2.0	(±1.0)		
ii. Operation average percent of calves forward priced:	1.6	(±0.9)		
iii. Percent of all calves forward priced:	5.0	(±1.3)		
iv. Percent of operations and percent of all calves using forward pricing by size of cow herd:				
<u>Cow Herd Size Group</u>	<u>Percent of Operations</u>	<u>Standard Error</u>	<u>Percent of All Calves</u>	<u>Standard Error</u>
1-4	0.0	(±0.0)	0.0	(±0.0)
5-9	0.0	(±0.0)	0.0	(±0.0)
10-49	2.1	(±1.7)	3.6	(±3.4)
50-99	1.7	(±1.5)	1.9	(±1.7)
100-299	3.3	(±1.2)	3.6	(±1.2)
300 or more	16.2	(±4.8)	20.9	(±5.1)
v. For operations using forward pricing, percent of calves forward priced:			82.5	(±9.9)

3. Operation records and marketing (continued)

vi. For operations using forward pricing, percent of operations and percent of calves by percent of calves forward priced:

Reported Percent of Calves Forward Priced (Interval)	Percent of Operations	Standard Error	Percent of Calves	Standard Error
1-24	3.9	(±2.8)	10.1	(±4.6)
25-49	10.5	(±9.9)	3.7	(±2.7)
50-74	10.1	(±6.3)	14.3	(±5.4)
75-99	13.3	(±11.7)	9.7	(±6.1)
100	<u>62.2</u>	(±19.9)	<u>62.2</u>	(±9.9)
Total	100.0		100.0	

Percent of Operations and Calves by Percent of Calves Forward Priced*



vii. For operations using forward pricing, operation average percent of forward priced contracts (and percent of calves on these operations) that were:

	Percent of Contracts	Standard Error	Percent of Calf Crop	Standard Error
Forward cash	37.4	(±18.6)	54.7	(±10.6)
Future contract	11.4	(±7.3)	13.8	(±5.0)
Options	48.5	(±24.9)	25.4	(±12.2)
Another technique	<u>2.7</u>	(±2.3)	<u>6.1</u>	(±4.0)
Total	100.0		100.0	

4. Nutrition

a. Nutritional analysis

i. Percent of operations (and cows on these operations) that calculate a winter feed schedule or ration based on the animals' requirements and the quality of the feedstuffs available:

	Percent	Standard Error
Percent of operations	48.7	(±3.1)
Number of cows on these operations as a percent of cows on all operations	56.7	(±3.0)


4. Nutrition (continued)

ii. Percent of operations (and cows on these operations) having a laboratory nutritional analysis completed on purchased or raised feed in the last 12 months:

	<u>Percent</u>	<u>Standard Error</u>
Percent of operations	8.0	(±1.3)
Number of cows on these operations as a percent of cows on all operations	17.9	(±2.0)

b. Supplements and feed fed

Laboratory Nutritional Analysis Completed on Purchased or Raised Feed



i. Percent of operations feeding the following to the cow herd in the previous 12 months:

<u>Compound/Element</u>	<u>Percent of Operations</u>			
	<u>Fall/Winter (10/91 - 3/92)</u>	<u>Stand. Error</u>	<u>Spring/Summer (4/92-9/92)</u>	<u>Stand. Error</u>
Salt	63.0	(±3.0)	62.2	(±3.0)
Trace mineral salt	82.1	(±2.3)	78.6	(±2.5)
Phosphorus	35.5	(±2.8)	31.0	(±2.6)
Magnesium	41.3	(±3.1)	46.1	(±3.0)

ii. Percent of operations where Vitamin A is supplied to the cow herd by:


<u>Method of Delivery</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
Feeding alfalfa hay	29.1	(±2.6)
Using a mineral mix containing Vitamin A	49.9	(±3.1)
Injecting Vitamin A	7.4	(±1.6)
Using a protein supplement containing Vitamin A	42.6	(±3.0)

4. Nutrition (continued)

iii. Percent of operations providing the cow herd with access to pasture or crop residue by month:

<u>Month</u>	<u>Percent of Operations</u>			
	<u>Pasture</u>	<u>Standard Error</u>	<u>Crop Residue</u>	<u>Standard Error</u>
January	83.1	(±1.9)	24.4	(±2.3)
March	84.1	(±1.8)	16.2	(±2.0)
May	97.6	(±1.1)	6.2	(±1.8)
July	98.8	(±0.8)	6.6	(±2.0)
September	99.6	(±0.2)	10.4	(±1.9)
November	91.2	(±1.5)	26.5	(±2.5)

Cow Herd Access to Pasture and Crop Residue by Month



iv. Percent of operations feeding the following to the cow herd during 1992 by month:

<u>Month</u>	<u>Percent of Operations Feeding</u>			
	<u>Hay</u>	<u>Silage</u>	<u>Supplements</u>	<u>Grain</u>
January	91.9	6.1	53.3	28.5
Stand. Error	(±1.4)	(±1.3)	(±3.0)	(±2.8)
March	89.7	5.9	51.6	27.3
Stand. Error	(±1.8)	(±1.3)	(±3.0)	(±2.8)
May	20.3	3.8	17.9	7.8
Stand. Error	(±2.4)	(±1.1)	(±2.3)	(±1.6)
July	4.9	2.6	13.2	4.5
Stand. Error	(±1.3)	(±1.0)	(±1.9)	(±1.3)
September	11.6	2.6	19.2	7.0
Stand. Error	(±2.0)	(±1.0)	(±2.6)	(±1.8)
November	66.2	3.4	44.6	18.9
Stand. Error	(±2.8)	(±1.0)	(±3.0)	(±2.5)

4. Nutrition (continued)

v. For operations feeding the following feedstuffs in the indicated month, average pounds fed per head per day to the cow herd during 1992 by month:

<u>Month</u>	<u>Average Pounds per Head per Day</u>			
	<u>Hay</u>	<u>Silage</u>	<u>Supplements</u>	<u>Grain</u>
January	23.7	14.8	2.9	4.0
Stand. Error	(±0.9)	(±2.2)	(±0.4)	(±0.3)
March	23.4	14.7	2.9	4.0
Stand. Error	(±0.9)	(±2.2)	(±0.4)	(±0.3)
May	12.7	10.9	2.1	3.2
Stand. Error	(±1.2)	(±2.7)	(±0.2)	(±0.3)
July	8.3	4.5	1.8	2.8
Stand. Error	(±2.4)	(±1.6)	(±0.2)	(±0.3)
September	14.9	5.7	1.8	3.1
Stand. Error	(±2.0)	(±2.5)	(±0.2)	(±0.5)
November	20.2	9.8	2.7	3.8
Stand. Error	(±1.2)	(±2.8)	(±0.4)	(±0.4)

vi. Percent of operations (and replacement heifers on these operations) that fed an ionophore to replacement heifers in the previous 12 months:

	<u>Percent</u>	<u>Standard Error</u>
Operations	7.3	(±1.4)
Number of replacement heifers on those operations as a percent of replacement heifers on all operations	17.7	(±2.7)


4. Nutrition (continued)

c. Implants

i. Percent of operations (and animals on these operations) that implant:

	<u>Percent</u>		<u>Percent</u>		<u>Percent</u>	
	<u>Suckling Calves</u>	<u>Stand. Error</u>	<u>Weaning Age Calves</u>	<u>Stand. Error</u>	<u>Either Suckling or Weaning Age Calves</u>	<u>Stand. Error</u>
Operations	18.3	(±2.1)	17.6	(±1.9)	27.1	(±2.4)
Number of calves on those operations as a percent of calves on all operations	37.9	(±3.1)	28.9	(±2.5)	47.9	(±3.0)

**Cow/Calf Health & Productivity Audit
Implant Practices**



*Number of calves on these operations as percent of calves on all operations.

ii. Of those operations that implant, the percent of operations that implant heifers for replacement purposes (and animals on those operations):

	<u>Percent</u>		<u>Percent</u>	
	<u>Suckling Calves</u>	<u>Standard Error</u>	<u>Weaning Age Calves</u>	<u>Standard Error</u>
Operations	38.7	(±6.0)	17.0	(±5.2)
Number of calves on these operations as a percent of calves on operations that implant	47.3	(±5.6)	14.3	(±3.0)

d. Creep feeding

i. Percent of operations providing unweaned calves with access to creep feed:

<u>Percent of Operations</u>	<u>Standard Error</u>
27.4	(±2.7)

4. Nutrition (continued)

ii. Operation average number of days unweaned calves had access to creep feed:

<u>Number of Days</u>	<u>Standard Error</u>
113.7	(±10.9)


5. Animal health

a. Mineral deficiencies

i. Percent of operations identifying the following minerals as deficient or a cause of health problems in the herd over the previous 5 years:

<u>Mineral</u>	<u>Percent of Operations</u>	<u>Standard Error</u>
Phosphorus	2.8	(±1.2)
Magnesium	5.0	(±1.3)
Cobalt	0.5	(±0.5)
Copper	2.1	(±0.8)
Iodine	1.3	(±0.6)
Manganese	1.9	(±0.8)
Selenium	3.9	(±1.1)
Zinc	0.6	(±0.5)

Identification of Minerals as Deficient or Cause of Health Problems in Herd Last 5 Years



ii. Percent of operations reporting a toxic level of selenium in tissue or body fluid samples from the herd in the last 5 years:

<u>Percent of Operations</u>	<u>Standard Error</u>
0.3	(±0.3)

b. Deworming

i. Percent of operations deworming one or more class of beef cattle:

<u>Percent of Operations</u>	<u>Standard Error</u>
77.4	(±2.6)

5. Animal health (continued)


ii. Operation average number of times animals were dewormed in the previous 12 months:

<u>Group</u>	<u>Number of Times</u>	<u>Standard Error</u>
Calves (1-12 months)	0.8	(±0.0) ¹
Yearlings (13-23 months)	0.9	(±0.1)
2 year-olds	0.9	(±0.0) ¹
Cows	1.0	(±0.0) ¹
Bulls	1.0	(±0.0) ¹

iii. Percent of operations by number of times animals were dewormed in the last 12 months:

<u>Group</u>	<u>Percent of Operations</u>				<u>Total</u>
	<u>0</u>	<u>1</u>	<u>2</u>	<u>3 or More</u>	
Calves (1-12 months)	33.5	50.9	14.4	1.2	100.0
Stand. Error	(±3.1)	(±3.1)	(±2.1)	(±0.5)	
Yearlings (13-23 months)	33.9	42.7	20.5	2.9	100.0
Stand. Error	(±3.0)	(±3.0)	(±2.5)	(±1.0)	
2 Year-olds	33.5	42.7	21.5	2.3	100.0
Stand. Error	(±2.9)	(±3.0)	(±2.5)	(±0.8)	
Cows	25.9	49.0	23.3	1.8	100.0
Stand. Error	(±2.8)	(±3.1)	(±2.7)	(±0.7)	
Bulls	27.5	47.7	22.9	1.9	100.0
Stand. Error	(±2.7)	(±3.1)	(±2.6)	(±0.7)	

Percent of Operations by Deworming Practice



¹ Standard error values rounded to zero.

5. Animal health (continued)

iv. Importance of factors in deworming cattle:

<u>Factors</u>	<u>Percent of Operations by Level of Importance</u>			
	<u>Not</u>	<u>Moderate</u>	<u>Very</u>	<u>Extreme</u>
Always have dewormed cattle (tradition)	18.3	13.6	35.8	32.3
Standard Error	(±2.4)	(±2.3)	(±3.3)	(±3.3)
Recommendation of veterinarian	31.0	20.5	28.3	20.2
Standard Error	(±3.2)	(±3.0)	(±3.2)	(±2.5)
Recommendation of another consultant or friend	55.7	25.4	14.5	4.4
Standard Error	(±3.3)	(±3.0)	(±2.8)	(±1.5)
Animals have loose feces (diarrhea)	38.1	16.1	24.3	21.5
Standard Error	(±3.3)	(±2.4)	(±3.0)	(±3.1)
Animals were looking poor (rough hair coat, weight loss, anemia, bottle jaw)	24.6	8.2	26.9	40.3
Standard Error	(±2.9)	(±1.7)	(±3.1)	(±3.5)
Fecal egg count	64.6	9.7	11.7	14.0
Standard Error	(±3.3)	(±2.0)	(±2.5)	(±2.2)

c. Treatment - grubs, ticks, lice, and flies

i. Percent of operations treating (using dips, sprays, insecticides, eartags, powders, injections, etc.) cattle over the previous 12 months for:


	<u>Percent of Operations</u>	<u>Standard Error</u>
Cattle grubs (warbles, hypoderma)	60.8	(±3.1)
Ticks	59.7	(±3.1)
Cattle lice	73.7	(±2.9)
Flies	84.1	(±2.5)

6. Sources of animal health or beef production information

a. Importance of sources of animal health information:

Source	Percent of Operations by Level of Importance			
	Not	Moderate	Very	Extreme
Cooperative Extension Service or university specialists	44.7	26.7	20.4	8.2
Standard Error	(±3.0)	(±2.6)	(±2.6)	(±1.6)
Veterinarian	10.1	12.6	34.6	42.7
Standard Error	(±2.1)	(±2.0)	(±2.9)	(±3.0)
Beef magazine or agricultural journal	34.2	38.8	22.0	5.0
Standard Error	(±2.9)	(±2.9)	(±2.5)	(±1.1)
Producer organization	60.4	28.8	9.5	1.3
Standard Error	(±3.0)	(±2.7)	(±1.8)	(±0.5)
Retail salespeople (feed, vaccines, etc.)	41.7	27.9	25.2	5.2
Standard Error	(±2.9)	(±2.8)	(±2.8)	(±1.3)
Family member	43.1	21.2	21.7	14.0
Standard Error	(±3.0)	(±2.6)	(±2.7)	(±2.1)
Private consultant	79.4	12.5	6.4	1.7
Standard Error	(±2.6)	(±2.0)	(±1.8)	(±0.8)
Other producers	39.8	32.6	23.3	4.3
Standard Error	(±3.1)	(±2.9)	(±2.7)	(±1.1)
Radio/television/newspaper	58.1	27.8	11.1	3.0
Standard Error	(±3.1)	(±2.8)	(±1.8)	(±1.1)
Other source	77.2	15.3	3.8	3.7
Standard Error	(±2.6)	(±2.0)	(±1.2)	(±1.2)

Sources of Animal Health Information




6. Sources of animal health or beef production information (continued)

b. Importance of sources of beef production information:

Source	Percent of Operations by Level of Importance			
	Not	Moderate	Very	Extreme
Cooperative Extension Service or university specialists	45.8	25.0	22.6	6.6
Standard Error	(±3.0)	(±2.6)	(±2.6)	(±1.4)
Veterinarian	25.7	18.2	29.3	26.8
Standard Error	(±2.8)	(±2.2)	(±2.7)	(±2.6)
Beef magazine or agricultural journal	36.0	29.7	27.7	6.6
Standard Error	(±2.9)	(±2.6)	(±2.7)	(±1.3)
Producer organization	56.8	26.8	12.7	3.7
Standard Error	(±3.0)	(±2.7)	(±1.8)	(±1.1)
Retail salespeople (feed, vaccines, etc.)	39.8	29.4	22.8	8.0
Standard Error	(±2.9)	(±2.9)	(±2.8)	(±1.6)
Family member	41.4	23.7	22.4	12.5
Standard Error	(±2.9)	(±2.7)	(±2.6)	(±2.0)
Private consultant	79.6	14.4	4.1	1.9
Standard Error	(±2.5)	(±2.3)	(±1.1)	(±0.9)
Other producers	44.4	30.8	19.6	5.2
Standard Error	(±3.0)	(±2.9)	(±2.4)	(±1.2)
Radio/television/newspaper	59.0	26.2	13.0	1.8
Standard Error	(±3.0)	(±2.8)	(±2.0)	(±0.8)
Other source	76.5	15.5	4.1	3.9
Standard Error	(±2.6)	(±2.3)	(±1.2)	(±1.3)

Sources of Beef Production Information



6. Sources of animal health or beef production information (continued)

c. Importance of sources of nutritional information:

<u>Source</u>	<u>Percent of Operations by Level of Importance</u>			
	<u>Not</u>	<u>Moderate</u>	<u>Very</u>	<u>Extremely</u>
Private nutritionist	80.9	11.1	6.2	1.8
Standard Error	(±2.6)	(±2.2)	(±1.4)	(±0.8)
Feed salesman or feed retailer	27.6	28.8	32.1	11.5
Standard Error	(±2.7)	(±2.7)	(±3.0)	(±2.1)
Extension agent	50.8	25.8	20.2	3.2
Standard Error	(±3.0)	(±2.8)	(±2.2)	(±1.1)
Veterinarian	26.6	19.1	30.7	23.6
Standard Error	(±2.9)	(±2.3)	(±2.7)	(±2.6)
Friend or neighbor	39.5	35.2	18.4	6.9
Standard Error	(±2.9)	(±3.0)	(±2.3)	(±1.6)
Producer magazine	48.2	35.1	15.0	1.7
Standard Error	(±3.0)	(±3.0)	(±2.0)	(±0.7)
Personal knowledge/education	6.4	7.0	32.9	53.7
Standard Error	(±1.5)	(±1.8)	(±2.9)	(±3.1)

d. Nutrition consultants

i. Percent of operations consulting an animal nutritionist in the previous 12 months:

<u>Percent of Operations</u>	<u>Standard Error</u>
10.6	(±1.7)

**National Animal Health Monitoring System
USDA:APHIS:VS
2150 Centre Ave., Bldg. B, MS 2E7
Fort Collins, Colorado 80526-8117
(970)494-7000**

N135.0194