

TABLE OF CONTENTS**VOLUME I**

	Page
FOREWORD	III
ABSTRACT	V
EXECUTIVE SUMMARY	VII
1.0 INTRODUCTION	1-1
2.0 MATERIAL AND METHODS	2-1
2.1 Access Point Survey Methods	2-1
2.1.1 Development of the Sampling Frame and Stratification.....	2-1
2.1.2 Scheduling of the Access Point Sampling	2-4
2.2 Aerial Survey Methods	2-7
2.2.1 Development of the Sampling Frame and Stratification.....	2-7
2.2.2 Scheduling of Angler Counts	2-7
2.3 Estimating equations	2-11
2.3.1 Catch and Harvest Rates and Mean Trip Length.....	2-11
2.3.2 Daytime Effort	2-13
2.3.3 Night effort.....	2-17
2.3.4 Total Catch and Harvest Estimates	2-18
3.0 RESULTS	3-1
3.1 Angler Fishing Effort.....	3-1
3.1.1 Daytime Fishing Effort	3-1
3.1.2 Night Fishing Effort.....	3-18
3.2 Angler Catch and Harvest Statistics	3-20
4.0 DISCUSSION	4-1
5.0 REFERENCES	5-1
6.0 ACKNOWLEDGEMENTS	6-1

**VOLUME II
APPENDICES**

A SAMPLING FRAME FOR THE ACCESS POINT SURVEY	A-1
B ACCESS POINT SAMPLING SCHEDULE	B-1

TABLE OF CONTENTS (CONT'D)

	Page
C SPECIES LIST AND NUMBER OF FISH CAUGHT BY ANGLERS INTERVIEWED	C-1
D CATCH AND HARVEST STATISTICS WITHIN AND ACROSS GEOGRAPHIC AND TEMPORAL STRATA	D-1
E FIELD DATA COLLECTIONS FORMS	E-1

LIST OF TABLES

Table	Page
2-1 Original Temporal Stratification.	2-2
2-2 Monthly Temporal Post-stratification.....	2-3
2-3 Number of sampling shifts that were completed in the access point survey by spatio-temporal strata.....	2-5
2-4 Delaware River Creel Survey 2002 - Scheduled and completed flights in the Estuary & Stratum 1.....	2-8
2-5 Delaware River Creel Survey 2002 - Scheduled and completed flights in Strata 2 and 3 (including East Branch)	2-9
2-6 Scheduled and completed flights for the second aerial survey in stratum 1. These flights were conducted during peak angler effort, using the Dauk and Schwarz (2001) flight scheduling method	2-10
2-7 Focus species and their Integrated Taxonomic Information System (ITIS) taxonomic serial number (ITSN), as recorded in the survey database.	2-12
3-1 Number of interviews completed within and across spatial and temporal strata. Stratum 3 includes the East Branch.	3-2
3-2 Aerial sampling effort (number of primary sampling units) by spatio-temporal strata. An instantaneous count was obtained for each PSU. Stratum 3 includes the East Branch.	3-4
3-3a Total daytime fishing effort for boat anglers for the non-tidal and tidal portions of the river estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Estimates based on original temporal stratification. .	3-5
3-3b Total daytime fishing effort for boat anglers for the non-tidal and tidal portions of the river estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Estimates based on monthly temporal post-stratification.	3-6
3-4a Total daytime fishing effort for shore anglers for the non-tidal and tidal portions of the river estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Estimates based on original temporal stratification.	3-7
3-4b Total daytime fishing effort for shore anglers for the non-tidal and tidal portions of the river estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Estimates based on monthly temporal post-stratification.	3-8
3-5a Total daytime fishing effort for boat and shore anglers for the non-tidal and tidal portions of the river estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Estimates based on original temporal stratification.	3-9

LIST OF TABLES (Cont'd)

Table	Page
3-5b Total daytime fishing effort for boat and shore anglers for the non-tidal and tidal portions of the river estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Estimates based on monthly temporal post-stratification.	3-10
3-6a Total daytime fishing effort by fishing mode in the non-tidal stratum 2 estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Based on original temporal stratification. Mode legend B= boat anglers, S= shore anglers, All = boat and shore anglers combined.	3-11
3-6b Total daytime fishing effort by fishing mode in the non-tidal stratum 3 estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Based on original temporal stratification. Mode legend B= boat anglers, S= shore anglers, All = boat and shore anglers combined.	3-12
3-7 Total night (9 PM to midnight) fishing effort (angler hours) for the non-tidal and tidal portions of the river estimated from the aerial survey, along with estimates of precision (relative standard error, RSE). Estimates based on monthly temporal stratification. Boat and shore anglers combined.	3-18
3-8 Total angler catch and harvest (daytime fishing) for the four key species and two species most frequently caught throughout the study period from March 17 to October 31, based on the original temporal strata. Boat and shore fishing modes are combined. ¹ 49 % were females.....	3-23
3-9 Total angler catch and harvest (daytime fishing) for the four key species and two species most frequently caught throughout the study period from March 17 to October 31, based on the monthly post- strata. Boat and shore fishing modes are combined. ¹ 49% were females	3-24
3-10a Total angler catch and harvest (daytime fishing) of American shad by original temporal strata. Boat and shore fishing modes are combined. ¹ 50% were females; ² 49% were females	3-24
3-10b Total angler catch and harvest (daytime fishing) of American shad by monthly post-strata. Boat and shore fishing modes are combined. ¹ 50% were females; ² 49% were females	3-25
3-11a Total angler catch and harvest (daytime fishing) of striped bass by original temporal strata. Boat and shore fishing modes are combined.	3-25
3-11b Total angler catch and harvest (daytime fishing) of striped bass by monthly post-strata. Boat and shore fishing modes are combined.....	3-25
3-12 Total angler catch and harvest (daytime fishing) for the non-tidal stratum 2 for selected species throughout the study period from March 17 to October 31, based on the original temporal stratification.	3-26

LIST OF TABLES (Cont'd)

Table		Page
3-13	Total angler catch and harvest (daytime fishing) for the non-tidal stratum 3 for the six species with highest catch throughout the study period from March 17 to October 31, based on the original temporal stratification. No catches of any species were reported in the East Branch, from Hancock, NY, to Downsville, NY.	3-26
3-14	Total angler catch and harvest during night (9 PM to midnight) for the four species with highest catch throughout the study period from March 17 to October 31. Boat and shore anglers combined.	3-27

LIST OF FIGURES

Figure	Page
3-1 Proportion of total daily fishing effort (angler hours) by 2-hour time intervals (military time) in the non-tidal portion of the river during weekdays for the original temporal strata 1-3, and 4-7.....	3-13
3-2 Proportion of total daily fishing effort (angler hours) by 2-hour time intervals (military time) in the non-tidal portion of the river during weekends for original temporal strata 1-3, and 4-7.	3-13
3-3 Proportion of total daily fishing effort (angler hours) by 2-hour time intervals (military time) in the tidal portion of the river during weekdays for original temporal strata 1-3, and 4-7.	3-14
3-4 Distribution of total daily fishing effort (angler hours) by 2-hour time intervals (military time) in the tidal portion of the river during weekends and holidays for original temporal strata 1-3, and 4-7.	3-14
3-5 Proportion of total fishing effort (angler hours) in the non-tidal portion of the river by fishing mode, based on the aerial survey and the access survey.	3-15
3-6 Proportion of total effort fishing (angler hours) in the tidal portion of the river by fishing mode, based on the aerial survey and the access survey.	3-16
3-7 Proportion of total effort fishing (angler hours) accounted for by boat anglers in the non-tidal stratum 3, based on the aerial survey and the access survey.	3-17
3-8 The proportion of total angler effort by daytime (7 AM to 9 PM) and night (9 PM to midnight) for the total sample period in the non-tidal portion of the River.....	3-19
3-9 The proportion of total angler effort by daytime (7 AM to 9 PM) and night (9 PM to midnight) for the total sample period in the tidal portion of the River.....	3-19
3-10 The species composition of catches (proportions) reported by anglers interviewed in the tidal portion of the river throughout the study period. Boat and shore anglers combined.....	3-28
3-11 The species composition of catches (proportions) reported by anglers interviewed in the non-tidal portion of the river (Strata 1 and 2) throughout the study period. Boat and shore anglers combined. It is likely that some blue gill was misidentified by anglers, and a large portion of the reported catch may represent redbreast sunfish.	3-28
3-12 The species composition of catches (proportions) reported by anglers interviewed in the non-tidal stratum 3 throughout the study period. Boat and shore anglers combined.	3-29
3-13 The species composition of catches (proportions) reported by anglers interviewed at night (9 PM to midnight) in the tidal portion of the river throughout the study period. Boat and shore anglers combined.	3-29
3-14 The species composition of catches (proportions) reported by anglers interviewed at night (9 PM to midnight) in the non-tidal portion of the river throughout the study period. Boat and shore anglers combined.	3-30

LIST OF FIGURES (Cont'd)

Figure	Page
3-15 Proportion of the total catch of American shad by fishing mode in the non-tidal portion of the river	3-31
3-16 Proportion of the total catch of American shad by fishing mode in the tidal portion of the river	3-31
3-17 Proportion of the total catch of striped bass by fishing mode in the non-tidal portion of the river	3-32
3-18 Proportion of the total catch of striped bass by fishing mode in the tidal portion of the river	3-32
3-19 Proportion of targeted species within anglers that target specific species for the non-tidal portion of the river from March to June.	3-33
3-20 Proportion of targeted species within anglers that target specific species for the non-tidal portion of the river from July to October.	3-33
3-21 Proportion of targeted species within anglers that target specific species for the tidal portion of the river from March to June.....	3-34
3-22 Proportion of targeted species within anglers that target specific species for the tidal portion of the river from July to October.....	3-34
3-23 Size distribution (total length by sex) of all harvested American shad measured by creel clerks throughout the study period.	3-35
3-24 Size distribution of all striped bass measured by creel clerks throughout the study period. Specimens below minimum size limits were generally released.	3-35
3-25 Demographic composition of state residency for anglers interviewed in the non-tidal portion of the river.	3-36
3-26 Demographic composition of state residency for anglers interviewed in the tidal portion of the river.	3-36
3-27 Demographic composition of Pennsylvania county residency for all anglers interviewed.....	3-37
3-28 Demographic composition of gender for all anglers interviewed.....	3-37
3-29 Demographic composition of age groups for all anglers interviewed.	3-38
4-1 Estimated effort by boat and shore anglers in the non-tidal portion of Delaware River from March 15 to May 31.	4-6
4-2 Daily mean streamflow value recorded at U.S. Geological Survey gauging station 01463500 at Trenton, NJ.	4-7
4-3 Estimated total catch and harvest of American shad in the non-tidal section of the Delaware River from March through May.	4-8