

**Climatological Data
for Clouds Over the
Globe from
Surface Observations,
1982 - 1991:**

THE TOTAL CLOUD EDITION

Carole J. Hahn, University of Colorado

Stephen G. Warren, University of Washington

Julius London, University of Colorado

CLIMATOLOGICAL DATA FOR CLOUDS OVER THE GLOBE
FROM SURFACE OBSERVATIONS, 1982-1991:
THE TOTAL CLOUD EDITION

Carole J. Hahn
Cooperative Institute for Research in Environmental Sciences
University of Colorado
Boulder, Colorado 80309

Stephen G. Warren
Department of Atmospheric Sciences
University of Washington
Seattle, Washington 98195

Julius London
Department of Astrophysical, Planetary and Atmospheric Sciences
University of Colorado
Boulder, Colorado 80309

✓ Environmental Sciences Division
Publication No. 4348

Date Published: October 1994


Printed for the
Global Change Research Program
Environmental Sciences Division
Office of Health and Environmental Research
U.S. Department of Energy
(KP 05 00 00 0)

Printed by the
Carbon Dioxide Information Analysis Center
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831-6335
managed by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under contract DE-AC05-84OR21400

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER



DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

TABLE OF CONTENTS

	<u>Page</u>
LIST OF TABLES	4
LIST OF FIGURES	5
ABSTRACT	7
1. INTRODUCTION	9
2. DATA SOURCE AND ANALYSIS	9
A. Data Sources	9
B. Data Analysis	9
1) Processing of Weather Reports	9
2) Determination of Cloudiness at Night	10
3) Averaging Methods	11
C. Grid Sizes	11
3. CONTENTS AND ORGANIZATION OF THE DATA ARCHIVE	12
A. General	12
B. Details of Organization	13
1) Map Groups and Data Formats	13
2) Group Headers	14
C. Details of Contents	15
1) The Missing Value Code	15
2) File 2: Grid Information; format 10	15
3) File 3: Land and Ocean Combined; formats 22,32	16
4) Files 4-7: Total Cloud Cover; format 22	16
5) Files 8-15: Weather Types; format 32	17
6) File 16: Harmonics, Interannual Variations and Trends; formats 40-42, 51-52	17
4. HOW TO OBTAIN THE DATA	19
ACKNOWLEDGEMENTS	19
REFERENCES	20
TABLES	21
FIGURES	31

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Cloud Information Contained in Synoptic Weather Reports	21
2. Cloud and Weather Type Definitions Used in Total Cloud Edition	21
3. Grid Box Sizes	22
4. File Information	23
5. Data Organization	24
6. List of Formats for Reading Data Records	26
7. Map Group Header Record Format and Codes	27
8. Terms and Abbreviations Used	28

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1. Flow chart of data selection and checking	31
2a. Annual Average Total Cloud Cover (%), Land & Ocean, 1982-1991	32
2b. Annual Precipitation Frequency (%), Land & Ocean, 1982-1991	33
3a. Phase of Annual Cycle in Total Cloud Cover for Ocean	34
3b. Phase of Annual Cycle in Precipitation Frequency for Ocean	35
3c. Phase of Annual Cycle in Fog Frequency for Ocean	36
4a. Diurnal Cycle in Total Cloud Cover for DJF (1982-1991) Ocean	37
4b. Diurnal Cycle in Total Cloud Cover for MAM (1982-1991) Ocean	38
4c. Diurnal Cycle in Total Cloud Cover for JJA (1982-1991) Ocean	39
4d. Diurnal Cycle in Total Cloud Cover for SON (1982-1991) Ocean	40
5a. Diurnal Cycle in Total Cloud Cover for DJF (1982-1991) Land, over China	41
5b. Diurnal Cycle in Total Cloud Cover for JJA (1982-1991) Land, over the U. S.	42

ABSTRACT

Hahn, C.J., S.G. Warren, and J. London, 1994: *Climatological Data for Clouds Over the Globe from Surface Observations, 1982-1991: The Total Cloud Edition*. NDP026A, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN. (Also available from Data Support Section, National Center for Atmospheric Research, Boulder, CO.)

Routine, surface synoptic weather reports from ships and land stations over the entire globe, for the ten-year period December 1981 through November 1991, were processed for total cloud cover and the frequencies of occurrence of clear sky, precipitation, and sky-obscured due to fog. Archived data, consisting of various annual, seasonal and monthly averages, are provided in grid boxes that are typically $2.5^\circ \times 2.5^\circ$ for land and $5^\circ \times 5^\circ$ for ocean. Day and nighttime averages are also given separately for each season. Several derived quantities, such as interannual variations and annual and diurnal harmonics, are provided as well. This data set incorporates an improved representation of nighttime cloudiness by utilizing only those nighttime observations for which the illuminance due to moonlight exceeds a specified threshold. This reduction in the night-detection bias increases the computed global average total cloud cover by about 2%. The impact on computed diurnal cycles is even greater, particularly over the oceans where is found, in contrast to previous surface-based climatologies, that cloudiness is often greater at night than during the day.

1. INTRODUCTION

This report describes a data archive that contains global coverage of analyzed cloud data. The cloud data contained in this archive (the "Total Cloud Edition") are for total cloud cover. In addition, the frequencies of occurrence of clear sky, precipitation, and sky-obscured due to fog (collectively referred to here as "weather types") are also included. Analyses of cloud types are not provided here. All data utilized here were obtained from routine, surface synoptic weather reports for the ten-year period December 1981 through November 1991. The specific contents of the archive are described in Section 3. Briefly, the archive contains various annual, seasonal and monthly averages for total cloud and the three weather types in grid boxes that are typically $2.5^\circ \times 2.5^\circ$ for land and $5^\circ \times 5^\circ$ for ocean. Several derived quantities, such as interannual variations and annual and diurnal harmonics, are also provided.

Previously the authors prepared a similar cloud data archive that spanned the years 1930 to 1981 (Hahn et al., 1988). Some of the data contained in that archive were presented in published atlases (Warren et al., 1986, 1988). The present archive represents not only an extension of the time period analyzed, but an improvement in the analysis scheme that results in more reliable estimates of cloudiness at night. This not only leads to more accurate daily averages, but, more significantly, leads to a more reliable determination of diurnal cycles than has been obtained previously from surface observations on a global scale.

2. DATA SOURCE AND ANALYSIS

A. Data Sources

For land stations, synoptic weather reports were obtained from the National Meteorological Center (NMC). Only those stations which have been assigned official station numbers by the World Meteorological Organization (WMO) were utilized. About 124 million reports were available for cloud analysis for the 10-year period December 1981 through November 1991 (referred to as 1982-91). Synoptic reports are recorded 8 times per day: 00, 03, 06, 09, 12, 15, 18, 21 GMT. However, many stations report only every 6 hours (notably those in the United States and Australia), some less often, and some only during the daytime.

Ship reports were obtained from the Comprehensive Ocean-Atmosphere Data Set (COADS), Interim Product CMR5 Reports (Woodruff et al., 1987). There were 14.4 million reports available for cloud analysis over the oceans.

B. Data Analysis

1) Processing of Weather Reports.

Synoptic weather reports are coded according to the system given by the World Meteorological Organization (WMO, 1988). The information in these reports that relates to cloud analysis is summarized in Table 1. For the total cloud and weather type analyses reported here, only N, ww, and

I_x are of direct relevance. However, N_h , C_L , C_M and C_H were used in error checking. Definitions of the cloud and weather types analyzed here are given in Table 2.

The flow chart in Figure 1 shows the processing and quality control checks performed on each weather report read from the original archives (NMC or COADS). The percentage of reports discarded at each stage of the processing is indicated. Land and ship reports required slightly different checks in the early stages of processing but were treated identically below the horizontal dashed line in the upper portion of the figure. If a land station did not have a WMO station number it was discarded (many of these were from the United States), thus ensuring more uniformity in reporting procedures. If a ship report was known to be from a buoy (from the "deck" number in the COADS data) it was discarded. Any report that had no cloud information ($N=/$) was discarded.

In 1982 WMO introduced several coding procedure changes (WMO, 1988). One of these changes now instructs observers to set $ww=/$ if present weather was either "not available" or "observed phenomena were not of significance" (ww codes 00-03 are considered to represent phenomena without significance). The present weather indicator, I_x , is used to distinguish these cases. Land station reports with I_x values of 4, 5 or 6 signify automatic weather stations and were discarded. Reports with $I_x=3$ (data not available) were also discarded because without ww it is not possible to interpret cases of $N=9$ (see Warren et al., 1986) or to evaluate the occurrence of precipitation. $I_x=2$ indicates that observed phenomena were not of significance, while I_x is coded as "1" when ww is given. Occasionally $I_x=1$ when $ww=/$. These inconsistent reports were also discarded.

Examination of the NMC data set showed that while land station reports conformed to this new coding procedure almost immediately, ship reports did not incorporate I_x coding consistently until 1985. The COADS data set does not even contain I_x . Thus some ship reports that should be discarded on the basis of I_x were kept. At the horizontal dashed line in Figure 1 there were 125 million land reports and 15.8 million ship reports remaining. The discard fractions below the line are fractions of these numbers.

If the sky was obscured due to fog (1.1% land, 2.5% ship), thunderstorms (0.05% land, 0.17% ship), or rain/snow (0.4% land, 1.1% ship), the sky was considered to be overcast ($N=8$). This source of "cloudiness" contributed about 1% to the total cloud cover globally, and much more in some locations and seasons (Hahn et al., 1992).

Other data consistency checks are indicated in the figure. The final one tests whether the reported latitude and longitude of a land station puts the station on water (rare) or whether reported latitude and longitude of a ship puts the ship on land (1.3%). The reports that survive these tests (124.2 million for land and 14.4 million for ships) are used to compute total cloud cover and the frequencies of occurrence of clear sky, fog, and precipitation. Cloud types were not analyzed further in this study.

2) Determination of Cloudiness at Night.

The ability of surface observers to adequately detect clouds at night has been questioned for many years (e.g. Riehl, 1947; Schneider et al., 1989). In an attempt to find a practical solution to this "night-detection-bias", the authors (Hahn et al., 1994) analyzed ten years of nighttime data for the zone 0-50°N and plotted reported cloud cover as a function of the illumination due to moonlight. The illuminance function used by the authors depends on the phase and altitude of the moon and on the distance of the moon from the earth. It was found that the amount of total cloud reported at night increased as the illuminance of the moon increased up to a certain threshold, after which reported

cloud amounts leveled off. This threshold is referred to as "the illuminance criterion" and corresponds to the light produced by the twilight sun at an altitude of about 9 degrees below the horizon. Thus the illuminance criterion is met when either the sun is at an altitude greater than -9° or the position of the moon is such that its illuminance exceeds the threshold. These conditions can be determined for each report with the use of an ephemeris and the latitude, longitude, and time of the report.

By using only reports for which the illuminance of the moon (or sun) exceeded the threshold illuminance, we can obtain more reliable estimates of nighttime cloudiness than have been previously obtained from surface observations. Application of the illuminance criterion increases the computed global average total cloudiness at night by about 4% and thus increases the daily average cloudiness by about 2%. There is also a significant effect on computed diurnal cycles which will be demonstrated in Section 3.

This illuminance criterion was applied in the analyses of total cloud cover and clear-sky frequency archived here, but not for fog and precipitation whose detection does not depend on illumination. (For comparative purposes, some analyses of total cloud and clear sky were also performed utilizing all observations as described in Section 3.) Application of the illuminance criterion caused 27% of the land reports and 24% of the ship reports to be discarded, leaving 90.4 million reports for land and 10.9 million reports from ships.

3) Averaging Methods.

An average for a synoptic hour, or for daytime or nighttime only, was obtained simply by averaging all the contributing reports, whether for a single year or a multi-year average. Because many nighttime reports are discarded due to the illuminance criterion, there are far fewer contributing reports at night than during the daytime. Therefore, to obtain the "daily" average, daytime and nighttime averages are first determined separately and then averaged together. For this purpose, daytime is considered to be 06-18 local time (determined from the longitude at the center of the grid box in which the observation was made) and nighttime is considered to be 18-06 local time. A daily average was obtained by this method if there were at least 50 observations contributing to both the day- and nighttime averages. If there were less than 50 observations at night (day) but 100 or more for the daytime (night), then only the daytime (nighttime) observations were used for the average. Otherwise averages were obtained by using all available observations, regardless of time of day. This method was applied uniformly over the globe, even though the method loses significance near the poles. (The poles themselves were considered to be on Greenwich Mean Time.) The particular method used in computing an average is coded in the data record which is described in Section 3.

It should be noted that in a single month at a single point on earth the moon will be above the horizon at night only for about 2 weeks. Thus a nighttime average for a single month, when the illuminance criterion has been applied, will not be fully representative of that month. Longer term averages will be statistically more reliable. For this reason, monthly mean daily averages are not provided in this data archive, although the data for obtaining them are available in the "monthly means by synoptic hour" that are provided (Section 3).

C. Grid Sizes

The globe was divided into grid boxes for which the various cloud quantities were computed. The three grid sizes used in these analyses are, nominally: $2.5^\circ \times 2.5^\circ$, $5^\circ \times 5^\circ$, and $10^\circ \times 20^\circ$ latitude x longitude. Because the area contained within a $5 \times 5^\circ$ box, for example, decreases with

increasing latitude, boxes poleward of 50° latitude were made to encompass a wider longitude range such as 5x10° or 5x20°, etc. A "c" is used to symbolize this condensation or contraction. Thus 5x5c (or 5c for short) means 5x5° between 50N and 50S but a larger longitude width poleward of 50° latitude. The three grid sizes used are described in Table 3.

Each grid box is assigned a number. The numbering goes from west to east (beginning with the Greenwich Meridian) and north to south. The west and south borders of a box are considered to be within the box (90°N is also considered to be within box 1). The latitude and longitude at the center of each numbered box for each grid are given in this archive (Section 3).

The 5c grid is used for most analyses over the ocean and for some land analyses. The 10c grid is used for some ocean analyses because relatively sparse ocean data make some analyses at the smaller grid size unreliable. The 2c grid is used only for analyses over land where finer resolution is practical.

3. CONTENTS AND ORGANIZATION OF THE DATA ARCHIVE

A. General

The data are divided into 15 files, numbered 2 to 16 as shown in Table 4. File 1 is a brief documentation of the archive, containing excerpts from this report. The organization of data into files is based on similarity of content and data format. Total cloud data are generally separated from weather type data. Land and ocean data are generally in separate files as well. There are many grid boxes that contain both land and ocean, and two separate values are retained in this way. The user can combine the two if desired, although land and ocean values are given at different grid scales making suitable averaging necessary. File 3 contains land and ocean data merged on a 5c grid for selected long-term averages (see below). The files listed with the same group cluster name would logically belong within the same file. However, monthly averages by synoptic hour require so much storage space that they are placed in separate files.

A detailed breakdown of the contents of each data file is given in Table 5. Each data file contains a series of "map groups", each of which consists of gridded data for total cloud or weather type averages for a particular season or year or time of day or for grid information (File 2) as indicated in the table. A map group is made up of the data records for a number of grid boxes over the globe and a header record which identifies the group:

- Header record identifying map group
- Data record for first reported box
- Data record for second reported box
- etc. for number of boxes specified in header.

This pattern is repeated throughout each file in the order indicated in Table 5. Data record formats shown in Table 6 and the header record described in Table 7 are discussed in the next section.

The number of data records within a map group depends on the grid size and whether it is for land or ocean. Since there would be no land data in an ocean-only grid box (and vice versa), data records for such boxes are uniformly not written. (Box numbers based on the full grid are provided in the data record itself.) The numbers of boxes given within each type of map group are listed in a

footnote to Table 5 (and are coded in the header record). Thus, in File 6, for example, each ocean 5c map group contains 1494 logical records - a header record and 1493 data records, while a land 5c map group in File 4 contains a total of 862 logical records. While there are actually 934 5c boxes with land fractions greater than 0.0001 (and 27 additional boxes with reporting stations on small islands), only 861 boxes have data for the period analyzed here and only those boxes are archived. Similarly there are 3027 2c boxes with land fractions >0.0001 (and 48 additional boxes with reporting stations on small islands) but only 2309 boxes have data. The 10c grid contains so few boxes that it is convenient to retain all 230 boxes, including the 16 land-only boxes. In file 3, where land and ocean are combined on a 5c grid, all 1820 boxes are retained.

B. Details of Organization

The use of Table 5, along with Tables 6 and 7, should enable the user to find any desired quantity, once a few conventions are understood. Abbreviations used are listed alphabetically in Table 8.

1) Map Groups and Data Formats.

Each data file shown in Table 5 is a series of map groups which are numbered consecutively within each group cluster. Each group contains the data relevant to the cloud or weather type quantities listed under the contents heading. These data are given for each reported grid box according to the indicated data format which is described in Table 6. For example, group 2 in File 6 contains data relevant to mean seasonal (DJF, 1982-91) total cloud cover over ocean on a 5c grid. These data are organized according to format 22. Table 6 shows that format 22 specifies the box number, the number of observations, the average amount of total cloud (given to hundredths of a percent), the standard deviation of the observations contributing to the average (given to tenths of a percent), and indicator telling whether contributing observations were from daytime, nighttime, or both (see IDN in Table 8), an indicator telling whether the observations were from land stations or ships, and the number of seasons contributing to the average (relevant only for annual averages).

Data format numbers are given as 2-digit integers. The tens digit distinguishes 5 data classes as shown in Table 6. The units digit is used to distinguish some small difference in the meaning of a variable represented. Format 40 is used to distinguish the fact that the phase of the annual harmonic is given in units of months as opposed to hour for the diurnal harmonic in formats 41 and 42. The differences between formats 41 and 42 (as well as 51 and 52) are simply that the data variables refer to amount or frequency, respectively. While these distinctions (and that between formats 22 and 32) are not essential in this "Total Cloud Edition", these numbers are retained to be consistent with our previous archive (Hahn et al., 1988) and a possible future archive containing cloud type data.

The order in which the groups follow each other, with respect to season, time of day, year or weather type, can also be determined from Table 5. Where there are simply 4 seasons (or 12 months) as with File 4 groups 2-5 (or 42-53), all the boxes for the first season (or month) are followed by the next group header and all the boxes for the second season (or month) and so on. The order in which the seasons are given is DJF, MAM, JJA, SON; months are given in the order Dec, Jan, Feb, . . . Nov. In cases such as for File 4 groups 10-41 or 98-137 where more than one time or year is given for each season, the convention adopted is to increment the parameter listed first while holding the parameter listed second constant. Thus the order for groups 10-41 would be 8 synoptic hours (in order of increasing hour) for DJF, then 8 synoptic hours for MAM, etc. The order for groups 98-137 would be 4 seasons for 1982, 4 seasons for 1983, etc. Thus the individual seasons follow each other

chronologically. The order in which weather type groups (as in File 8) follow each other is the numerical order of the numeric codes shown in Table 7. Using the convention of incrementing the leftmost group parameter first, it can be determined, for example, that the group number for precipitation frequency over land for MAM 1982-91 at 03 GMT is 69.

2) Group Headers.

The first record in each map group is a header record which identifies the group. The format of this header record is shown in Table 7. The first parameter of the header record gives the map group number. These numbers have no special significance other than that they run sequentially through a group cluster (Tables 4 and 5) and may aid in locating or identifying a map group.

The next three parameters specify the number of boxes reported in the group (the number of records to be read before reaching the next map group; this number may be less than the total number of boxes in a grid as discussed above), the grid size of the boxes, and whether the data are for land or ocean or both. This latter parameter, LO, differs from the parameter LOB in Table 6 in that LO specifies the *intent* of the map group while LOB indicates the *actual* condition for a particular box.

The IMOON parameter indicates whether the illuminance criterion was applied in the analysis of the data for the group. The last 5 parameters in the group header indicate the year, season (or month), time of day, cloud or weather type to which the data refer, and the data format.

Even without the map group number, the other 9 parameters together uniquely define each map group. The map group headers, along with example data records, that correspond to the examples given in the last section from Files 6 and 8 are:

21493	520829141-1	122				692309	211829142	3	332
1	{first data record}					24	{first data record}		
381	5182	7435	297	32-9		1444	50	200	-9 21-9
1809	{last data record}					7290	{last data record}		

The first example is for daily-average total cloud cover over the ocean with the illuminance criterion applied for DJF, 1982-91. There are 1493 data records in this 5c map group. The first data record is for box 1 and the last is for box 1809. The example box 381 (in the eastern North Pacific) has 5182 observations with an average of 74.35% total cloud cover. The standard deviation of the observations is 29.7%. The daily average was obtained by averaging daytime and nighttime averages. Only ship data contributed. The "-9" signifies that no value is reported for the last variable, NSN (see *The Missing Value Code* on page 15).

The second example is for the frequency of occurrence of precipitation over land at 03 GMT for MAM, 1982-91. The illuminance criterion was not applied. There are 2309 data records in this 2c map group. The first data record is for box 24 and the last is for box 7290. The example box 1444 (in the western United States) had only 50 observations during this time, giving an average precipitation frequency of 2.00%. The standard deviation is not given. These data are for nighttime over land.

C. Details of Contents

All data on this tape were written as integers. Floating point quantities were multiplied by a power of 10 and rounded off to give an integer. When read using the specified formats shown in Table 6, the proper floating point values are recovered. This was demonstrated with the examples given in the last section. Cloud cover and frequencies are given as percent. These and the units of other variables defined in the data formats are defined in Table 8.

After discussing the missing value code, details or peculiarities of the contents of each data file will be discussed. Refer to Tables 5, 6 and 7 for references to file contents, format numbers and header coding, respectively. Any non-standard terms not defined in the text can be found in Tables 1, 2, 3 or 8.

1) The Missing Value Code.

Any data variable for which no value is reported is assigned the "missing value code", which was chosen to be -9. Thus, when reading in the various floating point notations, the actual value obtained may vary but will always be less than 0. The trend (File 16, formats 51 & 52) is the only data variable for which a negative value is valid. Thus for trends the missing value code should be considered to be NYRS=0. Note that any time NOBS is zero, data variables such as AMT, FQ or SD will be assigned the missing value code, but it is possible for data variables, such as SD, to be assigned the missing value code even when NOBS is not zero.

2) File 2: Grid Information; format 10.

The three groups in this file contain data related to the grids used for dividing the globe (see Table 3). The parameters YEAR, SN, TIME and TYPE in the group header are set to -9. Inclusion of the box number in this data format is redundant since here all boxes are reported and data records are in box number order. The latitude (90 to -90 for N to S) and longitude (0 to 360E) of a box center are given to two decimal places. Since the gridded data in all subsequent files are identified only by box number, the information here allows those data to be located on a map. The inverse relation (converting latitude and longitude to box number) can be achieved with a simple mathematical relationship (Hahn et al., 1988). The fraction of each grid box that is land is given to four decimal places. Here "land" means "not ocean" since lakes and ice shelves are counted as land. The method for determining these fractions and a map for the 5c grid are given in Warren et al. (1986). Ocean fraction is 1 - land fraction.

The number of land stations in a 2c box was approximated by taking the number of observations over the 10-year period for a single season (MAM) for either 00 or 12Z, whichever was daytime for a box, and dividing by 10 (for the number of years) and by 3 (for the number of months in a season) to get the average number of reports per month at a single reporting time. If each station always reported, there should then be 30 reports per station. Since station reports are sometimes missing, the above number was divided by 25 to obtain an approximate value for the number of stations. This number may be of value in estimating the reliability or representativeness of data within a box. NLSTA is set to -9 in the 5c and 10c data records.

3) *File 3: Land and Ocean Combined; formats 22,32.*

This is the only file in which land and ocean values were merged onto a single grid, and this was done only for 10-year mean values (monthly, seasonal and annual). An average (total cloud cover or weather type frequency), for any grid box for which both land and ocean values contributed, was determined by weighting the contributing land and ocean values by their respective fractional area within the box. Only averages made from 100 or more observations were allowed to contribute. The variables SD and IDN (and NSN for seasonal and monthly averages) were set to -9 in the data record. LOB indicates whether land, ocean, both or neither contributed to the box. For the annual averages, NSN is the sum of the number of seasons that contributed to the land and ocean annual averages before merging, and so can have values 0 to 8 (in this file, -9 is used rather than 0). If NSN is 3 to 7 and LOB is 3, it cannot be known how many seasons were contributed by land or ocean without examining the annual map groups for land and ocean separately in their respective files.

NOBS is the sum of the number of observations contributed by land and ocean. For annual averages on a 5c grid, it is possible for NOBS to be greater than 999999 (the maximum allowed in the 16 format) since some boxes, notably in Europe, have a large number of land stations. NOBS in such cases was set to 999999. (The actual number can be retrieved, if desired, by adding up the number of observations in the contributing boxes and seasons given in the other files.)

Figures 2-5 are provided to illustrate some of the information available in this archive and serve as examples against which user output can be checked. The first map group of File 3 contains the data for the global distribution of annual average total cloud cover for the 1982-91 period with the illuminance criterion applied. This is shown in Figure 2a. The global distribution of annual average precipitation frequency (from all observations), a quantity not provided in our previous archive, is given in Figure 2b. These data are contained in map group 3 of File 3. Values are printed in these two figures only where there are at least 100 observations contributing.

4) *Files 4-7: Total Cloud Cover; format 22.*

Total cloud cover is given in percent (AMT in format 22). Ten-year means for daily average annual, seasonal and monthly total cloud cover are given on the 2c grid for land and on the 5c grid for ocean. Daytime means are given seasonally as well. Because some parts of the southern oceans are poorly sampled (see, for example, Figure 2a), mean seasonal values over the ocean are also given on the 10c grid. It is the 10c grid that is used for the ocean mean seasonal averages by synoptic hour, which are utilized in the diurnal cycle analysis given in File 16. Since land stations are at fixed locations, use of a smaller grid size does not pose the problem that it does for the ocean.

Since previous climatologies, such as our own (Warren et al., 1986; 1988), did not utilize the illuminance criterion, a discontinuity of about 2% would occur when comparing the present with the previous data. Therefore 10-year mean seasonal and yearly seasonal mean values were also computed using all observations (File 4 map groups 54-97 and File 6 map groups 58-101). These data also make it possible to analyze the effects of the illuminance criterion in the present data set. Grid sizes of 5c and 10c for land and ocean, respectively, were selected for ease of comparison with our previous data set (Hahn et al., 1988).

Seasonal means are given for daytime and nighttime values as well as for the daily averages to aid in analysis of possible differences in trends between day and night. Seasonal means for land are given at 5c as well as 2c to make merging with ocean values easier if desired.

The most basic units provided in this data set are the monthly means by synoptic hour. From these, all other averages given, and some not given, can be reconstructed. Since it requires 960 map groups for 10 years of 8 synoptic hours monthly, this unit is provided in a file separate from the rest of the group cluster. The IDN variable in the data record labels each GMT synoptic hour for each box as day or night.

The mean annual total cloud for a grid box was computed by averaging the mean seasonal values of the seasons with 100 or more observations. The number of seasons contributing to the annual average is entered in the NSN variable and may vary from 0 to 4. Note that if NSN is 1, as may be true for ocean boxes near the poles, the reported value may not be representative of the true annual value. NOBS is the sum of the number of observations in the contributing seasons. SD and IDN are set to -9. For seasonal and monthly averages, only NSN was uniformly set to -9.

5) Files 8-15: Weather Types; format 32.

Most of the comments made in the last section apply equally well here. A few differences are noted. Percent frequencies of occurrence of clear sky, precipitation and fog are given in these files. The SD variable in the data record is assigned the missing value code. Precipitation and fog are always computed from all observations and so do not have to be repeated in the "all" map groups. However, since fog is given on the 2c grid for land, it is repeated at 5c for ease of comparison with previous data sets. Daytime and nighttime seasonal means are not given here (but could be reconstructed from the monthly means by synoptic hour if desired). This is also true for seasonal means at 5c for land. Seasonal means for the ocean are given on the 10c grid.

6) File 16: Harmonics, Interannual Variations and Trends; formats 40-42, 51-52.

The unifying feature of the contents of this file is that these are quantities derived from data already given in the archive. Separate land and ocean values are included and two different format types are used to accommodate the harmonic analyses on the one hand and the interannual variations on the other.

The phase and amplitude of the annual harmonic of total cloud cover (or of the frequency of occurrence of a weather type) were computed from the mean monthly values if all 12 months had 100 or more observations. (NT in format 40 always gives the number of months with 100 observations but the other variables were only computed if NT=12.) The amplitude reported is the absolute amplitude so that, for example, if the mean value for a particular cloud amount is 25% and the maximum of the fitted cycle is 30% then the amplitude is reported as 5% (rather than 20% which would be the normalized amplitude). The phase is a numeric value that corresponds to a month such that 1.0 is the middle of January, 2.0 is the middle of February, etc. Phase values reported range from 0.5 to less than 12.5. The value 0.0 (rather than -9) was used for PHASE in cases in which the amplitude was exactly zero (thus distinguishing these from "missing value" cases). AVG is the average of the 12 months used in the analysis and may differ somewhat from the annual values given in the other files

because of the different averaging methods. VAF is the percent of the variance accounted for by the amplitude of the annual harmonic.

To exemplify some features of the annual harmonic, figures 3a-c give the geographic distribution of the phase of the annual harmonic for total cloud cover, precipitation frequency and fog frequency, respectively, from the ocean map groups 21, 23 and 24 of File 16. (A phase of "0", common for fog in Figure 3c, indicates that the amplitude is exactly zero.) It is interesting to note that maximum total cloud cover occurs during the summer months in the North Pacific (Figure 3a), while precipitation frequency reaches maximum values during the winter months (Figure 3b). Figure 3c shows that fog is at a maximum in this region during the summer. The amplitude of the annual cycle for fog (available in the data record but not shown here) reaches 10-20% in the western North Pacific, making a significant contribution to the computed total cloud cover in this region during the summer. By comparison, amplitudes for the fog annual cycle are generally near zero between 30N and 50S over the ocean.

The diurnal harmonic (formats 41-42) was computed from the mean seasonal by-synoptic-hour values if there were 8 or 4 evenly spaced hours with 100 or more observations. If these conditions were met then $NT=8$ or 4 , otherwise NT was set to 0 and the other variables were set to -9 . The amplitude reported is the absolute amplitude (not the normalized amplitude). The phase is the hour of day (mean solar time of the box center) and may range from 0 to less than 24 . Since zero is a valid phase here, PHASE was set to -9 for cases in which the amplitude was exactly zero. AVG is the average of the 8 or 4 hours used in the analysis and may differ somewhat from the seasonal values given in the other files because of the different averaging methods. VAF is the percent of the variance accounted for by the amplitude of the diurnal harmonic.

As might be expected, the utilization of the illuminance criterion had a profound effect on the outcome of the diurnal cycle analysis. The geographic distribution of the amplitude and phase of the diurnal harmonic for total cloud cover over the oceans is shown on a $10c$ grid between $60N$ and $60S$ for four seasons in Figures 4a-d. (The phase is printed in these figures only where the amplitude is not zero.) The effect of the illuminance criterion is apparent when comparing these figures to Maps 114-115 of Warren et al. (1988) which show near-noon phases dominating in the diurnal analysis presented for 1954-1983 data without application of the illuminance criterion. For example, for JJA, in Figure 4c, a band of midnight to 5 AM phases evident in the eastern Pacific replaces 4 AM to early afternoon phases in that same region shown in Map 115b of Warren et al. Early afternoon phases are still evident in the western North Pacific and the western North Atlantic, however.

Diurnal cycle analysis for land data is given on the $2c$ grid. Since it is difficult to display the entire $2c$ grid, examples are given for two selected mid-latitude regions in Figure 5. Figure 5a shows the diurnal cycle of total cloud cover for part of Asia for DJF. Here diurnal sampling tends to be 8 times per day and there tend to be many reporting stations, particularly in China in the southeast portion of the region. While phase differences between these data and those of Warren et al. (1986, Maps 18-19) tend not to be as dramatic as those over the ocean, the morning phases on the northwest and southeast portions of Figure 5a contrast with those of Warren et al. Where the phases are comparable (and near noon), however, the amplitudes tend to be smaller when the illuminance criterion is applied.

Figure 5b shows the diurnal cycle analysis for JJA for the region of North America that covers most of the United States, where diurnal sampling tends to be 4 times per day. In the eastern portion of the region phases tend to have early afternoon values, in agreement with those of Warren et al.

(1986, Maps 22-23), while in the western portion the phases are more variable. This illustrates another feature of the present data set. The $2.5 \times 2.5^\circ$ resolution is capable of resolving features that are missed at $5 \times 5^\circ$, such as the band of late night to early morning phases evident in the central plains of the United States.

The last map groups in this file contain the results of the analysis of interannual variations and trends. For each season, an individual year contributed to the computation of IAV, TRND and UNC (formats 51-52) for a grid box if there were at least 100 observations. The number of years contributing is given in the variable NYRS. SPAN gives the number of years between the first and last years contributing, including the first and last years. If NYRS (also SPAN) is zero, IAV, TRND and UNC are assigned the value -9. IAV is the standard deviation of the contributing yearly values about the multi-year mean. TRND is the slope of the straight line which was fit to the data points by least-squares analysis and is given in units of percent cloud amount (or percent frequency) per year. UNC is the uncertainty of the slope (Bevington, 1969, page 113) in the same units.

4. HOW TO OBTAIN THE DATA

This documentation and the data described herein are available from:

Carbon Dioxide Information Analysis Center
Oak Ridge National Laboratory
Post Office Box 2008
Oak Ridge, TN 37831-6335, U.S.A.
Telephone (615) 574-3645

or

Data Support Section
National Center for Atmospheric Research
Boulder, CO 80307, U.S.A.
Telephone (303) 497-1215.

The following citation should be used for referencing this archive and/or this documentation report:

Hahn, C.J., S.G. Warren, and J. London, 1994: *Climatological Data for Clouds Over the Globe from Surface Observations, 1982-1991: The Total Cloud Edition*. NDP026A, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN. (Also available from Data Support Section, National Center for Atmospheric Research, Boulder, CO.)

Note that the archive of our earlier climatology (Hahn et al., 1988), along with accompanying atlases (Warren et al., 1986, 1988), is available from the same sources listed above.

ACKNOWLEDGEMENTS

This work was supported by NASA Grant NAG-1-998 and by Battelle Pacific Northwest Laboratories, Atmospheric Radiation Measurement Program (Contract 144806-A-Q1). We would also like to acknowledge the long term cooperation and the provision of computing facilities made

available to us from the Scientific Computing Division of the National Center for Atmospheric Research.

REFERENCES

- Bevington, P.R. 1969: *Data Reduction and Error Analysis for the Physical Sciences*, McGraw-Hill, 336 pp.
- Hahn, C.J., S.G. Warren, J. London, and R.L. Jenne, 1988: *Climatological Data for Clouds Over the Globe from Surface Observations*. NDP-026, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN. (Also available from Data Support Section, National Center for Atmospheric Research, Boulder, CO.)
- Hahn, C.J., S.G. Warren, and J. London, 1992: The use of COADS ship observations in cloud climatologies. *Proceedings of the International COADS Workshop*, H.F. Diaz, K. Wolter, and S.D. Woodruff, Eds., NOAA/ERL, Boulder, CO, 271-280.
- Hahn, C.J., S.G. Warren and J. London, 1994: The effect of moonlight on observation of cloud cover at night, and application to cloud climatology. *Journal of Climate*, Accepted.
- Riehl, H., 1947: Diurnal variation of cloudiness over the subtropical Atlantic Ocean. *Bulletin of the American Meteorological Society*, **28**, 37-40.
- Schneider, G., P. Paluzzi and J.P. Oliver, 1989: Systematic error in the synoptic sky cover record of the South Pole. *Journal of Climate*, **2**, 295-302.
- Warren, S.G., C.J. Hahn, J. London, R.M. Chervin and R.L. Jenne, 1986: *Global distribution of total cloud cover and cloud type amounts over land*. NCAR Technical Note TN-273+STR, Boulder, CO, 29 pp. + 200 maps (also DOE/ER/60085-H1).
- Warren, S.G., C.J. Hahn, J. London, R.M. Chervin and R.L. Jenne, 1988: *Global distribution of total cloud cover and cloud type amounts over the ocean*. NCAR Technical Note TN-317+STR, Boulder, CO, 42 pp. + 170 maps (also DOE/ER-0406).
- Woodruff, S.D., R.J. Slutz, R.L. Jenne and P.M. Steurer, 1987: A comprehensive ocean-atmosphere data set. *Bulletin of the American Meteorological Society*, **68**, 1239-1250.
- World Meteorological Organization, 1988: *Manual on Codes, Volume 1*. (WMO Publ. No. 306), WMO, Geneva.

Table 1. Cloud Information Contained in Synoptic Weather Reports

Symbol	Meaning	Codes*
N	total cloud cover	0-8 eighths 9= sky obscured
N _h	lower cloud cover	0-8 eights
h	lower cloud base height	0-9
C _L	low cloud type	0-9
C _M	middle cloud type	0-9
C _H	high cloud type	0-9
ww	present weather	00-99
I _x	present weather indicator	1-6

* Any category for which information is lacking to the observer is coded as "/".

Table 2. Cloud and Weather Type Definitions Used in Total Cloud Edition

Shorthand notation	Meaning	Synoptic codes
Tcc	Total cloud cover	N= 0-9
Clr	Clear sky	N=0
Ppt (R) (T)	Precipitation rain or snow thunderstorm	ww= 50-75,77,79,80-99 50-75,77,79 80-99
Fog (F)	Sky obscured due to fog	N=9 with ww= 10-12 or 40-49

Table 3. Grid Box Sizes

Box size (shorthand)	Dimensions lat x lon degrees	Latitude range	Number of boxes in	
			zone	globe
2.5x2.5c (2c)	2.5x2.5	50N to 50S	144	7290
	2.5x5	50 to 70	72	
	2.5x10	70 to 80	36	
	2.5x20	80 to 85	18	
	2.5x40	85 to 87.5	8	
	2.5x360	87.5 to 90	1	
5x5c (5c)	5x5	50N to 50S	72	1820
	5x10	50 to 70	36	
	5x20	70 to 80	18	
	5x40	80 to 85	9	
	5x360	85 to 90	1	
10x20c (10c)	10x20	50N to 50S	18	230
	10x40	50 to 70	9	
	10x60	70 to 80	6	
	10x360	80 to 90	1	

Table 4. File Information (Total Cloud Edition)

File	Group cluster name	General contents	Logical records	Characters per record	Characters
1	RDME	README (brief documentation)	452	80	36,160
2	LLFR	Latitude, Longitude, Land-fraction & Number of land stations for grid boxes	9,343	24	224,232
3	TWLO	Land & Ocean Combined total cloud & weather types	123,828	"	2,971,872
4	TCCL	Land Total Cloud Cover	472,038	"	11,328,912
5	TCCL	Land Total Cloud Cover by synoptic hour, monthly	2,217,600	"	53,222,400
6	TCCO	Ocean Total Cloud Cover	229,134	"	5,499,216
7	TCCO	Ocean Total Cloud Cover by synoptic hour, monthly	1,434,240	"	34,421,760
8	WXTL	Land Weather Types	720,346	"	17,288,304
9	WXTL	Land Clear-sky Frequency by synoptic hour, monthly	2,217,600	"	53,222,400
10	WXTL	Land Precipitation Frequency by synoptic hour, monthly	2,217,600	"	53,222,400
11	WXTL	Land Fog(sky-obscured) Freq. by synoptic hour, monthly	2,217,600	"	53,222,400
12	WXTO	Ocean Weather Types	156,954	"	3,766,896
13	WXTO	Ocean Clear-sky Frequency by synoptic hour, monthly	1,434,240	"	34,421,760
14	WXTO	Ocean Precipitation Freq. by synoptic hour, monthly	1,434,240	"	34,421,760
15	WXTO	Ocean Fog(sky-obscured) Freq. by synoptic hour, monthly	1,434,240	"	34,421,760
16	HIAV	Harmonics & Interannual Variation, land and ocean, total cloud & weather types	96,528	"	2,316,672

Table 5. Data Organization
 Contents of Surface-based Cloud Climatology Archive, 1982-1991* Total Cloud Edition**

File	Number of map groups	Map group numbers	Contents (coded in group header, Table 7)	grid*	groups**	Data format (Table 6)
1			README			text
2-16	9003					
2	3	1-3	GRID LAT, LON, LAND FRACTION, NUM. LAND STATIONS			
	1	1	1820 5x5c boxes			10
	1	2	230 10x20c boxes			10
	1	3	7290 2.5x2.5c boxes			10
3	68	1-68	LAND+OCEAN			
	1	1	Mean Annual TC	5c	1ann	22
	3	2-4	Mean Annual WT	"	1ann, 3types	32
	4	5-8	Mean Seasonal TC	"	4sns	22
	12	9-20	Mean Seasonal WT	"	4sns,3types	32
	12	21-32	Mean Monthly TC	"	12mns	22
	36	33-68	Mean Monthly WT	"	12mns,3types	32
4	1217	1-1217	LAND TOTAL CLOUD			
	1	1	Mean Annual	2c	1ann	22
	4	2-5	Mean Seasonal	"	4sns	22
	4	6-9	daytime	"	4sns	22
	32	10-41	by synoptic hour	"	8hrs,4sns	22
	12	42-53	Mean Monthly	"	12mns	22
	4	54-57	Mean Seasonal (all)	5c	4sns	22
	40	58-97	Seasonal Means (all)	"	4sns,10yrs	22
	40	98-137	Seasonal Means	"	4sns,10yrs	22
	40	138-177	Seasonal Means	2c	4sns,10yrs	22
	40	178-217	daytime	"	4sns,10yrs	22
	40	218-257	nighttime	"	4sns,10yrs	22
			Monthly means			
5	960	258-1217	by synoptic hour	2c	8hrs,12mns,10yrs	22
6	1181	1-1181	OCEAN TOTAL CLOUD			
	1	1	Mean Annual	5c	1ann	22
	4	2-5	Mean Seasonal	"	4sns	22
	4	6-9	daytime	"	4sns	22
	4	10-13	Mean Seasonal	10c	4sns	22
	32	14-45	by synoptic hour	"	8hrs,4sns	22
	12	46-57	Mean monthly	5c	12mns	22
	4	58-61	Mean Seasonal (all)	10c	4sns	22
	40	62-101	Seasonal Means (all)	"	4sns,10yrs	22
	40	102-141	Seasonal Means	5c	4sns,10yrs	22
	40	142-181	daytime	"	4sns,10yrs	22
	40	182-221	nighttime	"	4sns,10yrs	22
			Monthly Means			
7	960	222-1181	by synoptic hour	5c	8hrs,12mns,10yrs	22

Table 5 continued. Data Organization

File	Number of map groups	Map group numbers	Contents (coded in group header, Table 7)			Data format (Table 6)
8	3247	1-3247	LAND WEATHER TYPES			
	3	1-3	Mean Annual	2c	lann,3types	32
	12	4-15	Mean Seasonal	"	4sns,3types	32
	12	16-27	daytime	"	4sns,3types	32
	96	28-123	by synoptic hour	"	8hrs,4sns,3types	32
	36	124-159	Mean Monthly	"	12mns,3 types	32
	8	160-167	Mean Seasonal (all)	5c	4sns,2types	32
	80	168-247	Seasonal Means (all)	"	4sns,10yrs,2types	32
	120	248-367	Seasonal means	2c	4sns,10yrs,3types	32
			Monthly Means			
9-11	2880	368-3247	by synoptic hour	"	8hr,12mn,10yr,3types	32
12	3215	1-3215	OCEAN WEATHER TYPES			
	3	1-3	Mean Annual	5c	lam, 3types	32
	12	4-15	Mean Seasonal	"	4sns,3types	32
	12	16-27	daytime	"	4sns,3types	32
	12	28-39	Mean Seasonal	10c	4sns,3types	32
	96	40-135	by synoptic hour	"	8hrs,4sns,3types	32
	36	136-171	Mean Monthly	5c	12ms,3types	32
	4	172-175	Mean Seasonal (all)	10c	4sns 1type	32
	40	176-215	Seasonal Means (all)	"	4sns:10yrs,1type	32
	120	216-335	Seasonal Means	"	4sns,10yrs,3types	32
			Monthly Means			
13-15	2880	336-3215	by synoptic hour	5c	8hr,12mn,10yr,3types	32
16	72	1-72	HARMONICS & IAV			
	1	1	Annual TC, land	2c	lann	40
	3	2-4	Annual WT, land	"	lann,3types	40
	4	5-8	Diurnal TC, land	"	4sns	41
	12	9-20	Diurnal WT, land	"	4sns,3types	42
	1	21	Annual TC, ocean	5c	lann	40
	3	22-24	Annual WT, ocean	5c	lann,3types	40
	4	25-28	Diurnal TC, ocean	10c	4sns	41
	12	29-40	Diurnal WT, ocean	"	4sns,3types	42
	4	41-44	IAV TC, land	2c	4sns	51
	12	45-56	IAV WT, land	"	4sns,3types	52
	4	57-60	IAV TC, ocean	10c	4sns	51
	12	61-72	IAV WT, ocean	"	4sns,3types	52

* Non-standard terms are defined in Table 8. Briefly:

TC= total cloud; WT= "weather types" (clear sky, precipitation, fog).

** Illuminance criterion applied to total cloud and clear sky unless "all" specified. Ppt and fog always determined from all observations.

* Grid sizes are described in Table 3.

Number of boxes archived for 2c land = 2309, for 5c land = 861, for 5c ocean = 1493,
for 5c land+ocean = 1820, and for 10c ocean = 230

** Months are given in the order: Dec, Jan, Feb,...Nov.

If "2types" are specified, they are clear sky and fog.

If "1type" is specified, it is clear sky.

Convention for the order of groups in a multigroup listing is:

increment left group qualifier while holding right qualifier constant.

Table 6. List of Formats for Reading Data Records*
(Total Cloud Edition)

<u>Data class</u>							
<u>Format number</u>	<u>Variables and Format</u>						
1__	Lat, Lon, Land-fraction, Number of land stations						
10	I4 BOX	F5.2 CLAT	F5.2 CLON	F5.4 FRL	I5 NLSTA		
2__	Total Cloud						
22	I4 BOX	I6 NOBS	F5.2 AMT	F4.1 SD	I2 IDN	I1 LOB	I2 NSN
3__	Weather Types						
32	I4 BOX	I6 NOBS	F5.2 FQ	F4.1 SD	I2 IDN	I1 LOB	I2 NSN
4__	Harmonic analyses						
40	I4 BOX	F5.2 PHASE	F5.2 AMP	F4.1 VAF	I2 NT	F4.1 AVG	
41	(for annual harmonic)						
42	(for diurnal AMT)						
	(for diurnal FQ)						
5__	Interannual variations and trends						
51	I4 BOX	I2 NYRS	I2 SPAN	F5.2 IAV	F6.3 TRND	F5.3 UNC	
52	(for AMT)						
	(for FQ)						

* Terms defined in Table 8.
Data records are 24 characters.

Table 7. Map Group Header Record Format and Codes*
(Total Cloud Edition)

Format	I4	I4	I2	I1	I1	I4	I2	I2	I2	I2
Parameter	MGRP	NBXS	SIZE	LO	IMOON	YEAR	SN	TIME	TYPE	FMT
Values	1	230	10	1=Land	0=Moon	1981	0=ann	-3night	1=TCC	10
		1820	5	2=Ocean	1=All			-2day	2=CLR	22
	3247	7290	2	3=Global		1991	1=Jan	-1daily	3=PPT	32
		861				8190		00GMT	11=FOG	40
		1493				8291	12=Dec	03		41
		2309						06		42
							41=DJF	09		51
							42=MAM	12		52
							43=JJA	15		
							44=SON	18		
								21		

Terms are defined in Tables 3 & 8.

Table 8. Terms and Abbreviations Used

<i>Term</i>	<i>Meaning and description</i>
AMP	Absolute amplitude of harmonic (not normalized).
AMT	Average amount of cloud cover, given in percent.
ann	Annual.
AVG	Average AMT or FQ. In formats 40-42 it is the average of NT values.
BOX	Box number specific to grid size. See Table 3.
Cb	Cumulonimbus cloud.
CLAT	Center latitude of grid box. Values +90 to -90 for N to S.
CLON	Center longitude of grid box. Values 0 to 360 East.
day (time)	Local time 06-18.
DJF	December (of the previous year), January, February.
FMT	Data format number (see Table 6).
FQ	Frequency of occurrence; given in percent.
FRL	Fraction (0. to 1.0000) of area in grid box that is land.
GMT	Greenwich Mean Time.
IAV	Interannual variation (standard deviation of contributing year averages).
IDN	Indicates whether reports contributing to box average were from day only (=1), night only (=2), both (=3) or had less than minimum observations (=4) {see Section 2B3}.
IMOON	Indicator for application of the illuminance criterion: 0= criterion applied ("Moon"), 1= not applied ("All").
JJA	June, July, August.
hrs	Hours.
lat	Latitude.
LO	Indicator that group data are intended to be for land (=1), ocean (=2), or both (=3).
LOB	Indicator that box data were from land only (=1), ocean only (=2), both (=3), or no data (=0).
lon	Longitude.

Table 8 continued. Terms and Abbreviations Used

<i>Term</i>	<i>Meaning and description</i>
MAM	March, April, May.
MGRP	Map group number. Increments serially through group cluster (see Tables 4 & 5).
missing value code	The integer -9. Inerted in data record where no legitimate value is reported. (In formats 51-52 use NYRS=0 for missing value code.)
night (time)	Local time 18-06.
NLSTA	Approximate average number of land stations reporting in a 2c grid box.
NOBS	Number of observations.
Ns	Nimbostratus cloud.
NSN	Number of seasons contributing to annual average.
NT	Number of synoptic hours used (4 or 8) for diurnal harmonic analysis or number of months used (12) for annual harmonic analysis.
NYRS	Number of years contributing to trends and IAV.
mean seasonal	Long-term average; average over several years for season.
Mns	Months.
PHASE	Phase of first harmonic. Diurnal: 0-24 hours mean solar time of box center (-9 if AMP=0). Annual: month (0.5 -12.4 [1.0 = middle of January, etc.]; 0 if AMP=0).
SD	Standard deviation; in units of variable (not normalized).
seasonal mean	Average for an individual year for a particular season.
SIZE	Grid box size indicator.
SN	Season or month indicator.
sns	Seasons (DJF, MAM, JJA, SON).
SON	September, October, November.
SPAN	Span of years contributing to trend and IAV (includes first and last years contributing).
TIME	Time of day for which group data apply.
TRND	Trend. Slope of least-squares fit (change in average/yr).

Table 8 continued. Terms and Abbreviations Used

<i>Term</i>	<i>Meaning and description</i>
TYPE	Cloud or weather type code.
UNC	Uncertainty of trend line; same units as TRND.
VAF	Percent variance accounted for by the first harmonic.
YEAR	Year or years for which group data apply. Coded as 19yr for single years where yr gives the last 2 digits of the year, and as yf1 for multi-year averages where yf=yr of the first year and 1=yr of the last year of the period of record.
yrs	Years.




Figure 1. Flow chart of data selection and checking. Abbreviations are defined in Tables 1, 2 & 8. (Discard fractions given in order "land, ship" where needed). See text for discussion.




Figure 2a. ANNUAL AVERAGE TOTAL CLOUD COVER (%), LAND & OCEAN, 1982-1991

from Map Group: 31820 5318291 0-1 332




Figure 2b. ANNUAL PRECIPITATION FREQUENCY (%), LAND & OCEAN, 1982-1991




Figure 3a. ANNUAL CYCLE in Total Cloud Cover for (1982-1991) Ocean




Figure 3b. ANNUAL CYCLE in Precipitation Frequency for (1982-1991) Ocean




Figure 3c. ANNUAL CYCLE in Fog (sky obscured) Frequency for (1982-1991) Ocean





Figure 4a. DIURNAL CYCLE in Total Cloud Cover for DJF (1982-1991) Ocean





Figure 4b. DIURNAL CYCLE in Total Cloud Cover for MAM (1982-1991) Ocean





Figure 4c. DIURNAL CYCLE in Total Cloud Cover for JJA (1982-1991) Ocean





Figure 4d. DIURNAL CYCLE in Total Cloud Cover for SON (1982-1991) Ocean





Figure 5a. DIURNAL CYCLE in Total Cloud Cover for DJF (1982-1991) Land over Asia





Figure 5b. DIURNAL CYCLE in Total Cloud Cover for JJA (1982-1991) Land over North America

INTERNAL DISTRIBUTION

1. L. D. Bates
2. B. A. Berven
3. T. A. Boden
4. J. B. Cannon
5. J. H. Cushman
6. R. M. Cushman
7. V. H. Dale
8. M. P. Farrell
9. D. E. Fowler
10. R. L. Graham
11. S. G. Hildebrand
12. S. B. Jones
13. P. Kanciruk
14. J. M. Loar
15. L. J. Morris
16. D. E. Reichle
17. M. J. Sale
18. F. E. Sharples
19. D. E. Shepherd
20. P. Singley
21. S. H. Stow
- 22-221. CDIAC
222. Central Research Library
- 223-224. ESD Library
- 225-226. Laboratory Records Department
227. Laboratory Records, RC
228. ORNL Patent Section
229. ORNL Y-12 Technical Library

EXTERNAL DISTRIBUTION

230. S. S. Alexander, Pennsylvania State University, Department of Geosciences, 503 Deike Building, University Park, PA 16802
231. J. H. Allen, National Oceanic and Atmospheric Administration, National Geophysical Data Center Code E/GC2, 325 Broadway, Boulder, CO 80303
232. R. C. Barry, University of Colorado, World Data Center A, Glaciology, CIRES, Campus Box 449, Boulder, CO 80309-0449
233. M. A. Chinnery, National Oceanic and Atmospheric Administration, National Geophysical Data Center Code E/GC2, 325 Broadway, Boulder, CO 80303

234. Roger C. Dahlman, Global Change Research Program, Environmental Sciences Division, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, Washington, DC 20585
235. W. Draeger, EROS Data Center, U.S. Geological Survey, Sioux Falls, SD 57198
236. M. Dryer, National Oceanic and Atmospheric Administration, Space Environmental Lab., ERL/OAR, R/E/SE, 320 Broadway, Boulder, CO 80303
237. R. N. Farvolden, Professor, Department of Earth Sciences, University of Waterloo, Waterloo, Ontario N2L 3G1 Canada
238. J. Filson, National Earthquake Information Center, U.S. Geological Survey, Denver Federal Center, P.O. Box 20546, Denver, CO 80225
239. Jerry F. Franklin, Bloedel Professor of Ecosystem Analysis, College of Forest Resources, University of Washington, Anderson Hall (AR-10), Seattle, WA 98195
240. Dr. Diana W. Freckman, Director, College of Natural Resources, 101 Natural Resources Building, Colorado State University, Fort Collins, CO 80523
241. S. Graves, National Aeronautics and Space Administration Headquarters Code SED, 600 Independence Avenue, Washington, DC 20546
242. J. L. Green, National Space Science Data Center, NASA Goddard Space Flight Center, Code 630.2, Greenbelt, MD 20771
243. K. D. Hadeen, National Oceanic and Atmospheric Administration, NESDIS/NCDC, Federal Building MC E/CC, Asheville, NC 28801
- 244-248. Carole J. Hahn, Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, Colorado, 80309
249. R. C. Harriss, Institute for the Study of Earth, Oceans, and Space, Science and Engineering Research Building, University of New Hampshire, Durham, NH 03824
250. Ann Henderson-Sellers, MacQuarie University, North Rude, NSW, Sydney 2109, Australia
251. W. J. Hinze, Purdue University, Department of Earth and Atmospheric Sciences, West Lafayette, IN 47907
252. R. L. Jenne, NCAR, Scientific Computing Division, PO Box 3000, 1850 Table Mesa Drive, Boulder, Colorado 80307-3000
253. G. Y. Jordy, Director, Office of Program Analysis, Office of Energy Research, ER-30, G-226, U.S. Department of Energy, Washington, DC 20585

254. Dennis Joseph, NCAR, Scientific Computing Division, PO Box 3000, 1850 Table Mesa Drive, Boulder, Colorado 80307-3000
255. Dr. Thomas R. Karl, National Climatic Data Center, Federal Building, 37 Battery Park Avenue, Asheville, NC 28801
256. D. Lauer, EROS Data Center, U.S. Geological Survey, Sioux Falls, SD 57198
257. S. Levitus, NOAA/National Oceanographic Data Center, 1825 Connecticut Avenue, NW, Washington, DC 20235
- 258-262. Julius London, Department of Astrophysical, Planetary and Atmospheric Sciences, University of Colorado, Boulder, Colorado 80309
263. M. S. Loughridge, National Oceanic and Atmospheric Administration, National Geophysical Data Center, Code E/GC3, 325 Broadway, Boulder, CO 80303
264. Peter Lunn, US Department of Energy, Environmental Sciences Division, Office of Energy Research, ER-74, Washington, DC 20585
265. H. M. McCammon, Acting Deputy Director, Environmental Sciences Division, Office of Health and Environmental Research, Office of Energy Research, ER-74, U.S. Department of Energy, Washington, DC 20585
266. R. H. Olsen, Vice President for Research, University of Michigan, Medical Science Building II, #5605, 1301 East Catherine Street, Ann Arbor, MI 48109-0620
267. J. T. Overpeck, National Oceanic and Atmospheric Administration, National Geophysical Data Center, Paleoclimatology Program, 325 Broadway E/EC, Boulder, CO 80303
268. Farn Parnungo, ARL/NOAA/ERL, Boulder CO 80303
269. Bobbi Parra, US Department of Energy, Environmental Sciences Division, Office of Health and Environmental Research, Washington, DC 20585
270. Ari Patrinos, Acting Director Office of Health and Environmental Research, ER-74, U.S. Department of Energy, Washington, DC 20585
271. S. Ichtiaque Rasool, IGBP Data and Information System Office, Universite Paris, Tour 26, 4 Etage, Aile 26-16, 4 Place Jussieu, 75230 Paris, Cedex 06, France
272. Michael R. Riches, Acting Director, Environmental Sciences Division, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, Washington, DC 20585
273. S. Ruttenberg, Univ. Corporation for Atmospheric Research, CSNET, P. O. Box 3000, Boulder, CO 80307-3000

274. G. S. Saylor, Professor, 10515 Research Drive, Suite 100, The University of Tennessee, Knoxville, TN 37932-2567
275. A. L. Shumbera, National Oceanic and Atmospheric Administration, WDC-A for Meteorology, National Climatic Data Center, Federal Building MC E/CC, Asheville, NC 28801
276. Mr. Peter M. Steurer, National Climatic Data Center, Federal Building, 37 Battery Park Avenue, Asheville, NC 28801
277. Jerry Stokes, Battelle Pacific Northwest Lab, Battelle, Blvd, MS K1-74, PO Box 999, Richland, Washington 99352
- 278-282. Stephen G. Warren, Department of Atmospheric Sciences, University of Washington, Seattle, Washington 98195
283. F. J. Wobber, Environmental Sciences Division, Office of Health and Environmental Research, Office of Energy Research, ER-74, U.S. Department of Energy, Washington, DC 20585
284. Office of Assistant Manager for Energy Research and Development, U.S. Department of Energy Oak Ridge Operations, P. O. Box 2001, Oak Ridge, TN 37831-8600
- 285-286. Office of Scientific and Technical Information, P. O. Box 62, Oak Ridge, TN 37831