

# Appendix D: Water Balance Model Procedures

Temperature and precipitation data were either the calculated historical values for the study area or the forecast-modified values under Special Report on Emissions Scenarios (SRES) scenario A1B. Reference evapotranspiration ( $ET_o$ ), was calculated by using the Turc (1961) model (Jensen et al., 1997; Fontenot, 2004). Turc was selected for use over the original Thornthwaite model (as described in Dingman 2002) because of its ability to more closely simulate FAO-56 Penman-Monteith  $ET_o$  with a limited set of meteorological data (Fontenot, 2004). Allen (2003) defined the Turc equation for operational use:

$$ET_o = a_T 0.013 \frac{T_{mean}}{T_{mean} + 15} \frac{23.8856R_s + 50}{\lambda} \quad (1)$$

where  $ET_o$  is evapotranspiration ( $\text{mm day}^{-1}$ ),  $T_{mean}$  is the mean daily air temperature ( $^{\circ}\text{C}$ ),  $R_s$  is solar radiation ( $\text{MJ m}^{-2} \text{day}^{-1}$ ), and  $\lambda$  is the latent heat of vaporization ( $\text{MJ kg}^{-1}$ ). The coefficient  $a_T$  is a humidity-based value. If the mean daily relative humidity ( $RH_{mean}$ ) is greater than or equal to 50 percent, then  $a_T = 1.0$ . If the mean daily relative humidity is less than 50 percent, then  $a_T$  has the value of:

$$a_T = 1 + \frac{50 - RH_{mean}}{70} \quad (2)$$

Humidity data (historical or forecasted) were not available for the study area, so the assumption was made that the dew point temperature was equal to the mean monthly minimum temperature. This procedure is recommended by Allen et al. (1998) for approximating daily humidity values when measured values are not available. Solar radiation ( $R_s$ ) was estimated by using the Hargreaves model as described by Allen et al. (1998):

$$R_s = k_{RS} \sqrt{(T_{MAX} - T_{MIN})} R_a \quad (3)$$

where  $R_s$  is the solar radiation as stated above,  $k_{RS}$  is an adjustment coefficient,  $T_{MAX}$  and  $T_{MIN}$  are the mean daily maximum and minimum air temperatures ( $^{\circ}\text{C}$ ), and  $R_a$  is extraterrestrial radiation ( $\text{MJ m}^{-2} \text{day}^{-1}$ ). A value of 0.19 was used for  $k_{RS}$  as suggested by Allen et al. (1998) for use in coastal locations. The Turc model was run by using the monthly temperature data and radiation data for the 15<sup>th</sup> – the midpoint – of each month. The values were then multiplied by the appropriate number of days in each month to create a monthly value for  $ET_o$ . For simplicity, leap days were not included.

After the basic input variables were prepared, the data were entered into the water balance model. First, by using the temperature data, the monthly precipitation was partitioned into rain and snow components, where:

$$RAIN_M = F_M \cdot P_M \quad (4)$$

$$SNOW_M = (1 - F_M) \cdot P_M \quad (5)$$

Where  $P_M$  is the monthly precipitation and  $F_M$  is a melt factor that is computed by using the following method:

$$\begin{aligned} \text{If } T_M \leq 0^\circ \text{ C: } & F_M = 0 \\ \text{If } 0^\circ \text{ C} < T_M < 6^\circ \text{ C: } & F_M = 0.167 \cdot T_M \\ \text{If } T_M \geq 6^\circ \text{ C: } & F_M = 1 \end{aligned} \quad (6)$$

where  $T_M$  is the mean monthly temperature (Dingman, 2002).  $F_M$  also is used to determine the monthly snowmelt amount:

$$MELT_M = F_M \cdot (PACK_{m-1} + SNOW_m) \quad (7)$$

with  $PACK_{m-1}$  being the water equivalent of the snow pack at the end of the previous month and  $SNOW_m$  being the snow fall total of the current month. The previous month's pack amount is calculated as:

$$PACK_m = (1 - F_M)^2 \cdot P_M + (1 - F_M) \cdot PACK_{m-1} \quad (8)$$

The overall hydrological input into the model is defined by  $W_M$  as:

$$W_M = RAIN_m + MELT_m \quad (9)$$

In this study, the probability of the study region having any significant snow amounts is low, but the variable was included to provide for the possibility in the forecasted model runs.

Changes in soil moisture are calculated by using the following logic. If  $W_M \geq ET_o$ , monthly evapotranspiration ( $ET_M$ ) occurs at the  $ET_o$  rate. If  $ET_M$  equals  $ET_o$ , then soil moisture would increase or remain steady if the soil moisture already is at field capacity (Dingman, 2002). For the purposes of this study, field capacity ( $SOIL_{MAX}$ ) has been set to 150 mm (5.9 in). The monthly value for soil moisture is therefore:

$$SOIL_M = \min\{[(W_M - ET_o) + SOIL_{m-1}], SOIL_{MAX}\} \quad (10)$$

where the soil moisture value is the lesser of the two values in the equation (Dingman, 2002). If  $W_M$  is less than  $ET_o$ , then  $ET_M$  is equal to the hydrological input ( $W_M$ ) and a drying factor:

$$ET_M = W_M + \left\{ SOIL_{m-1} \cdot \left[ 1 - \exp\left(-\frac{ET_{OM} - W_M}{SOIL_{MAX}}\right) \right] \right\} \quad (11)$$

where  $ET_{OM}$  is the monthly Turc  $ET_O$  value (Dingman, 2002).

After computing soil moisture change, any excess water in the budget was declared as surplus. The monthly surplus parameter is synonymous with runoff in these wetland environments, as long lags are not common between the generation of surplus water and the resultant streamflow. If  $W_M$  does not meet the environmental demand, then a deficit is created until  $W_M$  meets the environmental demand. In this study, we retained surplus as an index for runoff and dismissed the modeled runoff term as invalid.

## ■ References

- Allen, R.G.**, 2003: *REF-ET User's Guide*. University of Idaho Kimberly Research Stations: Kimberly, Idaho.
- Allen, R.G.**, L.S. Pereira, D. Raes, and M. Smith, 1998: *Crop Evapotranspiration – Guidelines for Computing Crop Water Requirements*. FAO Irrigation and Drainage Paper 56. Food and Agriculture Organization. Rome.
- Dingman, S.L.**, 2002: *Physical Hydrology, 2<sup>nd</sup> Ed.*: Upper Saddle River, New Jersey. Prentice Hall.
- Fontenot, R.**, 2004: *An Evaluation of Reference Evapotranspiration Models in Louisiana*. M.N.S. Thesis, Department of Geography and Anthropology, Louisiana State University.
- Jensen, D.T.**, G.H. Hargreaves, B. Temesgen, and R.G. Allen. 1997: Computation of  $ET_O$  under Nonideal Conditions. *Journal of Irrigation and Drainage Engineering* 123(5):394-400
- Turc, L.**, 1961: Evaluation des besoins en eau d'irrigation, evapotranspiration potentielle, formule climatique simpliffee et mise a jour. (In French). *Annales Agronomiques* 12(1):13-49.

