# **Drug Facts**

## Active ingredient (in each rectal suppository)

**Purposes** 

Acetaminophen 650 mg......Pain reliever/fever reducer

Uses temporarily # reduces fever # relieves minor aches, pains, and headache

## Warnings

## For rectal use only

**Alcohol warning**: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.

Do not use in children under 12 years

### Stop use and ask a doctor if

# fever lasts more than 3 days (72 hours), or recurs

# you need to use this product for pain for more than 10 days continuously Severe or recurrent pain, or high or continued fever may indicate a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children**. If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

## Directions<sup>1</sup>

#### # do not use more than directed

- # remove foil wrapper
- # insert suppository well up into rectum
- # adults and children 12 years and older:
  - # 1 suppository every 4 to 6 hours while symptoms persist
  - # do not exceed 6 suppositories in any 24-hour period
- # children under 12 years: do not use

#### Other information

# optional - tamper evident statement

<sup>&</sup>lt;sup>1</sup> The information for the two age groups may be presented in a table format in accord with 21 CFR 201.66(d)(9) with the first three bulleted statements appearing above the top line of the table.

# store at 8-25°C (46-77°F)<sup>2</sup> or refrigerate # optional - see [end or side] panel for lot number and expiration date

Inactive ingredients [list ingredients in alphabetical order]

Questions or comments? call toll free 1-800-XXX-XXXX

NOTE: The *Drug Facts (continued)* title should appear wherever the labeling continues onto another panel of the package.

\_

 $<sup>^2</sup>$  These temperatures reflect the storage conditions for acetaminophen suppositories, USP. Manufacturers who have stability data in their ANDA to support storage up to  $27^{\circ}$  C ( $80^{\circ}$  F) may use these temperatures in place of  $25^{\circ}$  C ( $77^{\circ}$  F).