

Table A2. Reported cases of sexually transmitted disease by sex and reporting source: United States, 2003

<i>Disease</i>	<i>Non-STD Clinic</i>			<i>STD Clinic</i>			<i>Total*</i>		
	<i>Male</i>	<i>Female</i>	<i>Total</i>	<i>Male</i>	<i>Female</i>	<i>Total</i>	<i>Male</i>	<i>Female</i>	<i>Total</i>
Chlamydia	114,814	555,957	672,073	66,354	88,444	154,944	190,244	685,017	877,478
Gonorrhea	82,627	131,780	214,819	69,335	30,766	100,173	160,106	174,230	335,104
Primary Syphilis	1,081	122	1,205	751	99	851	1,863	229	2,095
Secondary Syphilis	2,898	619	3,517	1,135	345	1,480	4,093	988	5,082
Early Latent Syphilis	3,692	1,695	5,393	1,892	962	2,854	5,664	2,691	8,361
Late and Late Latent Syphilis [†]	7,016	5,071	12,124	3,357	2,369	5,739	10,666	7,601	18,319
Neurosyphilis	387	117	504	25	3	28	420	121	541
Chancroid	8	7	15	17	18	35	25	29	54

*Totals include unknown sex and reporting source.

[†]Late and late latent syphilis includes cases of unknown duration, late syphilis with clinical manifestations, and neurosyphilis.

See Appendix (Reporting Sources).