

Table 3-152. Estimates of lost-time injuries by farm type for family workers in the Northeast region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	536.1	5.0	536.1	5.0
Field crops	1757.3	16.3	2293.4	21.3
Veg, fruit, nuts	1207.1	11.2	3500.5	32.5
Nursery	703.8	6.5	4204.3	39.0
Beef, hogs, sheep	1758.1	16.3	5962.4	55.3
Poultry	150.6	1.4	6113.0	56.7
Dairy	4462.4	41.4	10575.4	98.1
Other farms	200.9	1.9	10776.3	100.0

Table 3-153. Estimates of lost-time injuries by gender for family workers in the Northeast region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	8753.8	81.2	8753.8	81.2
Female	2022.5	18.8	10776.3	100.0

Table 3-154. Estimates of lost-time injuries by age group for family workers in the Northeast region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	729.3	6.8	729.3	6.8
20-29	2194.8	20.4	2924.1	27.1
30-39	1284.4	11.9	4208.5	39.1
40-49	2512.2	23.3	6720.7	62.4
50-59	1895.7	17.6	8616.4	80.0
60-69	697.4	6.5	9313.8	86.4
>=70	1462.5	13.6	10776.3	100.0

Table 3-155. Estimates of lost-time injuries by race or ethnic origin for family workers in the Northeast region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	10040.7	93.2	10040.7	93.2
Hispanic	735.6	6.8	10776.3	100.0

Table 3-156. Estimates of lost-time injuries by month for family workers in the Northeast region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	109.2	1.0	109.2	1.0
February	667.6	6.2	776.8	7.2
March	361.5	3.4	1138.3	10.6
April	749.8	7.0	1888.1	17.5
May	1295.0	12.0	3183.1	29.5
June	590.0	5.5	3773.1	35.0
July	1571.5	14.6	5344.6	49.6
August	575.0	5.3	5919.6	54.9
September	1033.9	9.6	6953.5	64.5
October	1491.7	13.8	8445.2	78.4
November	1231.5	11.4	9676.7	89.8
December	1099.6	10.2	10776.3	100.0

Table 3-157. Estimates of lost-time injuries by severity of the injury for family workers in the Northeast region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	10572.4	98.1	10572.4	98.1
Permanent	203.9	1.9	10776.3	100.0

Table 3-158. Estimates of lost-time injuries by whether professional medical attention was sought for family workers in the Northeast region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	203.9	1.9	203.9	1.9
Yes	8352.7	77.5	8556.6	79.4
No	2219.7	20.6	10776.3	100.0

Table 3-159. Estimates of lost-time injuries by body part injured for family workers in the Northeast region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	460.4	4.3	460.4	4.3
Eye	982.5	9.1	1442.9	13.4
Chest/trunk	446.0	4.1	1888.9	17.5
Back	840.4	7.8	2729.3	25.3
Arm/shoulder	749.4	7.0	3478.7	32.3
Finger	2805.9	26.0	6284.6	58.3
Hand/wrist	2031.6	18.9	8316.2	77.2
Leg/knee/hip	1279.5	11.9	9595.7	89.0
Toe	17.7	0.2	9613.4	89.2
Foot	670.1	6.2	10283.5	95.4
Mult. parts	134.2	1.2	10417.7	96.7
Other	358.6	3.3	10776.3	100.0

Table 3-160. Estimates of lost-time injuries by nature of injury for family workers in the Northeast region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	203.9	1.9	203.9	1.9
Bruise	2008.9	18.6	2212.8	20.5
Cut	1843.7	17.1	4056.5	37.6
Crush	521.6	4.8	4578.1	42.5
Fracture	1386.1	12.9	5964.2	55.3
Puncture	389.7	3.6	6353.9	59.0
Sprain/strain	2749.3	25.5	9103.2	84.5
Mult. injury	69.3	0.6	9172.5	85.1
Other	1603.8	14.9	10776.3	100.0

Table 3-161. Estimates of lost-time injuries by type of event for family workers in the Northeast region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	236.3	2.2	236.3	2.2
Caught between	1627.6	15.1	1863.9	17.3
Caught under	239.9	2.2	2103.8	19.5
Struck by/against	976.7	9.1	3080.5	28.6
Struck by falling	551.0	5.1	3631.5	33.7
Struck by flying	818.5	7.6	4450.0	41.3
Contact sharp obj.	1491.6	13.8	5941.6	55.1
Fall, same level	926.1	8.6	6867.7	63.7
Fall, elevation	1040.2	9.7	7907.9	73.4
Overexertion	1196.7	11.1	9104.6	84.5
Other	1671.7	15.5	10776.3	100.0

Table 3-162. Estimates of lost-time injuries by activity when worker was injured for family workers in the Northeast region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	1912.7	17.7	1912.7	17.7
Machine maint.	1346.0	12.5	3258.7	30.2
Field work	1507.4	14.0	4766.1	44.2
Crop handling	2360.5	21.9	7126.6	66.1
Livestock handling	2519.8	23.4	9646.4	89.5
Other	1129.9	10.5	10776.3	100.0

Table 3-163. Estimates of lost-time injuries by source of the injury for family workers in the Northeast region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	316.9	2.9	316.9	2.9
Machinery	1498.1	13.9	1815.0	16.8
Livestock	1990.6	18.5	3805.6	35.3
Hand tool	427.9	4.0	4233.5	39.3
Power tool	1465.2	13.6	5698.7	52.9
Plant/tree	326.6	3.0	6025.3	55.9
Working surface	1195.7	11.1	7221.0	67.0
Truck/auto	471.1	4.4	7692.1	71.4
Other	3084.2	28.6	10776.3	100.0

Table 3-164. Estimates of lost-time injuries by farm type for hired workers in the Northeast region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	59.7	1.7	59.7	1.7
Field crops	119.8	3.4	179.5	5.1
Veg, fruit, nuts	291.5	8.3	471.0	13.4
Nursery	703.8	20.0	1174.8	33.4
Beef, hogs, sheep	64.9	1.8	1239.7	35.2
Dairy	1365.1	38.8	2604.8	74.0
Other farms	916.7	26.0	3521.5	100.0

Table 3-165. Estimates of lost-time injuries by gender for hired workers in the Northeast region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	2086.3	59.2	2086.3	59.2
Female	1435.2	40.8	3521.5	100.0

Table 3-166. Estimates of lost-time injuries by age group for hired workers in the Northeast region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	554.5	15.7	554.5	15.7
20-29	1509.8	42.9	2064.3	58.6
30-39	449.3	12.8	2513.6	71.4
40-49	69.3	2.0	2582.9	73.3
50-59	307.0	8.7	2889.9	82.1
60-69	631.6	17.9	3521.5	100.0

Table 3-167. Estimates of lost-time injuries by race or ethnic origin for hired workers in the Northeast region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	3246.9	92.2	3246.9	92.2
Hispanic	274.6	7.8	3521.5	100.0

Table 3-168. Estimates of lost-time injuries by month for hired workers in the Northeast region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	0.0	0.0	0.0	0.0
February	0.0	0.0	0.0	0.0
March	36.1	1.0	36.1	1.0
April	696.5	19.8	732.6	20.8
May	206.6	5.9	939.2	26.7
June	156.1	4.4	1095.3	31.1
July	1395.6	39.6	2490.9	70.7
August	459.9	13.1	2950.8	83.8
September	414.6	11.8	3365.4	95.6
October	156.1	4.4	3521.5	100.0
November	156.1	4.4	3521.5	100.0
December	156.1	4.4	3521.5	100.0

Table 3-169. Estimates of lost-time injuries by severity of the injury for hired workers in the Northeast region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	3301.5	93.8	3301.5	93.8
Permanent	220.0	6.2	3521.5	100.0

Table 3-170. Estimates of lost-time injuries by whether professional medical attention was sought for hired workers in the Northeast region of the U.S., 1993.

Attention Required	Medical		Cumulative Frequency	Cumulative Percent
	Frequency	Percent		
Unknown	220.0	6.2	220.0	6.2
Yes	3232.2	91.8	3452.2	98.0
No	69.3	2.0	3521.5	100.0

Table 3-171. Estimates of lost-time injuries by body part injured for hired workers in the Northeast region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	310.2	8.8	310.2	8.8
Eye	56.0	1.6	366.2	10.4
Chest/trunk	220.0	6.2	586.2	16.6
Arm/shoulder	138.6	3.9	724.8	20.6
Finger	171.4	4.9	896.2	25.4
Hand/wrist	1824.3	51.8	2720.5	77.3
Leg/knee/hip	187.2	5.3	2907.7	82.6
Foot	613.8	17.4	3521.5	100.0

Table 3-172. Estimates of lost-time injuries by nature of injury for hired workers in the Northeast region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	221.2	6.3	221.2	6.3
Cut	1061.9	30.2	1283.1	36.4
Crush	102.1	2.9	1385.2	39.3
Fracture	1063.7	30.2	2448.9	69.5
Puncture	86.8	2.5	2535.7	72.0
Sprain/strain	985.8	28.0	3521.5	100.0

Table 3-173. Estimates of lost-time injuries by type of event for hired workers in the Northeast region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	376.1	10.7	376.1	10.7
Caught between	322.1	9.1	698.2	19.8
Caught under	69.3	2.0	767.5	21.8
Struck by/against	263.9	7.5	1031.4	29.3
Struck by falling	64.9	1.8	1096.3	31.1
Struck by flying	19.9	0.6	1116.2	31.7
Contact sharp obj.	1023.4	29.1	2139.6	60.8
Fall, same level	854.9	24.3	2994.5	85.0
Overexertion	289.3	8.2	3283.8	93.3
Other	237.7	6.7	3521.5	100.0

Table 3-174. Estimates of lost-time injuries by activity when worker was injured for hired workers in the Northeast region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	134.2	3.8	134.2	3.8
Machine maint.	1025.8	29.1	1160.0	32.9
Field work	1150.5	32.7	2310.5	65.6
Crop handling	476.1	13.5	2786.6	79.1
Livestock handling	734.9	20.9	3521.5	100.0

Table 3-175. Estimates of lost-time injuries by source of the injury for hired workers in the Northeast region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Machinery	1312.7	37.3	1312.7	37.3
Livestock	665.6	18.9	1978.3	56.2
Plant/tree	36.1	1.0	2014.4	57.2
Working surface	238.8	6.8	2253.2	64.0
Other vehicle	69.3	2.0	2322.5	66.0
Other	1199.0	34.0	3521.5	100.0

Table 3-176. Estimates of lost-time injuries by farm type for family workers in the Southern region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	2178.4	7.9	2178.4	7.9
Field crops	3702.2	13.4	5880.6	21.3
Veg, fruit, nuts	1860.0	6.7	7740.6	28.0
Nursery	148.8	0.5	7889.4	28.5
Beef, hogs, sheep	17594.3	63.6	25483.7	92.1
Poultry	670.7	2.4	26154.4	94.6
Dairy	404.6	1.5	26559.0	96.0
Other farms	1097.0	4.0	27656.0	100.0

Table 3-177. Estimates of lost-time injuries by gender for family workers in the Southern region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	23594.5	85.3	23594.5	85.3
Female	4061.5	14.7	27656.0	100.0

Table 3-178. Estimates of lost-time injuries by age group for family workers in the Southern region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	318.8	1.2	318.8	1.2
20-29	915.0	3.3	1233.8	4.5
30-39	4496.8	16.3	5730.6	20.7
40-49	5497.3	19.9	11227.9	40.6
50-59	6101.8	22.1	17329.7	62.7
60-69	6143.5	22.2	23473.2	84.9
>=70	4182.8	15.1	27656.0	100.0

Table 3-179. Estimates of lost-time injuries by race or ethnic origin for family workers in the Southern region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	27270.3	98.6	27270.3	98.6
American Indian	385.7	1.4	27656.0	100.0

Table 3-180. Estimates of lost-time injuries by month for family workers in the Southern region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	2378.2	8.6	2378.2	8.6
February	2799.6	10.1	5177.8	18.7
March	1583.9	5.7	6761.7	24.4
April	1407.3	5.1	8169.0	29.5
May	1608.4	5.8	9777.4	35.4
June	3100.9	11.2	12878.3	46.6
July	1340.7	4.8	14219.0	51.4
August	2718.3	9.8	16937.3	61.2
September	2550.7	9.2	19488.0	70.5
October	1233.7	4.5	20721.7	74.9
November	3059.2	11.1	23780.9	86.0
December	3875.1	14.0	27656.0	100.0

Table 3-181. Estimates of lost-time injuries by severity of the injury for family workers in the Southern region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	24828.7	89.8	24828.7	89.8
Permanent	2827.3	10.2	27656.0	100.0

Table 3-182. Estimates of lost-time injuries by whether professional medical attention was sought for family workers in the Southern region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	2827.3	10.2	2827.3	10.2
Yes	19400.1	70.1	22227.4	80.4
No	5428.6	19.6	27656.0	100.0

Table 3-183. Estimates of lost-time injuries by body part injured for family workers in the Southern region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	1296.2	4.7	1296.2	4.7
Eye	277.5	1.0	1573.7	5.7
Chest/trunk	4024.4	14.6	5598.1	20.2
Back	2053.3	7.4	7651.4	27.7
Arm/shoulder	2203.5	8.0	9854.9	35.6
Finger	1876.8	6.8	11731.7	42.4
Hand/wrist	3073.2	11.1	14804.9	53.5
Leg/knee/hip	6924.8	25.0	21729.7	78.6
Toe	398.6	1.4	22128.3	80.0
Foot	3235.4	11.7	25363.7	91.7
Mult. parts	2167.5	7.8	27531.2	99.5
Other	124.8	0.5	27656.0	100.0

Table 3-184. Estimates of lost-time injuries by nature of injury for family workers in the Southern region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	398.6	1.4	398.6	1.4
Bruise	3401.1	12.3	3799.7	13.7
Burn	398.6	1.4	4198.3	15.2
Cut	4665.5	16.9	8863.8	32.1
Crush	2718.6	9.8	11582.4	41.9
Fracture	9058.7	32.8	20641.1	74.6
Sprain/strain	3101.9	11.2	23743.0	85.9
Mult. injury	2671.2	9.7	26414.2	95.5
Other	1241.8	4.5	27656.0	100.0

Table 3-185. Estimates of lost-time injuries by type of event for family workers in the Southern region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	956.2	3.5	956.2	3.5
Caught between	666.6	2.4	1622.8	5.9
Caught under	950.6	3.4	2573.4	9.3
Struck by/against	8832.9	31.9	11406.3	41.2
Struck by falling	2627.1	9.5	14033.4	50.7
Struck by flying	905.3	3.3	14938.7	54.0
Contact sharp obj.	2775.1	10.0	17713.8	64.1
Fall, same level	4508.3	16.3	22222.1	80.4
Fall, elevation	3592.1	13.0	25814.2	93.3
Overexertion	527.1	1.9	26341.3	95.2
Other	1314.7	4.8	27656.0	100.0

Table 3-186. Estimates of lost-time injuries by activity when worker was injured for family workers in the Southern region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	5240.3	18.9	5240.3	18.9
Machine maint.	2109.2	7.6	7349.5	26.6
Field work	3864.7	14.0	11214.2	40.5
Crop handling	5168.5	18.7	16382.7	59.2
Livestock handling	10968.0	39.7	27350.7	98.9
Other	305.3	1.1	27656.0	100.0

Table 3-187. Estimates of lost-time injuries by source of the injury for family workers in the Southern region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	2781.4	10.1	2781.4	10.1
Machinery	3098.3	11.2	5879.7	21.3
Livestock	8152.6	29.5	14032.3	50.7
Hand tool	1569.0	5.7	15601.3	56.4
Power tool	805.9	2.9	16407.2	59.3
Pesticide/chem.	736.1	2.7	17143.3	62.0
Plant/tree	1123.4	4.1	18266.7	66.0
Working surface	2941.2	10.6	21207.9	76.7
Truck/auto	398.6	1.4	21606.5	78.1
Other vehicle	703.9	2.5	22310.4	80.7
Other	5345.6	19.3	27656.0	100.0

Table 3-188. Estimates of lost-time injuries by farm type for hired workers in the Southern region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	906.8	5.6	906.8	5.6
Field crops	3486.5	21.6	4393.3	27.2
Veg, fruit, nuts	3046.4	18.8	7439.7	46.0
Nursery	2221.6	13.7	9661.3	59.8
Beef, hogs, sheep	1867.0	11.5	11528.3	71.3
Poultry	285.0	1.8	11813.3	73.1
Dairy	3658.7	22.6	15472.0	95.7
Other farms	692.7	4.3	16164.7	100.0

Table 3-189. Estimates of lost-time injuries by gender for hired workers in the Southern region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	15718.3	97.2	15718.3	97.2
Female	446.4	2.8	16164.7	100.0

Table 3-190. Estimates of lost-time injuries by age group for hired workers in the Southern region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	1075.4	6.7	1075.4	6.7
20-29	6535.6	40.8	7611.0	47.5
30-39	5911.1	36.9	13522.1	84.4
40-49	1531.7	9.6	15053.8	94.0
50-59	662.1	4.1	15715.9	98.1
>=70	300.0	1.9	16015.9	100.0

(Frequency Missing = 148.8)

Table 3-191. Estimates of lost-time injuries by race or ethnic origin for hired workers in the Southern region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	8983.0	56.1	8983.0	56.1
Hispanic	5375.7	33.6	14358.7	89.7
Black	1258.6	7.9	15617.3	97.5
American Indian	398.6	2.5	16015.9	100.0

(Frequency Missing = 148.8)

Table 3-192. Estimates of lost-time injuries by month for hired workers in the Southern region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	494.5	3.1	494.5	3.1
February	1176.1	7.3	1670.6	10.4
March	266.1	1.7	1936.7	12.1
April	1684.3	10.5	3621.0	22.6
May	1317.9	8.2	4938.9	30.8
June	2351.0	14.7	7289.9	45.5
July	1058.7	6.6	8348.6	52.1
August	1963.7	12.3	10312.3	64.4
September	1143.5	7.1	11455.8	71.5
October	3037.3	19.0	14493.1	90.5
November	701.0	4.4	15194.1	94.9
December	821.8	5.1	16015.9	100.0

(Frequency Missing = 148.8)

Table 3-193. Estimates of lost-time injuries by severity of the injury for hired workers in the Southern region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	14508.0	89.8	14508.0	89.8
Permanent	1284.1	7.9	15792.1	97.7
Fatality	372.6	2.3	16164.7	100.0

Table 3-194. Estimates of lost-time injuries by whether professional medical attention was sought for hired workers in the Southern region of the U.S., 1993.

Attention Required	Medical		Cumulative Frequency	Cumulative Percent
	Frequency	Percent		
Unknown	1656.7	10.2	1656.7	10.2
Yes	13918.7	86.1	15575.4	96.4
No	589.3	3.6	16164.7	100.0

Table 3-195. Estimates of lost-time injuries by body part injured for hired workers in the Southern region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	909.9	5.6	909.9	5.6
Eye	1344.3	8.3	2254.2	13.9
Chest/trunk	1397.9	8.6	3652.1	22.6
Back	3231.3	20.0	6883.4	42.6
Arm/shoulder	324.3	2.0	7207.7	44.6
Finger	1874.1	11.6	9081.8	56.2
Hand/wrist	1994.6	12.3	11076.4	68.5
Leg/knee/hip	968.0	6.0	12044.4	74.5
Toe	566.2	3.5	12610.6	78.0
Foot	3067.8	19.0	15678.4	97.0
Mult. parts	486.3	3.0	16164.7	100.0

Table 3-196. Estimates of lost-time injuries by nature of injury for hired workers in the Southern region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	2526.9	15.6	2526.9	15.6
Burn	112.5	0.7	2639.4	16.3
Cut	3988.7	24.7	6628.1	41.0
Crush	1647.2	10.2	8275.3	51.2
Fracture	1444.7	8.9	9720.0	60.1
Puncture	1078.0	6.7	10798.0	66.8
Sprain/strain	3381.1	20.9	14179.1	87.7
Mult. injury	148.8	0.9	14327.9	88.6
Other	1836.8	11.4	16164.7	100.0

Table 3-197. Estimates of lost-time injuries by type of event for hired workers in the Southern region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	318.8	2.0	318.8	2.0
Caught between	2816.4	17.4	3135.2	19.4
Caught under	337.5	2.1	3472.7	21.5
Struck by/against	3038.8	18.8	6511.5	40.3
Struck by falling	1960.1	12.1	8471.6	52.4
Contact sharp obj.	2117.0	13.1	10588.6	65.5
Fall, same level	1103.9	6.8	11692.5	72.3
Fall, elevation	285.0	1.8	11977.5	74.1
Overexertion	2975.3	18.4	14952.8	92.5
Other	1211.9	7.5	16164.7	100.0

Table 3-198. Estimates of lost-time injuries by activity when worker was injured for hired workers in the Southern region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	3205.5	19.8	3205.5	19.8
Machine maint.	2305.8	14.3	5511.3	34.1
Field work	4910.8	30.4	10422.1	64.5
Crop handling	1949.2	12.1	12371.3	76.5
Livestock handling	3282.3	20.3	15653.6	96.8
Other	511.1	3.2	16164.7	100.0

Table 3-199. Estimates of lost-time injuries by source of the injury for hired workers in the Southern region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	1762.1	10.9	1762.1	10.9
Machinery	4465.1	27.6	6227.2	38.5
Livestock	1903.1	11.8	8130.3	50.3
Hand tool	1620.7	10.0	9751.0	60.3
Pesticide/chem.	150.0	0.9	9901.0	61.3
Plant/tree	1341.4	8.3	11242.4	69.5
Working surface	1170.0	7.2	12412.4	76.8
Truck/auto	933.0	5.8	13345.4	82.6
Other vehicle	367.7	2.3	13713.1	84.8
Other	2451.6	15.2	16164.7	100.0

Table 3-200. Estimates of lost-time injuries by farm type for family workers in the Midwest region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	20662.6	29.5	20662.6	29.5
Field crops	2471.4	3.5	23134.0	33.1
Veg, fruit, nuts	902.7	1.3	24036.7	34.3
Beef, hogs, sheep	28155.0	40.2	52191.7	74.6
Poultry	80.0	0.1	52271.7	74.7
Dairy	17079.2	24.4	69350.9	99.1
Other farms	640.0	0.9	69990.9	100.0

Table 3-201. Estimates of lost-time injuries by gender for family workers in the Midwest region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	61730.1	88.2	61730.1	88.2
Female	8260.8	11.8	69990.9	100.0

Table 3-202. Estimates of lost-time injuries by age group for family workers in the Midwest region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	6500.2	9.3	6500.2	9.3
20-29	5203.0	7.4	11703.2	16.7
30-39	17769.0	25.4	29472.2	42.1
40-49	12298.4	17.6	41770.6	59.7
50-59	13913.5	19.9	55684.1	79.6
60-69	10614.6	15.2	66298.7	94.7
>=70	3692.2	5.3	69990.9	100.0

Table 3-203. Estimates of lost-time injuries by race or ethnic origin for family workers in the Midwest region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	69374.0	99.1	69374.0	99.1
Hispanic	616.9	0.9	69990.9	100.0

Table 3-204. Estimates of lost-time injuries by month for family workers in the Midwest region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	3915.9	5.6	3915.9	5.6
February	1603.9	2.3	5519.8	7.9
March	1628.4	2.3	7148.2	10.2
April	6713.8	9.6	13862.0	19.8
May	4489.9	6.4	18351.9	26.2
June	10526.2	15.0	28878.1	41.3
July	9854.1	14.1	38732.2	55.3
August	7666.0	11.0	46398.2	66.3
September	9943.9	14.2	56342.1	80.5
October	5667.2	8.1	62009.3	88.6
November	5212.3	7.4	67221.6	96.0
December	2769.3	4.0	69990.9	100.0

Table 3-205. Estimates of lost-time injuries by severity of the injury for family workers in the Midwest region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	67307.1	96.2	67307.1	96.2
Permanent	1805.5	2.6	69112.6	98.7
Fatality	878.3	1.3	69990.9	100.0

Table 3-206. Estimates of lost-time injuries by whether professional medical attention was sought for family workers in the Midwest region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	2683.8	3.8	2683.8	3.8
Yes	55669.7	79.5	58353.5	83.4
No	11637.4	16.6	69990.9	100.0

Table 3-207. Estimates of lost-time injuries by body part injured for family workers in the Midwest region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	3345.2	4.8	3345.2	4.8
Eye	3422.8	4.9	6768.0	9.7
Chest/trunk	3332.1	4.8	10100.1	14.4
Back	9197.5	13.1	19297.6	27.6
Arm/shoulder	6556.2	9.4	25853.8	36.9
Finger	7165.8	10.2	33019.6	47.2
Hand/wrist	9197.7	13.1	42217.3	60.3
Leg/knee/hip	13090.6	18.7	55307.9	79.0
Toe	174.8	0.2	55482.7	79.3
Foot	7052.0	10.1	62534.7	89.3
Mult. parts	6440.2	9.2	68974.9	98.5
Other	1016.0	1.5	69990.9	100.0

Table 3-208. Estimates of lost-time injuries by nature of injury for family workers in the Midwest region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	174.8	0.2	174.8	0.2
Bruise	8426.1	12.0	8600.9	12.3
Burn	58.7	0.1	8659.6	12.4
Cut	13896.9	19.9	22556.5	32.2
Crush	4534.3	6.5	27090.8	38.7
Fracture	10102.0	14.4	37192.8	53.1
Puncture	4309.1	6.2	41501.9	59.3
Sprain/strain	17712.8	25.3	59214.7	84.6
Mult. injury	6394.3	9.1	65609.0	93.7
Other	4381.9	6.3	69990.9	100.0

Table 3-209. Estimates of lost-time injuries by type of event for family workers in the Midwest region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	1383.6	2.0	1383.6	2.0
Caught between	6532.7	9.3	7916.3	11.3
Caught under	5028.1	7.2	12944.4	18.5
Struck by/against	16823.0	24.0	29767.4	42.5
Struck by falling	2800.5	4.0	32567.9	46.5
Struck by flying	1871.5	2.7	34439.4	49.2
Contact sharp obj.	7622.6	10.9	42062.0	60.1
Fall, same level	10171.2	14.5	52233.2	74.6
Fall, elevation	6682.7	9.5	58915.9	84.2
Overexertion	6303.7	9.0	65219.6	93.2
Other	4771.3	6.8	69990.9	100.0

Table 3-210. Estimates of lost-time injuries by activity when worker was injured for family workers in the Midwest region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	11870.7	17.0	11870.7	17.0
Machine maint.	10045.7	14.4	21916.4	31.3
Field work	8261.1	11.8	30177.5	43.1
Crop handling	11561.9	16.5	41739.4	59.6
Livestock handling	23920.3	34.2	65659.7	93.8
Other	4331.2	6.2	69990.9	100.0

Table 3-211. Estimates of lost-time injuries by source of the injury for family workers in the Midwest region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	4167.1	6.0	4167.1	6.0
Machinery	12046.6	17.2	16213.7	23.2
Livestock	14859.2	21.2	31072.9	44.4
Hand tool	8587.0	12.3	39659.9	56.7
Power tool	2646.4	3.8	42306.3	60.4
Pesticide/chem.	271.0	0.4	42577.3	60.8
Plant/tree	1274.9	1.8	43852.2	62.7
Working surface	7183.5	10.3	51035.7	72.9
Truck/auto	2339.0	3.3	53374.7	76.3
Other vehicle	2082.2	3.0	55456.9	79.2
Other	14534.0	20.8	69990.9	100.0

Table 3-212. Estimates of lost-time injuries by farm type for hired workers in the Midwest region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	2409.5	20.6	2409.5	20.6
Field crops	838.3	7.2	3247.8	27.7
Veg, fruit, nuts	856.5	7.3	4104.3	35.0
Nursery	1749.5	14.9	5853.8	49.9
Beef, hogs, sheep	2685.5	22.9	8539.3	72.8
Dairy	3184.4	27.2	11723.7	100.0

Table 3-213. Estimates of lost-time injuries by gender for hired workers in the Midwest region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	11503.8	98.1	11503.8	98.1
Female	219.9	1.9	11723.7	100.0

Table 3-214. Estimates of lost-time injuries by age group for hired workers in the Midwest region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	864.6	7.4	864.6	7.4
20-29	3976.2	33.9	4840.8	41.3
30-39	3919.3	33.4	8760.1	74.7
40-49	1148.4	9.8	9908.5	84.5
50-59	1815.2	15.5	11723.7	100.0

Table 3-215. Estimates of lost-time injuries by race or ethnic origin for hired workers in the Midwest region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	10368.2	88.4	10368.2	88.4
Hispanic	1355.5	11.6	11723.7	100.0

Table 3-216. Estimates of lost-time injuries by month for hired workers in the Midwest region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	679.5	5.8	679.5	5.8
February	222.3	1.9	901.8	7.7
March	186.6	1.6	1088.4	9.3
April	426.1	3.6	1514.5	12.9
May	2561.2	21.8	4075.7	34.8
June	1124.7	9.6	5200.4	44.4
July	1483.8	12.7	6684.2	57.0
August	584.0	5.0	7268.2	62.0
September	33.3	0.3	7301.5	62.3
October	720.3	6.1	8021.8	68.4
November	1615.1	13.8	9636.9	82.2
December	2086.8	17.8	11723.7	100.0

Table 3-217. Estimates of lost-time injuries by severity of the injury for hired workers in the Midwest region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	11537.1	98.4	11537.1	98.4
Permanent	186.6	1.6	11723.7	100.0

Table 3-218. Estimates of lost-time injuries by whether professional medical attention was sought for hired workers in the Midwest region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	186.6	1.6	186.6	1.6
Yes	11118.2	94.8	11304.8	96.4
No	418.9	3.6	11723.7	100.0

Table 3-219. Estimates of lost-time injuries by body part injured for hired workers in the Midwest region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	860.4	7.3	860.4	7.3
Eye	186.6	1.6	1047.0	8.9
Chest/trunk	1803.6	15.4	2850.6	24.3
Back	279.1	2.4	3129.7	26.7
Arm/shoulder	397.1	3.4	3526.8	30.1
Finger	2105.1	18.0	5631.9	48.0
Hand/wrist	1169.8	10.0	6801.7	58.0
Leg/knee/hip	1948.1	16.6	8749.8	74.6
Toe	281.0	2.4	9030.8	77.0
Foot	1185.5	10.1	10216.3	87.1
Mult. parts	1507.4	12.9	11723.7	100.0

Table 3-220. Estimates of lost-time injuries by nature of injury for hired workers in the Midwest region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	186.6	1.6	186.6	1.6
Bruise	3490.9	29.8	3677.5	31.4
Burn	255.6	2.2	3933.1	33.5
Cut	2027.0	17.3	5960.1	50.8
Crush	222.3	1.9	6182.4	52.7
Fracture	1956.5	16.7	8138.9	69.4
Poisoning	33.3	0.3	8172.2	69.7
Puncture	408.9	3.5	8581.1	73.2
Sprain/strain	2216.8	18.9	10797.9	92.1
Mult. injury	703.5	6.0	11501.4	98.1
Other	222.3	1.9	11723.7	100.0

Table 3-221. Estimates of lost-time injuries by type of event for hired workers in the Midwest region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	612.7	5.2	612.7	5.2
Caught between	2463.1	21.0	3075.8	26.2
Struck by/against	1084.5	9.3	4160.3	35.5
Struck by falling	707.1	6.0	4867.4	41.5
Struck by flying	186.6	1.6	5054.0	43.1
Contact sharp obj.	1088.4	9.3	6142.4	52.4
Fall, same level	946.4	8.1	7088.8	60.5
Fall, elevation	2440.8	20.8	9529.6	81.3
Other	2194.1	18.7	11723.7	100.0

Table 3-222. Estimates of lost-time injuries by activity when worker was injured for hired workers in the Midwest region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	1434.2	12.2	1434.2	12.2
Machine maint.	1752.9	15.0	3187.1	27.2
Field work	2916.0	24.9	6103.1	52.1
Crop handling	3646.2	31.1	9749.3	83.2
Livestock handling	1323.8	11.3	11073.1	94.5
Other	650.6	5.5	11723.7	100.0

Table 3-223. Estimates of lost-time injuries by source of the injury for hired workers in the Midwest region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	1153.5	9.8	1153.5	9.8
Machinery	3834.1	32.7	4987.6	42.5
Livestock	1020.1	8.7	6007.7	51.2
Hand tool	1031.8	8.8	7039.5	60.0
Pesticide/chem.	66.6	0.6	7106.1	60.6
Plant/tree	120.7	1.0	7226.8	61.6
Working surface	242.9	2.1	7469.7	63.7
Truck/auto	982.6	8.4	8452.3	72.1
Other vehicle	584.0	5.0	9036.3	77.1
Liquids	493.3	4.2	9529.6	81.3
Other	2194.1	18.7	11723.7	100.0

Table 3-224. Estimates of lost-time injuries by farm type for family workers in the West region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	1251.0	7.0	1251.0	7.0
Field crops	2138.1	12.0	3389.1	19.0
Veg, fruit, nuts	5138.5	28.8	8527.6	47.7
Beef, hogs, sheep	7386.5	41.3	15914.1	89.1
Poultry	562.5	3.1	16476.6	92.2
Dairy	834.0	4.7	17310.6	96.9
Other farms	555.2	3.1	17865.8	100.0

Table 3-225. Estimates of lost-time injuries by gender for family workers in the West region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	15950.5	89.3	15950.5	89.3
Female	1915.3	10.7	17865.8	100.0

Table 3-226. Estimates of lost-time injuries by age group for family workers in the West region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	2051.4	11.5	2051.4	11.5
20-29	462.9	2.6	2514.3	14.1
30-39	3970.3	22.2	6484.6	36.3
40-49	4100.3	23.0	10584.9	59.2
50-59	2975.5	16.7	13560.4	75.9
60-69	2699.4	15.1	16259.8	91.0
>=70	1606.0	9.0	17865.8	100.0

Table 3-227. Estimates of lost-time injuries by race or ethnic origin for family workers in the West region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	16038.7	89.8	16038.7	89.8
Hispanic	1074.3	6.0	17113.0	95.8
Asian	752.8	4.2	17865.8	100.0

Table 3-228. Estimates of lost-time injuries by month for family workers in the West region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	1771.5	9.9	1771.5	9.9
February	758.9	4.2	2530.4	14.2
March	1995.8	11.2	4526.2	25.3
April	447.1	2.5	4973.3	27.8
May	462.1	2.6	5435.4	30.4
June	847.7	4.7	6283.1	35.2
July	2565.6	14.4	8848.7	49.5
August	2418.4	13.5	11267.1	63.1
September	1464.5	8.2	12731.6	71.3
October	457.5	2.6	13189.1	73.8
November	1260.5	7.1	14449.6	80.9
December	3416.2	19.1	17865.8	100.0

Table 3-229. Estimates of lost-time injuries by severity of the injury for family workers in the West region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	16387.2	91.7	16387.2	91.7
Permanent	1478.6	8.3	17865.8	100.0

Table 3-230. Estimates of lost-time injuries by whether professional medical attention was sought for family workers in the West region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	1478.6	8.3	1478.6	8.3
Yes	13888.1	77.7	15366.7	86.0
No	2499.1	14.0	17865.8	100.0

Table 3-231. Estimates of lost-time injuries by body part injured for family workers in the West region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	654.6	3.7	654.6	3.7
Eye	622.4	3.5	1277.0	7.1
Chest/trunk	1500.7	8.4	2777.7	15.5
Back	2274.2	12.7	5051.9	28.3
Arm/shoulder	1283.2	7.2	6335.1	35.5
Finger	3508.5	19.6	9843.6	55.1
Hand/wrist	1840.6	10.3	11684.2	65.4
Leg/knee/hip	3647.1	20.4	15331.3	85.8
Toe	392.6	2.2	15723.9	88.0
Foot	826.6	4.6	16550.5	92.6
Mult. parts	893.4	5.0	17443.9	97.6
Other	421.9	2.4	17865.8	100.0

Table 3-232. Estimates of lost-time injuries by nature of injury for family workers in the West region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	190.3	1.1	190.3	1.1
Bruise	1595.4	8.9	1785.7	10.0
Burn	422.0	2.4	2207.7	12.4
Cut	2933.5	16.4	5141.2	28.8
Crush	2027.5	11.3	7168.7	40.1
Fracture	3609.8	20.2	10778.5	60.3
Poisoning	356.5	2.0	11135.0	62.3
Puncture	1895.4	10.6	13030.4	72.9
Sprain/strain	2672.9	15.0	15703.3	87.9
Mult. injury	1311.2	7.3	17014.5	95.2
Other	851.3	4.8	17865.8	100.0

Table 3-233. Estimates of lost-time injuries by type of event for family workers in the West region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	455.7	2.6	455.7	2.6
Caught between	3031.7	17.0	3487.4	19.5
Caught under	511.7	2.9	3999.1	22.4
Struck by/against	1778.2	10.0	5777.3	32.3
Struck by falling	450.9	2.5	6228.2	34.9
Struck by flying	1895.4	10.6	8123.6	45.5
Contact sharp obj.	1904.6	10.7	10028.2	56.1
Fall, same level	2215.5	12.4	12243.7	68.5
Fall, elevation	2544.1	14.2	14787.8	82.8
Overexertion	1418.5	7.9	16206.3	90.7
Other	1659.5	9.3	17865.8	100.0

Table 3-234. Estimates of lost-time injuries by activity when worker was injured for family workers in the West region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	1668.8	9.3	1668.8	9.3
Machine maint.	4352.7	24.4	6021.5	33.7
Field work	1622.9	9.1	7644.4	42.8
Crop handling	964.0	5.4	8608.4	48.2
Livestock handling	5917.2	33.1	14525.6	81.3
Other	3340.2	18.7	17865.8	100.0

Table 3-235. Estimates of lost-time injuries by source of the injury for family workers in the West region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	151.9	0.9	151.9	0.9
Machinery	2729.4	15.3	2881.3	16.1
Livestock	3997.1	22.4	6878.4	38.5
Hand tool	3371.7	18.9	10250.1	57.4
Power tool	46.3	0.3	10296.4	57.6
Pesticide/chem.	115.8	0.6	10412.2	58.3
Plant/tree	1138.6	6.4	11550.8	64.7
Working surface	1121.1	6.3	12671.9	70.9
Truck/auto	396.3	2.2	13068.2	73.1
Liquids	734.4	4.1	13802.6	77.3
Other	4063.2	22.7	17865.8	100.0

Table 3-236. Estimates of lost-time injuries by farm type for hired workers in the West region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	347.4	0.8	347.4	0.8
Field crops	2551.6	6.2	2899.0	7.0
Veg, fruit, nuts	19325.3	46.8	22224.3	53.8
Nursery	9081.2	22.0	31305.5	75.8
Beef, hogs, sheep	2504.4	6.1	33809.9	81.8
Poultry	333.3	0.8	34143.2	82.7
Dairy	5480.8	13.3	39624.0	95.9
Other farms	1683.8	4.1	41307.8	100.0

Table 3-237. Estimates of lost-time injuries by gender for hired workers in the West region of the U.S., 1993.

Gender	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Male	40190.0	97.3	40190.0	97.3
Female	1117.8	2.7	41307.8	100.0

Table 3-238. Estimates of lost-time injuries by age group for hired workers in the West region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	779.2	1.9	779.2	1.9
20-29	16629.0	40.3	17408.2	42.1
30-39	11715.6	28.4	29123.8	70.5
40-49	9352.9	22.6	38476.7	93.1
50-59	2372.9	5.7	40849.6	98.9
>=70	458.2	1.1	41307.8	100.0

Table 3-239. Estimates of lost-time injuries by race or ethnic origin for hired workers in the West region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	6295.5	15.2	6295.5	15.2
Hispanic	33868.0	82.0	40163.5	97.2
Black	164.0	0.4	40327.5	97.6
American Indian	190.3	0.5	40517.8	98.1
Other	790.0	1.9	41307.8	100.0

Table 3-240. Estimates of lost-time injuries by month for hired workers in the West region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	2267.9	5.5	2267.9	5.5
February	267.2	0.6	2535.1	6.1
March	2117.6	5.1	4652.7	11.3
April	1426.9	3.5	6079.6	14.7
May	3876.0	9.4	9955.6	24.1
June	6632.3	16.1	16587.9	40.2
July	6567.6	15.9	23155.5	56.1
August	7241.7	17.5	30397.2	73.6
September	3010.8	7.3	33408.0	80.9
October	3064.2	7.4	36472.2	88.3
November	4122.8	10.0	40595.0	98.3
December	692.0	1.7	41287.0	100.0

(Frequency Missing = 20.8)

Table 3-241. Estimates of lost-time injuries by severity of the injury for hired workers in the West region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	40171.2	97.2	40171.2	97.2
Permanent	1136.6	2.8	41307.8	100.0

Table 3-242. Estimates of lost-time injuries by whether professional medical attention was sought for hired workers in the West region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	1136.6	2.8	1136.6	2.8
Yes	33961.2	82.2	35097.8	85.0
No	6210.0	15.0	41307.8	100.0

Table 3-243. Estimates of lost-time injuries by body part injured for hired workers in the West region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	2253.3	5.5	2253.3	5.5
Eye	6620.1	16.0	8873.4	21.5
Chest/trunk	4383.2	10.6	13256.6	32.1
Back	11724.8	28.4	24981.4	60.5
Arm/shoulder	3686.6	8.9	28668.0	69.4
Finger	4242.1	10.3	32910.1	79.7
Hand/wrist	2524.9	6.1	35435.0	85.8
Leg/knee/hip	2468.8	6.0	37903.8	91.8
Foot	954.4	2.3	38858.2	94.1
Mult. parts	346.6	0.8	39204.8	94.9
Other	2103.0	5.1	41307.8	100.0

Table 3-244. Estimates of lost-time injuries by nature of injury for hired workers in the West region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	5790.5	14.0	5790.5	14.0
Burn	1034.5	2.5	6825.0	16.5
Cut	6600.3	16.0	13425.3	32.5
Crush	1685.0	4.1	15110.3	36.6
Fracture	1767.6	4.3	16877.9	40.9
Puncture	1402.1	3.4	18280.0	44.3
Sprain/strain	18386.7	44.5	36666.7	88.8
Mult. injury	148.4	0.4	36815.1	89.1
Other	4492.7	10.9	41307.8	100.0

Table 3-245. Estimates of lost-time injuries by type of event for hired workers in the West region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	809.8	2.0	809.8	2.0
Caught between	2081.6	5.0	2891.4	7.0
Caught under	1232.7	3.0	4124.1	10.0
Struck by/against	4525.4	11.0	8649.5	20.9
Struck by falling	1457.0	3.5	10106.5	24.5
Struck by flying	1171.8	2.8	11278.3	27.3
Contact sharp obj.	4589.1	11.1	15867.4	38.4
Fall, same level	3268.1	7.9	19135.5	46.3
Fall, elevation	1630.2	3.9	20765.7	50.3
Overexertion	13272.0	32.1	34037.7	82.4
Other	7270.1	17.6	41307.8	100.0

Table 3-246. Estimates of lost-time injuries by activity when worker was injured for hired workers in the West region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	5725.9	13.9	5725.9	13.9
Machine maint.	3445.6	8.3	9171.5	22.2
Field work	13063.4	31.6	22234.9	53.8
Crop handling	4502.5	10.9	26737.4	64.7
Livestock handling	2248.7	5.4	28986.1	70.2
Other	12321.7	29.8	41307.8	100.0

Table 3-247. Estimates of lost-time injuries by source of the injury for hired workers in the West region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	432.0	1.0	432.0	1.0
Machinery	5341.4	12.9	5773.4	14.0
Livestock	3088.5	7.5	8861.9	21.5
Hand tool	5843.3	14.1	14705.2	35.6
Power tool	2275.8	5.5	16981.0	41.1
Pesticide/chem.	114.5	0.3	17095.5	41.4
Plant/tree	3287.9	8.0	20383.4	49.3
Working surface	2919.5	7.1	23302.9	56.4
Truck/auto	620.3	1.5	23923.2	57.9
Other vehicle	1034.3	2.5	24957.5	60.4
Other	16350.3	39.6	41307.8	100.0

Table 3-248. Estimates of lost-time injuries by farm type for male workers in the Northeast region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	595.8	5.4	595.8	5.4
Field crops	2182.9	19.9	2778.7	25.4
Veg, fruit, nuts	1297.2	11.8	4075.9	37.2
Nursery	739.9	6.8	4815.8	44.0
Beef, hogs, sheep	1156.4	10.6	5972.2	54.5
Poultry	122.8	1.1	6095.0	55.6
Dairy	4739.6	43.3	10834.6	98.9
Other farms	119.1	1.1	10953.7	100.0

Table 3-249. Estimates of lost-time injuries by status of the injured worker for male workers in the Northeast region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	8753.8	79.9	8753.8	79.9
Hired	2086.3	19.0	10840.1	99.0
Other	113.6	1.0	10953.7	100.0

Table 3-250. Estimates of lost-time injuries by age group for male workers in the Midwest region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	907.7	8.3	907.7	8.3
20-29	2797.3	25.5	3705.0	33.8
30-39	1127.5	10.3	4832.5	44.1
40-49	2113.1	19.3	6945.6	63.4
50-59	2045.4	18.7	8991.0	82.1
60-69	653.1	6.0	9644.1	88.0
>=70	1309.6	12.0	10953.7	100.0

Table 3-251. Estimates of lost-time injuries by race or ethnic origin for male workers in the Northeast region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	9961.2	90.9	9961.2	90.9
Hispanic	992.5	9.1	10953.7	100.0

Table 3-252. Estimates of lost-time injuries by month for male workers in the Northeast region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	109.2	1.0	109.2	1.0
February	376.1	3.4	485.3	4.4
March	361.5	3.3	846.8	7.7
April	592.5	5.4	1439.3	13.1
May	671.9	6.1	2111.2	19.3
June	524.7	4.8	2635.9	24.1
July	2949.4	26.9	5585.3	51.0
August	484.7	4.4	6070.0	55.4
September	1448.5	13.2	7518.5	68.6
October	1476.4	13.5	8994.9	82.1
November	876.9	8.0	9871.8	90.1
December	1081.9	9.9	10953.7	100.0

Table 3-253. Estimates of lost-time injuries by severity of the injury for male workers in the Northeast region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	10840.1	99.0	10840.1	99.0
Permanent	113.6	1.0	10953.7	100.0

Table 3-254. Estimates of lost-time injuries by whether professional medical attention was sought for male workers in the Northeast region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	113.6	1.0	113.6	1.0
Yes	9054.3	82.7	9167.9	83.7
No	1785.8	16.3	10953.7	100.0

Table 3-255. Estimates of lost-time injuries by body part injured for male workers in the Northeast region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	422.9	3.9	422.9	3.9
Eye	923.9	8.4	1346.8	12.3
Chest/trunk	511.5	4.7	1858.3	17.0
Back	590.4	5.4	2448.7	22.4
Arm/shoulder	888.0	8.1	3336.7	30.5
Finger	2503.9	22.9	5840.6	53.3
Hand/wrist	2968.2	27.1	8808.8	80.4
Leg/knee/hip	1244.5	11.4	10053.3	91.8
Foot	652.6	6.0	10705.9	97.7
Mult. parts	178.5	1.6	10884.4	99.4
Other	69.3	0.6	10953.7	100.0

Table 3-256. Estimates of lost-time injuries by nature of injury for male workers in the Northeast region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	69.3	0.6	69.3	0.6
Bruise	1470.4	13.4	1539.7	14.1
Cut	2905.6	26.5	4445.3	40.6
Crush	419.5	3.8	4864.8	44.4
Fracture	1478.1	13.5	6342.9	57.9
Puncture	361.9	3.3	6704.8	61.2
Sprain/strain	2769.2	25.3	9474.0	86.5
Mult. injury	113.6	1.0	9587.6	87.5
Other	1366.1	12.5	10953.7	100.0

Table 3-257. Estimates of lost-time injuries by type of event for male workers in the Northeast region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	392.4	3.6	392.4	3.6
Caught between	1432.0	13.1	1824.4	16.7
Struck by/against	1089.5	9.9	2913.9	26.6
Struck by falling	685.2	6.3	3599.1	32.9
Struck by flying	618.4	5.6	4217.5	38.5
Contact sharp obj.	2445.7	22.3	6663.2	60.8
Fall, same level	632.9	5.8	7296.1	66.6
Fall, elevation	1084.5	9.9	8380.6	76.5
Overexertion	1236.0	11.3	9616.6	87.8
Other	1337.1	12.2	10953.7	100.0

Table 3-258. Estimates of lost-time injuries by activity when worker was injured for male workers in the Northeast region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	2046.9	18.7	2046.9	18.7
Machine maint.	2371.8	21.7	4418.7	40.3
Field work	1767.1	16.1	6185.8	56.5
Crop handling	2036.8	18.6	8222.6	75.1
Livestock handling	2175.5	19.9	10398.1	94.9
Other	555.6	5.1	10953.7	100.0

Table 3-259. Estimates of lost-time injuries by source of the injury for male workers in the Northeast region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	316.9	2.9	316.9	2.9
Machinery	2386.9	21.8	2703.8	24.7
Livestock	1729.3	15.8	4433.1	40.5
Hand tool	427.9	3.9	4861.0	44.4
Power tool	1465.2	13.4	6326.2	57.8
Plant/tree	248.1	2.3	6574.3	60.0
Working surface	1133.7	10.3	7708.0	70.4
Truck/auto	471.1	4.3	8179.1	74.7
Other vehicle	69.3	0.6	8248.4	75.3
Other	2705.3	24.7	10953.7	100.0

Table 3-260. Estimates of lost-time injuries by farm type for female workers in the Northeast region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Field crops	610.9	17.7	610.9	17.7
Veg, fruit, nuts	201.4	5.8	812.3	23.5
Nursery	667.7	19.3	1480.0	42.8
Beef, hogs, sheep	666.6	19.3	2146.6	62.1
Poultry	27.8	0.8	2174.4	62.9
Dairy	1157.2	33.5	3331.6	96.4
Other farms	126.1	3.6	3457.7	100.0

Table 3-261. Estimates of lost-time injuries by status of the injured worker for female workers in the Northeast region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	2022.5	58.5	2022.5	58.5
Hired	1435.2	41.5	3457.7	100.0

Table 3-262. Estimates of lost-time injuries by age group for female workers in the Midwest region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	376.1	10.9	376.1	10.9
20-29	907.3	26.2	1283.4	37.1
30-39	606.2	17.5	1889.6	54.6
40-49	468.4	13.5	2358.0	68.2
50-59	201.6	5.8	2559.6	74.0
60-69	675.9	19.5	3235.5	93.6
>=70	222.2	6.4	3457.7	100.0

Table 3-263. Estimates of lost-time injuries by race or ethnic origin for female workers in the Northeast region of the U.S., 1993.

Race/Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	3440.0	99.5	3440.0	99.5
Hispanic	17.7	0.5	3457.7	100.0

Table 3-264. Estimates of lost-time injuries by month for female workers in the Northeast region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	44.3	1.3	44.3	1.3
February	291.5	8.4	335.8	9.7
March	36.1	1.0	371.9	10.8
April	853.8	24.7	1225.7	35.4
May	829.7	24.0	2055.4	59.4
June	221.4	6.4	2276.8	65.8
July	17.7	0.5	2294.5	66.4
August	550.2	15.9	2844.7	82.3
September	69.3	2.0	2914.0	84.3
October	171.4	5.0	3085.4	89.2
November	354.6	10.3	3440.0	99.5
December	17.7	0.5	3457.7	100.0

Table 3-265. Estimates of lost-time injuries by severity of the injury for female workers in the Northeast region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	3103.1	89.7	3103.1	89.7
Permanent	354.6	10.3	3457.7	100.0

Table 3-266. Estimates of lost-time injuries by whether professional medical attention was sought for female workers in the Northeast region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	354.6	10.3	354.6	10.3
Yes	2599.9	75.2	2954.5	85.4
No	503.2	14.6	3457.7	100.0

Table 3-267. Estimates of lost-time injuries by body part injured for female workers in the Northeast region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	347.7	10.1	347.7	10.1
Eye	114.6	3.3	462.3	13.4
Chest/trunk	154.5	4.5	616.8	17.8
Back	250.0	7.2	866.8	25.1
Arm/shoulder	69.3	2.0	936.1	27.1
Finger	473.4	13.7	1409.5	40.8
Hand/wrist	887.7	25.7	2297.2	66.4
Leg/knee/hip	222.2	6.4	2519.4	72.9
Toe	17.7	0.5	2537.1	73.4
Foot	631.3	18.3	3168.4	91.6
Other	289.3	8.4	3457.7	100.0

Table 3-268. Estimates of lost-time injuries by nature of injury for female workers in the Northeast region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	134.6	3.9	134.6	3.9
Bruise	759.7	22.0	894.3	25.9
Crush	204.2	5.9	1098.5	31.8
Fracture	1041.0	30.1	2139.5	61.9
Puncture	114.6	3.3	2254.1	65.2
Sprain/strain	965.9	27.9	3220.0	93.1
Other	237.7	6.9	3457.7	100.0

Table 3-269. Estimates of lost-time injuries by type of event for female workers in the Northeast region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	220.0	6.4	220.0	6.4
Caught between	517.7	15.0	737.7	21.3
Caught under	309.2	8.9	1046.9	30.3
Struck by/against	151.1	4.4	1198.0	34.6
Struck by flying	220.0	6.4	1418.0	41.0
Contact sharp obj.	69.3	2.0	1487.3	43.0
Fall, same level	1148.1	33.2	2635.4	76.2
Overexertion	250.0	7.2	2885.4	83.4
Other	572.3	16.6	3457.7	100.0

Table 3-270. Estimates of lost-time injuries by activity when worker was injured for female workers in the Northeast region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Field work	935.1	27.0	935.1	27.0
Crop handling	799.8	23.1	1734.9	50.2
Livestock handling	1148.5	33.2	2883.4	83.4
Other	574.3	16.6	3457.7	100.0

Table 3-271. Estimates of lost-time injuries by source of the injury for female workers in the Northeast region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Machinery	423.9	12.3	423.9	12.3
Livestock	926.9	26.8	1350.8	39.1
Plant/tree	114.6	3.3	1465.4	42.4
Working surface	345.1	10.0	1810.5	52.4
Other	1647.2	47.6	3457.7	100.0

Table 3-272. Estimates of lost-time injuries by farm type for male workers in the South region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	2485.2	6.3	2485.2	6.3
Field crops	6936.4	17.5	9421.6	23.8
Veg, fruit, nuts	4906.4	12.4	14328.0	36.2
Nursery	1775.2	4.5	16103.2	40.7
Beef, hogs, sheep	17034.4	43.0	33137.6	83.7
Poultry	955.7	2.4	34093.3	86.1
Dairy	3883.7	9.8	37977.0	95.9
Other farms	1613.3	4.1	39590.3	100.0

Table 3-273. Estimates of lost-time injuries by status of the injured worker for male workers in the South region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	23594.5	59.6	23594.5	59.6
Hired	15718.3	39.7	39312.8	99.3
Other	277.5	0.7	39590.3	100.0

Table 3-274. Estimates of lost-time injuries by age group for male workers in the South region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	1394.2	3.5	1394.2	3.5
20-29	7728.1	19.5	9122.3	23.0
30-39	9305.5	23.5	18427.8	46.5
40-49	6213.8	15.7	24641.6	62.2
50-59	4869.8	12.3	29511.4	74.5
60-69	6143.5	15.5	35654.9	90.1
>=70	3935.4	9.9	39590.3	100.0

Table 3-275. Estimates of lost-time injuries by race or ethnic origin for male workers in the South region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	32191.8	81.3	32191.8	81.3
Hispanic	5355.6	13.5	37547.4	94.8
Black	1258.6	3.2	38806.0	98.0
American Indian	784.3	2.0	39590.3	100.0

Table 3-276. Estimates of lost-time injuries by month for male workers in the South region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	2872.7	7.3	2872.7	7.3
February	3975.7	10.0	6848.4	17.3
March	1978.7	5.0	8827.1	22.3
April	2491.6	6.3	11318.7	28.6
May	2648.8	6.7	13967.5	35.3
June	4654.7	11.8	18622.2	47.0
July	2162.4	5.5	20784.6	52.5
August	4134.6	10.4	24919.2	62.9
September	3139.2	7.9	28058.4	70.9
October	4018.7	10.2	32077.1	81.0
November	3273.4	8.3	35350.5	89.3
December	4239.8	10.7	39590.3	100.0

Table 3-277. Estimates of lost-time injuries by severity of the injury for male workers in the South region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	35106.3	88.7	35106.3	88.7
Permanent	4111.4	10.4	39217.7	99.1
Fatality	372.6	0.9	39590.3	100.0

Table 3-278. Estimates of lost-time injuries by whether professional medical attention was sought for male workers in the South region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	4484.0	11.3	4484.0	11.3
Yes	29724.0	75.1	34208.0	86.4
No	5382.3	13.6	39590.3	100.0

Table 3-279. Estimates of lost-time injuries by body part injured for male workers in the South region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	1840.4	4.6	1840.4	4.6
Eye	1621.8	4.1	3462.2	8.7
Chest/trunk	5422.3	13.7	8884.5	22.4
Back	4737.2	12.0	13621.7	34.4
Arm/shoulder	2439.6	6.2	16061.3	40.6
Finger	3293.8	8.3	19355.1	48.9
Hand/wrist	4094.1	10.3	23449.2	59.2
Leg/knee/hip	7040.5	17.8	30489.7	77.0
Toe	964.8	2.4	31454.5	79.5
Foot	5904.6	14.9	37359.1	94.4
Mult. parts	2106.4	5.3	39465.5	99.7
Other	124.8	0.3	39590.3	100.0

Table 3-280. Estimates of lost-time injuries by nature of injury for male workers in the South region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	398.6	1.0	398.6	1.0
Bruise	6117.3	15.5	6515.9	16.5
Burn	112.5	0.3	6628.4	16.7
Cut	7856.6	19.8	14485.0	36.6
Crush	3967.2	10.0	18452.2	46.6
Fracture	9625.9	24.3	28078.1	70.9
Puncture	1078.0	2.7	29156.1	73.6
Sprain/strain	5509.3	13.9	34665.4	87.6
Mult. injury	2272.6	5.7	36938.0	93.3
Other	2652.3	6.7	39590.3	100.0

Table 3-281. Estimates of lost-time injuries by type of event for male workers in the South region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	1275.0	3.2	1275.0	3.2
Caught between	3025.9	7.6	4300.9	10.9
Caught under	1288.1	3.3	5589.0	14.1
Struck by/against	10749.3	27.2	16338.3	41.3
Struck by falling	4587.2	11.6	20925.5	52.9
Struck by flying	905.3	2.3	21830.8	55.1
Contact sharp obj.	4639.8	11.7	26470.6	66.9
Fall, same level	4585.9	11.6	31056.5	78.4
Fall, elevation	3728.3	9.4	34784.8	87.9
Overexertion	3224.9	8.1	38009.7	96.0
Other	1580.6	4.0	39590.3	100.0

Table 3-282. Estimates of lost-time injuries by activity when worker was injured for male workers in the South region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	8060.0	20.4	8060.0	20.4
Machine maint.	4016.4	10.1	12076.4	30.5
Field work	7774.4	19.6	19850.8	50.1
Crop handling	6413.9	16.2	26264.7	66.3
Livestock handling	12509.2	31.6	38773.9	97.9
Other	816.4	2.1	39590.3	100.0

Table 3-283. Estimates of lost-time injuries by source of the injury for male workers in the South region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	4394.7	11.1	4394.7	11.1
Machinery	7285.9	18.4	11680.6	29.5
Livestock	8503.9	21.5	20184.5	51.0
Hand tool	2937.4	7.4	23121.9	58.4
Power tool	805.9	2.0	23927.8	60.4
Pesticide/chem.	487.5	1.2	24415.3	61.7
Plant/tree	1716.0	4.3	26131.3	66.0
Working surface	3813.6	9.6	29944.9	75.6
Truck/auto	1331.6	3.4	31276.5	79.0
Other vehicle	1071.6	2.7	32348.1	81.7
Other	7242.2	18.3	39590.3	100.0

Table 3-284. Estimates of lost-time injuries by farm type for female workers in the South region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	600.0	13.3	600.0	13.3
Field crops	252.3	5.6	852.3	18.9
Nursery	595.2	13.2	1447.5	32.1
Beef,hogs,sheep	2704.4	60.0	4151.9	92.1
Dairy	179.6	4.0	4331.5	96.1
Other farms	176.4	3.9	4507.9	100.0

Table 3-285. Estimates of lost-time injuries by status of the injured worker for female workers in the South region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	4061.5	90.1	4061.5	90.1
Hired	446.4	9.9	4507.9	100.0

Table 3-286. Estimates of lost-time injuries by age group for female workers in the South region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
30-39	1102.4	25.3	1102.4	25.3
40-49	815.2	18.7	1917.6	44.0
50-59	1894.1	43.5	3811.7	87.4
>=70	547.4	12.6	4359.1	100.0
(Frequency Missing = 148.8)				

Table 3-287. Estimates of lost-time injuries by race or ethnic origin for female workers in the South region of the U.S., 1993.

Race/Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	4061.5	93.2	4061.5	93.2
Hispanic	297.6	6.8	4359.1	100.0
(Frequency Missing = 148.8)				

Table 3-288. Estimates of lost-time injuries by month for female workers in the South region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
March	148.8	3.4	148.8	3.4
April	600.0	13.8	748.8	17.2
May	277.5	6.4	1026.3	23.5
June	797.2	18.3	1823.5	41.8
July	237.0	5.4	2060.5	47.3
August	547.4	12.6	2607.9	59.8
September	555.0	12.7	3162.9	72.6
October	252.3	5.8	3415.2	78.3
November	486.8	11.2	3902.0	89.5
December	457.1	10.5	4359.1	100.0

(Frequency Missing = 148.8)

Table 3-289. Estimates of lost-time injuries by severity of the injury for female workers in the South region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	4507.9	100.0	4507.9	100.0

Table 3-290. Estimates of lost-time injuries by whether professional medical attention was sought for female workers in the South region of the U.S., 1993.

Attention Required	Medical		Cumulative Frequency	Cumulative Percent
	Frequency	Percent		
Yes	3872.3	85.9	3872.3	85.9
No	635.6	14.1	4507.9	100.0

Table 3-291. Estimates of lost-time injuries by body part injured for female workers in the South region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	365.7	8.1	365.7	8.1
Back	824.9	18.3	1190.6	26.4
Arm/shoulder	88.2	2.0	1278.8	28.4
Finger	457.1	10.1	1735.9	38.5
Hand/wrist	973.7	21.6	2709.6	60.1
Leg/knee/hip	852.3	18.9	3561.9	79.0
Foot	398.6	8.8	3960.5	87.9
Mult. parts	547.4	12.1	4507.9	100.0

Table 3-292. Estimates of lost-time injuries by nature of injury for female workers in the South region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	88.2	2.0	88.2	2.0
Burn	398.6	8.8	486.8	10.8
Cut	797.6	17.7	1284.4	28.5
Crush	398.6	8.8	1683.0	37.3
Fracture	877.5	19.5	2560.5	56.8
Sprain/strain	973.7	21.6	3534.2	78.4
Mult. injury	547.4	12.1	4081.6	90.5
Other	426.3	9.5	4507.9	100.0

Table 3-293. Estimates of lost-time injuries by type of event for female workers in the South region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught between	457.1	10.1	457.1	10.1
Struck by/against	1399.9	31.1	1857.0	41.2
Contact sharp obj.	252.3	5.6	2109.3	46.8
Fall, same level	1026.3	22.8	3135.6	69.6
Fall, elevation	148.8	3.3	3284.4	72.9
Overexertion	277.5	6.2	3561.9	79.0
Other	946.0	21.0	4507.9	100.0

Table 3-294. Estimates of lost-time injuries by activity when worker was injured for female workers in the South region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	385.8	8.6	385.8	8.6
Machine maint.	398.6	8.8	784.4	17.4
Field work	1001.1	22.2	1785.5	39.6
Crop handling	703.8	15.6	2489.3	55.2
Livestock handling	2018.6	44.8	4507.9	100.0

Table 3-295. Estimates of lost-time injuries by source of the injury for female workers in the South region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	148.8	3.3	148.8	3.3
Machinery	277.5	6.2	426.3	9.5
Livestock	1829.3	40.6	2255.6	50.0
Hand tool	252.3	5.6	2507.9	55.6
Pesticide/chem.	398.6	8.8	2906.5	64.5
Plant/tree	748.8	16.6	3655.3	81.1
Working surface	297.6	6.6	3952.9	87.7
Other	555.0	12.3	4507.9	100.0

Table 3-296. Estimates of lost-time injuries by farm type for male workers in the Midwest region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	23043.1	31.2	23043.1	31.2
Field crops	3179.7	4.3	26222.8	35.5
Veg, fruit, nuts	1581.3	2.1	27804.1	37.6
Nursery	1716.2	2.3	29520.3	40.0
Beef, hogs, sheep	27362.6	37.0	56882.9	77.0
Dairy	16970.4	23.0	73853.3	100.0

Table 3-297. Estimates of lost-time injuries by status of the injured worker for male workers in the Midwest region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	61730.1	83.6	61730.1	83.6
Hired	11503.8	15.6	73233.9	99.2
Other	619.4	0.8	73853.3	100.0

Table 3-298. Estimates of lost-time injuries by age group for male workers in the Midwest region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	6354.7	8.6	6354.7	8.6
20-29	9340.3	12.6	15695.0	21.3
30-39	19528.9	26.4	35223.9	47.7
40-49	10524.4	14.3	45748.3	61.9
50-59	13401.1	18.1	59149.4	80.1
60-69	10836.9	14.7	69986.3	94.8
>=70	3867.0	5.2	73853.3	100.0

Table 3-299. Estimates of lost-time injuries by race or ethnic origin for male workers in the Midwest region of the U.S., 1993.

Race/Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	71706.1	97.3	71706.1	97.3
Hispanic	1972.4	2.7	73678.5	100.0
(Frequency Missing = 174.8)				

Table 3-300. Estimates of lost-time injuries by month for male workers in the Midwest region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	4199.6	5.7	4199.6	5.7
February	1506.2	2.0	5705.8	7.7
March	1815.0	2.5	7520.8	10.2
April	6339.6	8.6	13860.4	18.8
May	5660.8	7.7	19521.2	26.5
June	10475.1	14.2	29996.3	40.7
July	9983.3	13.5	39979.6	54.3
August	7786.1	10.6	47765.7	64.8
September	10166.2	13.8	57931.9	78.6
October	6257.5	8.5	64189.4	87.1
November	6056.8	8.2	70246.2	95.3
December	3432.3	4.7	73678.5	100.0

(Frequency Missing = 174.8)

Table 3-301. Estimates of lost-time injuries by severity of the injury for male workers in the Midwest region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	70808.1	96.1	70808.1	96.1
Permanent	1992.1	2.7	72800.2	98.8
Fatality	878.3	1.2	73678.5	100.0

(Frequency Missing = 174.8)

Table 3-302. Estimates of lost-time injuries by whether professional medical attention was sought for male workers in the Midwest region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Known	3045.2	4.1	3045.2	4.1
Yes	59315.1	80.3	62360.3	84.4
No	11493.0	15.6	73853.3	100.0

Table 3-303. Estimates of lost-time injuries by body part injured for male workers in the Midwest region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	3237.2	4.4	3237.2	4.4
Eye	3609.4	4.9	6846.6	9.3
Chest/trunk	5135.7	7.0	11982.3	16.2
Back	8588.4	11.6	20570.7	27.9
Arm/shoulder	5532.3	7.5	26103.0	35.3
Finger	9063.5	12.3	35166.5	47.6
Hand/wrist	9330.6	12.6	44497.1	60.3
Leg/knee/hip	12980.6	17.6	57477.7	77.8
Toe	455.8	0.6	57933.5	78.4
Foot	7573.5	10.3	65507.0	88.7
Mult. parts	7330.3	9.9	72837.3	98.6
Other	1016.0	1.4	73853.3	100.0

Table 3-304. Estimates of lost-time injuries by nature of injury for male workers in the Midwest region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	361.4	0.5	361.4	0.5
Bruise	10951.8	14.8	11313.2	15.3
Burn	314.3	0.4	11627.5	15.7
Cut	13375.0	18.1	25002.5	33.9
Crush	4570.0	6.2	29572.5	40.0
Fracture	10617.9	14.4	40190.4	54.4
Puncture	4940.3	6.7	45130.7	61.1
Sprain/strain	18087.5	24.5	63218.2	85.6
Mult. injury	6310.0	8.5	69528.2	94.1
Other	4325.1	5.9	73853.3	100.0

Table 3-305. Estimates of lost-time injuries by type of event for male workers in the Midwest region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	1996.3	2.7	1996.3	2.7
Caught between	7691.5	10.4	9687.8	13.1
Caught under	4018.0	5.4	13705.8	18.6
Struck by/against	15382.6	20.8	29088.4	39.4
Struck by falling	3427.6	4.6	32516.0	44.0
Struck by flying	2280.4	3.1	34796.4	47.1
Contact sharp obj.	7288.4	9.9	42084.8	57.0
Fall, same level	10297.4	13.9	52382.2	70.9
Fall, elevation	8742.3	11.8	61124.5	82.8
Overexertion	6075.8	8.2	67200.3	91.0
Other	6653.0	9.0	73853.3	100.0

Table 3-306. Estimates of lost-time injuries by activity when worker was injured for male workers in the Midwest of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	12679.4	17.2	12679.4	17.2
Machine maint.	11146.6	15.1	23826.0	32.3
Field work	9805.3	13.3	33631.3	45.5
Crop handling	14805.4	20.0	48436.7	65.6
Livestock handling	21637.3	29.3	70074.0	94.9
Other	3779.3	5.1	73853.3	100.0

Table 3-307. Estimates of lost-time injuries by source of the injury for male workers in the Midwest region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	4911.4	6.7	4911.4	6.7
Machinery	15110.1	20.5	20021.5	27.1
Livestock	12973.2	17.6	32994.7	44.7
Hand tool	9411.4	12.7	42406.1	57.4
Power tool	2646.4	3.6	45052.5	61.0
Pesticide/chem.	304.3	0.4	45356.8	61.4
Plant/tree	1395.6	1.9	46752.4	63.3
Working surface	7309.7	9.9	54062.1	73.2
Truck/auto	2820.2	3.8	56882.3	77.0
Other vehicle	2240.1	3.0	59122.4	80.1
Liquids	493.3	0.7	59615.7	80.7
Other	14237.6	19.3	73853.3	100.0

Table 3-308. Estimates of lost-time injuries by farm type for female workers in the Midwest region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	648.4	7.6	648.4	7.6
Field crops	130.0	1.5	778.4	9.2
Veg, fruit, nuts	177.9	2.1	956.3	11.3
Nursery	33.3	0.4	989.6	11.7
Beef, hogs, sheep	3477.9	41.0	4467.5	52.7
Poultry	80.0	0.9	4547.5	53.6
Dairy	3293.2	38.8	7840.7	92.5
Other farms	640.0	7.5	8480.7	100.0

Table 3-309. Estimates of lost-time injuries by status of the injured worker for female workers in the Midwest region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	8260.8	97.4	8260.8	97.4
Hired	219.9	2.6	8480.7	100.0

Table 3-310. Estimates of lost-time injuries by age group for female workers in the Midwest region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	1010.1	11.9	1010.1	11.9
20-29	61.2	0.7	1071.3	12.6
30-39	2159.4	25.5	3230.7	38.1
40-49	2922.4	34.5	6153.1	72.6
50-59	2327.6	27.4	8480.7	100.0

Table 3-311. Estimates of lost-time injuries by race or ethnic origin for female workers in the Midwest region of the U.S., 1993.

Race/Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	8480.7	100.0	8480.7	100.0

Table 3-312. Estimates of lost-time injuries by month for female workers in the Midwest region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	395.8	4.7	395.8	4.7
February	320.0	3.8	715.8	8.4
March	0.0	0.0	715.8	8.4
April	800.3	9.4	1516.1	17.9
May	1390.3	16.4	2906.4	34.3
June	1398.1	16.5	4304.5	50.8
July	1354.6	16.0	5659.1	66.7
August	463.9	5.5	6123.0	72.2
September	33.3	0.4	6156.3	72.6
October	130.0	1.5	6286.3	74.1
November	770.6	9.1	7056.9	83.2
December	1423.8	16.8	8480.7	100.0

Table 3-313. Estimates of lost-time injuries by severity of the injury for female workers in the Midwest region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	8480.7	100.0	8480.7	100.0

Table 3-314. Estimates of lost-time injuries by whether professional medical attention was sought for female workers in the Midwest region of the U.S., 1993.

Attention Required	Medical		Cumulative Frequency	Cumulative Percent
	Frequency	Percent		
Yes	7917.4	93.4	7917.4	93.4
No	563.3	6.6	8480.7	100.0

Table 3-315. Estimates of lost-time injuries by body part injured for female workers in the Midwest region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	968.4	11.4	968.4	11.4
Back	1063.0	12.5	2031.4	24.0
Arm/shoulder	1421.0	16.8	3452.4	40.7
Finger	652.0	7.7	4104.4	48.4
Hand/wrist	1036.9	12.2	5141.3	60.6
Leg/knee/hip	2058.1	24.3	7199.4	84.9
Foot	664.0	7.8	7863.4	92.7
Mult. parts	617.3	7.3	8480.7	100.0

Table 3-316. Estimates of lost-time injuries by nature of injury for female workers in the Midwest region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	965.2	11.4	965.2	11.4
Cut	2548.9	30.1	3514.1	41.4
Crush	186.6	2.2	3700.7	43.6
Fracture	1440.6	17.0	5141.3	60.6
Poisoning	33.3	0.4	5174.6	61.0
Sprain/strain	2016.9	23.8	7191.5	84.8
Mult. injury	1010.1	11.9	8201.6	96.7
Other	279.1	3.3	8480.7	100.0

Table 3-317. Estimates of lost-time injuries by type of event for female workers in the Midwest region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught between	1304.3	15.4	1304.3	15.4
Caught under	1010.1	11.9	2314.4	27.3
Struck by/against	2524.9	29.8	4839.3	57.1
Struck by falling	302.3	3.6	5141.6	60.6
Contact sharp obj.	1422.6	16.8	6564.2	77.4
Fall, same level	820.2	9.7	7384.4	87.1
Fall, elevation	381.2	4.5	7765.6	91.6
Overexertion	402.7	4.7	8168.3	96.3
Other	312.4	3.7	8480.7	100.0

Table 3-318. Estimates of lost-time injuries by activity when worker was injured for female workers in the Midwest region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	800.3	9.4	800.3	9.4
Machine maint.	874.3	10.3	1674.6	19.7
Field work	1594.1	18.8	3268.7	38.5
Crop handling	402.7	4.7	3671.4	43.3
Livestock handling	3606.8	42.5	7278.2	85.8
Other	1202.5	14.2	8480.7	100.0

Table 3-319. Estimates of lost-time injuries by source of the injury for female workers in the Midwest region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	584.0	6.9	584.0	6.9
Machinery	770.6	9.1	1354.6	16.0
Livestock	2906.1	34.3	4260.7	50.2
Hand tool	652.0	7.7	4912.7	57.9
Pesticide/chem.	33.3	0.4	4946.0	58.3
Working surface	116.7	1.4	5062.7	59.7
Truck/auto	501.4	5.9	5564.1	65.6
Other vehicle	426.1	5.0	5990.2	70.6
Other	2490.5	29.4	8480.7	100.0

Table 3-320. Estimates of lost-time injuries by farm type for male workers in the West region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	1464.2	2.6	1464.2	2.6
Field crops	6038.4	10.7	7502.6	13.2
Veg, fruit, nuts	23544.2	41.5	31046.8	54.8
Nursery	9081.2	16.0	40128.0	70.8
Beef, hogs, sheep	9494.5	16.8	49622.5	87.6
Poultry	333.3	0.6	49955.8	88.1
Dairy	6120.4	10.8	56076.2	98.9
Other farms	596.8	1.1	56673.0	100.0

Table 3-321. Estimates of lost-time injuries by status of the injured worker for male workers in the West region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	15950.5	28.1	15950.5	28.1
Hired	40190.0	70.9	56140.5	99.1
Other	532.5	0.9	56673.0	100.0

Table 3-322. Estimates of lost-time injuries by age group for male workers in the West region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
10-19	2830.6	5.0	2830.6	5.0
20-29	16977.4	30.0	19808.0	35.0
30-39	14872.4	26.2	34680.4	61.2
40-49	12434.9	21.9	47115.3	83.1
50-59	5356.6	9.5	52471.9	92.6
60-69	2136.9	3.8	54608.8	96.4
>=70	2064.2	3.6	56673.0	100.0

Table 3-323. Estimates of lost-time injuries by race or ethnic origin for male workers in the West region of the U.S., 1993.

Race/Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	21151.7	37.3	21151.7	37.3
Hispanic	34186.7	60.3	55338.4	97.6
Black	164.0	0.3	55502.4	97.9
Asian	190.3	0.3	55692.7	98.3
American Indian	190.3	0.3	55883.0	98.6
Other	790.0	1.4	56673.0	100.0

Table 3-324. Estimates of lost-time injuries by month for male workers in the West region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	4039.4	7.1	4039.4	7.1
February	1026.1	1.8	5065.5	8.9
March	3863.4	6.8	8928.9	15.8
April	1827.7	3.2	10756.6	19.0
May	4143.7	7.3	14900.3	26.3
June	6801.5	12.0	21701.8	38.3
July	9133.2	16.1	30835.0	54.4
August	9545.6	16.8	40380.6	71.3
September	4323.4	7.6	44704.0	78.9
October	3140.0	5.5	47844.0	84.5
November	5308.8	9.4	53152.8	93.8
December	3499.4	6.2	56652.2	100.0

(Frequency Missing = 20.8)

Table 3-325. Estimates of lost-time injuries by severity of the injury for male workers in the West region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	54209.7	95.7	54209.7	95.7
Permanent	2463.3	4.3	56673.0	100.0

Table 3-326. Estimates of lost-time injuries by whether professional medical attention was sought for male workers in the West region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Known	2463.3	4.3	2463.3	4.3
Yes	45500.6	80.3	47963.9	84.6
No	8709.1	15.4	56673.0	100.0

Table 3-327. Estimates of lost-time injuries by body part injured for male workers in the West region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	2659.2	4.7	2659.2	4.7
Eye	7128.0	12.6	9787.2	17.3
Chest/trunk	5883.9	10.4	15671.1	27.7
Back	13952.7	24.6	29623.8	52.3
Arm/shoulder	4393.7	7.8	34017.5	60.0
Finger	7914.6	14.0	41932.1	74.0
Hand/wrist	3756.7	6.6	45688.8	80.6
Leg/knee/hip	5544.9	9.8	51233.7	90.4
Toe	392.6	0.7	51626.3	91.1
Foot	1413.8	2.5	53040.1	93.6
Mult. parts	855.3	1.5	53895.4	95.1
Other	2777.6	4.9	56673.0	100.0

Table 3-328. Estimates of lost-time injuries by nature of injury for male workers in the West region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Amputation	190.3	0.3	190.3	0.3
Bruise	6330.8	11.2	6521.1	11.5
Burn	1456.5	2.6	7977.6	14.1
Cut	9683.3	17.1	17660.9	31.2
Crush	3712.5	6.6	21373.4	37.7
Fracture	4618.0	8.1	25991.4	45.9
Poisoning	162.1	0.3	26153.5	46.1
Puncture	3297.5	5.8	29451.0	52.0
Sprain/strain	20760.6	36.6	50211.6	88.6
Mult. injury	1269.3	2.2	51480.9	90.8
Other	5192.1	9.2	56673.0	100.0

Table 3-329. Estimates of lost-time injuries by type of event for male workers in the West region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught in	1265.5	2.2	1265.5	2.2
Caught between	5113.3	9.0	6378.8	11.3
Caught under	1592.5	2.8	7971.3	14.1
Struck by/against	5863.3	10.3	13834.6	24.4
Struck by falling	1065.4	1.9	14900.0	26.3
Struck by flying	3319.9	5.9	18219.9	32.1
Contact sharp obj.	6032.1	10.6	24252.0	42.8
Fall, same level	4393.1	7.8	28645.1	50.5
Fall, elevation	4290.1	7.6	32935.2	58.1
Overexertion	14690.5	25.9	47625.7	84.0
Other	9047.3	16.0	56673.0	100.0

Table 3-330. Estimates of lost-time injuries by activity when worker was injured for male workers in the West region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	7291.5	12.9	7291.5	12.9
Machine maint.	7914.1	14.0	15205.6	26.8
Field work	13204.2	23.3	28409.8	50.1
Crop handling	5466.5	9.6	33876.3	59.8
Livestock handling	7134.8	12.6	41011.1	72.4
Other	15661.9	27.6	56673.0	100.0

Table 3-331. Estimates of lost-time injuries by source of the injury for male workers in the West region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Tractor	583.9	1.0	583.9	1.0
Machinery	7936.6	14.0	8520.5	15.0
Livestock	7140.1	12.6	15660.6	27.6
Hand tool	8947.8	15.8	24608.4	43.4
Power tool	2486.1	4.4	27094.5	47.8
Pesticide/chem.	230.3	0.4	27324.8	48.2
Plant/tree	3864.0	6.8	31188.8	55.0
Working surface	3787.9	6.7	34976.7	61.7
Truck/auto	1016.6	1.8	35993.3	63.5
Other vehicle	1034.3	1.8	37027.6	65.3
Liquids	734.4	1.3	37762.0	66.6
Other	18911.0	33.4	56673.0	100.0

Table 3-332. Estimates of lost-time injuries by farm type for female workers in the West region of the U.S., 1993.

Farm Type	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Cash grain	250.0	8.2	250.0	8.2
Field crops	190.3	6.3	440.3	14.5
Veg, fruit, nuts	919.6	30.3	1359.9	44.8
Beef, hogs, sheep	649.1	21.4	2009.0	66.2
Poultry	562.5	18.5	2571.5	84.8
Dairy	194.4	6.4	2765.9	91.2
Other farms	267.2	8.8	3033.1	100.0

Table 3-333. Estimates of lost-time injuries by status of the injured worker for female workers in the West region of the U.S., 1993.

Status of Injured Worker	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Family	1915.3	63.1	1915.3	63.1
Hired	1117.8	36.9	3033.1	100.0

Table 3-334. Estimates of lost-time injuries by age group for female workers in the West region of the U.S., 1993.

Age Group (Years)	Frequency	Percent	Cumulative Frequency	Cumulative Percent
20-29	114.5	3.8	114.5	3.8
30-39	1093.3	36.0	1207.8	39.8
40-49	1018.3	33.6	2226.1	73.4
50-59	244.5	8.1	2470.6	81.5
60-69	562.5	18.5	3033.1	100.0

Table 3-335. Estimates of lost-time injuries by race or ethnic origin for female workers in the West region of the U.S., 1993.

Race/ Ethnic Origin	Frequency	Percent	Cumulative Frequency	Cumulative Percent
White	1551.0	51.1	1551.0	51.1
Hispanic	919.6	30.3	2470.6	81.5
Asian	562.5	18.5	3033.1	100.0

Table 3-336. Estimates of lost-time injuries by month for female workers in the West region of the U.S., 1993.

Month	Frequency	Percent	Cumulative Frequency	Cumulative Percent
January	0.0	0.0	0.0	0.0
February	0.0	0.0	0.0	0.0
March	250.0	8.2	250.0	8.2
April	299.0	9.9	549.0	18.1
May	194.4	6.4	743.4	24.5
June	842.5	27.8	1585.9	52.3
July	0.0	0.0	1585.9	52.3
August	114.5	3.8	1700.4	56.1
September	151.9	5.0	1852.3	61.1
October	381.7	12.6	2234.0	73.7
November	190.3	6.3	2424.3	79.9
December	608.8	20.1	3033.1	100.0

Table 3-337. Estimates of lost-time injuries by severity of the injury for female workers in the West region of the U.S., 1993.

Severity of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Temporary	2881.2	95.0	2881.2	95.0
Permanent	151.9	5.0	3033.1	100.0

Table 3-338. Estimates of lost-time injuries by whether professional medical attention was sought for female workers in the West region of the U.S., 1993.

Medical Attention Required	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Unknown	151.9	5.0	151.9	5.0
Yes	2881.2	95.0	3033.1	100.0

Table 3-339. Estimates of lost-time injuries by body part injured for female workers in the West region of the U.S., 1993.

Body Part Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Head/neck	364.5	12.0	364.5	12.0
Eye	114.5	3.8	479.0	15.8
Back	46.3	1.5	525.3	17.3
Arm/shoulder	576.1	19.0	1101.4	36.3
Hand/wrist	608.8	20.1	1710.2	56.4
Leg/knee/hip	571.0	18.8	2281.2	75.2
Foot	367.2	12.1	2648.4	87.3
Mult. parts	384.7	12.7	3033.1	100.0

Table 3-340. Estimates of lost-time injuries by nature of injury for female workers in the West region of the U.S., 1993.

Nature of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Bruise	1055.1	34.8	1055.1	34.8
Cut	267.2	8.8	1322.3	43.6
Fracture	875.2	28.9	2197.5	72.5
Poisoning	194.4	6.4	2391.9	78.9
Sprain/strain	299.0	9.9	2690.9	88.7
Mult. injury	190.3	6.3	2881.2	95.0
Other	151.9	5.0	3033.1	100.0

Table 3-341. Estimates of lost-time injuries by type of event for female workers in the West region of the U.S., 1993.

Type of Event	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Caught under	151.9	5.0	151.9	5.0
Struck by/against	440.3	14.5	592.2	19.5
Struck by falling	842.5	27.8	1434.7	47.3
Contact sharp obj.	461.6	15.2	1896.3	62.5
Fall, same level	1090.5	36.0	2986.8	98.5
Other	46.3	1.5	3033.1	100.0

Table 3-342. Estimates of lost-time injuries by activity when worker was injured for female workers in the West region of the U.S., 1993.

Activity When Injured	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Farm maint.	267.2	8.8	267.2	8.8
Field work	1482.1	48.9	1749.3	57.7
Livestock handling	1283.8	42.3	3033.1	100.0

Table 3-343. Estimates of lost-time injuries by source of the injury for female workers in the West region of the U.S., 1993.

Source of Injury	Frequency	Percent	Cumulative Frequency	Cumulative Percent
Machinery	250.0	8.2	250.0	8.2
Livestock	198.2	6.5	448.2	14.8
Hand tool	267.2	8.8	715.4	23.6
Plant/tree	562.5	18.5	1277.9	42.1
Working surface	252.7	8.3	1530.6	50.5
Other	1502.5	49.5	3033.1	100.0