

2006 National Youth Tobacco Survey and Key Prevalence Indicators

TABLE 1. Percentage of students in middle school (grades 6-8) who were current users\* of any tobacco product, by product type, sex, and race/ethnicity — United States, 2004 and 2006

Characteristic	Any tobacco†		Cigarettes		Cigars		Smokeless tobacco		Pipes		Bidis		Kreteks	
	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)	%	(95% CI§)
<b>Middle school, 2006</b>														
<b>Sex</b>														
Male	10.9	(9.2-12.6)	6.3	(5.1-7.5)	5.3¶	(4.4-6.2)	4.1	(3.1-5.1)	3.0	(2.4-3.6)	1.9**	(1.4-2.4)	1.7	(1.3-2.1)
Female	8.2¶	(6.7-9.7)	6.4¶	(5.0-7.8)	2.7¶	(2.2-3.2)	1.2¶	(0.9-1.5)	1.3¶	(1.0-1.6)	1.5	(1.2-1.8)	1.0	(0.6-1.4)
<b>Race/Ethnicity</b>														
White, non-Hispanic	9.2	(7.3-11.1)	6.5	(4.9-8.1)	3.1¶	(2.4-3.8)	2.8	(2.0-3.6)	1.9	(1.3-2.5)	1.2¶	(0.8-1.6)	1.0	(0.7-1.3)
Black, non-Hispanic	9.8	(7.6-12.0)	5.5	(3.9-7.1)	5.8	(4.2-7.4)	1.7	(1.0-2.4)	1.5	(0.9-2.1)	2.3	(1.4-3.2)	1.7	(1.0-2.4)
Hispanic	10.9**	(9.3-12.5)	6.8**	(5.6-8.0)	6.1¶	(4.9-7.3)	3.4	(2.6-4.2)	4.3	(3.5-5.1)	3.3	(2.6-4.0)	2.6	(1.9-3.3)
Asian	4.9	(3.2-6.6)	2.6	(1.3-3.9)	1.6	(0.5-2.7)	2.0	(0.7-3.3)	2.2	(0.9-3.5)	1.9	(0.6-3.2)	1.2	(0.4-2.0)
<b>Total</b>	<b>9.5¶</b>	<b>(8.0-11.0)</b>	<b>6.3¶</b>	<b>(5.1-7.5)</b>	<b>4.0¶</b>	<b>(3.4-4.6)</b>	<b>2.6</b>	<b>(2.0-3.2)</b>	<b>2.2</b>	<b>(1.8-2.6)</b>	<b>1.7¶</b>	<b>(1.4-2.0)</b>	<b>1.4</b>	<b>(1.2-1.6)</b>
<b>Middle school, 2004</b>														
<b>Sex</b>														
Male	12.7	(11.2-14.2)	7.9	(6.7-9.1)	6.7	(5.8-7.6)	3.8	(3.0-4.6)	3.5	(2.8-4.2)	3.0	(2.5-3.5)	2.0	(1.6-2.4)
Female	10.9	(9.5-12.3)	8.8	(7.4-10.2)	3.8	(3.2-4.4)	1.9	(1.5-2.3)	1.9	(1.5-2.3)	1.8	(1.4-2.2)	1.2	(0.9-1.5)
<b>Race/Ethnicity</b>														
White, non-Hispanic	11.3	(9.7-12.9)	8.5	(7.0-10.0)	4.4	(3.7-5.1)	3.0	(2.3-3.7)	2.3	(1.7-2.9)	1.9	(1.5-2.3)	1.3	(1.0-1.6)
Black, non-Hispanic	12.4	(10.2-14.6)	7.6	(5.9-9.3)	6.9	(5.2-8.6)	2.0	(1.2-2.8)	2.2	(1.5-2.9)	2.9	(2.0-3.8)	1.6	(1.0-2.2)
Hispanic	15.1	(13.0-17.2)	9.9	(8.4-11.4)	8.2	(6.9-9.5)	3.8	(2.9-4.7)	5.3	(4.1-6.5)	4.3	(3.5-5.1)	2.9	(2.2-3.6)
Asian	5.1	(2.8-7.4)	2.7	(1.0-4.4)	1.2	(0.1-2.3)	1.0	(0.3-1.7)	1.5	(0.3-2.7)	1.1	(0.0-2.2)	1.5	(0.2-2.8)
<b>Total</b>	<b>11.8</b>	<b>(10.5-13.1)</b>	<b>8.4</b>	<b>(7.3-9.5)</b>	<b>5.3</b>	<b>(4.6-6.0)</b>	<b>2.8</b>	<b>(2.3-3.3)</b>	<b>2.7</b>	<b>(2.2-3.2)</b>	<b>2.4</b>	<b>(2.0-2.8)</b>	<b>1.6</b>	<b>(1.3-1.9)</b>

\* Used tobacco on on  $\geq 1$  of the 30 preceding days.

† Cigarettes, cigars, smokeless tobacco, pipes, bidis (leaf-wrapped, flavored cigarettes from India), or kreteks (clove cigarettes).

§ Confidence interval.

¶ Significant difference ( $p < 0.05$ ), 2006 versus 2004.

\*\* Significant difference ( $p < 0.01$ ), 2006 versus 2004.