

8. REFERENCES

- *Abou-Donia MB, Hu ZH, Lapadula DM, et al.1991. Mechanisms of joint neurotoxicity of *n*-hexane, methyl isobutyl ketone and O-ethyl O-4-nitrophenyl phenylphosphonothioate in hens. *J pharmacol Exp Ther* 257(1):282-289.
- *Abou-Donia MB, Lapadula DM.1990. Mechanisms of organophosphorus ester-induced delayed neurotoxicity: Type I and type II. *Annu Rev Pharmacol Toxicol* 30:405-440.
- *Abou-Donia MB, Lapadula DM, Campbell G, et al.1985 The synergism of *n*-hexane-induced neurotoxicity by methyl isobutyl ketone following subchronic (90 days) inhalation in hens: Induction of hepatic microsomal cytochrome P-450. *Toxicol Appl Pharmacol* 81(1):1-16.
- *Abou-Donia MB, Lapadula DM, Suwita E.1988. Cytoskeletal proteins as targets for organophosphorous compound and aliphatic hexacarbon-induced neurotoxicity. *Toxicology* 49:469-477.
- *Abou-Donia MB, Makkawy HA, Graham DG.1982. The relative neurotoxicities of *n*-hexane, methyl *n*-butyl ketone, 2,5-hexanediol, and 2,5-hexanedione following oral or intraperitoneal administration in hens. *Toxicol Appl Pharmacol* 62(3):369-389.
- *ACGIH.1996. Threshold limit values for chemical substances and physical agents and biological exposure indices for 1995-1996. American Conference of Governmental Industrial Hygienists, Cincinnati, OH.
- *Adinolfi M.1985. The development of the human blood-CSF-brain barrier. *Developmental Medicine & Child Neurology* 27:532-537.
- *Ahearn DG, Crow SA, Simmons RB, et al.1996. Fungal colonization of fiberglass insulation in the air distribution system of a multi-story office building: VOC production and possible relationship to a sick building syndrome. *J Ind Microbial* 16:280-285.
- *Allen N, Mendell JR, Billmaier DJ, et al.1975. Toxic polyneuropathy due to methyl *n*-butyl ketone. *Arch Neurol* 32:209-218.
- *Altenkirch H, Mager J, Stoltenburg G, et al.1977. Toxic polyneuropathies after sniffing a glue thinner. *J Neurol* 214:137-152.
- *Altenkirch H, Wagner HM, Stoltenburg G, et al.1982. Nervous system responses of rats to subchronic inhalation of *n*-hexane and *n*-hexane + methyl-ethyl-ketone mixture. *J Neurol Sci* 57:209-219.
- *Altenkirch H, Wagner HM.1982. Potentiation of hexacarbon-neurotoxicity by methyl-ethyl-ketone (MEK) and other substances: Clinical and experimental aspects. *Neurobehav Toxicol Teratol* 4:623-627.
- *Altman PK, Dittmer DS.1974. In: *Biological handbooks: Biology data book, Volume III*, second edition. Bethesda, MD: Federation of American Societies for Experimental Biology, 1987-2008, 2041.

* Cited in text

8. REFERENCES

- *Amoore JE, Hautala E.1983. Odor as an aid to chemical safety: Odor thresholds compared with threshold limit values and volatilities for 214 industrial chemicals in air and water dilution. *J Appl Toxicol* 3(6):272-290.
- *Andersen ME, Krishnan K.1994. Relating *in vitro* to *in vivo* exposures with physiologically-based tissue dosimetry and tissue response models. In: H. Salem, ed. Current concepts and approaches on animal test alternatives. U.S. Army Chemical Research Development and Engineering Center, Aberdeen Proving Ground, Maryland.
- *Andersen ME, MacNaughton MG, Clewell HJ, et al.1987. Adjusting exposure limits for long and short exposure periods using a physiological pharmacokinetic model. *Am Ind Hyg Assoc J* 48(4):335-343.
- *Anlauf KG, Bottenheim JW, Brice KA.1985. Measurement of atmospheric aerosols and photochemical products at a rural site in SW Ontario. *Atmos Environ* 19(11):1859-1870.
- *Ashby J.1985. Fundamental structural alerts to potential carcinogenicity or non-carcinogenicity. *Environ Mutagen* 7:919-921.
- *Ashley DL, Bonin MA, Cardinali FL.1992. Determining volatile organic compounds in human blood from a large sample production by using purge and trap gas chromatography-mass spectrometry. *Anal Chem* 64(9):1021-1029.
- *Ashley DL, Bonin MA, Cardinali FL, et al.1994. Blood concentration of volatile organic compounds *in* a nonoccupationally exposed US population and in groups with suspected exposure. *Clin Chem* 40(7):1401-1409.
- *ASTER.1995. ASTER Ecotoxicity Profile: *n*-Hexane. Duluth, MN: U.S. EPA Office of Research and Development, National Health and Environmental Effects Research Laboratory, Mid-Continent Ecology Division.
- *Atkinson R.1985. Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions. *Chem Rev* 85:69-201.
- *Atkinson R, Aschman SM, Carter WPL.1982. Kinetics of the reactions of OH radicals with *n*-alkanes at 299K. *Int J Chem Kinetics* 14:781-788.
- *Atkinson R, Carter WPL.1984. Kinetics and mechanisms of the gas phase reactions of ozone with organic compounds under atmospheric conditions. *Chem Rev* 84:437-470.
- *ATSDR.1989. Decision guide for identifying substance-specific data needs related to toxicological profiles. Agency for Toxic Substances and Disease Registry, Division of Toxicology, Atlanta, GA.
- *ATSDR.1991. Toxicological profile for 2-hexanone. Agency for Toxic Substances and Disease Registry. Public Health Service. U.S. Department of Health and Human Services. Atlanta, GA. TP-91/18.
- *ATSDR/CDC.1990. Subcommittee report on biological indicators of organ damage. Agency for Toxic Substances and Disease Registry, Centers for Disease Control and Prevention, Atlanta, GA.

8. REFERENCES

- *Bachmann MO, De Beer Z, Myers JE.1993. *n*-Hexane neurotoxicity in metal can manufacturing workers. *Occup Med* 43:149-154.
- *Baelum J, Molhave L, Hansen SH, et al.1998. Metabolic interaction between toluene, trichloroethylene and *n*-hexane in humans. *Stand J Work Environ Health* 24:30-37.
- *Baker TS Rickert DE.1981. Dose-dependent uptake, distribution, and elimination of inhaled *n*-hexane in the Fischer-344 rat. *Toxicol Appl Pharmacol* 61(3):414-422.
- *Barnes DG, Dourson M.1988. Reference dose (RfD): Description and use in health risk assessments. U.S. Environmental Protection Agency. *Regul Toxicol Pharmacol* 8:471-486.
- *Barrefors G, Petersson G.1993. Assessment of ambient volatile hydrocarbons from tobacco smoke and from vehicle emissions. *J Chromatogr* 643:71-76.
- *Bartolucci GB, Perbellini L, Gori GP, et al.1986. Occupational exposure to solvents: Field comparison of active and passive samplers and biological monitoring of exposed workers. *Ann Occup Hyg* 30(3):295-306.
- *Bavazzano P, Apostoli P, Balducci C, et al.1998. Determination of urinary 2,5-hexanedione in the general Italian population. *Int Arch Occup Environ Health* 71(4):284-288.
- *Bayer C.1995. Beware of fungus among us: Emissions from mold and fungus may be culprits in indoor air problems. *Georgia Tech Research News*, August 3, 1995. Atlanta GA: Institute of Technology.
- *Belkin F, Hable MA.1988. Analysis of gasoline in water using a stripping preconcentration procedure. *Bull Environ Contam Toxicol* 40:244-248.
- *Bhagya S, Srinivas H.1992. Extraction of soybean with hexane-acetic acid: Effect on oil quality. *Food Chem* 44:123-125.
- *Bianchi AP, Varnoy MS, Phillips J.1991. Analysis of volatile organic compounds in estuarine sediments using dynamic headspace and gas chromatography mass spectrometry. *J Chromatogr* 542:413-450.
- *Bio/Dynamics.1995a. An inhalation oncogenicity study of commercial hexane in rats and mice--1. Rats, abridged final report. American Petroleum Institute Publication TR 404 41-332319
- *Bio/Dynamics.1995b. An inhalation oncogenicity study of commercial hexane in rats and mice--2. Mice, abridged final report. American Petroleum Institute Publication TR 405 41-33232.
- *Bio/Dynamics Inc.1978. 26 Week inhalation toxicity study of *n*-hexane in the rat. Unpublished report prepared for the American Petroleum Institute, Washington, DC. EPA -FYI-AX-1081-0137.
- *Bishop GA, Zhang Y, McLaren SE, et al.1994. Enhancement of remote sensing for vehicle emissions in tunnels. *Air & Waste* 44:169-75.
- *Biziuk M, Namiesnik J, Czerwinski J et al.1996. Occurrence and determination of organic pollutants in tap and surface waters of the Gdansk district. *J Chromatogr A* 733:171-183.

8. REFERENCES

- *Bottenheim JW, Shepherd MF.1995. C2-C6 Hydrocarbon measurements at four rural locations across Canada. *Atmospheric Environment* 29(6):647-664.
- *Bouhamra WS.1995. Estimation of volatile organic compounds in Kuwaiti houses after the Gulf War. *Environmental Pollution* 90(1):121-125.
- *Brosseau J, Heitz M.1994. Trace gas compound emissions from municipal landfill sanitary sites. *Atmos Environ* 28(2):285-293.
- *Brugnone F, Maranelli G, Romeo L, et al.1991. Ubiquitous pollution by *n*-hexane and reference biological levels in the general population. *Int Arch Occup Environ Health* 63(3):157-160.
- *Brugnone F, Perbellini L, Giulaiari C, et al.1994. Blood and urine concentrations of chemical pollutants in the general population. *Med Lav* 85:370-389.
- *Bus JS, Deyo D, Cox M.1982. Dose-dependent disposition of *n*-hexane in F-344 rats after inhalation exposure. *Fund Appl Toxicol* 2(5):226-229.
- *Bus JS, White EL, Gillies PJ, et al.1981. Tissue distribution of *n*-hexane, methyl *n*-butyl ketone, and 2,5-hexanedione in rats after single or repeated inhalation exposure to *n*-hexane. *Drug Metab Dispos* 9(4):386-387.
- *Bus JS, White EL, Tyl RW, et al.1979. Perinatal toxicity and metabolism of *n*-hexane in Fischer-344 rats after inhalation exposure during gestation. *Toxicol Appl Pharmacol* 51(2):295-302.
- *Cao X-L, Hewitt CN.1993. Passive sampling and gas chromatographic determination of low concentrations of reactive hydrocarbons in ambient air with reduction gas detector. *J Chromatogr* 648:191-197.
- *Cardona A, Marhuenda D, Marti J, et al.1993. Biological monitoring of occupational exposure to *n*-hexane by measurement of urinary 2,5-hexanedione. *Int Arch Occup Environ Health* 65:71-74.
- *Cardona A, Marhuenda D, Prieto MJ, et al.1996. Behaviour of urinary 2,5-hexanedione in occupational co-exposure to *n*-hexane and acetone. *Int Arch Occup Environ Health* 68(2):88-93.
- *Cavender FL, Casey HW, Salem H, et al.1984. A 13-week vapor inhalation study of *n*-hexane in rats with emphasis on neurotoxic effects. *Fund Appl Toxicol* 4:191-201.
- *Chan C, Liu S, Her G.1994. Office workers exposure to volatile organic compounds while commuting and working in Taipei City. *Atmos Environ* 28(14):2351-2359.
- *Chang AP, England JD, Garcia CA, et al.1998. Focal conduction block in *n*-hexane polyneuropathy. *Muscle and Nerve* 21:964-969.
- *Chang CM, Yu CW, Fong KY, et al.1993. *n*-Hexane neuropathy in offset printers. *J Neurol Neurosurg Psychiatry* 56:538-542.
- *Chapin RE, Norton RM, Popp JA, et al.1982. The effects of 2,5-hexanedione on reproductive hormones and testicular enzyme activities in the F-344 rat. *Toxicol Appl Pharmacol* 62:262-272.

8. REFERENCES

- CIIT.1977. Current status report No.1. *n*-Hexane. Chemical Industry Institute of Toxicology, Research Triangle Park, NC., Feb. 4, 1977.
- CIIT.1982. Ninety-day vapor inhalation toxicity study on *n*-hexane in albino rats. Unpublished report prepared for the Chemical Industry Institute of Toxicology by ToxiGenics Inc. CIIT Docket No.10242. Chemical Industry Institute of Toxicology, Research Triangle Park, NC.
- *Clewell HJ III, Andersen M.1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.
- *Coates M, Connell DW, Barron DM.1985. Aqueous solubility and octanol to water partition coefficients of aliphatic hydrocarbons. *Environ Sci Technol* 19:628-632.
- Yonkerton EJ, Wan PJ, Richard OA.1995. Hexane and heptane as extraction solvents for cottonseed: A laboratory-scale study. *J Am Oil Chem Soc* 72:963-965.
- *Cooper DA, Peterson K, Simpson D.1996. Hydrocarbon PAH and PCB emissions from ferries: A case study in the Skagerak Kattégatt oresund region. *Atmos Environ* 30(14):2463-2473.
- *Couri D, Abdel-Rahman MS, Hetland LB.1978. Biotransformation of *n*-hexane and methyl *n*-butyl ketone in guinea pigs and mice. *Am Ind Hyg Assoc J* 39(4):295-300.
- *Crawford DW, Bonnevie NL, Wenning RJ.1995. Sources of pollution and sediment contamination in Newark Bay, New Jersey. *Ecotoxicol Environ Safety* 30:85-100.
- *Creaser C, Welch J, Bishop K.1983. Letters to the editor. *n*-Hexane in petroleum ether fractions. *Ann Occup Hyg* 27(2):229-230.
- *Crosbie SJ, Blain PG, Williams FM.1997. Metabolism of *n*-hexane by rat liver and extrahepatic tissues and the effect of cytochrome P-450 inducers. *Hum Exp Toxicol* 16:131-137.
- *Daisey JM, Hodgson AT, Fisk WJ, et al.1994. Volatile Organic Compounds in Twelve California Office Buildings: Classes, Concentrations and Sources. *Atmospheric Environment* 28(22):3557-3562.
- *De Martino C, Malorni W, Amantini MC. et al.1987. Effects of respiratory treatment with *n*-hexane on rat testis morphology. I. A light microscopic study. *Exp Mol Pathol* 46(2):199-216.
- *DeCaprio AP, O'Neill EA.1985. Alterations in rat axonal cytoskeletal proteins induced by *in vitro* and *in vivo* 2,5-hexanedione exposure. *Toxicol Appl Pharmacol* 78:235-247.
- *DeCaprio AP, Olajos EJ, Weber P, et al.1982. Covalent binding of a neurotoxic *n*-hexane metabolite: conversion of primary amines to substituted pyrrole adducts by 2,5-hexanedione. *Toxicol Appl Pharmacol* 65:440-450.
- *DeLuchi MA.1993. Emissions from the production, storage, and transport of crude oil and gasoline. *Air & Waste* 43:1486-1495.
- Dominguez H, Nunez MJ, Lema JM.1995. Enzyme-assisted hexane extraction of soya bean oil. *Food Chem* 54:223-231.

8. REFERENCES

- *DOT.1990. Department of Transportation. Code of Federal Regulations. 49 CFR 172.101.
- *Dunnick JK, Graham DG, Yang RS, et al.1989. Thirteen-week toxicity study of *n*-hexane in B6C3F1 mice after inhalation exposure. *Toxicology* 57(2):163-172.
- *Eiceman GA,McConnon JT, Zaman M, et al.1986. Hydrocarbons and aromatic hydrocarbons in groundwater surrounding an earthen waste disposal pit for produced water in the Duncan oil field of New Mexico. Groundwater Hazardous Waste Bureau, Environmental Improvement Division, Environment and Health Dept, State of New Mexico, Santa Fe.
- *EPA.1980. Ambient water quality criteria for chlorinated benzenes. U.S. Environmental Protection Agency, Office of Water, Regulations and Standards, Washington, DC. EPA 440/5-80-028, NTIS No. PB81-117392.
- *EPA.1985. Standards for the management of specific hazardous wastes and specific types of hazardous waste management facilities. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 266.
- *EPA.1987a. Exposure analysis modeling system (EXAMSII). Center for Exposure Assessment Modeling, Ecosystems Research Division, National Exposure Research Laboratory, Office of Research and Development. U. S. Environmental Protection Agency. Athens, GA.
- *EPA.1987b. Bulk organic chemicals. Applicability; description of the bulk organic chemicals subcategory. U. S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 414.70.
- *EPA.1988. Compendium of methods for the determination of toxic organic compounds in ambient air. U. S. Environmental Protection Agency, U. S Department of Commerce, National Technical Information Service . Springfield VA.
- *EPA.1990. Standards of performance for volatile organic compounds (VOC) emissions from synthetic organic chemical manufacturing industry (SOCMI) distillation operation. U. S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 60.667.
- *EPA.1991. Alpha 2y-globulin: association with chemically induced renal toxicity and neoplasia in the male rat. Risk Assessment Forum U.S. Environmental Protection Agency, Washington DC. EPA/625/3-91/019F.
- *EPA.1993a. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 302.4.
- *EPA.1993b. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 60.707, Subpart RRR.
- *EPA.1994a. U.S. Environmental Protection Agency. Federal Register Notice. 59 FR 1788.
- *EPA.1994b. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 63.100, subpart F.
- *EPA.1994~. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 63.106.

8. REFERENCES

- *EPA.1994d. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 63, Subpart G, Appendix, Tables 8,9, and 34.
- *EPA.1994e. U.S. Environmental Protection Agency. Federal Register Notice. 59 FR 61432.
- *EPA.1994f. U.S. Environmental Protection Agency. Federal Register Notice. 59 FR 9808.
- *EPA .1994g. Health Effects Notebook for Hazardous Air Pollutants. Durham, NC: U.S. EPA, Office of Air Quality Planning and Standards, Air Risk Information Support Center.
- *EPA.1994h. Automobile Emissions: an Overview. EPA 400-F-92-007. Ann Arbor, MI: U.S. EPA, Office of Mobile Sources, National Vehicle and Fuel Emissions Laboratory.
- *EPA.1994i. Method 0030. Volatile organic sampling train. Method Status Table SW-846, third edition updates I, II, IIA and IIB. U. S. Environmental Protection Agency.
- *EPA.1994j. Health effects notebook for hazardous air pollutants. U.S. Environmental Protection Agency.
- *EPA.1995a. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 63, Subpart CC, Appendix, Tables 1, 5, and 7.
- *EPA.1995b. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 63.800, Subpart JJ.
- *EPA.1995c. U.S. Environmental Protection Agency. Federal Register Notice. 60 FR 21592.
- *EPA.1995d. Universal Treatment Standards. U. S. Environment Protection Agency. 60 FR 242.
- *EPA.1995e. Hazadous waste management system; carbamate production identification and listing quantities. Final Rule. U.S. Environmental Protection Agency. Federal Register. 60 FR 7824. February 9, 1995.
- *EPA.1995f. Toxics Chemical Release Inventory Reporting Form R and instructions: Revised 1994 version. EPA 745-K-95-051. Washington DC: U. S. EPA. Office of Pollution Prevention and Toxics.
- *EPA.1995g. Toxics release inventory: reporting modifications beginning with 1995 reporting year. U.S. Environmental Protection Agency. EPA 745-R-95-009.
- *EPA.1996a. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 63.506, Subpart U.
- *EPA.1996b. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 63.698, Subpart DD.
- *EPA.1996c. Proposed air toxics regulation for the printing and publishing industry. Air Toxics Regulations: Printing and Publishing. U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. Durham, NC.

8. REFERENCES

- *EPA.1996d. Understanding the hazardous waste rule: A handbook for small business. EPA530-K-95-001 (1996 Update). Washington, DC: U.S. Environmental Protection Agency.
- *EPA.1996e. National emission standards for hazardous air pollutant emissions: group I polymers and resins. Final Rule. Environmental Protection Agency. Federal Register. 61 FR 46906. September 5, 1997.
- *EPA.1996f. Drinking water regulations and health advisories. U.S. Environmental Protection Agency. Office of Water.
- *Ervin ME, Manske MG.1990. Petroleum distillates and turpentine. In: Haddad LM, Winchester JF, eds. Clinical management of poisoning and drug overdose. 2nd edition. Philadelphia, PA: W.B. Saunders Company, 1177-1186.
- *Farmer TH.1996. Occupational hygiene limits for hydrocarbon solvents. *Ann Occup Hyg* 40(2):237-242.
- *Fedtke N, Bolt HM.1986a. Detection of 2,5-hexanedione in the urine of persons not exposed to *n*-hexane. *Int Arch Occup Environ Health* 57:143-148.
- *Fedtke N, Bolt HM.1986b. Methodological investigations on the determination of *n*-hexane metabolites in urine. *Int Arch Occup Environ Health* 57:149-158.
- *Fedtke N, Bolt HM.1987. The relevance of 4,5-dihydroxy-2-hexanone in the excretion kinetics of *n*-hexane metabolites in rat and man. *Arch Toxicol* 61:131-137.
- *Feenstra S, Mackay DM, Cherry JA.1991. A method for assessing residual NAPL based on organic chemical concentrations in soil samples. *Ground Water Monitoring Review* 11(2):128-136.
- *Fellin P, Otson R.1994. Assessment of the influence of climatic factors on concentration levels of volatile organic compounds (VOCs) in Canadian homes. *Atmos Environ* 28(22):3581-3586.
- *Filser JG, Bolt HM, Muliawan H, et al, 1983. Quantitative evaluation of ethane and *n*-pentane as indicators of lipid peroxidation *in vivo*. *Arch Toxicol* 52:135-147.
- *Filser JG, Peter H, Bolt HM, et al.1987. Pharmacokinetics of the neurotoxin *n*-hexane in rat and man. *Arch Toxicol* 60:77-80.
- *Firestone D.1997. Personal communication.
- *Fisher J, Mahle D, Bankston L, et al.1997. Lactational transfer of volatile chemicals in breast milk. *Am Ind Hyg Assoc J* 58(6):425-431.
- *Foman SJ.1966. Body composition of the infant (Part I: The male reference infant). In: Falkner F, editor. Human development. Philadelphia, PA: WB Saunders, 239-246.
- *Fornan, SJ, Haschke F, Ziegler EE et al.1982. Body composition of reference children from birth to age 10 years. *American Journal of Clinical Nutrition* 35 :1169-1175.

8. REFERENCES

- *Frontali N, Amantini MC, Spagnolo AM, et al.1981. Experimental neurotoxicity and urinary metabolites of the C5-C7 aliphatic hydrocarbons used as glue solvents in shoe manufacture. Clin Toxicol 18(12):1357-1367.
- *FSTRAC.1990. Summary of state and federal drinking water standards and guidelines. U.S. Environmental Protection Agency. Chemical Communication Subcommittee, Federal State Toxicology and Regulatory Alliance Committee (FSTRAC).
- *FSTRAC.1995. Summary of state and federal drinking water standards and guidelines. U.S. Environmental Protection Agency. Chemical Communications Subcommittee, Federal State Toxicology and Regulatory Alliance Committee.
- *Fukui Y, Doskey PV.1996. An enclosure technique for measuring nonmethane organic compound emissions from grasslands. Journal of Environmental Quality. 25:601-610.
- *Cargas ML, Burgess RJ, Voisard DE, et al.1989. Partition coefficients of low-molecular-weight volatile chemicals in various liquids and tissues. Toxicol Appl Pharmacol 98:87-99.
- *Gentry SJ, Walsh PT.1987. Eight hour TWA personal monitoring using a diffusive sampler and short-term stain tube. Am Ind Hyg Assoc J 48(3):287-292.
- *Gillies PJ, Norton RM, Baker TS, et al.1981. Altered lipid metabolism in 2,5-hexanedione-induced testicular atrophy and peripheral neuropathy in the rat. Toxicol Appl Pharmacol 59:293-299.
- *Goodgold J, Eberstein A.1983. Motor and sensory nerve conduction measurements. In: Electrodiagnosis of neuromuscular diseases. 3rd ed. Baltimore, MD: Williams and Wilkins, 104-153.
- *Gori G, Bartolucci GB, Sturaro A, et al.1995. High performance liquid chromatographic determination of urinary 2,5- hexanedione as mono 2,4-dinitrophenylhydrazone using ultraviolet detection. J Chromatogr B 673:165-172.
- *Graham DG, Amarnath V, Valentine WM, et al.1995. Pathogenetic studies of hexane and carbon disulfide neurotoxicity. Crit Rev Toxicol 25(2):91-112.
- *Graham DG, Anthony DC, Boekelheide K, et al.1982. Studies of the molecular pathogenesis of hexane neuropathy. II. Evidence that pyrrole derivatization of lysyl residues leads to protein crosslinking. Toxicol Appl Pharmacol 64:415-422.
- *Guzelian PS, Henry CJ, Olin SS.1992. Similarities and differences between children and adults: Implications for risk assessment. International Life Sciences Institute Press, Washington, D.C.
- *Hakkola M, Saarinen L.1996. Exposure of tanker drivers to gasoline and some of its components. Ann Occup Hyg 40(1):1-10.
- *Hamilton, Bruce.1996. Gasoline frequently asked questions. INTERNET USENET, Archive Name: autos/gasoline-faq/part 1 (Document Version 1.09) <http://www.paranoia.com/~filipg/HTML>
- *Hautfenne A, Pocklington WD, Wolfe JP.1987. Determination of hexane residues in oils: Results of a collaborative study and the standardised method. Pure Appl Chem 59(11):1561-1570.

8. REFERENCES

- *HazDat.1998. Agency for Toxic Substances and Disease Registry (ATSDR). Atlanta, GA.
- *Heath JS, Koblis K, Sager SL, et al.1993. Risk assessment for total petroleum hydrocarbons. In: Kostecki PT, Calabrese EJ, eds. Hydrocarbon contaminated soils and groundwater. Vol. 3, 267-301. Chelsea, MI: Lewis Publishers.
- *Herbert R, Gerr F, Luo J, et al.1995. Peripheral neurological abnormalities among roofing workers: Sentinel case and clinical screening. Arch Environ Health 50(5):349-354.
- *Heringa JW, Huybregtse R, Van Der Linden AC.1961. *n*-Alkane oxidation by a Pseudomonas formation and B-oxidation of intermediate fatty acids. Antonie Van Leeuwenhoek 2751-58.
- *Hernandez JE, Machado LT, Corbella R, et al.1995. *n*-Alkanes and polynuclear aromatic hydrocarbons in fresh-frozen and pre-cooked-frozen mussels. Bull Environ Contam Toxicol 55:461-468.
- *Herskowitz A, Ishii N, Schaumburg H.1971. *n*-Hexane neuropathy: A syndrome occurring as a result of industrial exposure. N Engl J Med 285:82-85.
- *Hine CH, Zuidema HH.1970. The toxicological properties of hydrocarbon solvents. Ind Med 39:215-220.
- *Houk VS, Schaikowsky S, Claxton LD.1989. Development and validation of the spiral Salmonella assay: An automated approach to bacterial mutagenicity testing. Mutat Res 223:49-64.
- *Howd RA, Rebert CS, Dickinson J, et al.1983. A comparison of the rates of development of functional hexane neuropathy in weanling and young adult rats. Neurobehav Toxicol Teratol 5:63-68.
- *HSDB.1996. Hazardous Substances Database: *n*-Hexane. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.
- *Huang C, Shih T, Cheng S, et al.1991. *n*-Hexane polyneuropathy in a ball-manufacturing factory. J Occup Med 33(2):139-142.
- *Huang J, Kato K, Shibata E, et al.1989. Effects of chronic *n*-hexane exposure on nervous system-specific and muscle-specific proteins. Arch Toxicol 63(5):381-385.
- *Hunt JR, Sitar N, Udell KS.1988. Nonaqueous phase liquid transport and cleanup:1. Analysis of mechanisms. Water Resources Research 24(8):1247-1258.
- *IARC.1979b. International Agency for Research on Cancer (IARC). Monograph on the evaluation of the carcinogenic risk of chemicals to humans: some halogenated hydrocarbons. Vol. 20, Lyon France, 155-178.
- *IARC.1987. IARC monographs on the evaluation of carcinogenic risks to humans. Supplement 7, Overall evaluation of carcinogenicity: An updating of IARC monographs (Volumes 1 to 42) Lyon, France.
- Iida M.1982. Neurophysiological studies of *n*-hexane polyneuropathy in the sandal factory. Kyoto Symposia (Elsevier Biomedical Press Supplement) 36:671-81.

8. REFERENCES

- *Imbriani M, Ghittori S, Pezzagno G, et al.1984. *n*-Hexane urine elimination and weighted exposure concentration. *Int Arch Occup Environ Health* 55:33-41.
- *IRDC.1981. Six-month continuous inhalation exposure of rats to hexane mixtures-Phase II. Unpublished study by International Research and Development Corporation, Mattawan MI. Prepared for the American Petroleum Institute, Washington DC. EPA-FYI-AX-0282-0166.
- *IRIS.1998. Integrated Risk Information System, Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH. May 1998.
- *Ishidate M, Sofuni T, Yoshikawa K, et al.1984. Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 22(8):623-636.
- *Isidorov VA, Zenkevich IG, Ioffe BV.1985. Volatile organic compounds in the atmosphere of forest. *Atmos Environ* 19(1):1-8.
- *Iwata M, Takeuchi Y, Hisanaga N, et al.1983. A study on biological monitoring of *n*-hexane exposure. *Int Arch Occup Environ Health* 51:253-260.
- *Jackson JR, deKechoue G, Karakaya A.1997. Letter. *Hum Exp Toxicol* 16:284-285.
- *Johanson CE.1980. Permeability and vascularity of the developing brain: Cerebellum vs cerebral cortex. *Brain Research* 190:3-16.
- *Johnson BL, Boyd J, Burg JR, et al.1983. Effects on the peripheral nervous system of workers' exposure to carbon disulfide. *Neurotoxicology* 4(1):53-66.
- *Johnson DJ, Lack L, Ibrahim S, et al.1995. Protein-bound pyrroles in rat hair following subchronic intraperitoneal injections of 2,5-hexanedione. *J Toxicol Environ Health* 45:313-324.
- *Jorgensen NK, Chor KH.1981. *n*-Hexane and its toxicologic effects: A review. *Stand J Work Environ Health* 7:157-68.
- *Kanatharana P, Chindarasamme C, Kaewnarong B.1993. Extracting solvents for pesticide residues. *J Environ Sci Health A28(10):2323-2332*.
- *Karakaya A, Yucesoy B, Burgaz S, et al.1996. Some immunological parameters in workers occupationally exposed to *n*-hexane. *Hum Exp Toxicol* 15:56-58
- *Kawai T Mizunuma K, Yasugi T.1990. The method choice for the determination of 2,5-hexanedione as an indicator of occupational exposure to *n*-hexane. *Int Arch Occup Environ Health* 62:403-408.
- *Kawai T, Yasugi T, Mizunuma K.1992. Urinalysis vs. blood analysis as a tool for biological monitoring of solvent exposure. *Toxicol Letters* 63:333-343.
- *Kawai T, Yasugi T, Mizunuma K, et al.1993. Comparative evaluation of blood and urine analysis as a tool for biological monitoring of *n*-hexane and toluene. *Int Arch Occp Environ Health* 65:S123-S126.

8. REFERENCES

- *Kim IH, Yoon SH.1990. Effect of extraction solvents on oxidative stability of crude soybean oil. J Am Oil Chem Soc 67(3):165-167.
- *Kimura ET, Ebert DM, Dodge PW.1971. Acute toxicity and limits of solvent residue of sixteen organic solvents. Toxicol Appl Pharmacol 19:699-704.
- *Klein BL, Simon JE.1986. Hydrocarbon poisonings. Pediatric Clinics of North America 33:411-419.
- *Komori M, Nishio K, Kitada M et al.1990. Fetus-specific expression of a form of cytochrome P-450 in human livers. Biochemistry 29:4430-4433.
- *Kopczynski SL, Lonneman WA, Sutterfield FD, et al.1972. Photochemistry of atmospheric samples in Los Angeles. Environmental Science & Technology 6(4):342-347.
- *Kozloski RP.1985. Simple method for concentrating volatiles in water for gas chromatographic analysis by vacuum distillation. J Chromotgr 346:408-412.
- *Krasavage WJ, O'Donoghue JL, DiVincenzo GD, et al.1980. The relative neurotoxicity of methyl-*n*-butyl ketone, *n*-hexane and their metabolites. Toxicol Appl Pharmacol 52(3):433-441.
- *Krishnan K, Andersen ME.1994. Physiologically-based pharmacokinetic modeling in toxicology. In: Wallace Hayes, ed. Principles and methods of toxicology . 3rd edition. New York, NY: Raven Press Ltd.
- *Krishnan K, Andersen ME, Clewell HJ III, et al.1994. Physiologically-based pharmacokinetic modeling of chemical mixtures. In: RSA Yang, ed. Toxicology of chemical mixtures. New York, NY: Academic Press.
- *Krost KJ, Pellizzari ED, Walburn SG, et al.1982. Collection and analysis of hazardous organic emissions. Anal Chem 54:810-817.
- *Krotoszynski BK, Bruneau GM, OWeill HJ.1979. Measurement of chemical inhalation exposure in urban population in the presence of endogenous effluents. J Anal Toxicol 3(6):225-234.
- *Ladefoged O, Hass U, Simonsen L.1989. Neurophysiological and behavioural effects of combined exposure to 2,5-hexanedione and acetone or ethanol in rats. Pharmacol Toxicol 65:372-375.
- *Ladefoged O, Perbellini L.1986. Acetone-induced changes in the toxicokinetics of 2,5-hexanedione in rabbits. Stand J Work Environ Health 12:627-629.
- *Ladefoged O, Roswall K, Larsen J-J.1994. Acetone potentiation and influence on the reversibility of 2,5-hexanedione-induced neurotoxicity studied with behavioral and morphometric methods in rats. Pharmacol Toxicol 74:294-299.
- *Lam C-W, Galen TJ, Boyd JF, et al.1990. Mechanism of transport and distribution of organic solvents in blood. Toxicol Appl Pharmacol 104:117-129.
- *Lapadula DM, Habig C, Gupta RP, et al.1991. Induction of cytochrome P450 isozymes by simultaneous inhalation exposure of hens to *n*-hexane and methyl iso-butyl ketone (MiBK). Biochem Pharmacol 41(6-7):877-883.

8. REFERENCES

- *Lapadula DM, Irwin RD, Suwita E et al.1986. Cross-linking of neurofilament proteins of rat spinal cord *in vivo* after administration of 2,5-hexanedione. J Neurochem 46:1843-1850.
- *Latimer JS, Hoffman EJ, Hoffman G, et al.1990. Sources of petroleum hydrocarbons in urban runoff. Water Air and Soil Pollution 52:1-21.
- *Lawson H.1995. Food oils and fats: Technology, utilization, and nutrition. New York, NY: Chapman Jz Hall.
- *Leahy GL, Colwell RR.1990. Microbial degradation of hydrocarbons in the environment. Microbial Rev 54(3):305-315.
- *Leeder JS, Kearns GL.1997. Pharmacogenetics in pediatrics: Implications for practice. Pediatric Clinics of North America 44:55-77.
- *Leung H.1993. Physiologically-based pharmacokinetic modeling. In: Ballantine B, Marro T, Turner T, eds. General and applied toxicology, Vol. I. New York, NY: Stockton Press, 153-164.
- *Lide DR.1994. CRC handbook of chemistry and physics. 74th edition. Boca Raton, FL: CRC Press Inc.
- *Linder RE, Strader LF, Slott VL, et al.1992. Endpoints of spermatotoxicity in the rat after short duration exposures to fourteen reproductive toxicants. Reprod Toxicol 6(6):491-505.
- *Litton Bionetics.1979. Teratology study in rats *n*-hexane. Unpublished study by Litton Bionetics Inc., Kensington MD. Prepared for the American Petroleum Institute, Washington DC. EPA-FYI-AX-O 183-0231.
- *Litton Bionetics.1980. Mutagenicity evaluation of *n*-hexane in the mouse dominant lethal assay Final report. Unpublished study by Litton Bionetics Inc., Kensington MD. Prepared for the American Petroleum Institute, Washington DC. EPA-FYI-AX-01 83-0231.
- *Loden M.1986. The *in vitro* permeability of human skin to benzene, ethylene glycol, formaldehyde, and *n*-hexane. Acta Pharmacol Toxicol 58(5):382-389.
- *Lungarella G, Barni-Comparini I, Fonzi L.1984. Pulmonary changes induced in rabbits by long-term exposure to *n*-hexane. Arch Toxicol 55:224-228.
- *Lyman WJ, Reehl WF, Rosenblatt DH.1982. Handbook of chemical property estimation methods: Environmental behavior of organic compound. New York, NY: McGraw-Hill Book Company.
- *Marks TA, Fisher PW, Staples RE.1980. Influence of *n*-hexane on embryo and fetal development in mice. Drug Chem Toxicol 3(4):393-406.
- *Mast TA, Decker JR, Clark ML, et al.1987. Inhalation developmental toxicology studies: Teratology study of *n*-hexane in rats: Final report. Department of Energy, Washington DC. NTIS No.-DE88006812. PNL-6453.

8. REFERENCES

- *Mast TA, Hackett PL, Decker JR, et al.1989a. Inhalation reproductive toxicology studies: Sperm morphology study of *n*-hexane in B6C3F1 mice. Department of Energy, Washington DC. NTIS No.-DE89000262. PNL-6672.
- *Mast TA, Rommerein RL, Evanoff JJ, et al.1989b. Inhalation reproductive toxicology studies: Male dominant lethal study of *n*-hexane in Swiss (CD-1) mice. Department of Energy, Washington DC. NTIS No.-DE89000271. PNL-6679.
- *Mast TJ, Decker JR, Stoney KH, et al.1988. Inhalation developmental toxicology studies: Teratology study of *n*-hexane in mice. Department of Energy, NTIS No. DE88012680 (PNL-6590).
- *Mathews JM, Raymer JH, Etheridge As, et al.1997. Do endogenous volatile organic chemicals measured in breath reflect and maintain CYP2E1 levels *in vivo*? *Toxicol Appl Pharmacol* 146:255-260.
- *Mayer VW, Goin CJ.1994. Induction of chromosome loss in yeast by combined treatment with neurotoxic hexacarbons and monoketones. *Mutat Res* 341:83-91.
- *McCann MF.1992. Occupational and environmental hazards in art. *Environ Res* 59:139-144.
- *McCarroll NE, Keech BH, Piper CE.1981a. A microsuspension adaptation of the *Bacillus subtilis* "ret" assay. *Environ Mutagen* 3:607-616.
- *McCarroll NE, Piper CE, Keech BH. 1981 b. An *E. coli* microsuspension assay for the detection of DNA damage induced by direct-acting agents and promutagens. *Environ Mutagen* 3:429-444.
- *McClay K, Streger SH, Steffan RJ.1995. Induction of toluene oxidation activity in *Pseudomonas mendocina* KRI and *Pseudomonas sp.* Strain ENVPCS by chlorinated solvents and alkanes. *Appl Environ Microbiol* 61(9):3479-3481.
- *McKay WA, Turner MF, Jones BMR, et al.1996. Emission of hydrocarbons from marine phytoplankton - Some results from controlled laboratory experiments. *Atmos Environ* 30(14):2583-2593.
- *McLaren R, Singleton DL, Lai JYK, et al.1996. Analysis of motor vehicle sources and their contribution to ambient hydrocarbon distributions at urban sites in Toronto during the Southern Ontario oxidants study. *Atmos Environ* 30(12):2219-2232.
- *Merck.1989. Merck index: An encyclopedia of chemicals, drugs, and biologicals.11 th ed. Budavari S, ed. Rahway NJ: Merck & Co., Inc.
- *Michael LC, Erickson MD, Parks SP, et al.1980. Volatile environmental pollutants in biological matrices with a headspace purge technique. *Anal Chem* 52:1836-1841.
- *Michael LC, Pellizzari ED, Wiseman RW.1988. Development and evaluation of a procedure for determining volatile organics in water. *Environ Sci Technol* 22(5):565-570
- .
*Montgomery JH.1991. Groundwater chemicals desk reference. Volume 2. Ann Arbor: Lewis Publishers.
- *Morgenroth E, Schroeder ED, Chang DPY, et al.1996. Nutrient limitation in a compost biofilter degrading hexane. *J Air & Waste Management Association* 46:300-308.

8. REFERENCES

- *Morsehi PL, France-Morselli R, Bossi L.1980. Clinical pharmacokinetics in newborns and infants. *Clinical Pharmacokinetics* 5:485-527.
- *Mortelmans K, Haworth S, Lawlor T, et al.1986. Salmonella mutagenicity tests. II. Results from the testing of 270 chemicals. *Environ Mutagen* 8(Suppl 7):1-119.
- *Moschonas N, Glavas S.1996. C3-C 10 Hydrocarbons in the atmosphere of Athens, Greece. *Atmos Environ* 30(15):2769-7272.
- *Mutti A, Bergamaschi E, Ghittori S, et al.1993. On the need of a sampling strategy in biological monitoring: The example of hexane exposure. *Int Arch Occup Environ Health* 65(1):S172-S176.
- *Mutti A, Cavatorta A, Lommi G , et al.1982a. Neurophysiological effects of long-term exposure to hydrocarbon mixtures. *Arch Toxicol Suppl* 5:120-124.
- *Mutti A, Falzoi M, Lucertini S, et al.1984. *n*-Hexane metabolism in occupationally exposed workers. *Br J Ind Med* 41(4):533-538.
- *Mutti A, Ferri F, Lommi G, et al.1982b. *n*-Hexane-induced changes in nerve conduction velocities and somatosensory evoked potentials. *Int Arch Occup Environ Health* 51(1):45-54.
- *Nakajima T, Elovaara E, Park SS, et al.1991. Immunochemical detection of cytochrome P450 isozymes induced in rat liver by *n*-hexane, 2-hexanone and acetyl acetone. *Arch Toxicol* 65:542-547.
- *NAS/NRC.1980. Drinking water and health: Volume3. Washington, DC: National Academy Press.
- *NAS/NRC.1989. Biologic markers in reproductive toxicology. National Academy of Sciences/National Research Council. Washington, DC: National Academy Press, 15-35.
- *NAS/NRC.1989. Biological markers in reproductive toxicology. National Research Council. Board of Environmental Studies and Toxicology. Committee on Biological Markers, pp.15-35.
- *NATICH.1992. National Air Toxics Information Clearing House (NATICH) data base report of federal, state, and local air toxics activities. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. EPA-453\R-92-008.
- *Neeper-Bradley TL .1989a. Developmental toxicity study of commercial hexane vapor in CD(Sprague-Dawley) rats. Prepared by Bushy Run Research Center (Export, PA) for the American Petroleum Institute (API), Project Report 52-605, October 27, 1989.
- *Neeper-Bradley TL .1989b. Developmental toxicity study of commercial hexane vapor in CD-1 mice. Prepared by Bushy Run Research Center (Export, PA) for the American Petroleum Institute (API), Project Report 52-604, October 27, 1989.
- *Neligan RE.1962. Hydrocarbons in the Los Angeles atmosphere. *Arch Environ Health* 5:581-91.
- *Nelson KW, Ege JF Jr, Ross M, et al.1943. Sensory response to certain industrial solvent vapors. *J Ind Hyg Toxicol* 25:282-285.

8. REFERENCES

- *NFPA.1994. Fire protection guide to hazardous materials.11th edition. National Fire Protection Association. One Batterymarch Park, Quincy, MA 02269.
- *NIOSH.1992. NIOSH Recommendations for occupational safety and health--compendium of policy documents and statements. National Institute for Occupational Safety and Health. Department of Health and Human Services. Publication No. 92-100. Cincinnati, Ohio.
- *NIOSH.1995. NIOSH Report to Congress on workers' home contamination study conducted under the Workers' Family Protection Act. National Institute for Occupational Safety and Health. Department of Health and Human Services. Publication No. 96-192-000. Cincinnati, Ohio.
- *NOES.1991. *n*-Hexane: Numbers of potentially exposed employees from the National Occupational Exposure Database. Cincinnati, OH: National Institute for Occupational Safety.
- *NRC.1993. Pesticides in the diets of infants and children. National Research Council, Washington DC: National Academy Press.
- *NTP 1991. Toxicity studies of *n*-hexane in B6C3F1 (inhalation studies) National Toxicology Program. Research Triangle Park, NC. U.S. Department of Health and Human Services. Public Health Service. National Institutes of Health. Publication No. 91-3121.
- *Nylen P, Ebendal T, Eriksson Nilsson M, et al.1989. Testicular atrophy and loss of nerve growth factor-immunoreactive germ cell line in rats exposed to *n*-hexane and a protective effect of simultaneous exposure to toluene or xylene. Arch Toxicol 63:296-307.
- *Nylen P, Hagman M.1994. Function of the auditory and visual systems, and of peripheral nerve, in rats after long-term combined exposure to *n*-hexane and methylated benzene derivatives. II. Xylene. Pharmacol Toxicol 74(2):124-9.
- *Nylen P, Hagman M, Johnson AC.1994. Function of the auditory and visual systems, and of peripheral nerve, in rats after long-term combined exposure to *n*-hexane and methylated benzene derivatives. I. Toluene. Pharmacol Toxicol 74(2):116-23.
- *O'Leary P, Walsh P.1995. Solid waste landfills. Madison, WI: University of Wisconsin at Madison.
- *Oliver KD, Adams JR, Daughtrey EH, et al.1996. Technique for monitoring ozone precursor hydrocarbons in air at photochemical assessment monitoring stations: Sorbent preconcentration, closed-cycle cooler cryofocusing, and GC-FID analysis. Atmos Environ 30(15):2751-2757.
- *Ong CN, Koh D, Foo SC, et al.1993. Volatile organic solvents in correction fluids: Identification and potential hazards. Bull Environ Contam Toxicol 50:787-793.
- *Ono Y, Takeuchi Y, Hisanaga N.1981. A comparative study on the toxicity of *n*-hexane and its isomers on the peripheral nerve. Int Arch Occup Environ Health 48:289-294.
- *OSHA.1974. U.S. Occupational Safety and Health Administration. Code of Federal Regulations 29 CFR 1910.1000.
- *OTA.1990. Neurotoxicology: Identifying and controlling poisons of the nervous system. Office of Technology Assessment, Washington, DC. OTA-BA-438.

8. REFERENCES

- *Otson R, Fellin P, Tran Q.1994. VOCs in representative Canadian residences. *Atmos Environ* 28(22):3563-3569.
- *Owen GM, Brozek J.1966. Influence of age, sex, and nutrition on body composition during childhood and adolescence. In: Falkner F, ed. *Human development*. Philadelphia, PA: Saunders, 222-238.
- *Pastore C, Marhuenda D, Marti J, et al.1994. Early diagnosis of *n*-hexane-caused neuropathy. *Muscle Nerve* 17(9):981-6.
- *Patten CJ, Ning SM, Lu AYH, et al.1986. Acetone-inducible cytochrome P-450: Purification, catalytic activity, and interaction with cytochrome b5. *Arch Biochem Biophys* 251:629-638.
- *Pellizzari ED, Hartwell TD, Harris BSH, et al.1982. Purgeable organic compounds in mother's milk. *Bull Environ Contam Toxicol* 28:322-328.
- *PennWell.1994. *Energy statistics sourcebook*. 9th edition. Tulsa, OK: PennWell Publishing Company.
- *Perbellini L, Brugnone F, Caretta D, et al.1985. Partition coefficients of some industrial aliphatic hydrocarbons (C5-C7) in blood and human tissues. *Br J Ind Med*. 42(3):162-167.
- *Perbellini L, Brugnone F, Faggionato G.1981. Urinary excretion of the metabolites of *n*-hexane and its isomers during occupational exposure. *Br J Ind Med*. 38(1):20-26.
- *Perbellini L, Brugnone F, Pavan I.1980. Identification of the metabolites of *n*-hexane, cyclohexane, and their isomers in men's urine. *Toxicol Appl Pharmacol* 53:220-229.
- *Perbellini L, Marhuenda Amoros D, Cardona Llorens A, et al.1990a. An improved method of analysing 2,5-hexanedione in urine. *Brit J Ind Med* 47:421-424.
- *Perbellini L, Mozzo P, Brugnone F, et al.1986. Physiologicomathematical model for studying human exposure to organic solvents: Kinetics of blood/tissue *n*-hexane concentrations and of 2,5-hexanedione in urine. *Brit J Ind Med* 43(11):760-768.
- *Perbellini L, Mozzo P, Olivato D, et al.1990b. Dynamic biological exposure indexes for *n*-hexane and 2,5-hexanedione, suggested by a physiologically based pharmacokinetic model. *Am Ind Hyg Assoc J* 51(7):356-362.
- *Perbellini L, Pezzoli G, Brugnone F, et al.1993. Biochemical and physiological aspects of 2,5-hexanedione: Endogenous or exogenous product? *Int Arch Occup Environ Health* 65:49-52.
- *Periago FJ, Luna A, Morente A, et al.1991. Design and evaluation of an exhaled breath sampler for biological monitoring of organic solvents. *J Appl Toxicol* 12(2):91-96.
- *Periago JF, Cardona A, Marhuenda D, et al.1993. Biological monitoring of occupational exposure to *n*-hexane by exhaled air analysis and urinalysis. *Int Arch Occup Environ Health* 65:275-278.
- *Perocco P, Bolognesi S, Alberghini W.1983. Toxic activity of seventeen industrial solvents and halogenated compounds on human lymphocytes cultured *in vitro*. *Toxicol Lett* 16:69-75.

8. REFERENCES

- *Pezzoli G, Antonini A, Barbieri S, et al.1995. *n*-Hexane-induced Parkinsonism: Pathogenetic hypothesis. *Movement Disorders* 10:279-282.
- *Potter TL.1993. Analysis of petroleum contaminated soil and water: An overview. In: Calabrese EJ, Kostecki PT, eds. *Principles and practices for petroleum contaminated soils*. Ann Arbor, MI: Lewis Publishers, 1-14.
- *Pryor GT, Rebert CS.1992. Interactive effects of toluene and hexane on behavior and neurophysiologic responses in Fischer-344 rats. *Neurotoxicology* 13:225-234.
- *Pyle SJ, Amarnath V, Graham DG et al.1993. Decreased levels of the high molecular weight subunit of neurofilaments and accelerated neurofilament transport during the recovery phase of 2,5-hexanedione exposure. *Cell Motil Cytoskeleton* 26(2):133-143.
- *Rastogi SC.1993. Organic solvent levels in model and hobby glues. *Bull Environ Contam Toxicol* 51:501-507.
- *Raymer JH, Pellizzari ED.1996. Estimation of recent exposures to volatile organic chemicals using alveolar air measurements. *Toxicol Ind Health* 12:201-20.
- *Raymer JH, Thomas KW, Pellizzari ED, et al.1994. A breath sampling device for measuring human exposure to volatile organic compounds in microgravity. *Aviation, Space and Environmental Medicine* 353-360.
- *Rebert C, Sorenson.1983. Concentration-related effects of *n*-hexane on evoked responses from brain and peripheral nerve of the rat. *Neurobehav Toxicol Teratol* 5:69-76.
- *Roberts MT, Etherington D.1996. *Bookbinding and the conservation of books: A dictionary of descriptive terminology*. Menlo Park, CA: Electronic Edition, Stanford University
<http://palimnset.stanford.edu/don/don.html>
- *Roberts NJ, Burton HR.1994. Volatile compounds in Meromictic Antarctic lakes and basins *Chemosphere* 29(8):1627-1637.
- *Robertson P, White EL, Bus JS.1989. Effects of methyl ethyl ketone pretreatment on hepatic mixed-function oxidase activity and on *in vivo* metabolism of *n*-hexane. *Xenobiotica* 19(7):721-729.
- *Rosenberg E, Legmann R, Kushmaro A, et al.1992. Petroleum bioremediation: A multiphase problem. *Biodegradation* 3:337-350.
- *RTECS.1997. *Registry of Toxic Effects of Chemical Substances (RTECS)*. National Institute for Occupational Safety and Health (NIOSH). Computer database online.
- *Sack TM, Steele DH, Hammerstrom K, et al.1992. A survey of household products for volatile organic compounds. *Atmos Environ* 26A:1063-1070.
- *Sanagi S, Seki Y, Sugimoto K, et al.1980. Peripheral nervous system functions of workers exposed to *n*-hexane at a low level. *Int Arch Occup Environ Health* 47(1):69-79.

8. REFERENCES

- *Sauer TC, Brown JS, Boehm PD, et al. 1993. Hydrocarbon source identification and weathering characterization of intertidal and subtidal sediments along the Saudi Arabian coast after the Gulf War Oil Spill. *Marine Pollution Bulletin* 27:117-134.
- *Schaumburg HH, Spencer PS. 1976. Degeneration in central and peripheral nervous systems produced by pure *n*-hexane: An experimental study. *Brain* 99:183-192.
- *Schuberth J. 1994. Joint use of retention index and mass spectrum in postmortem test for volatile organics by headspace capillary gas chromatography with ion-trap detection. *J Chromatogr A* 674:63-71.
- *Seppalainen AM, Raitta C, Huuskonen MS. 1979. *n*-Hexane-induced changes in visual evoked potentials and electroretinograms of industrial workers. *Electroencephal Clin Neurophysiol* 47:492-498.
- *Setchell BP, Waites GMH. 1975. The blood testis barrier. In: Creep RO, Astwood EB, eds., Geiger SR, executive ed. *Handbook of physiology: Endocrinology V* (Chapter 6). Washington DC: American Physiological Society.
- *Seto Y. 1994. Determination of volatile substances in biological samples by headspace gas chromatography. *J Chromatogr A* 674:25-62.
- *Shaw DG, Hogan TE, McIntosh DJ. 1986. Hydrocarbons in the sediments of Port Valdez, Alaska: Consequences of five years' permitted discharge. *Estuarine, Coastal and Shelf Science* 21:131-144.
- *Sieber WK, Sundin DS, Frazier TM, et al. 1991. Development, use, and availability of a job exposure matrix based on national occupational hazard survey data. *Amer J Ind Med* 20:163-174.
- *Sittig M. 1994. *World-wide limits for toxic and hazardous chemicals in air, water, and soil*. Park Ridge, NJ. Noyes Publications.
- *Speight JG. 1991. *Classification. The chemistry and technology of petroleum*. Second edition, 197-527.
- *Spencer PS, Schaumburg HH. 1977a. Ultrastructural studies of the dying-back process. III. The evolution of experimental peripheral giant axonal degeneration. *J Neuropathol Exp Neurol* 36(2):276-299.
- *Spencer PS, Schaumburg HH. 1977b. Ultrastructural studies of the dying-back process. IV. Differential vulnerability of PNS and CNS fibers in experimental central-peripheral distal axonopathies. *J Neuropathol Exp Neurol* 36(2):300-320.
- *Spencer PS, Schaumburg HH, Sabri MI, et al. 1980. The enlarging view of hexacarbon neurotoxicity. *CRC Crit Rev Toxicol* 17(4):279-356.
- *SRC. 1994a. Syracuse Research Center. Henry's Law Constant Program (HENRYWIN, version 2,50 serial H0142). Chemical Hazard Assessment Division, Environmental Chemistry Center, Syracuse, NY.
- *SRC. 1994b. Fate and transport models: Model outputs for the chemical *n*-hexane. Syracuse, NY: Syracuse Research Corporation.

8. REFERENCES

- *SRC.1995. Syracuse Research Center. Octanol-Water Partition Coefficient Program (KOWWIN, version 137, serial L0148). Chemical Hazard Assessment Division, Environmental Chemistry Center, Syracuse, NY.
- *SRI.1988.1988 Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- *SRI.1990.1990 Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- *SRI.1992.1992 Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- *SRI.1994.1994 Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- *SRI.1995.1995 Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- *SRI.1996.1996 Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- *Srinivas H, Swamylingappa B, Chand N.1992. Secondary extraction of soybeans using hexane-acetic acid: Effect on beany flavor removal and physicochemical properties. *J Agric Food Chem* 40:276-279.
- *St. Clair MB, Amarnath V, Moody MA, et al.1988. Pyrrole oxidation and protein cross-linking as necessary steps in the development of gamma-diketone neuropathy. *Chem Res Toxicol* 1:179-185.
- *Stelljes ME, Watkin GE.1993. Comparison of environmental impacts posed by different hydrocarbon mixtures: A need for site-specific composition analyses. In: Kostecki PT, Calabrese EJ, eds. *Hydrocarbon contaminated soils and groundwater*, Vol. 3. Chelsea, MI: Lewis Publishers, 549-569.
- *Stevens P.1988. *Oil and gas dictionary*. New York, NY: Nichols Publishing.
- *Stoltenburg-Didinger G, Altenkirch H, Wagner M.1990. Neurotoxicity of organic solvent mixtures: Embryotoxicity and fetotoxicity. *Neurotoxicol Teratol* 12:585-589.
- *SUCRON.1996. *The SUCRON wood and cane to gasoline and hydrogen process*. Clinton, LA: SUCRON, Inc.
- *Sundell J.1996. What we know, and don't know, about sick building syndrome. *ASHRAE Journal* 38(6):51-57.
- *Swarm RL, Laskowski DA, McCall PJ, et al.1983. A rapid method for the estimation of the environmental parameters octanol/water partition coefficient, soil sorption constant, water to air ratio, and water solubility. *Residue Reviews* 85:1.
- *Takeuchi Y, Hisanaga N, Ono Y, et al.1993. Modification of metabolism and neurotoxicity of hexane by co-exposure of toluene. *Int Arch of Occup Environ Health* 65(1):S227-S230.

8. REFERENCES

- *Takeuchi Y, Ono Y, Hisanaga N, et al.1980. A comparative study on the neurotoxicity of *n*-pentane, *n*-hexane, and *n*-heptane in the rat. *Brit J Ind Med* 37:241-247.
- *Tang Y-Z, Cheng WK, Fellin P, et al.1996. Laboratory evaluation of sampling methods for C 1 to C4 hydrocarbons. *Amer Ind Hyg Assoc J* 57:245-250.
- *Thomas KW, Pellizzari ED, Cooper SD.199 I. Canister based method for collection and GC/MS analysis of volatile organic compounds in human breath. *J Anal Toxicol* 15:54-59.
- *Toftgard R, Haaparanta T, Eng L, et al.1986. Rat lung and liver microsomal cytochrome P-450 isozymes involved in the hydroxylation of *n*-hexane. *Biochem Pharmacol* 35(21):3733-3738.
- *Toftgard R, Halpert J, Gustafsson JA.1983. Xylene induces a cytochrome P-450 isozyme in rat liver similar to the major isozyme induced by phenobarbital. *Mol Pharmacol* 23(1):265-271.
- *TRI96.1998. Toxic Chemical Release Inventory. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.
- *U.S. Congress.1986. Superfund amendments and reauthorization act of 1986. Title III--Emergency Planning and Community Right-to-Know. Ninety-ninth Congress of the United States of America.
- *U.S. Congress.1990. Clean air amendments. Title III, Hazardous Air Pollutants, Section 112(b), Hazardous Air Pollutants as Amended, October 26, 1990. One Hundred and First Congress of the United States of America, 2nd Session Report 101-952.
- *USITC.1994. Synthetic organic chemical: United States production and sales, 1992. USITC Publication 2720, February, 1994. Washington, DC: United States International Trade Commission.
- *van Engelen JG, Kezie S, de Haan W.1995. Determination of 2,5 hexanedione a metabolite of *n*-hexane, in urine evaluation and application of three analytical methods. *J Chromatogr B* 667:233-240.
- *van Engelen JGM, Rebel-deHaan W, Opdam JJG, et al.1997. Effect of coexposure to methyl ethyl ketone (MEK) on *n*-hexane in human volunteers. *Toxicol Appl Pharmacol* 144:385-395
- *Van Ert MD, Arp EW, Harris RL.1980. Worker exposures to chemical agents in the manufacture of rubber tires: Solvent vapor studies. *Am Ind Hyg Assoc J* 41(3):212-219.
- *Verma DK, Shaw DS, McLean JD.1992. An evaluation of airborne concentrations of selected hydrocarbons at an oil refinery land reclamation operation. *Ann Occup Hyg* 36(3):307-315.
- *Veulemans H, Groeseneken D, Masschelein R, et al.1987. Survey of ethylene glycol ether exposure in Belgian industries and workshops. *Am Ind Hyg Assoc J* 48(8):671-676.
- *Veulemans H, Van Vlem E, Janssens H, et al.1982. Experimental human exposure to *n*-hexane. Study of the respiratory uptake and elimination, and of *n*-hexane concentrations in peripheral venous blood. *Int Arch Occup Environ Health*. 49(3-4):251-263.

8. REFERENCES

- *Vieira I, Sonnier M, Cresteil T.1996. Developmental expression of CYP2E1 in the human liver: hypermethylation control of gene expression during the neonatal period. *European Journal of Biochemistry* 238:476-483.
- *Wadden RA, Scheff PA, Franke JE, et al.1995. VOC emission rates and emission factors for a sheetfed offset printing shop. *Am Ind Hygiene Assoc J* 56:368-376.
- *Wahlberg JE.1984. Erythema-inducing effects of solvents following epicutaneous administration to man-studied by laser doppler flowmetry. *Stand J Work Environ Health* 10:159-162.
- *Wahlberg JE, Boman A.1979. Comparative percutaneous toxicity of ten industrial solvents in the guinea pig. *Stand J Work Environ & Health* 5:345-351.
- *Wanasundara JPD, Shahidi F.1994. Alkanol-ammonia-water/hexane extraction of flaxseed. *Food Chem* 49:39-44.
- *Wang JD, Chang YC, Kao KP, et al.1986. An outbreak of *n*-hexane induced polyneuropathy among press proofing workers in Taipei. *Am J Ind Med* 10(2):111-118.
- *Watts JO, Holswade W.1967. Gas chromatographic determination of residual hydrocarbon solvents in solvent-extracted edible oils. *J AOAC* 50(3):718-726.
- *West JR, Smith HW, Chasis H.1948. Glomerular filtration rate, effective renal blood flow, and maximal tubular excretory capacity in infancy. *J of Pediatrics* 32a:10-18.
- *White EL, Bus JS, Heck H d'A.1979. Simultaneous determination of *n*-hexane, 2-hexanone and 2,5-hexanedione in biological tissues by gas chromatography mass spectrometry. *Biomedical Mass Spectrometry* 6(4):169-172.
- *WHO.1984. Guidelines for drinking-water quality: Volume I: Recommendations. Geneva, Switzerland: World Health Organization, 203-228.
- * WHO.1991. *n*-Hexane. Environmental health criteria 122. World Health Organization. Geneva, Switzerland.
- *Widdowson EM, Dickerson JWT.1964. Chapter 17: Chemical composition of the body. In: Comar CL and Bronner F, eds. *Mineral metabolism: An advanced treatise volume II - the elements part A*. New York, NY: Academic Press.
- *Wirier AM, Arey J, Atkinson R, et al.1992. Emission rates of organics from vegetation in California's Central valley. *Atmos Environ* 26A(14):2647-2659.
- *Wang P SH, Cooks RG, Cisper ME, et al.1995. *On-line in situ* analysis with membrane introduction MS. *Environ Sci Technol* 29(5):215A-218A.
- *Xu YH, Mitra S.1994. Continuous monitoring of volatile organic compounds in water using *on-line* membrane extraction and microtrap gas chromatography system. *J Chromatogr* 688:171-180.

8. REFERENCES

- *Yamamura Y.1969. *n*-Hexane polyneuropathy. *Folia Psychiatrica et Neurologica Japonica* 23(1):45-57.
- *Yang Yu, Hawthorne B, Miller DJ.1995. Comparison of sorbent and solvent trapping after supercritical fluid extraction of volatile petroleum hydrocarbons from soil. *J Chromatogr A* 699:265-276.
- *Zhao WY, Misumi J, Yasui T, et al.1998. Effects of methyl ethyl ketone, acetone, or toluene coadministration on 2,5-hexanedione concentration in the sciatic nerve, serum, and urine of rats. *Int Arch Occup Environ Health* 71:236-244.
- *Zhu M, Spink DC, Yan B, et al.1995. Inhibition of 2,5-hexanedione-induced protein cross-linking by biological thiols: Chemical mechanisms and toxicological implications. *Chem Res Toxicol* 8:764-771.
- *Ziegler EE, Edwards BB, Jensen RL et al.1978. Absorption and retention of lead by infants. *Pediatr Res* 12:29-34.
- *Zielinska B, Sagebiel JC, Harshfield G, et al.1996. Volatile organic compounds up to C20 emitted from motor vehicles: Measurement methods. *Atmos Environ* 30(12):2269-2286.

