

8. REFERENCES

- Abdel-Rahman MS, Hetland LB, Couri D. 1976. Toxicity and metabolism of methyl n-butyl ketone. *Am Ind Hyg Assoc J* 37:95-102.
- *Abdel-Rahman MS, Saladin JJ, Bohman CE, et al. 1978. The effect of 2-hexanone and 2-hexanone metabolites on pupillomotor activity and growth. *Am Ind Hyg Assoc J* 39:94-99.
- *Abdo KM, Graham DG, Timmons PR, et al. 1982. Neurotoxicity of continuous (90 days) inhalation of technical grade methyl butyl ketone in hens. *J Toxicol Environ Health* 9:199-215.
- Abe K, Misumi J, Kawakami M, et al. 1980. Effects of n-hexane, methyl n-butyl ketone, and 2,5-hexanedione on the excitability of sweat glands in rats to mecholy. *Jap J Ind Health* 22:380-381.
- Abou-Donia MB. 1983. Interaction between neurotoxicities induced by organophosphorus and long-chain hexacarbon compounds. *Neurotoxicity* 4:117-135.
- Abou-Donia MB, Nomeir AA. 1986. The role of pharmacokinetic and metabolism in species sensitivity to neurotoxic agents. *Fundam Appl Toxicol* 6:190-207.
- *Abou-Donia MB, Makkawy H-AM, Graham DB. 1982. The relative neurotoxicities of n-hexane, methyl n-butyl ketone, 2,5-hexanediol, and 2,5-hexanedione following oral or intraperitoneal administration in hens. *Toxicol Appl Pharmacol* 62:369-389.
- *Abou-Donia MB, Lapadula DM, Campbell G, et al. 1985a. The joint neurotoxic action of inhaled methyl butyl ketone vapor and dermally applied 0-ethyl 0-4-nitrophenyl phenylphosphonothioate in hens: Potentiating effect. *Toxicol Appl Pharmacol* 79:69-82.
- *Abou-Donia MB, Makkawy H-AM, Campbell GM. 1985b. Pattern of neurotoxicity of n-hexane, methyl n-butyl ketone, 2,5-hexanediol, and 2,5-hexanedione alone and in combination with 0-ethyl 0-4-nitrophenyl phenylphosphonothioate in hens. *J Toxicol Environ Health* 16:85-100.
- *Abou-Donia MB, Lapadula DM, Campbell G, et al. 1985c. The synergism of n-hexane-induced neurotoxicity by methyl isobutyl ketone following subchronic (90 days) inhalation in hens: Induction of hepatic microsomal cytochrome P-450. *Toxicol Appl Pharmacol* 81:1-16.

* Cited in text

8. REFERENCES

- *ACGIH. 1986. Documentation of the threshold limit values and biological exposure indices. 5th ed. American Conference of Governmental Industrial Hygienists. Cincinnati, OH.
- Allen N, Mendell JR, Billmaier D, et al. 1974. An outbreak of a previously undescribed toxic polyneuropathy due to industrial solvent. Transactions of the American Neurological Association 99:74-79.
- *Allen N, Mendell JR, Billmaier DJ, et al. 1975. Toxic polyneuropathy due to methyl n-butyl ketone: An industrial outbreak. Arch Neurol 32:209-218.
- Altenkirch H, Mager J, Stoltenburg G, et al. 1977. Toxic polyneuropathies after sniffing a glue thinner. J Neurol 214:137-152.
- *Ambrose D, Ellender JH, Lees EB, et al. 1975. Thermodynamic properties of organic oxygen compounds. XXXVIII. Vapour pressures of some aliphatic ketones. J Chem Thermodynamics 7:453-472.
- *Amoore JE, Hautala E. 1983. Odor as an aid to chemical safety: Odor thresholds compared with threshold limit values and volatilities for 214 industrial chemicals in air and water dilution. J Appl Toxicol 3:272-290.
- *Anderson RA, Harland WA. 1980. The analysis of volatiles in blood from fire fatalities. In: Proceedings of the Forensic Toxicology European Meeting, Int Assoc Toxicol, 279-292.
- Anger WK, Lynch DW. 1977. The effect of methyl n-butyl ketone on response rates of rats performing on a multiple schedule of reinforcement. Environ Res 14:204-211.
- Anonymous. 1974. Solvent causes motor neuropathy in workers at clothing factory. J Am Med Assoc 229:247-248.
- Anonymous. 1979. Hexacarbon neuropathy. The Lancet 2(November 3):942-943.
- Anthony DC, Boekelheide K, Anderson CW, et al. 1983. The effect of 3,4-dimethyl substitution on the neurotoxicity of 2,5-hexanedione. II. Dimethyl substitution accelerates pyrrole formation and protein crosslinking. [Abstract] Toxicol Appl Pharmacol 71:372-382.
- Asbury AK. 1979. Pathology of industrial toxic neuropathies. Acta Neurol Stand Suppl 60:52-53.
- ASTM. 1987a. Standard practice for sampling atmospheres to collect organic compound vapors (activated charcoal tube adsorption method) - method D 3686-84. In: 1987 annual book of ASTM standards. Vol. 11.03. Atmospheric analysis; occupational health and safety. Philadelphia, PA: American Society for Testing and Materials, 326-336.

8. REFERENCES

ASTM. 1987b. Standard practice for analysis of organic compound vapors collected by the activated charcoal tube adsorption method - method D 3687-84. In: 1987 annual book of ASTM standards. Vol. 11.03. Atmospheric analysis; occupational health and safety. Philadelphia, PA: American Society for Testing and Materials, 337-344.

ASTM. 1988. Standard practice for measuring volatile organic matter in water by aqueous-injection gas chromatography - method D 2908-87. In: 1988 annual book of ASTM standards. Vol. 11.02. Water II. Philadelphia, PA: American Society for Testing and Materials, 46-51.

*Atkinson R, Carter WP, Aschmann SM, et al. 1985. Atmospheric fates of organic chemicals: Prediction of ozone and hydroxyl radical reaction rates and mechanisms. Report to U.S. Environmental Protection Agency, Office of Research and Development, Research Triangle Park, NC, by Statewide Air Pollution Research Center, University of California, Riverside, CA. EPA/600/3-85/063. NTIS Publication No. PB85-241529.

*Babeu L, Vaishnav DD. 1987. Prediction of biodegradability for selected organic chemicals. J Ind Microbiol 2:107-115.

*Badings HT, De Jong C, Dooper RP. 1985. Automatic system for rapid analysis of volatile compounds by purge-and-cold trapping/capillary gas chromatography. J High Resolut Chromatogr Commun 8:755-763.

Baker EL Jr. 1983. Neurological disorders. In: Rom WN et al., eds. Environmental and Occupational Medicine. Boston, MA: Little, Brown and Company, 313-327.

Baker EL, Fine LJ. 1986. Solvent neurotoxicity: The current evidence. J Occup Med 28:126-129.

*Barnes D, Bellin J, DeRosa C, et al. 1988. Reference dose (RfD): Description and use in health risk assessments. Vol. I. Appendix A: Integrated risk information system supportive documentation. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. EPA/600/8-88/032a.

Bierbaum PJ. 1973. Survey of Columbus Coated Fabrics, Newark, California. Cincinnati, OH: National Institute for Occupational Safety and Health, Division of Field Studies and Clinical Investigations. NTIS No. PB81-229601.

*Bierbaum PJ, Marceleno T. 1973. Survey of Brammer Manufacturing Company, Davenport, Iowa. Cincinnati, OH: National Institute for Occupational Safety and Health, Division of Field Studies and Clinical Investigations. NTIS No. PB81-231318.

8. REFERENCES

- Bierbaum PJ, Parnes WD. 1974. Survey of Electrical Division of Bristol Brass Corporation, South Windsor, Connecticut. Cincinnati, OH: National Institute for Occupational Safety and Health, Division of Field Studies and Clinical Investigations. NTIS No. PB-82-110073.
- *Billmaier D, Yee HT, Allen N, et al. 1974. Peripheral neuropathy in a coated fabrics plant. *J Occup Med* 16:665-671.
- *Boekelheide K. 1987. 2,5-Hexanedione alters microtubule assembly. I. Testicular atrophy, not nervous system toxicity, correlates with enhanced tubulin polymerization. *Toxicol Appl Pharmacol* 88:370-382.
- *Branchflower RV, Pohl IX. 1981. Investigation of the mechanism of the potentiation of chloroform-induced hepatotoxicity and nephrotoxicity by methyl *n*-butyl ketone. *Toxicol Appl Pharmacol* 61:407-413.
- Branchflower RV, Schulick RD, George JW, et al. 1983. Comparison of the effects of methyl-*n*-butyl ketone and phenobarbital on rat liver cytochromes P-450 and the metabolism of chloroform to phosgene. *Toxicol Appl Pharmacol* 71:414-421.
- *Bronstein AC, Currance PL. 1988. Emergency care for hazardous materials exposure. St. Louis, MO: The C.V. Mosby Company, 65, 215-216.
- *Brown EM, Hewitt WR. 1984. Dose-response relationships in ketone-induced potentiation of chloroform hepato- and nephrotoxicity. *Toxicol Appl Pharmacol* 76:437-453.
- *Brown KW, Donnelly KC. 1988. An estimation of the risk associated with the organic constituents of hazardous and municipal waste landfill leachates. *Hazardous Wastes and Hazardous Materials* 5:1-30.
- Browning E. 1965. Toxicity and metabolism of industrial solvents. London, England: Elsevier Publishing Company, 428-429.
- *Calvert JG, Pitts JN. 1966. Photochemistry of the polyatomic molecules. In: *Photochemistry*. New York, NY: John Wiley & Sons, Inc.
- *CLPSD. 1989, Contract Laboratory Program Statistical Database. Vitar and Company, Management Services Division, Alexandria, VA. June 1986.
- Couri D, Milks M. 1982. Toxicity and metabolism of the neurotoxic hexacarbonyl *n*-hexane, 2-hexanone, and 2,5-hexanedione. *Ann Rev Pharmacol Toxicol* 22:145-166.

8. REFERENCES

- Couri D, Hetland LB, O'Neill JJ, et al. 1974. Comments on a plastics industry neurotoxicity in relationship to methylbutyl ketone. Wright-Patterson Air Force Base, OH: Aerospace Medical Research Laboratory. NTIS No. ADA-011857.
- *Couri D, Hetland LB, Abdel-Rahman MS, et al. 1977. The influence of inhaled ketone solvent vapors on hepatic microsomal biotransformation activities. *Toxicol Appl Pharmacol* 41:285-289.
- Couri D, Abdel-Rahman MS, Hetland LB. 1978. Biotransformation of n-hexane and methyl n-butyl ketone in guinea pigs and mice. *Am Ind Hyg Assoc J* 39:295-300.
- *Cowlen MS, Hewitt WR, Schroeder F. 1984a. 2-Hexanone potentiation of [¹⁴C]chloroform hepatotoxicity: Covalent interaction of a reactive intermediate with rat liver phospholipid. *Toxicol Appl Pharmacol* 73:478-491.
- *Cowlen MS, Hewitt WR, Schroeder F. 1984b. Mechanisms in 2-hexanone potentiation of chloroform hepatotoxicity. *Toxicol Lett* 22:293-299.
- Craft BF. 1973. Summary status report of an investigation of an incident of toxic polyneuropathy at the Borden Chemical Company's Columbus Coated Fabrics Division, Columbus, Ohio. Cincinnati, OH: National Institute for Occupational Safety and Health. NTIS No. PB89-122915.
- Davenport JG, Farrell DF, Sumi SM. 1976. Giant axonal neuropathy caused by industrial chemicals: Neurofilamentous axonal masses in man. *Neurology* 26:919-923.
- *Day EA, Anderson DF. 1965. Gas chromatographic and mass spectral identification of natural components of the aroma fraction of blue cheese. *J Agric Food Chem* 13:2-4.
- *DeCaprio AP, Olajos EJ, Weber P. 1982. Covalent binding of a neurotoxic n-hexane metabolite: Conversion of primary amines to substituted pyrrole adducts by 2,5-hexanedione. *Toxicol Appl Pharmacol* 65:440-450.
- *DeCaprio AP, Briggs RG, Jackowski SJ, et al. 1988. Comparative neurotoxicity and pyrrole-forming potential of 2,5-hexanedione and perdeuterio-2,5-hexanedione in the rat. *Toxicol Appl Pharmacol* 92:75-85.
- DeJesus C, Pleasure DF, Asbury AK, et al. 1977. Effects of methyl butyl ketone on peripheral nerves and its mechanism of action. Report to National Institute of Occupational Safety and Health, Cincinnati, OH, by Yale University of Medicine, New Haven, CT and Veterans Administration Hospital, West Haven, CT. NTIS No. PB83-105148.

8. REFERENCES

- *DiVincenzo GD, Kaplan CJ, Dedinas J. 1976. Characterization of the metabolites of methyl n-butyl ketone, methyl iso-butyl ketone, and methyl ethyl ketone in guinea pig serum and their clearance. *Toxicol Appl Pharmacol* 36:511-522.
- *DiVincenzo GD, Hamilton ML, Kaplan CJ, et al. 1977. Metabolic fate and disposition of ¹⁴C-labeled methyl n-butyl ketone in the rat. *Toxicol Appl Pharmacol* 41:547-560.
- *DiVincenzo GD, Hamilton ML, Kaplan CJ, et al. 1978. Studies on the respiratory uptake and excretion and the skin absorption of methyl n-butyl ketone in humans and dogs. *Toxicol Appl Pharmacol* 44:593-604.
- DiVincenzo GD, Hamilton ML, Kaplan CJ, et al. 1980. Characterization of the metabolites of methyl n-butyl ketone. In: Spencer PS, Schaumburg HH, eds. *Experimental and clinical neurotoxicology*. Baltimore, MD: Williams and Wilkins Company, 846-855.
- Duckett S, Williams N, Francis S. 1974. Peripheral neuropathy associated with inhalation of methyl-n-butyl ketone. *Experientia* 30:1283-1284.
- *Duckett S, Streletz IJ, Chambers RA, et al. 1979. 50 ppm MnBK subclinical neuropathy in rats. *Experientia* 35:1365-1367.
- Dumdei BE, Henderson R. 1988. Air toxics monitoring program during bioremediation of a Superfund site. Proceedings of the 81st Annual Meeting of Air Pollution Control Association, Dallas, TX. June, 1988.
- *Dumont JP, Adda J. 1978. Occurrence of sesquiterpenes in mountain cheese volatiles. *J Agric Food Chem* 26:364-367.
- Durham HD, Pena SD J, Ecobichon DJ. 1988. Hexahydrocarbon effects on intermediate filament organization in human fibroblasts. *Muscle Nerve* 11:160-165.
- *Eben A, Flucke W, Mihail F, et al. 1979. Toxicological and metabolic studies of methyl n-butylketone, 2,5-hexanedione, and 2,5-hexanediol in male rats. *Ecotoxicol Environ Safety* 3:204-217.
- *Egan G, Spencer P, Schaumburg H, et al. 1980. n-Hexane-"free" hexane mixture fails to produce nervous system damage. *Neurotoxicology* 1:515-524.
- *Ellenhorn MJ, Barceloux DG. 1988. *Medical toxicology: Diagnosis and treatment of human poisoning*. New York, NY: Elsevier, 999-1000.
- *EPA. 1981. Chemical hazard information profile: Draft report methyl n-butyl ketone. Washington, DC: U.S. Environmental Protection Agency, Office of Pesticides and Toxic Substances.

8. REFERENCES

- *EPA. 1986. Gas chromatography/mass spectrometry for volatile organics - method 8240. In: Test methods for evaluating solid waste. 3rd ed. SW-846. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.
- *EPA. 1987a. U.S. Environmental Protection Agency: Part II. Federal Register 52:25942-25953.
- *EPA. 1987b. U.S. Environmental Protection Agency. Federal Register 52:11823-11826.
- *EPA. 1989. Interim methods for development of inhalation reference doses. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. EPA 600/8-88/066F.
- *Fedtke N, Bolt HM. 1986. Methodological investigations on the determination of E-hexane metabolites in urine. Int Arch Occup Environ Health 57:149-158.
- Garman JR, Freund T, Lawless EW. 1987. Testing for groundwater contamination at hazardous waste sites. J Chromatogr Sci 25:328-337.
- Genter MB, Szakal-Quin G, Anderson W, et al. 1986. Evidence that pyrrole formation is a pathogenetic step in gamma-diketone neuropathy. [Abstract] Toxicol Appl Pharmacol 87:351-362.
- Goldberg AM. 1980. Mechanisms of neurotoxicity as studied in tissue culture systems. Toxicology 17:201-208.
- Goldstein MN. 1981. Effect of methyl-n-butyl ketone and other ketone homologues on mammalian cells in culture. Report to National Institute of Occupational Health and Safety, Cincinnati, OH, by Washington University, St. Louis, MO. NTIS No. PB81-225997.
- Gosselin RE, Smith RP, Hodge HC, et al. 1984. Clinical toxicology of commercial products. 5th ed. Baltimore, MD: Williams and Wilkins, 11-185-11-186.
- Graedel TE. 1978. Aliphatic ketones. In: Chemical compounds in the atmosphere. New York, NY: Academic Press, 181, 184-185.
- *Green DR, Le Pape D. 1987. Stability of hydrocarbon samples on solid-phase extraction columns. Anal Chem 59:699-703.
- *Grey TC, Shrimpton DH. 1967. Volatile components of raw chicken breast muscle. Br Poult Sci 8:23-33.

8. REFERENCES

- *Hassett JJ, Banwart WL, Griffin RA. 1983. Correlation of compound properties with sorption characteristics of nonpolar compounds by soils and sediments: Concepts and limitations. In: Francis CW, Auerbach SI, Jacobs VA, eds. Environment and solid wastes: Characterization, treatment, and disposal. Boston, MA: Butterworths, 161-176.
- *Hawthorne SB, Sievers RE, Barkley RM. 1985. Organic emissions from shale oil wastewaters and their implications for air quality. Environ Sci Technol 19:992-997.
- *Hewitt WR, Miyajima H, Cote MG, et al. 1980a. Acute alteration of chloroform-induced hepato- and nephrotoxicity by n-hexane, methyl n-butyl ketone and 2,5-hexanedione. Toxicol Appl Pharmacol 53:230-248.
- *Hewitt WR, Miyajima H, Cote MG, et al. 1980b. Modification of haloalkane-induced hepatotoxicity by exogenous ketones and metabolic ketosis. Fed Proc 39:3118-3123.
- *Hewitt WR, Brown EM, Plaa GL. 1983. Relationship between the carbon skeleton length of ketonic solvents and potentiation of chloroform-induced hepatotoxicity in rats. Toxicol Lett 16:297-304.
- *Hewitt LA, Ayotte P, Plaa GL. 1986. Modifications in rat hepatobiliary function following treatment with acetone, 2-butanone, 2-hexanone, mirex, or chlordecone and subsequently exposed to chloroform. Toxicol Appl Pharmacol 83:465-473.
- *HSDB. 1989. Hazardous Substances Data Bank. National Library of Medicine, National Toxicology Information Program, Bethesda, MD. September 5, 1989.
- IRPTC. 1989. International Register of Potentially Toxic Chemicals. United Nations Environment Programme, Geneva, Switzerland. September 1989.
- *Johnson BL, Setzer JV, Lewis TR, et al. 1977. Effects of methyl n-butyl ketone on behavior and the nervous system. Am Ind Hyg Assoc J 38:567-579.
- Johnson BL, Setzer JV, Lewis TR, et al. 1978. An electrodiagnostic study of the neurotoxicity of methyl n-amyl ketone. Am Ind Hyg Assoc J 39:866-872.
- Johnson BL, Anger WK, Setzer JV, et al. 1979. Neurobehavioral effects of methyl n-butyl ketone and methyl n-amyl ketone in rats and monkeys: A summary of NIOSH investigations. J Environ Pathol Toxicol 2:113-133.
- *Katz GV, O'Donoghue JL, DiVincenzo GD, et al. 1980. Comparative neurotoxicity and metabolism of ethyl n-butyl ketone and methyl n-butyl ketone in rats. Toxicol Appl Pharmacol 52:153-158.

8. REFERENCES

- *King JW. 1989. Fundamentals and applications of supercritical fluid extraction in chromatographic science. *J Chromatogr Sci* 27:355-364.
- *Kinlin TE, Muralidhara R, Pittet AO, et al. 1972. Volatile components of roasted filberts. *J Agr Food Chem* 20:1021-1028.
- Konasewich D, Traversy W, Zar H. 1978. Status report on organic and heavy metal contaminants in the Lakes Erie, Michigan, Huron and Superior basins. Great Lakes Water Quality Board.
- *Krasavage WJ, O'Donoghue JL, DiVincenzo GD, et al. 1980. The relative neurotoxicity of methyl-n-butyl ketone, n-hexane and their metabolites. *Toxicol Appl Pharmacol* 52:433-441.
- *Laity JL, Burstain IG, Appel BR. 1973. Photochemical smog and the atmospheric reactions of solvents. In: Tess RW, ed. *Solvents theory and practice*. Advances in Chemistry Series 124, Washington, DC: American Chemical Society, 95-112.
- *Lande SS, Durkin PR, Christopher DE, et al. 1976. Investigation of selected potential environmental contaminants: Ketonic solvents. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA-560/2-76-003. NTIS No. TR 76-500.
- *Lapin EP, Weissbarth S, Maker HS, et al. 1982. The sensitivities of creatine and adenylate kinases to the neurotoxins acrylamide and methyl n-butyl ketone. *Environ Res* 28:21-31.
- Leo A, Hansch C, Elkins D. 1971. Partition coefficients and their uses. *Chem Rev* 71:525,563.
- Lloyd AC. 1979. Tropospheric chemistry of aldehydes. Washington, DC: National Bureau of Standards. Special Publication No. 557.
- *Lowery CE Jr., Foster JW, Jurtschuk P. 1968. The growth of various filamentous fungi and yeasts on n-alkanes and ketones: I. Studies on substrate specificity. *Arch Mikrobiol* 60:246-254.
- *Lucas SV. 1984, GC/MS analysis of organics in drinking water concentrates and advanced waste treatment concentrates. Vol. 1. Analysis results for 17 drinking water, 16 advanced waste treatment and 3 process blank concentrates. Report to U.S. Environmental Protection Agency, Office of Research and Development, Research Triangle Park, NC, Battelle Columbus Laboratories, Columbus, OH. EPA-600/1-84-020a. NTIS No. PB85-128221.
- *Lukins HB, Foster JW. 1963. Methyl ketone metabolism in hydrocarbonutilizing mycobacteria. *J Bacterial* 85:1074-1087.

8. REFERENCES

- *Mabey WR, Smith JH, Pod011 RT, et al. 1982. Aquatic fate process data for organic priority pollutants. Report to U.S. Environmental Protection Agency, Office of Water Regulations and Standards, Washington, DC, by SRI International, Menlo Park, CA. EPA 440/4-81-014. NTIS No. PB87-169090.
- *MacLeod H, Jourdain JL, Poulet G, et al. 1984. Kinetic study of reactions of some organic sulfur compounds with OH radicals. Atmos Environ 18:2621-2626.
- Makkawy HM, Graham DG, Abou-Donia MB. 1981. Differential neuro toxicity of n - hexane methyl-E-butyl ketone, 2,5-hexanediol and 2,5-hexanedione following subchronic 90 days oral administration in hens. Fed Proc 40:678.
- Mallov JS. 1976. MBK neuropathy among spray painters. J Am Med Assoc 235:1455-1457.
- Mallov JS, Marceleno T, Fontaine R. 1975. Methyl n-butyl ketone and peripheral neuropathy: A summary report. Cincinnati, OH: U.S. Department of Health, Education, and Welfare, Division of Field Studies and Clinical Investigations. NTIS No. PB81-229981.
- *Marceleno T, Bierbaum PJ, Mallov JS. 1974. Survey of Premium Finishes, Incorporated, Cincinnati, OH. Cincinnati, OH: National Institute for Occupational Safety and Health, Division of Field Studies and Clinical Investigations. NTIS No. PB81-242729.
- McDonough JR. 1974. Possible neuropathy from methyl butyl ketone. N Engl J Med 290:695.
- Mendell JR, Duckett S, Williams N, et al. 1974a. Neuropathy and methyl n-butyl ketone. N Engl J Med 290:1263-1264.
- *Mendell JR, Saida K, Ganansia MF, et al. 1974b. Toxic polyneuropathy produced by methyl n-butyl ketone. Science 185:787-789.
- Mendell JR, Sahenk Z, Saida K, et al. 1977. Alterations of fast axoplasmic transport in experimental methyl n-butyl ketone neuropathy. Brain Res 133:107-118.
- Michael LC, Pellizzari ED, Wiseman RW. 1988. Development and evaluation of a procedure for determining volatile organics in water. Environ Sci Technol 22:565-570.
- Misumi J, Nagano M. 1984. Neurophysiological studies on the relation between the structural properties and neurotoxicity of aliphatic hydrocarbon compounds in rats. Br J Ind Med 41:526-532.

8. REFERENCES

Misumi J, Nagano M. 1985. Experimental study on the enhancement of the neurotoxicity of methyl n-butyl ketone by non-neurotoxic aliphatic monoketones. *Br J Ind Med* 42:155-161.

*Morrison RT, Boyd RN. 1974. *Organic chemistry*. 3rd ed. Boston, MA: Allyn and Bacon, Inc., 620.

*Myers VB. 1983. Remedial activities at the Miami Drum site, Florida. In: National conference on management of uncontrolled hazardous waste sites. Silver Springs, MD: Hazardous Materials Control Research Institute, 354-357.

Nachtman JP, Couri D. 1984. An electrophysiological study of 2-hexanone and 2,5-hexanedione neurotoxicity in rats. *Toxicol Lett* 23:141-145.

*NAS/NRC. 1989. *Biologic markers in reproductive toxicology*. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press.

*NATICH. 1989. National Air Toxics Information Clearinghouse: NATICH database report on state, local and EPA air toxics activities. Report to U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards, Research Triangle Park, NC, by Radian Corporation, Austin, TX. EPA-450/3-89-29.

*Neely WB, Branson DR, Blau GE. 1974. Partition coefficient to measure bioconcentration potential of organic chemicals in fish. *Environ Sci Technol* 8:1113-1115.

Nelson BK. 1986. Developmental neurotoxicology of in utero exposure to industrial solvents in experimental animals. *Neurotoxicology* 7:441-448.

NIOSH. 1978a. Criteria for a recommended standard: Occupational exposure to ketones. Cincinnati, OH: U.S. Department of Health, Education and Welfare, National Institute for Occupational Safety and Health. DHEW (NIOSH) Publication No. 78-173.

*NIOSH. 1978b. Criteria for a recommended standard: Occupational exposure in coal gasification plants. Cincinnati, OH: National Institute for Occupational Safety and Health. DHEW (NIOSH) Pub No. 78-191.

*NIOSH. 1984. Ketones I - method 1300. In: NIOSH manual of analytical methods. 3rd ed. Vol. 2. Cincinnati, OH: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health.

*NIOSH. 1985. Pocket guide to chemical hazards. Washington, DC: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health. DHHS (NIOSH) Publication No. 85-114.

8. REFERENCES

NIOSH. 1987. Organic solvent neurotoxicity. Cincinnati, OH: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health. Current Intelligence Bulletin 48. DHHS (NIOSH) Publication No. 87-104.

*NIOSH. 1989. National Occupational Exposure Survey. National Institute of Occupational Safety and Health, Cincinnati, OH. October 18, 1989.

*NLM, 1989. Chemline. National Library of Medicine, Bethesda, MD. September 1989.

*NOHS. 1989. National Occupational Hazard Survey. National Institute of Occupational Safety and Health, Cincinnati, OH. October 18, 1989.

*Nomeir AA, Abou-Donia MB. 1985. Analysis of D-hexane, 2-hexanone, 2,5-hexanedione, and related chemicals by capillary gas chromatography and high performance liquid chromatography. Anal Biochem 151:381-388.

*O'Donoghue, JL. 1985. Alkanes, alcohols, ketones, and ethylene oxide. In: O'Donoghue JL, ed. Neurotoxicity of industrial and commercial chemicals. Vol. II. Boca Raton, FL: CRC Press, Inc. 61-97.

O'Donoghue JL, Krasavage WJ, DiVincenzo GD, et al. 1984. Further studies on ketone neurotoxicity and interactions. Toxicol Appl Pharmacol 72:201-209.

O'Donoghue JL, Haworth SR, Curren RD, et al. 1988. Mutagenicity studies on ketone solvents: Methyl ethyl ketone, methyl isobutyl ketone, and isophorone. Mutat Res 206:149-161.

*OSHA. 1989. Occupational Safety and Health Administration: Part III. Federal Register 54:2940.

*Pankow JF, Rosen ME. 1988. Determination of volatile compounds in water by purging directly to a capillary column with whole column cryotrapping. Environ Sci Technol 22:398-405.

*Pellizzari ED, Castillo NP, Willis S, et al. 1979. Identification of organic components in aqueous effluents from energy-related processes. In: Van Hall CE, ed. Measurement of organic pollutants in water and wastewater. Philadelphia, PA: American Society for Testing and Materials, 256-274. ASTM Special Technical Publication No. 686.

*Perry JJ. 1968. Substrate specificity in hydrocarbon utilizing microorganisms. Antonie Van Leeuwenhoek 34:27-36.

8. REFERENCES

- *Perry DL, Chuang CC, Jungclaus GA, et al. 1979. Identification of organic compounds in industrial effluent discharges. Athens, GA: U.S. Environmental Protection Agency, Office of Research and Development, Environmental Research Laboratory. EPA-600/4-79-016. PB-294794.
- *Peters MA, Hudson PM, Dixon RL. 1981. The effect of gestational exposure to methyl n-butyl ketone has on postnatal development and behavior. *Ecotoxicol Environ Safety* 5:291-306.
- Pilon D, Charbonneau M, Brodeur J, et al. 1986. Metabolites and ketone body production following methyl n-butyl ketone exposure as possible indices of MnBK potentiation of carbon tetrachloride hepatotoxicity. *Toxicol Appl Pharmacol* 85:49-59.
- Politis MJ, Pellegrino RG, Spencer PS. 1980. Ultrastructural studies of the dying-back process. 5. Axonal neurofilaments accumulate at sites of 2,5-hexanedione application: Evidence for nerve fibre dysfunction in experimental hexacarbon neuropathy. [Abstract] *J Neurocytol* 9:505-516.
- Prockop L, Couri D. 1977. Nervous system damage from mixed organic solvents. In: Sharp CW, ed. *Review of inhalants: Euphoria to dysfunction*. Washington, DC: U.S. Government Printing Office, 185-198.
- Proctor NH, Hughes JP, Fischman ML. 1988. *Chemical hazards of the workplace*. 2nd edition. Philadelphia, PA: J. B. Lippincott Company, 324-325.
- Puskar MA, Levine SP, Lowry SR. 1987. Qualitative screening of hazardous waste mixtures. *Environ Sci Technol* 21:90-96.
- Raisbeck MF. 1984. Effects of 2-hexanone upon hepato- and nephrotoxicity of several haloalkanes and cephaloridine. *Diss Abstr Int B* 46:1083.
- Raisbeck MF, Brown EM, Hewitt WR. 1986. Renal and hepatic interactions between 2-hexanone and carbon tetrachloride in F-344 rats. *Toxicol Lett* 31:15-21.
- Raleigh RL, Spencer PS, Schaumburg HH. 1975. Toxicity of methyl- n-butyl ketone. *Arch Environ Health* 30:317-318.
- Roy WR, Griffin RA. 1985. Mobility of organic solvents in water-saturated soil materials. *Environ Geol Water Sci* 7:241-247.
- *Sabri MI. 1984. In vitro effect of n-hexane and its metabolites on selected enzymes in glycolysis, pentose phosphate pathway and citric acid cycle. *Brain Res* 297:145-150.

8. REFERENCES

- Sabri MI, Moore CL, Spencer PS. 1979a. Studies on the biochemical basis of distal axonopathies-I. Inhibition of glycolysis by neurotoxic hexacarbon compounds. *J Neurochem* 32:683-689.
- *Sabri MI, Ederle K, Holdsworth CE, et al. 1979b. Studies on the biochemical basis of distal axonopathies. II. Specific inhibition of fructose-6-phosphate kinase by 2,5-hexanedione and methyl-butyl ketone. *Neurotoxicology* 1:285-297.
- *Saida K, Mendell JR, Weiss HS. 1976. Peripheral nerve changes induced by methyl E-butyl ketone and potentiation by methyl ethyl ketone. *Neuropathol Exp Neurol* 35:207-225.
- *Sate A, Nakajima T. 1979. Partition coefficients of some aromatic hydrocarbons and ketones in water, blood and oil. *Br J Ind Med* 36:231-234.
- *Sax NI. 1984. Dangerous properties of industrial materials. 6th ed. New York: Van Nostrand Reinhold Company, 1526-1527.
- *Sax NI, Lewis RJ Sr. 1987. Hawley's condensed chemical dictionary. 11th ed. New York: Van Nostrand Reinhold Company, 762.
- Sayre LM, Shearson CM, Wongmongkolrit T, et al. 1986. Structural basis of gamma-diketone neurotoxicity: Non-neurotoxicity of 3,3-dimethyl-2,5-hexanedione, a gamma-diketone incapable of pyrrole formation. [Abstract] *Toxicol Appl Pharmacol* 84:36-44.
- Schrenk HH, Yant WP, Patty FA. 1936. Acute response of guinea pigs to vapors of some new commercial organic compounds. *Public Health Rep* 51:624-631.
- Selkoe DJ, Luckenbill-Edds L, Shelanski ML. 1978. Effects of neurotoxic industrial solvents on cultured neuroblastoma cells: Methyl n-butyl ketone, n-hexane and derivatives. *J Neuropathol Exp Neurol* 37:768-789.
- *Shackelford WM, Keith LH. 1976. Frequency of organic compounds identified in water. Athens, GA: U.S. Environmental Protection Agency, Office of Research and Development. EPA-600/4-76-062. NTIS No. PB-165470.
- Sharkawi M, Elfassy B. 1985. Inhibition of mouse liver alcohol dehydrogenase by methyl n-butyl ketone. *Toxicol Lett* 25:185-189.
- *Smyth HF, Carpenter CP, Weil CS, et al. 1954. Range-finding toxicity data: List V. *Arch Ind Hyg Occup Med* 10:61-68.
- Spector WS, ed. 1956. Lethal doses of solid and liquid compounds: Laboratory animals. In: Spector WS, ed. *Handbook of Toxicology*. Vol. I. Philadelphia, PA: W.B. Saunders and Company 5, 156-157, 321, 338-339.

8. REFERENCES

Spencer PS, Schaumburg HH. 1976. Feline nervous system response to chronic intoxication with commercial grades of methyl n-butyl ketone, methyl isobutyl ketone, and methyl ethyl ketone. *Toxicol Appl Pharmacol* 37:301-311.

Spencer PS, Schaumburg HH. 1977a. Ultrastructural studies of the dying-back process. III. The evolution of experimental peripheral giant axonal degeneration. *J Neuropathol Exp Neurol* 36:276-299.

*Spencer PS, Schaumburg HH. 1977b. Ultrastructural studies of the dying-back process. IV. Differential vulnerability of PNS and CNS fibers in experimental central-peripheral distal axonopathies. *J Neuropathol Exp Neurol* 36:300-320.

*Spencer PS, Schaumburg HH, Raleigh RL, et al. 1975. Nervous system degeneration produced by the industrial solvent methyl n-butyl ketone. *Arch Neurol* 32:219-222.

*Stutz DR, Janusz SJ. 1988. Hazardous materials injuries: A handbook for pre-hospital care. 2nd ed. Beltsville, MD: Bradford Communications Corporation, 312-313.

*Takeoka GR, Flath RA, Guntert M, et al. 1988. Nectarine volatiles: Vacuum steam distillation versus headspace sampling. *J Agric Food Chem* 36:553-560.

Thomas RD, ed. 1986. Drinking water and health. Vol. 6. Washington, DC: National Academy Press, 130.

Thorn P, Gardener H, Hrutfiord B. 1977. Odorous compounds from magnesite pulping. *Pulp and Paper Canada* 78:93-95.

TRI. 1989. Toxic Chemical Release Inventory. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.

*Upreti RK, Shanker S. 1987. 2,5-Hexanedione-induced immunomodulatory effect in mice. *Environ Res* 43:48-59.

*Upreti RK, Singh KP, Saxena AK, et al. 1986. Effect of 2,5-hexanedione on lymphoid organs of rats: A preliminary report. *Environ Res* 39:188-198.

Vaishnav DD, Boethling RS, Babeu L. 1987. Quantitative structure-biodegradability relationships for alcohols, ketones and alicyclic compounds. *Chemosphere* 16:695-703.

Veronesi B, Peterson ER, Bornstein MB, et al. 1983. Ultrastructural studies of the dying-back process. VI. Examination of nerve fibers undergoing giant axonal degeneration in organotypic culture. *J Neuropathol Exp Neurol* 42:153-165. [Abstract]

8. REFERENCES

*Verschueren K. 1983. Handbook of environmental data on organic chemicals. 2nd ed. New York: Van Nostrand Reinhold Company, 320.

*View Database. 1989. Agency for Toxic Substances and Disease Registry (ATSDR), Office of External Affairs, Exposure and Disease Registry Branch, Atlanta GA. September 25, 1989.

*Weast RC, ed. 1985. CRC Handbook of chemistry and physics. Boca Raton, FL: CRC Press, Inc., C-307.

*West C. 1990. Written communication (March 5) to Barry L. Johnson, Agency for Toxic Substances and Disease Registry, regarding regulatory information of hazardous substances. Department of Environmental Quality Engineering, The Commonwealth of Massachusetts, Boston, MA.

*White EL, Bus JS, Heck HD. 1979. Simultaneous determination of n-hexane, 2-hexanone and 2,5-hexanedione in biological tissues by gas chromatography mass spectrometry. Biomed Mass Spectrum 6:169-172.

*Wieboldt RC, Adams GE, Later DW. 1988. Sensitivity improvement in infrared detection for supercritical fluid chromatography. Anal Chem 60:2422-2427.

*Windholz M, ed. 1983. The Merck index: An encyclopedia of chemicals, drugs, and biologicals. 10th ed. Rahway, NJ: Merck and Company, Inc., 5909.