

Table 4-1. Youth reporting ever having used marijuana, by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting marijuana use ever													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13	6.1	(4.7,7.9)	5.3	(3.9,7.3)	5.7	(4.5,7.1)	5.4	(4.1,7.2)	5.0	(3.2,7.8)	-1.1	(-3.8,1.6)	-0.7	(-3.6,2.2)
14 to 15	15.4	(12.4,18.8)	19.5	(16.6,22.7)	18.8	(16.4,21.5)	18.5	(16.7,20.5)	17.3	(14.9,20.0)	1.9	(-2.4,6.3)	-1.6	(-5.1,1.9)
16 to 18	40.4	(36.2,44.7)	40.3	(36.7,43.9)	41.4	(38.2,44.6)	37.5	(35.2,39.9)	37.6	(34.2,41.1)	-2.8	(-8.4,2.7)	-3.8	(-8.6,1.1)
14 to 16	20.6	(18.2,23.3)	22.4	(19.7,25.5)	24.1	(22.1,26.1)	22.3	(20.5,24.3)	21.8	(19.6,24.1)	1.1	(-2.3,4.6)	-2.3	(-5.1,0.5)
14 to 18	29.1	(26.6,31.7)	30.9	(28.1,33.8)	31.8	(29.7,34.0)	29.4	(27.8,31.2)	29.3	(27.0,31.7)	0.2	(-3.3,3.7)	-2.5	(-5.6,0.6)
12.5 to 18	23.6	(21.7,25.7)	24.8	(22.7,27.1)	25.5	(23.9,27.2)	23.7	(22.3,25.1)	23.5	(21.7,25.5)	-0.1	(-2.9,2.8)	-2.0	(-4.5,0.5)
Youth aged 12.5 to 13														
Gender														
Males	6.8	(5.0,9.1)	6.2	(4.2,9.2)	4.6	(3.2,6.5)	5.1	(3.4,7.5)	4.4	(2.8,7.0)	-2.3	(-5.3,0.6)	-0.2	(-2.9,2.6)
Females	5.4	(3.7,7.8)	4.3	(2.8,6.7)	6.7	(5.0,9.1)	5.8	(3.9,8.5)	5.6	(3.0,10.2)	0.2	(-3.9,4.3)	-1.2	(-5.7,3.3)
Race/ethnicity														
White	5.1	(3.5,7.4)	4.4	(3.0,6.6)	5.2	(3.7,7.3)	4.9	(3.6,6.8)	3.5	(2.4,5.2)	-1.6	(-3.8,0.5)	-1.7	(-3.9,0.5)
African American	6.8	(3.3,13.7)	3.5	(1.5,8.0)	4.8	(2.3,9.6)	5.4	(2.4,11.6)	S	(S)	S	(S)	S	(S)
Hispanic	8.5	(4.6,15.0)	10.9	(5.7,19.7)	6.3	(3.8,10.2)	7.3	(4.0,13.0)	4.0	(1.5,10.3)	-4.4	(-11.8,2.9)	-2.3	(-7.1,2.5)
Risk score														
Higher risk	36.6	(27.4,47.0)	24.8	(14.9,38.3)	34.0	(25.1,44.2)	30.2	(20.8,41.6)	41.0	(28.7,54.5)	4.4	(-10.8,19.5)	7.0	(-8.9,22.8)
Lower risk	2.6	(1.6,4.2)	2.7	(1.8,4.3)	2.0	(1.2,3.3)	2.2	(1.2,3.7)	2.1	(0.9,4.8)	-0.5	(-2.5,1.5)	0.1	(-1.9,2.1)
Sensation seeking														
High	11.4	(8.6,15.1)	9.6	(6.5,14.1)	10.0	(7.6,13.1)	10.0	(7.3,13.6)	11.2	(7.0,17.2)	-0.3	(-6.2,5.6)	1.1	(-5.1,7.4)
Low	2.0	(1.2,3.6)	2.1	(1.2,3.6)	2.4	(1.3,4.2)	1.6	(0.9,2.8)	0.7	(0.2,2.1)	-1.4	(-2.8,0.0)	-1.7	*(-3.3,-0.1)

Table 4-1. Youth reporting ever having used marijuana, by age, gender, race/ethnicity, risk score, and sensation seeking (continued)

Characteristics	Percent reporting marijuana use ever													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 14 to 18														
Gender														
Males	30.8	(26.8,35.2)	32.7	(28.8,36.9)	32.1	(29.7,34.7)	29.4	(27.2,31.7)	29.1	(26.1,32.2)	-1.8	(-7.4,3.9)	-3.1	(-6.8,0.7)
Females	27.2	(24.0,30.7)	28.9	(25.4,32.7)	31.4	(28.1,34.9)	29.5	(27.0,32.2)	29.5	(26.1,33.1)	2.2	(-2.9,7.4)	-1.9	(-6.8,2.9)
Race/ethnicity														
White	30.3	(27.2,33.6)	32.2	(28.7,35.9)	32.9	(30.6,35.4)	30.2	(28.1,32.3)	29.5	(27.0,32.1)	-0.8	(-4.7,3.1)	-3.5	*(-6.7,-0.3)
African American	25.0	(19.6,31.3)	24.6	(18.7,31.6)	26.3	(21.9,31.3)	25.0	(20.9,29.7)	26.3	(20.6,32.8)	1.3	(-6.7,9.3)	0.0	(-7.1,7.0)
Hispanic	28.5	(22.0,36.1)	31.8	(24.0,40.9)	33.6	(27.6,40.2)	29.9	(24.5,35.9)	33.8	(27.4,40.8)	5.3	(-2.7,13.2)	0.2	(-6.1,6.5)
Risk score														
Higher risk	49.3	(45.5,53.0)	52.5	(48.4,56.5)	51.8	(48.7,54.9)	49.0	(46.6,51.4)	49.1	(45.8,52.3)	-0.2	(-5.5,5.1)	-2.7	(-7.3,1.8)
Lower risk	5.3	(3.8,7.4)	7.2	(5.6,9.3)	8.7	(6.7,11.1)	8.8	(7.5,10.4)	9.6	(7.5,12.4)	4.3	*(1.5,7.1)	1.0	(-2.4,4.4)
Sensation seeking														
High	39.1	(35.7,42.6)	38.6	(34.9,42.5)	41.0	(38.4,43.6)	36.9	(34.6,39.3)	39.1	(36.2,42.0)	0.0	(-4.3,4.3)	-1.9	(-5.4,1.6)
Low	14.7	(11.5,18.7)	18.0	(15.1,21.4)	18.1	(15.2,21.5)	18.5	(16.5,20.7)	15.6	(12.6,19.1)	0.9	(-3.9,5.7)	-2.6	(-7.1,2.0)

¹The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-2. Youth reporting using marijuana in the past year, by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting marijuana use in the past year													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13	3.9	(2.9,5.3)	3.5	(2.2,5.4)	3.9	(3.0,5.1)	4.2	(3.0,6.1)	3.2	(2.0,5.1)	-0.7	(-2.7,1.2)	-0.7	(-2.8,1.4)
14 to 15	11.6	(9.0,14.7)	13.8	(11.6,16.5)	12.6	(10.6,14.9)	13.6	(12.0,15.3)	13.3	(11.1,15.8)	1.8	(-2.2,5.8)	0.7	(-2.3,3.8)
16 to 18	29.2	(25.7,33.0)	27.1	(24.0,30.4)	29.1	(26.4,32.0)	27.5	(25.2,29.9)	26.2	(23.1,29.4)	-3.0	(-7.7,1.6)	-3.0	(-7.0,1.1)
14 to 16	15.4	(13.2,17.9)	15.8	(13.7,18.3)	17.2	(15.5,19.1)	16.6	(15.0,18.4)	16.0	(14.0,18.2)	0.5	(-2.6,3.6)	-1.3	(-3.6,1.0)
14 to 18	21.2	(19.2,23.4)	21.1	(18.8,23.5)	22.1	(20.2,24.1)	21.6	(20.0,23.2)	20.9	(18.9,23.1)	-0.3	(-3.2,2.6)	-1.2	(-3.8,1.5)
12.5 to 18	17.1	(15.5,18.9)	16.9	(15.2,18.8)	17.7	(16.3,19.3)	17.4	(16.2,18.7)	16.7	(15.1,18.4)	-0.4	(-2.6,1.9)	-1.0	(-3.1,1.1)
Youth aged 12.5 to 13														
Gender														
Males	4.1	(2.9,5.8)	3.5	(1.9,6.4)	2.8	(1.8,4.3)	3.9	(2.4,6.2)	3.4	(1.9,6.0)	-0.7	(-3.1,1.7)	0.6	(-1.7,3.0)
Females	3.7	(2.4,5.8)	3.5	(2.1,5.8)	5.1	(3.6,7.0)	4.6	(2.9,7.4)	2.9	(1.4,6.2)	-0.8	(-3.6,2.0)	-2.1	(-5.2,1.0)
Race/ethnicity														
White	2.6	(1.5,4.4)	3.1	(1.8,5.2)	3.6	(2.4,5.3)	3.8	(2.6,5.6)	2.3	(1.3,4.2)	-0.3	(-2.3,1.7)	-1.3	(-3.3,0.8)
African American	5.7	(2.6,12.0)	1.4	(0.3,5.9)	2.5	(1.1,5.7)	4.9	(2.1,10.9)	4.2	(1.7,10.1)	-1.5	(-7.0,4.0)	1.7	(-2.8,6.1)
Hispanic	6.1	(3.3,11.2)	7.0	(2.7,16.8)	4.7	(2.5,8.7)	5.6	(2.5,11.8)	4.0	(1.5,10.3)	-2.1	(-7.2,3.0)	-0.7	(-5.9,4.6)
Risk score														
Higher risk	24.6	(17.5,33.4)	19.0	(10.8,31.4)	25.1	(17.5,34.5)	22.6	(14.0,34.3)	32.6	(20.1,48.2)	8.0	(-8.4,24.4)	7.5	(-10.4,25.5)
Lower risk	1.7	(1.0,3.0)	1.9	(1.0,3.5)	1.1	(0.6,2.0)	1.8	(0.9,3.4)	0.7	(0.3,1.7)	-1.0	(-2.1,0.1)	-0.4	(-1.3,0.5)
Sensation seeking														
High	7.7	(5.5,10.7)	6.9	(4.2,11.4)	7.8	(5.7,10.7)	8.3	(5.6,12.0)	6.8	(4.1,11.1)	-0.9	(-5.1,3.3)	-1.0	(-5.8,3.7)
Low	1.0	(0.5,1.8)	0.8	(0.4,1.8)	0.9	(0.4,2.2)	0.8	(0.4,2.0)	0.7	(0.2,2.1)	-0.3	(-1.3,0.8)	-0.2	(-1.3,0.9)

Table 4-2. Youth reporting using marijuana in the past year, by age, gender, race/ethnicity, risk score, and sensation seeking (continued)

Characteristics	Percent reporting marijuana use in the past year													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 14 to 18														
Gender														
Males _____	23.8	(20.1,27.8)	21.7	(18.6,25.1)	21.5	(19.4,23.8)	21.8	(19.8,23.9)	21.7	(18.9,24.9)	-2.1	(-6.8,2.7)	0.2	(-3.1,3.5)
Females _____	18.6	(16.0,21.4)	20.5	(17.4,23.9)	22.7	(19.9,25.7)	21.4	(19.1,23.9)	20.1	(17.5,22.9)	1.5	(-2.5,5.5)	-2.7	(-6.5,1.2)
Race/ethnicity														
White _____	23.1	(20.5,26.0)	22.6	(19.6,25.9)	24.2	(22.0,26.5)	22.5	(20.7,24.4)	21.2	(18.9,23.8)	-1.9	(-5.1,1.4)	-2.9	(-6.0,0.1)
African American _____	17.4	(12.7,23.4)	16.1	(11.8,21.6)	14.5	(11.0,18.9)	17.3	(13.6,21.7)	18.8	(14.0,24.7)	1.3	(-6.2,8.9)	4.2	(-2.1,10.5)
Hispanic _____	17.0	(12.4,22.9)	19.4	(14.5,25.4)	21.2	(16.9,26.4)	20.9	(16.6,26.1)	24.0	(18.4,30.7)	7.0	(-0.6,14.6)	2.8	(-2.8,8.3)
Risk score														
Higher risk _____	37.1	(33.8,40.5)	36.1	(32.4,40.0)	36.5	(33.5,39.6)	36.6	(34.2,39.1)	35.9	(32.4,39.5)	-1.3	(-6.2,3.7)	-0.6	(-4.8,3.6)
Lower risk _____	3.6	(2.3,5.5)	4.9	(3.5,6.9)	5.6	(4.1,7.7)	6.2	(5.1,7.6)	6.9	(5.0,9.3)	3.3	*(0.8,5.7)	1.3	(-1.6,4.2)
Sensation seeking														
High _____	30.4	(27.4,33.5)	27.7	(24.6,31.0)	30.1	(27.4,32.9)	28.4	(26.2,30.8)	29.5	(26.8,32.3)	-0.9	(-4.6,2.8)	-0.6	(-4.2,2.9)
Low _____	8.1	(5.8,11.0)	10.1	(7.7,13.1)	10.3	(8.2,12.8)	11.5	(9.5,13.7)	8.7	(6.6,11.4)	0.7	(-2.8,4.1)	-1.6	(-4.8,1.6)

¹The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-3. Youth reporting using marijuana in the past month, by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting marijuana use in the past month													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13	1.7	(1.1,2.6)	1.6	(0.8,3.4)	1.1	(0.7,1.8)	1.3	(0.7,2.4)	1.2	(0.7,2.1)	-0.5	(-1.5,0.5)	0.1	(-0.9,1.0)
14 to 15	3.9	(2.5,5.9)	7.2	(5.4,9.5)	6.4	(5.0,8.0)	6.2	(5.0,7.6)	6.8	(5.3,8.8)	3.0	*(0.5,5.4)	0.5	(-1.9,2.9)
16 to 18	14.6	(12.3,17.2)	13.8	(11.3,16.7)	16.7	(14.6,19.0)	14.3	(12.6,16.1)	12.9	(10.8,15.3)	-1.7	(-5.1,1.7)	-3.8	*(-6.7,-0.8)
14 to 16	6.0	(4.6,7.7)	7.5	(5.8,9.7)	8.5	(7.3,10.0)	8.1	(6.9,9.5)	7.8	(6.5,9.3)	1.8	(-0.1,3.7)	-0.8	(-2.7,1.2)
14 to 18	9.7	(8.3,11.5)	10.8	(9.1,12.8)	12.3	(10.9,13.9)	10.8	(9.7,12.0)	10.4	(9.0,12.1)	0.7	(-1.6,3.0)	-1.9	(-3.8,0.1)
12.5 to 18	7.8	(6.7,9.2)	8.6	(7.3,10.1)	9.6	(8.5,10.8)	8.5	(7.7,9.4)	8.2	(7.1,9.5)	0.4	(-1.4,2.2)	-1.3	(-2.8,0.1)
Youth aged 12.5 to 13														
Gender														
Males	2.2	(1.3,3.7)	2.0	(0.8,5.1)	0.6	(0.3,1.3)	1.7	(0.8,3.8)	1.2	(0.6,2.4)	-1.0	(-2.4,0.4)	0.6	(-0.4,1.6)
Females	1.1	(0.5,2.4)	1.2	(0.5,2.9)	1.6	(0.8,3.2)	0.8	(0.3,2.0)	1.1	(0.4,3.2)	0.0	(-1.5,1.5)	-0.5	(-2.3,1.4)
Race/ethnicity														
White	1.5	(0.8,2.8)	0.9	(0.3,2.5)	1.2	(0.6,2.4)	0.8	(0.3,1.8)	0.9	(0.4,2.1)	-0.6	(-1.8,0.6)	-0.3	(-1.4,0.8)
African American	1.7	(0.4,6.5)	1.4	(0.3,5.9)	1.0	(0.4,2.4)	2.5	(0.6,9.9)	1.6	(0.4,7.0)	-0.1	(-3.3,3.2)	0.6	(-2.0,3.2)
Hispanic	2.2	(0.8,5.7)	4.4	(1.2,15.5)	1.2	(0.4,3.6)	2.0	(0.8,4.8)	0.1	(0.0,1.1)	-2.0	(-4.2,0.1)	-1.0	(-2.6,0.5)
Risk score														
Higher risk	12.0	(6.8,20.4)	7.1	(2.5,18.3)	7.7	(4.1,14.2)	7.1	(2.9,16.6)	12.6	(6.6,22.7)	0.6	(-9.9,11.0)	4.8	(-5.0,14.7)
Lower risk	0.4	(0.2,0.9)	1.0	(0.4,2.5)	0.3	(0.1,0.7)	0.5	(0.2,1.4)	0.3	(0.1,1.2)	-0.2	(-0.7,0.3)	0.0	(-0.5,0.5)
Sensation seeking														
High	3.0	(1.7,5.1)	3.1	(1.3,7.0)	2.2	(1.2,4.1)	2.0	(0.9,4.2)	2.4	(1.4,4.4)	-0.5	(-2.7,1.7)	0.2	(-2.0,2.4)
Low	0.6	(0.3,1.3)	0.3	(0.1,1.3)	0.3	(0.1,0.9)	0.6	(0.2,1.8)	0.3	(0.0,2.2)	-0.3	(-1.1,0.6)	0.0	(-0.7,0.7)

Table 4-3. Youth reporting using marijuana in the past month, by age, gender, race/ethnicity, risk score, and sensation seeking (continued)

Characteristics	Percent reporting marijuana use in the past month													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 14 to 18														
Gender														
Males_____	11.8	(9.2,15.0)	10.9	(8.6,13.7)	12.1	(10.3,14.2)	11.3	(9.9,12.9)	11.5	(9.4,13.9)	-0.3	(-4.1,3.5)	-0.7	(-3.6,2.3)
Females_____	7.6	(5.7,10.2)	10.7	(8.6,13.3)	12.4	(10.4,14.8)	10.3	(8.6,12.3)	9.3	(7.5,11.5)	1.7	(-1.2,4.6)	-3.1	*(-6.0,-0.2)
Race/ethnicity														
White_____	9.9	(8.1,12.0)	12.0	(9.9,14.5)	13.4	(11.6,15.3)	10.8	(9.5,12.3)	10.1	(8.4,12.2)	0.2	(-2.4,2.9)	-3.2	*(-5.6,-0.8)
African American__	8.5	(5.1,13.8)	8.4	(5.2,13.3)	9.1	(6.2,13.3)	9.0	(6.6,12.2)	12.0	(8.4,16.8)	3.4	(-2.3,9.2)	2.8	(-1.8,7.5)
Hispanic_____	10.4	(6.9,15.4)	7.6	(4.3,13.1)	12.2	(9.2,16.0)	12.6	(9.2,17.0)	10.9	(7.3,15.9)	0.5	(-5.3,6.4)	-1.3	(-5.9,3.3)
Risk score														
Higher risk_____	17.8	(15.0,21.0)	18.9	(16.0,22.2)	21.0	(18.6,23.6)	19.6	(17.5,21.9)	18.1	(15.4,21.1)	0.3	(-4.2,4.7)	-2.9	(-6.3,0.4)
Lower risk_____	1.2	(0.7,2.1)	2.3	(1.3,3.9)	2.4	(1.7,3.5)	2.1	(1.5,2.9)	3.1	(2.0,4.7)	1.8	*(0.5,3.2)	0.7	(-1.0,2.3)
Sensation seeking														
High_____	14.6	(12.1,17.4)	14.1	(11.7,16.8)	17.1	(15.0,19.5)	14.9	(13.2,16.8)	14.7	(12.7,16.9)	0.1	(-3.5,3.7)	-2.4	(-5.2,0.4)
Low_____	2.7	(1.6,4.5)	5.5	(3.8,7.9)	5.3	(4.0,6.9)	4.9	(3.7,6.3)	4.2	(2.7,6.5)	1.6	(-0.8,3.9)	-1.0	(-3.4,1.4)

¹The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-4. Youth reporting regular marijuana use,¹ by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting regular marijuana use													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ²	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13	0.6	(0.3,1.3)	0.4	(0.1,1.2)	0.6	(0.3,1.2)	0.7	(0.3,1.7)	0.8	(0.3,2.2)	0.2	(-0.8,1.2)	0.2	(-0.7,1.1)
14 to 15	2.3	(1.5,3.4)	5.4	(4.0,7.3)	4.4	(3.5,5.6)	3.9	(3.0,5.0)	3.9	(2.9,5.2)	1.6	*(0.2,2.9)	-0.5	(-2.1,1.1)
16 to 18	12.6	(10.4,15.0)	11.6	(9.3,14.5)	12.4	(10.5,14.7)	11.5	(9.9,13.2)	10.6	(8.7,12.9)	-1.9	(-5.3,1.4)	-1.8	(-4.7,1.1)
14 to 16	4.0	(2.9,5.5)	5.9	(4.8,7.4)	5.6	(4.7,6.7)	5.8	(4.8,7.0)	5.1	(4.1,6.3)	1.0	(-0.6,2.7)	-0.5	(-2.1,1.0)
14 to 18	7.9	(6.7,9.3)	8.8	(7.3,10.6)	9.0	(7.8,10.5)	8.2	(7.3,9.3)	7.9	(6.7,9.3)	-0.1	(-2.0,1.9)	-1.2	(-2.9,0.6)
12.5 to 18	6.2	(5.3,7.3)	6.8	(5.6,8.2)	7.0	(6.1,8.1)	6.4	(5.7,7.2)	6.2	(5.3,7.3)	0.0	(-1.5,1.5)	-0.8	(-2.1,0.5)
Youth aged 14 to 18														
Gender														
Males	10.0	(7.9,12.5)	10.8	(8.5,13.6)	9.4	(7.9,11.1)	9.0	(7.8,10.5)	9.4	(7.6,11.5)	-0.6	(-3.8,2.6)	0.0	(-2.6,2.5)
Females	5.8	(4.1,8.1)	6.7	(5.2,8.6)	8.6	(6.8,11.0)	7.4	(6.0,9.2)	6.3	(4.7,8.4)	0.5	(-1.9,2.9)	-2.4	(-4.8,0.1)
Race/ethnicity														
White	9.1	(7.4,11.1)	10.4	(8.4,12.8)	9.9	(8.3,11.9)	8.6	(7.4,10.0)	7.9	(6.3,9.8)	-1.2	(-3.6,1.2)	-2.0	(-4.3,0.3)
African American	4.0	(2.1,7.4)	3.6	(2.0,6.5)	6.5	(4.2,9.9)	5.4	(3.5,8.2)	6.6	(3.7,11.4)	2.6	(-1.3,6.5)	0.1	(-4.1,4.4)
Hispanic	5.8	(3.2,10.5)	6.8	(3.7,12.4)	8.6	(6.0,12.0)	9.8	(6.7,14.2)	9.6	(6.9,13.2)	3.7	(-1.3,8.7)	1.0	(-2.7,4.8)
Risk score														
Higher risk	14.8	(12.5,17.4)	14.9	(12.4,17.9)	15.8	(13.5,18.4)	15.0	(13.1,17.1)	14.8	(12.4,17.7)	0.1	(-3.6,3.8)	-1.0	(-4.3,2.3)
Lower risk	0.5	(0.2,1.2)	1.6	(0.8,3.1)	1.4	(0.9,2.3)	1.3	(0.7,2.3)	1.4	(0.8,2.3)	0.8	(0.0,1.6)	0.0	(-1.1,1.0)
Sensation seeking														
High	11.9	(9.9,14.2)	12.5	(10.3,15.2)	13.0	(11.1,15.1)	11.7	(10.2,13.4)	12.0	(10.1,14.1)	0.1	(-3.3,3.4)	-1.0	(-3.7,1.7)
Low	2.4	(1.2,4.6)	2.7	(1.7,4.1)	3.0	(2.1,4.3)	3.1	(2.2,4.4)	1.9	(1.2,2.9)	-0.5	(-2.2,1.2)	-1.1	(-2.4,0.2)

¹Regular use = Used 10 or more times in past year.²The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-5. Youth never receiving offers of marijuana, by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting never receiving offers of marijuana													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13	78.8	(76.0,81.4)	79.3	(76.7,81.7)	80.1	(77.2,82.8)	80.9	(78.1,83.5)	82.9	(79.7,85.7)	4.1	(-0.1,8.3)	2.8	(-1.2,6.7)
14 to 15	53.9	(50.4,57.3)	54.1	(49.9,58.2)	54.2	(51.0,57.3)	58.6	(56.3,60.9)	59.8	(56.8,62.7)	5.9	*(1.4,10.4)	5.6	*(1.5,9.7)
16 to 18	29.1	(25.9,32.6)	28.9	(26.0,31.9)	29.8	(27.2,32.5)	34.6	(32.2,37.1)	34.8	(31.5,38.2)	5.6	*(0.9,10.4)	5.0	*(1.0,9.0)
14 to 16	48.7	(45.6,51.9)	48.3	(44.9,51.7)	49.1	(46.3,51.9)	52.3	(50.3,54.3)	54.5	(51.8,57.1)	5.8	*(1.7,9.8)	5.4	*(1.9,8.8)
14 to 18	40.3	(37.6,43.1)	40.3	(37.5,43.0)	40.1	(37.9,42.4)	44.8	(43.1,46.5)	45.0	(42.4,47.6)	4.7	*(1.2,8.1)	4.9	*(1.9,7.9)
12.5 to 18	49.4	(47.2,51.6)	49.5	(47.3,51.7)	49.7	(47.6,51.8)	53.5	(51.8,55.1)	54.0	(51.8,56.2)	4.6	*(1.8,7.4)	4.3	*(1.9,6.6)
Youth aged 12.5 to 13														
Gender														
Males	78.8	(74.7,82.4)	80.1	(76.1,83.5)	80.9	(77.1,84.2)	81.5	(78.4,84.2)	83.6	(79.9,86.8)	4.8	(0.0,9.7)	2.7	(-2.5,8.0)
Females	78.8	(75.1,82.1)	78.4	(73.6,82.6)	79.4	(75.7,82.6)	80.4	(75.6,84.4)	82.2	(76.1,87.0)	3.4	(-3.5,10.3)	2.8	(-3.7,9.3)
Race/ethnicity														
White	81.6	(78.6,84.3)	81.3	(78.2,84.2)	83.2	(80.4,85.6)	82.4	(78.9,85.4)	85.7	(82.9,88.1)	4.1	*(0.0,8.1)	2.5	(-1.0,6.1)
African American	74.0	(65.1,81.3)	81.9	(74.3,87.6)	69.7	(58.5,79.0)	81.1	(70.8,88.3)	77.5	(63.0,87.4)	3.5	(-11.9,18.8)	7.8	(-7.0,22.6)
Hispanic	70.2	(60.6,78.2)	66.5	(60.2,72.2)	79.2	(73.3,84.1)	74.5	(67.7,80.3)	77.6	(68.6,84.5)	7.4	(-4.1,18.8)	-1.7	(-10.2,6.9)
Risk score														
Higher risk	34.1	(25.3,44.0)	33.3	(21.8,47.2)	40.4	(31.6,49.7)	40.4	(30.4,51.4)	39.5	(28.5,51.6)	5.4	(-8.9,19.7)	-0.9	(-15.8,14.0)
Lower risk	83.6	(80.6,86.2)	85.2	(82.8,87.3)	85.2	(82.4,87.6)	86.5	(84.3,88.5)	87.8	(84.8,90.3)	4.2	*(0.2,8.3)	2.7	(-0.9,6.2)
Sensation seeking														
High	67.2	(62.6,71.5)	68.4	(63.1,73.3)	70.9	(66.4,75.0)	70.7	(65.9,75.1)	73.9	(67.5,79.4)	6.7	(-1.5,14.8)	3.0	(-4.2,10.2)
Low	87.7	(84.7,90.2)	87.8	(84.7,90.3)	87.1	(83.8,89.9)	90.4	(87.5,92.7)	90.1	(86.6,92.7)	2.3	(-2.0,6.6)	2.9	(-1.3,7.1)

Table 4-5. Youth never receiving offers of marijuana, by age, gender, race/ethnicity, risk score, and sensation seeking (continued)

Characteristics	Percent reporting never receiving offers of marijuana													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 14 to 18														
Gender														
Males_____	36.7	(33.3,40.1)	36.2	(32.3,40.3)	38.9	(36.2,41.8)	43.7	(41.1,46.2)	43.0	(40.0,46.0)	6.3	*(2.1,10.5)	4.0	*(0.4,7.6)
Females_____	44.1	(39.6,48.8)	44.5	(40.4,48.6)	41.4	(38.1,44.7)	46.0	(43.7,48.3)	47.2	(43.5,50.9)	3.0	(-2.8,8.9)	5.8	*(1.1,10.5)
Race/ethnicity														
White_____	41.4	(38.5,44.3)	41.3	(38.2,44.6)	41.0	(38.7,43.4)	44.8	(42.6,47.0)	46.1	(43.0,49.3)	4.7	*(1.4,8.0)	5.1	*(1.4,8.8)
African American_	37.8	(32.0,44.0)	39.3	(32.8,46.3)	37.1	(30.9,43.7)	46.2	(40.9,51.7)	46.3	(40.0,52.7)	8.5	(-1.2,18.2)	9.2	*(3.0,15.4)
Hispanic_____	33.0	(24.8,42.4)	34.8	(27.7,42.6)	36.6	(31.4,42.2)	41.4	(37.2,45.6)	36.5	(31.0,42.3)	3.4	(-7.7,14.6)	-0.2	(-7.0,6.6)
Risk score														
Higher risk_____	20.6	(17.8,23.7)	18.5	(16.0,21.3)	20.5	(18.0,23.1)	23.4	(21.2,25.7)	23.3	(20.7,26.0)	2.7	(-1.3,6.7)	2.8	(-0.7,6.3)
Lower risk_____	63.3	(58.5,67.9)	64.2	(59.9,68.4)	62.1	(58.8,65.3)	66.4	(63.8,68.9)	66.8	(63.2,70.3)	3.5	(-2.3,9.4)	4.8	*(0.7,8.9)
Sensation seeking														
High_____	31.1	(27.8,34.5)	30.2	(27.2,33.5)	30.7	(28.3,33.3)	35.4	(33.3,37.7)	35.1	(32.2,38.1)	4.0	(0.0,8.0)	4.3	*(0.9,7.7)
Low_____	53.0	(48.0,58.0)	56.9	(51.8,61.9)	53.6	(50.3,56.9)	58.1	(55.6,60.5)	58.4	(54.7,62.1)	5.4	(-0.6,11.5)	4.8	*(0.2,9.5)

¹The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-6. Youth receiving offers of marijuana one or more times in the past 30 days, by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting receiving offers of marijuana one or more times in past 30 days													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13	11.1	(9.4,13.1)	10.8	(9.0,13.1)	10.9	(9.0,13.1)	11.0	(8.8,13.7)	9.0	(7.0,11.6)	-2.1	(-4.8,0.6)	-1.9	(-4.9,1.1)
14 to 15	26.8	(23.2,30.7)	28.2	(24.7,31.9)	28.0	(25.4,30.8)	25.4	(23.2,27.6)	25.5	(22.8,28.4)	-1.3	(-5.9,3.3)	-2.5	(-6.3,1.4)
16 to 18	46.7	(42.9,50.5)	47.0	(43.1,50.8)	45.8	(42.7,49.0)	43.1	(40.2,45.9)	40.7	(37.7,43.8)	-5.9	*(-10.6,-1.3)	-5.1	*(-9.5,-0.6)
14 to 16	30.9	(28.2,33.7)	32.5	(29.3,35.7)	30.7	(28.3,33.2)	30.1	(28.0,32.4)	28.8	(26.3,31.3)	-2.1	(-5.9,1.7)	-1.9	(-5.3,1.4)
14 to 18	37.7	(35.1,40.4)	38.5	(35.6,41.4)	38.2	(36.1,40.4)	35.5	(33.6,37.5)	34.5	(32.1,36.9)	-3.2	*(-6.4,0.0)	-3.7	*(-6.8,-0.7)
12.5 to 18	31.4	(29.3,33.6)	31.9	(29.7,34.3)	31.7	(29.8,33.6)	29.6	(28.1,31.3)	28.5	(26.6,30.5)	-2.9	*(-5.5,-0.4)	-3.2	*(-5.5,-0.8)
Youth aged 12.5 to 13														
Gender														
Males	10.9	(8.4,14.1)	8.9	(6.5,12.2)	9.1	(6.9,11.8)	9.6	(7.2,12.6)	8.7	(6.3,11.9)	-2.2	(-6.0,1.6)	-0.4	(-4.3,3.6)
Females	11.4	(9.2,14.1)	13.0	(9.9,17.1)	12.7	(10.3,15.7)	12.5	(9.1,16.9)	9.4	(6.1,14.0)	-2.0	(-6.5,2.5)	-3.4	(-8.0,1.2)
Race/ethnicity														
White	9.1	(7.2,11.4)	9.6	(7.3,12.4)	9.4	(7.6,11.6)	9.3	(6.9,12.4)	6.9	(5.1,9.3)	-2.2	(-5.1,0.6)	-2.6	(-5.2,0.1)
African American	12.3	(7.8,19.0)	7.6	(4.1,13.9)	15.0	(9.5,22.8)	15.8	(8.9,26.4)	13.1	(5.8,26.8)	0.8	(-10.8,12.3)	-1.9	(-13.9,10.1)
Hispanic	19.1	(13.5,26.2)	20.1	(14.9,26.6)	14.1	(10.4,18.8)	14.9	(10.5,20.8)	11.8	(7.7,17.8)	-7.2	(-14.9,0.5)	-2.2	(-9.0,4.5)
Risk score														
Higher risk	40.1	(32.2,48.6)	42.9	(31.0,55.7)	38.1	(30.2,46.6)	40.1	(29.7,51.4)	42.5	(30.2,55.7)	2.3	(-13.2,17.9)	4.4	(-11.6,20.4)
Lower risk	7.9	(6.1,10.0)	8.1	(6.5,10.0)	7.1	(5.5,9.2)	6.9	(5.4,8.7)	6.1	(4.4,8.5)	-1.7	(-4.4,0.9)	-1.0	(-3.7,1.7)
Sensation seeking														
High	17.6	(14.5,21.2)	18.4	(14.5,23.0)	17.4	(14.5,20.7)	17.1	(13.2,21.9)	16.1	(11.8,21.6)	-1.5	(-7.3,4.3)	-1.3	(-6.9,4.3)
Low	6.2	(4.5,8.5)	5.4	(3.9,7.4)	6.0	(4.2,8.4)	5.0	(3.3,7.4)	4.1	(2.7,6.0)	-2.1	(-4.6,0.3)	-1.9	(-4.5,0.6)

Table 4-6. Youth receiving offers of marijuana one or more times in the past 30 days, by age, gender, race/ethnicity, risk score, and sensation seeking (continued)

Characteristics	Percent reporting receiving offers of marijuana one or more times in past 30 days													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 14 to 18														
Gender														
Males_____	40.6	(36.9,44.3)	39.0	(34.6,43.7)	39.0	(36.3,41.7)	34.6	(32.0,37.3)	34.5	(31.2,37.9)	-6.1	*(-11.1,-1.1)	-4.5	*(-8.7,-0.3)
Females_____	34.7	(30.8,38.9)	37.9	(34.6,41.3)	37.5	(34.2,40.8)	36.5	(34.0,39.0)	34.5	(31.4,37.7)	-0.2	(-4.9,4.5)	-2.9	(-7.5,1.6)
Race/ethnicity														
White_____	37.4	(34.2,40.6)	37.6	(33.9,41.5)	38.0	(35.4,40.6)	35.6	(33.4,37.8)	33.1	(30.3,35.9)	-4.3	*(-7.8,-0.8)	-4.9	*(-8.8,-1.0)
African American__	38.9	(33.4,44.7)	34.1	(28.4,40.4)	40.7	(35.7,46.0)	31.9	(26.1,38.2)	35.1	(29.3,41.5)	-3.7	(-12.1,4.6)	-5.6	(-12.0,0.7)
Hispanic_____	40.5	(33.5,48.0)	47.5	(39.7,55.4)	41.7	(37.1,46.5)	41.0	(36.6,45.6)	41.8	(35.8,48.1)	1.3	(-8.6,11.2)	0.1	(-6.6,6.7)
Risk score														
Higher risk_____	55.0	(50.7,59.2)	57.0	(52.7,61.1)	55.5	(52.2,58.8)	53.1	(50.0,56.1)	49.8	(46.0,53.5)	-5.2	(-10.7,0.2)	-5.7	*(-10.4,-1.1)
Lower risk_____	17.9	(14.6,21.7)	18.8	(15.7,22.3)	19.2	(17.1,21.5)	17.6	(15.3,20.1)	19.5	(16.9,22.4)	1.6	(-2.4,5.7)	0.3	(-3.2,3.8)
Sensation seeking														
High_____	47.6	(44.3,51.0)	46.2	(42.4,50.2)	47.1	(44.1,50.1)	44.4	(41.9,46.9)	43.5	(40.3,46.8)	-4.1	(-8.5,0.3)	-3.5	(-7.5,0.4)
Low_____	23.7	(20.0,27.8)	25.7	(21.7,30.1)	25.3	(22.2,28.6)	23.0	(20.6,25.6)	22.0	(19.3,24.9)	-1.7	(-6.1,2.7)	-3.3	(-7.5,0.9)

¹The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-7. Youth reporting ever having used inhalants, by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting inhalant use ever													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13	1.5	(0.9,2.2)	1.6	(1.0,2.6)	1.7	(1.2,2.4)	2.3	(1.4,3.9)	1.9	(1.2,3.0)	0.5	(-0.5,1.4)	0.2	(-0.9,1.3)
14 to 15	5.9	(3.9,8.6)	3.6	(2.6,5.0)	3.2	(2.4,4.2)	3.5	(2.6,4.6)	3.4	(2.3,4.9)	-2.5	(-5.2,0.2)	0.2	(-1.3,1.6)
16 to 18	8.0	(6.4,9.9)	6.0	(4.7,7.7)	5.7	(4.4,7.4)	4.4	(3.4,5.6)	5.0	(3.8,6.6)	-3.0	*(-5.4,-0.7)	-0.7	(-2.5,1.2)
14 to 16	6.6	(5.0,8.6)	3.4	(2.5,4.6)	3.5	(2.7,4.4)	3.4	(2.7,4.3)	3.4	(2.5,4.5)	-3.2	*(-5.3,-1.2)	-0.1	(-1.4,1.1)
14 to 18	7.0	(5.8,8.5)	4.9	(4.0,6.1)	4.6	(3.8,5.7)	4.0	(3.3,4.9)	4.3	(3.5,5.3)	-2.7	*(-4.4,-1.0)	-0.3	(-1.4,0.9)
12.5 to 18	5.7	(4.8,6.8)	4.2	(3.4,5.0)	3.9	(3.2,4.8)	3.6	(3.0,4.3)	3.8	(3.1,4.5)	-2.0	*(-3.3,-0.6)	-0.1	(-1.1,0.8)
Youth aged 12.5 to 13														
Gender														
Males	1.6	(1.0,2.8)	1.2	(0.5,2.8)	1.1	(0.7,1.9)	2.1	(1.1,4.1)	2.5	(1.4,4.4)	0.8	(-0.9,2.6)	1.3	(-0.1,2.8)
Females	1.3	(0.6,2.5)	2.1	(1.1,3.9)	2.2	(1.4,3.5)	2.6	(1.4,4.8)	1.3	(0.6,3.1)	0.1	(-1.2,1.4)	-0.9	(-2.4,0.7)
Race/ethnicity														
White	1.6	(1.0,2.7)	1.6	(0.9,2.8)	1.8	(1.1,2.7)	3.0	(1.7,5.0)	1.6	(0.8,3.1)	0.0	(-1.4,1.3)	-0.2	(-1.7,1.4)
African American	0.3	(0.0,1.7)	1.4	(0.3,5.9)	0.3	(0.0,3.6)	S	(S)	3.0	(1.0,8.8)	2.8	(-0.6,6.1)	2.7	(-0.7,6.1)
Hispanic	1.9	(0.8,4.8)	2.1	(0.9,5.1)	2.9	(1.3,6.1)	2.6	(0.7,9.3)	0.7	(0.1,4.8)	-1.3	(-3.3,0.8)	-2.2	(-5.2,0.8)
Risk score														
Higher risk	8.5	(4.7,15.0)	11.0	(5.7,20.2)	7.5	(4.1,13.3)	11.1	(5.4,21.6)	8.8	(3.5,20.2)	0.3	(-8.1,8.6)	1.3	(-7.4,10.1)
Lower risk	0.7	(0.3,1.3)	0.6	(0.3,1.4)	0.7	(0.4,1.2)	1.3	(0.6,2.5)	1.1	(0.6,2.2)	0.5	(-0.4,1.4)	0.4	(-0.4,1.2)
Sensation seeking														
High	3.1	(2.0,4.8)	2.6	(1.5,4.3)	3.4	(2.3,5.0)	4.3	(2.4,7.9)	4.1	(2.7,6.2)	1.1	(-1.0,3.1)	0.7	(-1.5,2.9)
Low	0.1	(0.0,0.6)	0.7	(0.3,2.0)	0.2	(0.1,1.0)	0.7	(0.3,1.8)	0.4	(0.1,2.5)	0.3	(-0.4,1.0)	0.1	(-0.8,1.1)

Table 4-7. Youth reporting ever having used inhalants, by age, gender, race/ethnicity, risk score, and sensation seeking (continued)

Characteristics	Percent reporting inhalant use ever													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 14 to 18														
Gender														
Males_____	9.1	(6.8,12.0)	5.1	(3.6,7.1)	4.5	(3.4,6.0)	3.7	(2.7,5.0)	3.9	(2.7,5.6)	-5.2	*(-8.5,-1.9)	-0.6	(-2.4,1.2)
Females_____	4.9	(3.6,6.6)	4.8	(3.5,6.5)	4.7	(3.6,6.2)	4.3	(3.4,5.5)	4.8	(3.5,6.5)	-0.1	(-2.2,2.0)	0.1	(-1.7,1.9)
Race/ethnicity														
White_____	7.8	(6.2,9.9)	5.5	(4.4,6.7)	5.5	(4.3,6.9)	4.7	(3.8,5.8)	4.8	(3.8,6.1)	-3.0	*(-5.3,-0.7)	-0.7	(-2.2,0.8)
African American__	1.5	(0.5,4.0)	1.7	(0.8,3.6)	1.0	(0.5,2.1)	1.0	(0.4,2.1)	0.4	(0.1,1.8)	-1.0	(-2.7,0.6)	-0.6	(-1.5,0.3)
Hispanic_____	8.4	(5.3,13.1)	6.2	(3.5,10.9)	5.2	(3.2,8.2)	4.1	(2.4,6.8)	6.5	(4.3,9.7)	-1.9	(-6.0,2.2)	1.3	(-1.6,4.2)
Risk score														
Higher risk_____	12.6	10.4,15.2)	9.3	(7.5,11.4)	7.6	(6.1,9.4)	7.1	(5.7,8.7)	7.1	(5.7,8.7)	-5.6	*(-8.5,-2.7)	-0.5	(-2.6,1.5)
Lower risk_____	1.3	(0.7,2.4)	0.7	(0.5,1.1)	1.0	(0.5,2.2)	0.7	(0.4,1.3)	1.0	(0.5,1.8)	-0.3	(-1.4,0.7)	-0.1	(-1.0,0.9)
Sensation seeking														
High_____	11.4	(9.4,13.8)	6.6	(5.3,8.3)	6.8	(5.6,8.3)	5.5	(4.4,6.7)	6.2	(4.8,8.1)	-5.2	*(-8.1,-2.3)	-0.6	(-2.5,1.3)
Low_____	0.6	(0.3,1.3)	2.2	(1.2,3.9)	1.5	(0.9,2.5)	1.7	(1.0,2.9)	1.4	(0.8,2.5)	0.8	(0.0,1.7)	-0.1	(-1.3,1.1)

¹The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-8. Youth reporting using inhalant in the past year, by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting inhalant use in the past year													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13 _____	1.0	(0.5,1.8)	1.0	(0.5,1.9)	1.1	(0.7,1.7)	1.7	(0.9,3.1)	1.5	(0.9,2.6)	0.6	(-0.4,1.5)	0.5	(-0.5,1.5)
14 to 15 _____	2.8	(1.8,4.3)	1.9	(1.1,3.1)	1.9	(1.3,2.8)	2.0	(1.4,2.9)	2.2	(1.4,3.4)	-0.6	(-2.1,1.0)	0.3	(-0.9,1.4)
16 to 18 _____	3.2	(2.1,4.9)	2.2	(1.3,3.7)	1.6	(1.0,2.6)	1.7	(1.1,2.6)	2.1	(1.4,3.3)	-1.1	(-2.8,0.7)	0.5	(-0.8,1.8)
14 to 16 _____	3.0	(2.2,4.2)	1.6	(1.0,2.5)	1.7	(1.2,2.4)	2.0	(1.4,2.8)	2.1	(1.5,3.0)	-0.9	(-2.3,0.4)	0.4	(-0.5,1.3)
14 to 18 _____	3.0	(2.2,4.1)	2.1	(1.4,3.0)	1.7	(1.3,2.4)	1.8	(1.3,2.5)	2.2	(1.6,2.9)	-0.8	(-2.0,0.4)	0.4	(-0.5,1.3)
12.5 to 18 _____	2.5	(1.9,3.3)	1.8	(1.3,2.5)	1.6	(1.2,2.1)	1.8	(1.3,2.4)	2.0	(1.5,2.7)	-0.5	(-1.4,0.4)	0.4	(-0.3,1.2)
Youth aged 12.5 to 13														
Gender														
Males _____	1.0	(0.4,2.3)	0.7	(0.3,2.1)	0.5	(0.2,1.1)	1.0	(0.4,2.5)	1.8	(0.8,3.8)	0.8	(-0.8,2.4)	1.2	(-0.1,2.6)
Females _____	1.0	(0.5,2.3)	1.3	(0.6,2.8)	1.7	(1.0,2.9)	2.4	(1.2,4.7)	1.3	(0.6,3.1)	0.3	(-0.9,1.5)	-0.3	(-1.8,1.1)
Race/ethnicity														
White _____	1.2	(0.6,2.4)	1.0	(0.4,2.1)	1.3	(0.8,2.3)	2.0	(1.1,3.9)	1.0	(0.4,2.7)	-0.2	(-1.5,1.1)	-0.3	(-1.7,1.1)
African American _____	S	(S)	0.8	(0.1,6.4)	S	(S)	S	(S)	3.0	(1.0,8.8)	S	(S)	S	(S)
Hispanic _____	1.6	(0.6,4.3)	1.6	(0.6,4.2)	1.4	(0.5,4.4)	2.3	(0.5,9.5)	0.7	(0.1,4.8)	-0.9	(-2.8,1.0)	-0.8	(-2.9,1.3)
Risk score														
Higher risk _____	5.4	(2.2,12.2)	6.5	(2.6,15.3)	5.2	(2.4,10.9)	9.4	(4.2,19.8)	7.7	(2.8,19.3)	2.4	(-5.8,10.5)	2.6	(-5.7,10.8)
Lower risk _____	0.5	(0.2,1.2)	0.3	(0.1,0.8)	0.4	(0.2,0.8)	0.7	(0.3,1.8)	0.8	(0.4,1.8)	0.3	(-0.5,1.1)	0.4	(-0.2,1.1)
Sensation seeking														
High _____	2.1	(1.2,3.8)	1.7	(0.8,3.4)	2.0	(1.2,3.4)	3.2	(1.6,6.5)	3.3	(2.0,5.3)	1.2	(-0.7,3.1)	1.2	(-0.7,3.1)
Low _____	0.0	(0.0,1.1)	0.3	(0.1,1.3)	0.2	(0.1,1.0)	0.4	(0.1,1.6)	0.4	(0.1,2.5)	0.4	(-0.3,1.0)	0.1	(-0.8,1.1)

Table 4-8. Youth reporting using inhalants in the past year, by age, gender, race/ethnicity, risk score, and sensation seeking (continued)

Characteristics	Percent reporting inhalant use in the past year													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 14 to 18														
Gender														
Males_____	3.5	(2.2,5.5)	2.3	(1.3,3.9)	1.9	(1.3,3.0)	1.8	(1.1,2.9)	1.8	(1.2,2.8)	-1.7	(-3.5,0.2)	-0.1	(-1.2,1.0)
Females_____	2.5	(1.6,3.7)	1.8	(1.2,2.8)	1.5	(1.0,2.3)	1.8	(1.3,2.5)	2.5	(1.6,3.9)	0.0	(-1.5,1.5)	1.0	(-0.3,2.2)
Race/ethnicity														
White_____	3.5	(2.4,5.0)	2.6	(1.7,3.9)	2.1	(1.4,3.0)	2.2	(1.5,3.1)	2.7	(1.8,3.8)	-0.8	(-2.5,0.8)	0.6	(-0.6,1.8)
African American_____	0.1	(0.0,0.9)	0.5	(0.1,2.5)	0.7	(0.2,2.7)	0.4	(0.1,1.5)	0.4	(0.1,1.8)	0.3	(-0.4,1.0)	-0.3	(-1.4,0.9)
Hispanic_____	2.1	(1.0,4.4)	1.3	(0.6,2.7)	1.5	(0.7,2.9)	1.6	(0.8,3.2)	2.0	(1.0,4.1)	-0.1	(-2.2,2.0)	0.6	(-1.3,2.5)
Risk score														
Higher risk_____	5.7	(4.1,7.8)	3.9	(2.6,5.7)	2.7	(1.9,3.8)	3.3	(2.3,4.7)	4.1	(2.9,5.7)	-1.6	(-4.0,0.8)	1.4	(-0.3,3.1)
Lower risk_____	0.5	(0.2,1.2)	0.6	(0.3,1.0)	0.7	(0.3,1.8)	0.3	(0.1,0.6)	0.5	(0.2,1.3)	0.0	(-0.6,0.7)	-0.2	(-1.0,0.6)
Sensation seeking														
High_____	4.8	(3.5,6.5)	2.9	(1.9,4.4)	2.6	(1.9,3.6)	2.6	(1.9,3.6)	3.2	(2.2,4.6)	-1.6	(-3.6,0.5)	0.6	(-1.0,2.1)
Low_____	0.6	(0.3,1.2)	0.7	(0.4,1.3)	0.5	(0.2,1.1)	0.5	(0.2,1.4)	0.8	(0.4,1.6)	0.2	(-0.5,0.9)	0.3	(-0.4,1.1)

¹The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-9. Youth reporting using inhalants in the past month, by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting inhalant use in the past month													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13 _____	0.3	(0.1,0.7)	0.5	(0.2,1.2)	0.4	(0.2,0.8)	0.7	(0.2,1.7)	1.0	(0.5,2.2)	0.8	(0.0,1.6)	0.6	(-0.1,1.4)
14 to 15 _____	0.3	(0.1,0.7)	0.8	(0.3,1.8)	0.8	(0.4,1.4)	0.6	(0.3,1.1)	1.2	(0.7,2.1)	0.9	*(0.2,1.6)	0.4	(-0.3,1.2)
16 to 18 _____	0.9	(0.4,2.0)	0.4	(0.2,1.0)	0.6	(0.3,1.1)	0.6	(0.3,1.3)	1.2	(0.6,2.3)	0.3	(-0.8,1.3)	0.6	(-0.2,1.4)
14 to 16 _____	0.5	(0.2,1.1)	0.6	(0.3,1.2)	0.6	(0.4,1.1)	0.5	(0.3,0.9)	1.1	(0.7,1.7)	0.6	(-0.1,1.2)	0.4	(-0.1,1.0)
14 to 18 _____	0.6	(0.3,1.2)	0.6	(0.3,1.0)	0.7	(0.4,1.0)	0.6	(0.4,1.0)	1.2	(0.8,1.8)	0.5	(-0.1,1.2)	0.5	(0.0,1.1)
12.5 to 18 _____	0.5	(0.3,1.0)	0.6	(0.4,0.9)	0.6	(0.4,0.9)	0.6	(0.4,1.0)	1.1	(0.8,1.7)	0.6	*(0.1,1.1)	0.6	*(0.1,1.0)
Youth aged 12.5 to 13														
Gender														
Males _____	0.1	(0.0,0.9)	0.5	(0.2,1.8)	0.3	(0.1,0.8)	0.2	(0.1,1.0)	1.8	(0.8,3.8)	1.6	*(0.2,3.0)	1.4	*(0.1,2.8)
Females _____	0.4	(0.2,1.2)	0.5	(0.1,1.6)	0.5	(0.2,1.2)	1.1	(0.3,3.4)	0.3	(0.0,2.0)	-0.2	(-0.9,0.6)	-0.2	(-1.0,0.5)
Race/ethnicity														
White _____	0.2	(0.1,0.5)	0.5	(0.2,1.6)	0.6	(0.3,1.1)	0.5	(0.2,1.4)	0.6	(0.2,2.1)	0.4	(-0.3,1.2)	0.1	(-0.8,0.9)
African American _____	S	(S)	S	(S)	S	(S)	S	(S)	3.0	(1.0,8.8)	S	(S)	S	(S)
Hispanic _____	1.1	(0.2,4.4)	1.0	(0.2,4.4)	0.5	(0.1,3.2)	2.1	(0.4,9.9)	S	(S)	S	(S)	S	(S)
Risk score														
Higher risk _____	S	(S)	3.2	(0.8,11.6)	1.4	(0.5,3.8)	4.6	(1.4,14.1)	5.2	(1.5,16.5)	S	(S)	3.8	(-2.5,10.1)
Lower risk _____	0.4	(0.1,0.9)	0.2	(0.0,0.8)	0.2	(0.1,0.6)	0.1	(0.0,1.0)	0.6	(0.2,1.9)	0.3	(-0.5,1.0)	0.4	(-0.3,1.1)
Sensation seeking														
High _____	0.6	(0.3,1.5)	0.8	(0.3,2.5)	0.7	(0.3,1.5)	1.4	(0.5,3.7)	2.0	(0.9,4.3)	1.4	(-0.3,3.1)	1.4	(-0.3,3.0)
Low _____	0.0	(0.0,1.1)	0.3	(0.1,1.3)	0.1	(0.0,0.2)	0.0	(0.0,1.1)	0.4	(0.1,2.5)	0.4	(-0.3,1.0)	0.3	(-0.4,1.0)

Table 4-9. Youth reporting using inhalants in the past month, by age, gender, race/ethnicity, risk score, and sensation seeking (continued)

Characteristics	Percent reporting inhalant use in the past month													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ¹	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 14 to 18														
Gender														
Males _____	0.6	(0.2,1.8)	0.4	(0.1,1.2)	1.0	(0.6,1.7)	0.4	(0.2,0.9)	1.0	(0.6,1.6)	0.3	(-0.5,1.1)	0.0	(-0.8,0.7)
Females _____	0.6	(0.3,1.2)	0.8	(0.4,1.3)	0.3	(0.1,0.7)	0.9	(0.5,1.5)	1.4	(0.7,2.7)	0.8	(-0.2,1.8)	1.1	*(0.2,2.1)
Race/ethnicity														
White _____	0.6	(0.3,1.1)	0.7	(0.4,1.2)	0.7	(0.4,1.2)	0.6	(0.4,1.1)	1.5	(0.9,2.4)	0.9	*(0.0,1.7)	0.8	(0.0,1.5)
African American _____	0.1	(0.0,0.9)	0.5	(0.1,2.5)	0.3	(0.0,2.4)	0.4	(0.1,1.5)	0.4	(0.1,1.8)	0.3	(-0.4,1.0)	0.1	(-0.7,1.0)
Hispanic _____	0.0	(0.0,2.5)	0.3	(0.1,0.9)	0.9	(0.4,2.4)	0.9	(0.3,2.6)	0.8	(0.2,2.9)	0.8	(-0.3,1.9)	-0.1	(-1.6,1.3)
Risk score														
Higher risk _____	1.3	(0.7,2.4)	1.1	(0.6,1.9)	0.9	(0.5,1.5)	1.0	(0.6,1.7)	2.4	(1.5,3.9)	1.2	(-0.2,2.5)	1.5	*(0.3,2.8)
Lower risk _____	0.1	(0.0,0.4)	0.1	(0.1,0.4)	0.2	(0.1,0.7)	0.1	(0.0,0.4)	0.1	(0.0,0.6)	0.1	(-0.2,0.3)	-0.1	(-0.4,0.2)
Sensation seeking														
High _____	1.0	(0.5,1.9)	0.8	(0.4,1.6)	0.9	(0.6,1.5)	0.8	(0.5,1.3)	1.8	(1.1,2.9)	0.8	(-0.3,1.9)	0.9	(-0.1,1.8)
Low _____	0.1	(0.0,0.4)	0.2	(0.0,0.8)	0.2	(0.1,0.7)	0.2	(0.0,0.5)	0.3	(0.1,0.8)	0.2	(0.0,0.5)	0.1	(-0.3,0.5)

¹The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.

Table 4-10. Youth reporting regular inhalant use,¹ by age, gender, race/ethnicity, risk score, and sensation seeking

Characteristics	Percent reporting regular inhalant use													
	Year 2000 (Average for Waves 1 and 2)		Year 2001 (Average for Waves 3 and 4)		Year 2002 (Average for Waves 5 and 6)		Year 2003 (Average for Waves 7 and 8)		Year 2004 (Wave 9 only)		Change from Year 2000 to Year 2004		Change from Year 2002 to Year 2004 ²	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	Est	95% CI	Est	95% CI
Youth aged 12.5 to 18														
12.5 to 13	0.1	(0.0,0.8)	0.0	(0.0,0.7)	0.1	(0.0,0.5)	0.0	(0.0,0.6)	0.0	(0.0,1.0)	-0.1	(-0.4,0.1)	-0.1	(-0.3,0.1)
14 to 15	0.2	(0.0,1.1)	0.2	(0.1,0.9)	0.3	(0.1,0.8)	0.0	(0.0,0.4)	0.3	(0.1,1.2)	0.1	(-0.5,0.7)	0.0	(-0.5,0.5)
16 to 18	0.5	(0.2,1.0)	0.4	(0.2,0.9)	0.1	(0.0,0.6)	0.3	(0.1,0.8)	0.1	(0.0,0.4)	-0.4	*(-0.8,-0.1)	-0.1	(-0.3,0.1)
14 to 16	0.3	(0.1,0.6)	0.2	(0.1,0.7)	0.2	(0.1,0.5)	0.0	(0.0,0.3)	0.3	(0.1,0.8)	0.0	(-0.4,0.4)	0.0	(-0.3,0.4)
14 to 18	0.4	(0.2,0.6)	0.3	(0.2,0.7)	0.2	(0.1,0.5)	0.2	(0.1,0.4)	0.2	(0.1,0.5)	-0.2	(-0.5,0.1)	-0.1	(-0.3,0.2)
12.5 to 18	0.3	(0.2,0.5)	0.2	(0.1,0.5)	0.2	(0.1,0.4)	0.1	(0.0,0.3)	0.1	(0.0,0.4)	-0.2	(-0.4,0.0)	-0.1	(-0.3,0.1)
Youth aged 14 to 18														
Gender														
Males	0.6	(0.3,1.1)	0.4	(0.1,1.0)	0.4	(0.2,0.8)	0.3	(0.1,0.8)	0.1	(0.0,1.1)	-0.5	*(-0.9,0.0)	-0.2	(-0.6,0.2)
Females	0.1	(0.1,0.3)	0.3	(0.1,0.7)	0.1	(0.0,0.4)	0.1	(0.0,0.5)	0.2	(0.1,0.7)	0.1	(-0.2,0.3)	0.1	(-0.2,0.4)
Race/ethnicity														
White	0.6	(0.3,1.0)	0.4	(0.2,0.9)	0.2	(0.1,0.6)	0.1	(0.0,0.4)	0.2	(0.1,0.7)	-0.4	(-0.8,0.1)	0.0	(-0.3,0.3)
African American	0.0	(0.0,2.5)	0.0	(0.0,2.6)	0.2	(0.1,0.6)	0.0	(0.0,1.4)	0.0	(0.0,2.2)	0.0	(N/A)	-0.2	(-0.4,0.0)
Hispanic	0.0	(0.0,2.5)	0.4	(0.1,1.1)	0.4	(0.1,1.9)	0.6	(0.1,2.6)	0.2	(0.0,1.4)	0.2	(-0.2,0.6)	-0.2	(-1.0,0.6)
Risk score														
Higher risk	0.7	(0.4,1.3)	0.6	(0.3,1.3)	0.4	(0.2,0.9)	0.2	(0.1,0.5)	0.3	(0.1,1.1)	-0.4	(-1.0,0.2)	-0.1	(-0.6,0.4)
Lower risk	0.1	(0.0,0.3)	0.1	(0.0,0.3)	0.0	(0.0,0.5)	0.0	(0.0,0.4)	0.0	(0.0,0.7)	-0.1	(-0.2,0.0)	0.0	(N/A)
Sensation seeking														
High	0.6	(0.4,1.1)	0.5	(0.3,1.1)	0.4	(0.2,0.8)	0.1	(0.0,0.4)	0.3	(0.1,0.9)	-0.3	(-0.8,0.1)	-0.1	(-0.5,0.4)
Low	0.0	(0.0,1.0)	0.0	(0.0,0.9)	0.0	(0.0,0.1)	0.0	(0.0,0.5)	0.0	(0.0,0.8)	0.0	(N/A)	0.0	*(-0.1,0.0)

¹Regular use = Used 10 or more times in past year.²The 2002 to 2004 time period is used to cover the period of the Campaign from the launch of the marijuana initiative.

NOTE: All estimates presented in this report were calculated using weights that have been adjusted to reflect 2000 Census data, while weights used in prior reports were based on 1990 Census data. Also, estimates in this report are based on youth aged 12.5 to 18, while estimates in prior reports were based on youth aged 12 to 18. Consequently, results in this report may differ from those presented earlier.