



THE CENTER FOR
FOOD SAFETY



June 2, 2003

Re: Docket #03-019N

03-019N 03-019N-1 Peter T. Jenkins Wenonah Hauter
--

Greetings:

The Center for Food Safety (CFS) and Public Citizen are pleased to submit this public comment on the above-referenced Draft Revised Codex General Standard for Irradiated Foods, which is due to be considered by the Codex Alimentarius Commission at its 26th meeting, 30 June – 7 July in Rome.

CFS is a national, non-profit, membership organization established in 1997 to use science and the law to address increasing concerns over the impacts of the global food production system on human health, animal welfare, and the environment. Public Citizen is a national, non-profit, membership organization established in 1971 that advocates for consumer protection and for government and corporate accountability.

CFS and Public Citizen **oppose** the proposed revision of the Codex standards that would remove the existing 10 kiloGray (kGy) irradiation maximum average absorbed dose limit. Important new information indicates that critical concerns remain unresolved as to the safety of irradiated foods.

The European Union in 1999 commissioned a detailed assessment of the toxicity of several “unique radiolytic products” that have been found to be toxic in various contexts. Our earlier comments disclosed some of these toxicity concerns for 2-alkylcyclobutanones (2-ACBs). The recent EU report, entitled “Toxicological Study to Assess the Risks Associated with the Consumption of Irradiated, Fat-containing Foods,” was prepared over the last four years by a consortium of German and French scientists from recognized institutions. This report, by Burnouf et al, contains major new findings.¹

For example, tumor promotion, which has never been assessed in any other irradiated food animal or human feeding studies, represents a new area of toxicity that cannot be dismissed as already covered:

In an experiment with rats treated with a specific colon carcinogen, it was shown that 2-tDCB and 2-tDeCB have a promoter effect on the development of colon tumors. In this experiment, we found a larger number of aberrant crypts and development of more and larger tumors in the animals that received 2-ACBs in combination with the carcinogen azoxymethane (AOM). Although we did not observe initiation of tumor development by 2-ACBs alone, both the in vitro tests

[S]ince our results point to toxic, genotoxic and even tumor-promoting activity of several 2-ACBs, we consider it necessary that further research, including confirmation of our results by other laboratories, be conducted to permit an assessment of the possible risks associated with consumption of irradiated, fat-containing foods. Unfortunately, these potential risks cannot be assessed at present due to the lack of studies in various areas. In order to characterize the potential risks, the hazards must be identified, and further research is required to precisely determine exposure to these substances, the precise dose-response relationship, and in particular the kinetics and metabolism of 2-ACBs in the living organism. All of this research is necessary to gain insight into the mechanisms of the toxic effects. Numerous questions still remain to be answered, and much research must still be done, before an informed risk assessment can be conducted. However, a start has been made, and we hope to be able to provide answers in the near future.

In response to a review of their report by the EU Scientific Committee on Food in July 2002, the report's authors, Burnouf et al, made a statement to clarify the significance of their work:

[O]ur new data which will be published in peer-reviewed journals, raise some doubts or at least suggest that caution should be exercised before any risk to consumers by exposure to these compounds is denied. At present, knowledge about the potential toxicity of the 2-ACBs (including possible metabolites) and their toxic potency is very limited. Since these compounds are uniquely formed by irradiation and are not inherent in food, in our opinion, complementary studies are needed to make a qualified risk assessment. It needs to be shown that despite the presence of potentially cyto- and genotoxic radiation-induced agents, the consumption of irradiated fat-containing food is safe for consumers.⁴

As the leading researchers to have done any irradiation toxicity assessment in recent decades – and with representation from the well-known food irradiation research program of the Federal Research Center for Nutrition in Karlsruhe, Germany – it is extraordinarily significant that they say that current knowledge is inadequate to show the food is “safe for consumers” and that, pending further research, “risk to consumers” should not be “denied”

Because these new results raise many more questions than they answered, a path of caution must be taken. These statements give no assurance of safety at all, rather they are a clear call for more studies before safety from now clearly-proven potential risks can be assured.

Here is a summary of 2-ACBs studied and the properties detected:

Chapter of Burnouf et al	Particular 2-ACB				
	2-DCB	2-dDCB	2-tDCB	2-dDeCB	2-tDeCB
2.5 – found in ground beef	X	X	X*	X	X
2.6.1 – cyto- and genotoxic to human cells		X	X		X
2.6.2 – cytotoxic/oxidative damage to DNA in human cells	X	X	X		X
2.6.3 – cytotoxic to bacteria	X	X	X		

Comment on Draft Revised Codex Standard for Irradiated Foods

seasonings, where doses up to 30 kGy may be needed to ensure a product in a satisfactory hygienic condition.

On the basis of the information presently supplied to it, the Committee is still of the opinion, that it is appropriate to specify a maximum dose for the treatment of certain food products by ionising radiation and that irradiated foodstuffs should continue to be evaluated individually taking into account the technological need and their safety.⁵

Thus, it is not merely scientists and consumer groups who have raised concerns, but also the EU itself.

Public Citizen and CFS made an earlier joint comment to the Codex CCFAC dated May 14, 2001 (at www.centerforfoodsafety.org/li/commcodx.htm) that raised mutagenicity concerns that go far beyond this recent cyclobutanone debate, based on careful review of decades of scientific articles. That comment demonstrated mistakes in the 1999 FAO/WHO/IAEA Technical Report #890, *High-Dose Irradiation: Wholesomeness of Foods Irradiated Above 10 kGy*. At least 10 positive in vivo published studies that found mutagenic effects in mammals – including one in humans – were misclassified or ignored in that 1999 report, upon which the Codex CCFAC explicitly relied in its preliminary approval of removing the 10 kGy limit. These 10 positive studies compare to only 17 published in vivo studies that were reportedly negative for mutagenicity. Similarly, for published in vitro studies, 5 mutagenicity studies were positive and 8 were negative. Overall, more than one-third of published studies indicate mutagenicity of irradiated food substances. This is hardly a record upon which Codex can assert safety.

Further, evidence suggesting that irradiated foods may not be safe for human consumption has been dismissed and misrepresented by the World Health Organization, the International Atomic Energy Agency and the United Nations' Food and Agriculture Organization.


In 1994, 1995 and 1999, the WHO published the three most important documents since international deliberations over food irradiation policy began in earnest in 1961. These documents culminated in an endorsement that any food could be irradiated at any dose. The agencies arrived at this decision after taking research that revealed health problems in animals that ate irradiated foods, and stating that the research actually revealed no health problems that could be attributed to irradiation. In addition to reclassifying studies that found “adverse effects” as “negative,” many studies that found negative effects were omitted from key reports published later. These discrepancies occurred 52 times.

In 1994, the WHO published a report entitled *Safety and Nutritional Adequacy of Irradiated Food*.⁶ The report stemmed from an FAO/IAEA/WHO meeting held in Geneva two years earlier. The document lists about 150 studies conducted on the safety of irradiated foods, including those involving monkeys, dogs, rabbits, pigs, hamsters, mice, rats and fruit flies. Among these studies, the report lists a wide range of adverse health effects.

In the 1994 report, 11 studies classified as yielding adverse effects were re-classified as negative in an FAO/IAEA/WHO report published in 1999, *High-Dose Irradiation of Food*.⁷ Among these studies, the 1994 report lists a wide range of adverse health effects in animals that ate irradiated


Comment on Draft Revised Codex Standard for Irradiated Foods

Sincerely,



Peter T. Jenkins
Attorney/Policy Analyst
Center for Food Safety

Center for Food Safety
660 Pennsylvania Ave., S.E.
Suite 302
Washington, DC 20003 USA



Wenonah Haute
Director, Critical Mass Energy and
Environment Program

Public Citizen
215 Pennsylvania Ave., S.E.
Third Floor
Washington, DC 20003 USA

¹ Translation was done by William Freese Translations of Mt. Rainier, MD. Mr. Freese has a degree in chemistry and more than 13 years experience translating medical and scientific texts. The translated report is online at: www.bfa-ernaehrung.de/Bfe-Deutsch/Information/bfeber91.htm (2nd 2002 paper). The full citation is: D. Burnouf, H. Delincée, A. Hartwig, E. Marchioni, M. Miesch, F. Raul, D. Werner (2001), Etude toxicologique transfrontalière destinée à évaluer le risque encouru lors de la consommation d'aliments gras ionisés - Toxikologische Untersuchung zur Risikobewertung beim Verzehr von bestrahlten fetthaltigen Lebensmitteln - Eine französisch-deutsche Studie im Grenzraum Oberrhein, Rapport final d'étude Interreg II, projet N° 3.171. BFE-R--02-02, Federal Research Centre for Nutrition, Karlsruhe, Germany.

² Raul, F. et al. "Food-borne radiolytic compounds (2-alkylcyclobutanones) may promote experimental colon carcinogenesis." *Nutrition and Cancer*, 44(2):189-91, 2002.

³ Horvatovich P. et al. "Detection of 2-alkylcyclobutanones, markers for irradiated foods, in adipose tissues of animals fed with these substances." *Journal of Food Protection*, 65(10):1610-3, 2002.

⁴ D. Burnouf, H. Delincée, A. Hartwig, E. Marchioni, M. Miesch, F. Raul, D. Werner. Comment on a statement of the SCF on a report on 2-alkylcyclobutanones.

⁵ "Revision of the opinion of the Scientific Committee on Food on the irradiation of food." Scientific Committee on Food, Health and Consumer Protection Directorate-General, European Commission, SCF/CS/NF/IRR/24 Final, 24 April 2003.

< http://europa.eu.int/comm/food/fs/sc/scf/out193_en.pdf >

⁶ *Review of Data on High Dose (10-70 kGy) Irradiation of Food*. Report of a Consultation, Karlsruhe, Germany, 29 August - 2 September 1994. Geneva: World Health Organization, 1995.

⁷ *High-Dose Irradiation: Wholesomeness of Food Irradiated with Doses Above 10 kGy*. Report of a Joint FAO/IAEA/WHO Study Group, Geneva, 15-20 September 1997. Geneva: World Health Organization, 1999.

⁸ *Review of Data on High Dose (10-70 kGy) Irradiation of Food*. Report of a Consultation, Karlsruhe, Germany, 29 August - 2 September 1994. Geneva: World Health Organization, 1995.