

Estimated Number and Proportion* of AIDS Cases among Female Adults and Adolescents 1985–2006—United States and Dependent Areas








Estimated Number of AIDS Cases and Rates for Female Adults and Adolescents, by Race/Ethnicity 2006—50 States and DC

Race/Ethnicity	Cases	Rate (cases per 100,000 population)
White, not Hispanic	1,659	1.9
Black, not Hispanic	6,391	40.4
Hispanic	1,516	9.5
Asian/Pacific Islander	95	1.6
American Indian/Alaska Native	37	3.6
Total*	9,801	7.8





Proportion of AIDS Cases among Female Adults and Adolescents, by Transmission Category 2006—United States and Dependent Areas





Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.

‡Includes sex with a bisexual male, a person with hemophilia, and HIV-infected transfusion recipient, or an HIV-infected person with an unspecified risk factor.



Proportion of AIDS Cases among Female Adults and Adolescents, by Transmission Category and Age at Diagnosis, 2006—United States and Dependent Areas

Age at diagnosis (in years)

Transmission category	13–19	20–24	25–34	35–44	≥45
	%	<u></u> %		<u></u> %	%
Injection drug use	10	15	19	27	28
High-risk					
heterosexual contact*	52	82	80	72	71
Other/not identified [†]	38	3	1	1	2




Note. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.



^{*} Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

† Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.


AIDS Rates for Female Adults and Adolescents Reported in 2006—United States and Dependent Areas









AIDS Cases among Female Adults and Adolescents Attributed to Injection Drug Use or High-Risk Heterosexual Contact, by Region, 2002–2006—50 States and DC





* Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


Reported AIDS Cases among Female Adults and Adolescents, by Region and Race/Ethnicity, 2006 50 States and DC








Female Adults and Adolescents 15 to 44 Years of Age Reported to be Living with HIV Infection (not AIDS), 2006—50 Areas






*Includes 94 persons reported from areas with confidential name-based HIV infection reporting, but who were residents of areas without confidential name-based HIV infection reporting. Includes 183 persons whose state of residence is unknown or missing.




Female Adults and Adolescents 15 to 44 Years of Age Reported to be Living with AIDS, 2006—United States and Dependent Areas





Proportion of HIV/AIDS Cases among Female Adults and Adolescents, by Transmission Category 2006—33 States






Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection. †Includes blood transfusion, perinatal exposure, and risk factor not reported or not identified.

Proportion of HIV/AIDS Cases and Population among Female Adults and Adolescents, by Race/Ethnicity 2006—33 States







Proportion of HIV/AIDS Cases among Female Adults and Adolescents, by Transmission Category and Age at Diagnosis, 2006—33 States

Age at diagnosis (in years)

Transmission category	13–19	20–24	25–34	35–44	≥45
	%	 %	%	 %	%
Injection drug use	13	16	16	20	21
High-risk heterosexual contact*	86	83	83	78	78
Other/not identified [†]	1	1	1	1	2




Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential named-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and cases without risk factor information were proportionally redistributed.

*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

tlncludes blood transfusion, perinatal exposure, and risk factor not reported or not identified.



Diagnosis Rates of HIV/AIDS for Female Adults and Adolescents, 2006—33 States







Estimated Number of HIV/AIDS Cases and Rates for Female Adults and Adolescents, by Race/Ethnicity 2006—33 States

Race/Ethnicity	Cases	(cases per 100,000 population)
White, not Hispanic	1,664	2.9
Black, not Hispanic	6,033	56.2
Hispanic	1,400	15.1
Asian/Pacific Islander	79	3.2
American Indian/ Alaska Native	35_	4.6
Total*	9,252	11.5



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 33 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.

* Includes 41 female adults and adolescents of unknown race or multiple races.



Rata