

TOXICOLOGICAL PROFILE FOR  
ACRYLONITRILE

Agency for Toxic Substances and Disease Registry  
U.S. Public Health Service

December 1990

**DISCLAIMER**

The use of company or product name(s) is for identification only and does not imply endorsement by the Agency for Toxic Substances and Disease Registry.

## FOREWORD

The Superfund Amendments and Reauthorization Act (SARA) of 1986 (Public Law 99-499) extended and amended the Comprehensive Environmental Response, Compensation, and Liability Act of 1980 (CERCLA or Superfund). This public law directed the Agency for Toxic Substances and Disease Registry (ATSDR) to prepare toxicological profiles for hazardous substances which are most commonly found at facilities on the CERCLA National Priorities List and which pose the most significant potential threat to human health, as determined by ATSDR and the Environmental Protection Agency (EPA). The lists of the 250 most significant hazardous substances were published in the Federal Register on April 17, 1987, on October 20, 1988, on October 26, 1989, and on October 17, 1990.

Section 104(i)(3) of CERCLA, as amended, directs the Administrator of ATSDR to prepare a toxicological profile for each substance on the list. Each profile must include the following content:

- (A) An examination, summary, and interpretation of available toxicological information and epidemiological evaluations on the hazardous substance in order to ascertain the levels of significant human exposure for the substance and the associated acute, subacute, and chronic health effects,
- (B) A determination of whether adequate information on the health effects of each substance is available or in the process of development to determine levels of exposure which present a significant risk to human health of acute, subacute, and chronic health effects, and
- (C) Where appropriate, an identification of toxicological testing needed to identify the types or levels of exposure that may present significant risk of adverse health effects in humans.

This toxicological profile is prepared in accordance with guidelines developed by ATSDR and EPA. The original guidelines were published in the Federal Register on April 17, 1987. Each profile will be revised and republished as necessary, but no less often than every three years, as required by CERCLA, as amended.

The ATSDR toxicological profile is intended to characterize succinctly the toxicological and adverse health effects information for the hazardous substance being described. Each profile identifies and reviews the key literature (that has been peer-reviewed) that describes a hazardous substance's toxicological properties. Other pertinent literature is also presented but described in less detail than the key studies. The profile is not intended to be an exhaustive document; however, more comprehensive sources of specialty information are referenced.

*Foreword*

Each toxicological profile begins with a public health statement, which describes in nontechnical language a substance's relevant toxicological properties. Following the public health statement is information concerning significant health effects associated with exposure to the substance. The adequacy of information to determine a substance's health effects is described. Data needs that are of significance to protection of public health will be identified by ATSDR, the National Toxicology Program (NTP) of the Public Health Service, and EPA. The focus of the profiles is on health and toxicological information; therefore, we have included this information in the beginning of the document.

The principal audiences for the toxicological profiles are health professionals at the federal, state, and local levels, interested private sector organizations and groups, and members of the public.

This profile reflects our assessment of all relevant toxicological testing and information that has been peer reviewed. It has been reviewed by scientists from ATSDR, the Centers for Disease Control, the NTP, and other federal agencies. It has also been reviewed by a panel of nongovernment peer reviewers and is being made available for public review. Final responsibility for the contents and views expressed in this toxicological profile resides with ATSDR.



William L. Roper, M.D., M.P.H.  
Administrator  
Agency for Toxic Substances and  
Disease Registry

## CONTENTS

FOREWORD . . . . .	iii
LIST OF FIGURES . . . . .	ix
LIST OF TABLES . . . . .	xi
1. PUBLIC HEALTH STATEMENT . . . . .	1
1.1 WHAT IS ACRYLONITRILE? . . . . .	1
1.2 HOW MIGHT I BE EXPOSED TO ACRYLONITRILE? . . . . .	2
1.3 HOW CAN ACRYLONITRILE ENTER AND LEAVE MY BODY? . . . . .	2
1.4 HOW CAN ACRYLONITRILE AFFECT MY HEALTH? . . . . .	2
1.5 WHAT LEVELS OF EXPOSURE HAVE RESULTED IN HARMFUL HEALTH EFFECTS? . . . . .	3
1.6 IS THERE A MEDICAL TEST TO DETERMINE WHETHER I HAVE BEEN EXPOSED TO ACRYLONITRILE? . . . . .	4
1.7 WHAT RECOMMENDATIONS HAS THE FEDERAL GOVERNMENT MADE TO PROTECT HUMAN HEALTH? . . . . .	4
1.8 WHERE CAN I GET MORE INFORMATION? . . . . .	9
2. HEALTH EFFECTS . . . . .	11
2.1 INTRODUCTION . . . . .	11
2.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE . . . . .	11
2.2.1 Inhalation Exposure . . . . .	12
2.2.1.1 Death . . . . .	12
2.2.1.2 Systemic Effects . . . . .	18
2.2.1.3 Immunological Effects . . . . .	21
2.2.1.4 Neurological Effects . . . . .	21
2.2.1.5 Developmental Effects . . . . .	22
2.2.1.6 Reproductive Effects . . . . .	22
2.2.1.7 Genotoxic Effects . . . . .	22
2.2.1.8 Cancer . . . . .	23
2.2.2 Oral Exposure . . . . .	24
2.2.2.1 Death . . . . .	24
2.2.2.2 Systemic Effects . . . . .	33
2.2.2.3 Immunological Effects . . . . .	36
2.2.2.4 Neurological Effects . . . . .	36
2.2.2.5 Developmental Effects . . . . .	37
2.2.2.6 Reproductive Effects . . . . .	37
2.2.2.7 Genotoxic Effects . . . . .	38
2.2.2.8 Cancer . . . . .	38
2.2.3 Dermal Exposure . . . . .	39
2.2.3.1 Death . . . . .	39
2.2.3.2 Systemic Effects . . . . .	39
2.2.3.3 Immunological Effects . . . . .	39
2.2.3.4 Neurological Effects . . . . .	41
2.2.3.5 Developmental Effects . . . . .	41
2.2.3.6 Reproductive Effects . . . . .	41
2.2.3.7 Genotoxic Effects . . . . .	41

	2.2.3.8 Cancer . . . . .	41
2.3	TOXICOKINETICS . . . . .	41
	2.3.1 Absorption . . . . .	41
	2.3.1.1 Inhalation Exposure . . . . .	41
	2.3.1.2 Oral Exposure . . . . .	41
	2.3.1.3 Dermal Exposure . . . . .	42
	2.3.2 Distribution . . . . .	42
	2.3.2.1 Inhalation Exposure . . . . .	42
	2.3.2.2 Oral Exposure . . . . .	42
	2.3.2.3 Dermal Exposure . . . . .	43
	2.3.3 Metabolism . . . . .	43
	2.3.4 Excretion . . . . .	46
	2.3.4.1 Inhalation Exposure . . . . .	46
	2.3.4.2 Oral Exposure . . . . .	46
	2.3.4.3 Dermal Exposure . . . . .	46
2.4	RELEVANCE TO PUBLIC HEALTH . . . . .	47
2.5	BIOMARKERS OF EXPOSURE AND EFFECT . . . . .	54
	2.5.1 Biomarkers Used to Identify and/or Quantify Exposure to Acrylonitrile . . . . .	55
	2.5.2 Biomarkers Used to Characterize Effects Caused by Acrylonitrile . . . . .	55
2.6	INTERACTIONS WITH OTHER CHEMICALS . . . . .	56
2.7	POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE . . . . .	57
2.8	ADEQUACY OF THE DATABASE . . . . .	57
	2.8.1 Existing Information on the Health Effects of Acrylonitrile . . . . .	57
	2.8.2 Identification of Data Needs . . . . .	59
	2.8.3 On-going Studies . . . . .	62
3.	CHEMICAL AND PHYSICAL INFORMATION . . . . .	67
	3.1 CHEMICAL IDENTITY . . . . .	67
	3.2 PHYSICAL AND CHEMICAL PROPERTIES . . . . .	67
4.	PRODUCTION, IMPORT, USE, AND DISPOSAL . . . . .	71
	4.1 PRODUCTION . . . . .	71
	4.2 IMPORT . . . . .	71
	4.3 USE . . . . .	71
	4.4 DISPOSAL . . . . .	72
5.	POTENTIAL FOR HUMAN EXPOSURE . . . . .	73
	5.1 OVERVIEW . . . . .	73
	5.2 RELEASES TO THE ENVIRONMENT . . . . .	73
	5.2.1 Air . . . . .	73
	5.2.2 Water . . . . .	75
	5.2.3 Soil . . . . .	75
	5.3 ENVIRONMENTAL FATE . . . . .	75
	5.3.1 Transport and Partitioning . . . . .	75
	5.3.2 Transformation and Degradation . . . . .	76
	5.3.2.1 Air . . . . .	76
	5.3.2.2 Water . . . . .	77
	5.3.2.3 Soil . . . . .	78


5.4	LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT . . . . .	78
5.4.1	Air . . . . .	78
5.4.2	Water . . . . .	78
5.4.3	Soil . . . . .	79
5.4.4	Other Media . . . . .	79
5.5	GENERAL POPULATION AND OCCUPATIONAL EXPOSURE . . . . .	80
5.6	POPULATIONS WITH POTENTIALLY HIGH EXPOSURES . . . . .	80
5.7	ADEQUACY OF THE DATABASE . . . . .	80
5.7.1	Identification of Data Needs . . . . .	82
5.7.2	On-going Studies . . . . .	83
6.	ANALYTICAL METHODS . . . . .	85
6.1	BIOLOGICAL MATERIALS . . . . .	85
6.2	ENVIRONMENTAL SAMPLES . . . . .	85
6.3	ADEQUACY OF THE DATABASE . . . . .	87
6.3.1	Identification of Data Needs . . . . .	87
6.3.2	On-going Studies . . . . .	89
7.	REGULATIONS AND ADVISORIES . . . . .	91
8.	REFERENCES . . . . .	95
9.	GLOSSARY . . . . .	125
	APPENDIX . . . . .	129





LIST OF FIGURES

2-1	Levels of Significant Exposure to Acrylonitrile - Inhalation . . . .	16
2-2	Levels of Significant Exposure to Acrylonitrile - Oral . . . . .	30
2-3	Metabolic Scheme for Acrylonitrile . . . . .	44
2-4	Existing Information on Health Effects of Acrylonitrile . . . . .	58
5-1	Frequency of Sites with Acrylonitrile Contamination . . . . .	74



## LIST OF TABLES

1-1	Human Health Effects from Breathing Acrylonitrile . . . . .	5
1-2	Animal Health Effects from Breathing Acrylonitrile . . . . .	6
1-3	Human Health Effects from Eating or Drinking Acrylonitrile . . . . .	7
1-4	Animal Health Effects from Eating or Drinking Acrylonitrile . . . . .	8
2-1	Levels of Significant Exposure to Acrylonitrile - Inhalation . . . . .	13
2-2	Levels of Significant Exposure to Acrylonitrile - Oral . . . . .	25
2-3	Levels of Significant Exposure to Acrylonitrile - Dermal . . . . .	40
2-4	Genotoxicity of Acrylonitrile <u>In Vitro</u> . . . . .	51
2-5	Genotoxicity of Acrylonitrile <u>In Vivo</u> . . . . .	53
2-6	On-going Studies on Acrylonitrile . . . . .	63
3-1	Chemical Identity of Acrylonitrile . . . . .	68
3-2	Physical and Chemical Properties of Acrylonitrile . . . . .	69
5-1	Estimated Levels of Human Exposure to Acrylonitrile for Nonoccupational and Occupational Exposure . . . . .	81
6-1	Analytical Methods for Determining Acrylonitrile in Biological Materials . . . . .	86
6-2	Analytical Methods for Determining Acrylonitrile in Environmental Samples . . . . .	88
7-1	Regulations and Guidelines Applicable to Acrylonitrile . . . . .	92



## 1. PUBLIC HEALTH STATEMENT

This Statement was prepared to give you information about acrylonitrile and to emphasize the human health effects that may result from exposure to it. The Environmental Protection Agency (EPA) has identified 1,177 sites on its National Priorities List (NPL). Acrylonitrile has been found at 3 of these sites. However, we do not know how many of the 1,177 NPL sites have been evaluated for acrylonitrile. As EPA evaluates more sites, the number of sites at which acrylonitrile is found may change. The information is important for you to know because acrylonitrile may cause harmful health effects and because these sites are potential or actual sources of human exposure to acrylonitrile.

When a chemical is released from a large area, such as an industrial plant, or from a container, such as a drum or bottle, it enters the environment as a chemical emission. This emission, which is also called a release, does not always lead to exposure. You can be exposed to a chemical only when you come into contact with the chemical. You may be exposed to it in the environment by breathing, eating, or drinking substances containing the chemical or from skin contact with it.

If you are exposed to a hazardous substance such as acrylonitrile, several factors will determine whether harmful health effects will occur and what the type and severity of those health effects will be. These factors include the dose (how much), the duration (how long), the route or pathway by which you are exposed (breathing, eating, drinking, or skin contact), the other chemicals to which you are exposed, and your individual characteristics such as age, sex, nutritional status, family traits, life style, and state of health.

### 1.1 WHAT IS ACRYLONITRILE?

Acrylonitrile is a colorless, liquid, man-made chemical with a sharp, onion- or garlic-like odor. Acrylonitrile is used mostly to make plastics, acrylic fibers, and synthetic rubber. Because acrylonitrile evaporates quickly, it is most likely to be found in the air around chemical plants where it is made. Acrylonitrile breaks down quickly in the air. It has been found in small amounts in the water and soil near manufacturing plants and hazardous waste sites. In water, acrylonitrile usually breaks down in about 1 to 2 weeks, although this can vary depending on conditions. For example, high concentrations of acrylonitrile (such as might occur after a spill) tend to be broken down more slowly. In one case, measurable amounts of acrylonitrile were found in nearby wells 1 year after a spill. Further information on the properties and uses of acrylonitrile and how it behaves in the environment may be found in Chapters 3, 4, and 5.

## 1. PUBLIC HEALTH STATEMENT

### 1.2 HOW MIGHT I BE EXPOSED TO ACRYLONITRILE?

Unless you live near a factory where acrylonitrile is made or near a hazardous waste site that contains acrylonitrile, you are unlikely to be exposed to acrylonitrile in the air you breathe or the water you drink. Concentrations of acrylonitrile in average air samples are too low to be measured, and most water samples also have no measurable acrylonitrile. Measurable amounts of acrylonitrile are found primarily near factories and hazardous waste sites. Concentrations in the air near a factory producing or using acrylonitrile average less than 1 part per billion (ppb). Extremely small amounts of acrylonitrile may be found in water near some factories that make or use it, but acrylonitrile rapidly breaks down and disappears from water. Plastic food containers that are made from acrylonitrile are regulated by the Food and Drug Administration such that only 0.17 ppb can enter food; therefore, acrylonitrile intake from food packaging would be extremely low. Because acrylonitrile has been found in water and soil in some hazardous waste sites that contain this chemical, residents living very close to waste sites might possibly be exposed to acrylonitrile by breathing the air or drinking contaminated groundwater.

Further information on how you might be exposed to acrylonitrile is given in Chapter 5.

### 1.3 HOW CAN ACRYLONITRILE ENTER AND LEAVE MY BODY?

Acrylonitrile can enter your body if you breathe its vapors or eat or drink acrylonitrile-contaminated food or water. Acrylonitrile can pass through your skin, but how much gets through is not known. Inside the body, acrylonitrile is broken down into other chemicals, including cyanide. Most of these breakdown products are removed from the body in the urine. Overall, most acrylonitrile is removed from the body within 24 hours, but approximately 25% of what is taken in becomes attached to materials inside cells of the body. More information on how acrylonitrile enters and leaves the body is given in Chapter 2.

### 1.4 HOW CAN ACRYLONITRILE AFFECT MY HEALTH?

The effects of acrylonitrile on your health depend on how much you take into your body and whether you are exposed for a short or long period of time. If the levels of acrylonitrile are high enough, or if the exposure is for a long enough period of time, acrylonitrile can cause death. Small children are more likely to be affected than adults. In several cases, children died following exposures that adults found only mildly irritating. It should be noted that specific levels of acrylonitrile causing death were not reported. Exposure to large amounts of acrylonitrile for a short period of time, as might occur in the case of an industrial accident, results mainly in effects on the

## 1. PUBLIC HEALTH STATEMENT

nervous system. Symptoms can include headache and nausea. At higher concentrations of acrylonitrile there may be temporary damage to red blood cells and the liver. These symptoms disappear when the exposure is stopped.

Direct contact of your skin with acrylonitrile will damage the skin so that it may blister and peel. Exposure of the skin to high concentrations of acrylonitrile in the air may irritate the skin and cause it to turn red. The redness may last for several days.

Long-term exposure to acrylonitrile in air or water may increase your chances of getting cancer. Humans who are repeatedly exposed to acrylonitrile in the workplace for many years may have a higher-than-average chance of developing lung cancer, although this is not clearly established. In animals, exposure to acrylonitrile in the air or in drinking water has been found to increase the number of tumors occurring in the brain, salivary glands, and intestines.

Birth defects have been seen in animals exposed to high concentrations of acrylonitrile in the air or drinking water. Reproductive effects have been seen in animals given acrylonitrile in drinking water for three generations. However, no birth defects or effects on reproduction have been reported in humans.

Further information on the health effects of acrylonitrile in humans and animals can be found in Chapter 2.

### 1.5 WHAT LEVELS OF EXPOSURE HAVE RESULTED IN HARMFUL HEALTH EFFECTS?

In humans, breathing acrylonitrile at a concentration of 16 parts of acrylonitrile per million parts of air (ppm) causes headaches, nausea, and disorientation (Table 1-1). This concentration is close to that at which acrylonitrile can be smelled in air (about 21 ppm). Breathing acrylonitrile in air for long periods of time and at high concentrations can cause death. The actual concentrations of acrylonitrile and breathing times which cause death have not been measured. There is no information on human health effects from eating or drinking acrylonitrile. Acrylonitrile can be smelled at a concentration of 19 ppm when dissolved in water.

In animals, drinking water that contains 142 ppm of acrylonitrile has caused nervous system disorders leading to death (Table 1-4). Birth defects and effects on reproduction have occurred in animals that breathed acrylonitrile in air at levels of 80 ppm or drank it in water at 180 ppm.

## 1. PUBLIC HEALTH STATEMENT

Tables 1-1 through 1-4 show the relationship between exposure to acrylonitrile and known health effects. Short-term and longer-term Minimal Risk Levels (MRLs) are also included in Tables 1-1 and 1-3. These MRLs were derived from animal and human data for both short-term and long-term exposure, as described in Chapter 2 and in Tables 2-1 and 2-2. The MRLs provide a basis for comparison with levels that people might encounter either in the air or in food or drinking water. If a person is exposed to acrylonitrile at an amount below the MRL, it is not expected that harmful (noncancer) health effects will occur. Because these levels are based only on information currently available, some uncertainty is always associated with them. Also, because the method for deriving MRLs does not use any information about cancer, an MRL does not imply anything about the presence, absence, or level of risk for cancer.

Additional information on the levels of exposure associated with harmful effects can be found in Chapter 2.

### **1.6 IS THERE A MEDICAL TEST TO DETERMINE WHETHER I HAVE BEEN EXPOSED TO ACRYLONITRILE?**

There is a test that can detect acrylonitrile in blood. In addition, the major breakdown products of acrylonitrile by the body (termed metabolites) can be measured in urine. Some breakdown products that can be measured are specific to acrylonitrile. However, one breakdown product of the body (cyanide) that is commonly measured is not specific to acrylonitrile exposure, and the results can be affected by cigarette smoking. Because special equipment is needed, these tests cannot be performed routinely in your doctor's office. There is not enough information at present to use the results of such tests to predict the nature or severity of any health effects that may result from exposure to acrylonitrile. Further information on how acrylonitrile can be measured in exposed humans is presented in Chapters 2 and 6.

### **1.7 WHAT RECOMMENDATIONS HAS THE FEDERAL GOVERNMENT MADE TO PROTECT HUMAN HEALTH?**

The federal government has developed regulations and advisories to protect individuals from the potential health effects of acrylonitrile in the environment. The U.S. Environmental Protection Agency (EPA) recommends that acrylonitrile levels in water not exceed 0.058 ppb. Any release to the environment of more than 100 lb must be reported to the federal government. The Occupational Safety and Health Administration (OSHA) has established a legally enforceable maximum limit of 2 ppm in workplace air for an 8-hour exposure over a 40-hour work week.



## 1. PUBLIC HEALTH STATEMENT

TABLE 1-1. Human Health Effects from Breathing Acrylonitrile\*

Short-term Exposure (less than or equal to 14 days)		
<u>Levels in Air (ppm)</u>	<u>Length of Exposure</u>	<u>Description of Effects**</u>
0.1		Minimal Risk Level (based on human studies; see Section 1.5 for discussion)
16	20-45 min	Headaches, nausea, diarrhea, apprehension and redness of the skin.
Long-term Exposure (greater than 14 days)		
<u>Levels in Air</u>	<u>Length of Exposure</u>	<u>Description of Effects</u>
		The health effects resulting from long-term exposure of humans to air containing specific levels of acrylonitrile are not known.

\*See Section 1.2 for a discussion of exposures encountered in daily life.

\*\*These effects are listed at the lowest level at which they were first observed. They may also be seen at higher levels.

## 1. PUBLIC HEALTH STATEMENT

TABLE 1-2. Animal Health Effects from Breathing Acrylonitrile

Short-term Exposure (less than or equal to 14 days)		
<u>Levels in Air (ppm)</u>	<u>Length of Exposure</u>	<u>Description of Effects*</u>
30	4 hours	Excessive watering of the mouth in dogs.
65	4 hours	Death in dogs.
80	10 days	Birth defects in rats.
90	4 hours	Reddened skin in monkeys.
Long-term Exposure (greater than 14 days)		
<u>Levels in Air (ppm)</u>	<u>Length of Exposure</u>	<u>Description of Effects*</u>
20	2 years	Premature death in some rats.
80	2 years	Brain tumors in rats.

\*These effects are listed at the lowest level at which they were first observed. They may also be seen at higher levels.

## 1. PUBLIC HEALTH STATEMENT

TABLE 1-3. Human Health Effects from Eating or Drinking Acrylonitrile\*

Short-term Exposure (less than or equal to 14 days)		
<u>Levels in Food</u>	<u>Length of Exposure</u>	<u>Description of Effects</u>
		The health effects resulting from short-term exposure of humans to food containing specific levels of acrylonitrile are not known.
<u>Levels in Water (ppm)</u> 3.0		Minimal Risk Level (based on animal studies; see Section 1.5 for discussion).
Long-term Exposure (greater than 14 days)		
<u>Levels in Food</u>	<u>Length of Exposure</u>	<u>Description of Effects</u>
		The health effects resulting from long-term exposure of humans to food containing specific levels of acrylonitrile are not known.
<u>Levels in Water (ppm)</u> 1.4		Minimal Risk Level (based on animal studies; see Section 1.5 for discussion).

\*See Section 1.2 for a discussion of exposures encountered in daily life.

## 1. PUBLIC HEALTH STATEMENT

TABLE 1-4. Animal Health Effects from Eating or Drinking Acrylonitrile

Short-term Exposure (less than or equal to 14 days)		
<u>Levels in Food</u>	<u>Length of Exposure</u>	<u>Description of Effects*</u>
		The health effects resulting from short-term exposure of animals to food containing specific levels of acrylonitrile are not known.
<u>Levels in Water (ppm)</u>		
142	1 day	Death and nervous system disorders in mice.
180	10 days	Birth defects in rats.
Long-term Exposure (greater than 14 days)		
<u>Levels in Food</u>	<u>Length of Exposure</u>	<u>Description of Effects*</u>
		The health effects resulting from short-term exposure of animals to food containing specific levels of acrylonitrile are not known.
<u>Levels in Water (ppm)</u>		
35	2 years	Premature death in rats.
52	60 days	Low sperm count in mice.
100	19 months	Low red blood cell counts.
200	6 months	Ulcers in the throat and premature death in dogs.
500	220 days	Reduced reproductive capability in rats.

\*These effects are listed at the lowest level at which they were first observed. They may also be seen at higher levels.

## 1. PUBLIC HEALTH STATEMENT

Additional information on governmental regulations can be found Chapter 7.

**1.8 WHERE CAN I GET MORE INFORMATION?**

If you have any more questions or concerns not covered here, please contact your State Health or Environmental Department or:

Agency for Toxic Substances and Disease Registry  
Division of Toxicology  
1600 Clifton Road, E-29  
Atlanta, Georgia 30333

This agency can also give you information on the location of the nearest occupational and environmental health clinics. Such clinics specialize in recognizing, evaluating, and treating illnesses that result from exposure to hazardous substances.



## 2. HEALTH EFFECTS

### 2.1 INTRODUCTION

This chapter contains descriptions and evaluations of studies and interpretation of data on the health effects associated with exposure to acrylonitrile. Its purpose is to present levels of significant exposure for acrylonitrile based on toxicological studies, epidemiological investigations, and environmental exposure data. This information is presented to provide public health officials, physicians, toxicologists, and other interested individuals and groups with (1) an overall perspective of the toxicology of acrylonitrile and (2) a depiction of significant exposure levels associated with various adverse health effects.

### 2.2 DISCUSSION OF HEALTH EFFECTS BY ROUTE OF EXPOSURE

To help public health professionals address the needs of persons living or working near hazardous waste sites, the data in this section are organized first by route of exposure -- inhalation, oral, and dermal -- and then by health effect -- death, systemic, immunological, neurological, developmental, reproductive, genotoxic, and carcinogenic effects. These data are discussed in terms of three exposure periods -- acute, intermediate, and chronic.

Levels of significant exposure for each exposure route and duration (for which data exist) are presented in tables and illustrated in figures. The points in the figures showing no-observed-adverse-effect levels (NOAELs) or lowest-observed-adverse-effect levels (LOAELs) reflect the actual doses (levels of exposure) used in the studies. LOAELs have been classified into "less serious" or "serious" effects. These distinctions are intended to help the users of the document identify the levels of exposure at which adverse health effects start to appear, determine whether or not the intensity of the effects varies with dose and/or duration, and place into perspective the possible significance of these effects to human health.

The significance of the exposure levels shown on the tables and figures may differ depending on the user's perspective. For example, physicians concerned with the interpretation of clinical findings in exposed persons or with the identification of persons with the potential to develop such disease may be interested in levels of exposure associated with "serious" effects. Public health officials and project managers concerned with response actions at Superfund sites may want information on levels of exposure associated with more subtle effects in humans or animals (LOAELs) or exposure levels below which no adverse effects (NOAELs) have been observed. Estimates of levels posing minimal risk to humans (Minimal Risk Levels, MRLs) are of interest to health professionals and citizens alike.

## 2. HEALTH EFFECTS

For certain chemicals, levels of exposure associated with carcinogenic effects may be indicated in the figures. These levels reflect the actual doses associated with the tumor incidences reported in the studies cited. Because cancer effects could occur at lower exposure levels, the figures also show estimated excess risks, ranging from a risk of one in 10,000 to one in 10,000,000 ( $10^{-4}$  to  $10^{-7}$ ), as developed by EPA.

Estimates of exposure levels posing minimal risk to humans (MRLs) have been made, where data were believed reliable, for the most sensitive noncancer end point for each exposure duration. MRLs include adjustments to reflect human variability and, where appropriate, the uncertainty of extrapolating from laboratory animal data to humans. Although methods have been established to derive these levels (Barnes et al. 1987; EPA 1989a), uncertainties are associated with the techniques. Furthermore, ATSDR acknowledges additional uncertainties inherent in the application of these procedures to derive less than lifetime MRLs. As an example, acute inhalation MRLs may not be protective for health effects that are delayed in development or are acquired following repeated acute insults, such as hypersensitivity reactions, asthma, or chronic bronchitis. As these kinds of health effects data become available and methods to assess levels of significant human exposure improve, these MRLs will be revised.

### 2.2.1 Inhalation Exposure

Table 2-1 and Figure 2-1 summarize quantitative data on the health effects observed in humans and laboratory animals exposed to acrylonitrile by inhalation.

#### 2.2.1.1 Death

In humans, the death of a child (age 3) who was exposed by sleeping in a room that had been fumigated with acrylonitrile has been described by Grunske (1949). Respiratory malfunction, lip cyanosis and tachycardia were among the symptoms described prior to death. Five adults who spent the night in the same room complained only of eye irritation or showed no signs of acrylonitrile poisoning. The concentrations of acrylonitrile in the air were not reported. Several other instances of death in children with only mild irritation in adults were reported by Grunske (1949), but not described in detail.

Data on the NOAELS and LOAELS for death are presented in Table 2-1 for several animal species. The data presented indicate that species differences exist with respect to acute lethal effects. Dogs appear to be the most susceptible species, but this is based on studies involving only a few animals. The cause of death varied among test species. In guinea pigs, death resulted from pulmonary irritation (see



TABLE 2-1. Levels of Significant Exposure to Acrylonitrile - Inhalation

Figure Key	Species	Exposure Frequency/ Duration	Effect	NOAEL (ppm)	LOAEL (Effect)		Reference
					Less Serious (ppm)	Serious (ppm)	
ACUTE EXPOSURE							
Death							
1	Rat	4 hr		130		315	Dudley and Neal 1942
2	Gn Pig	4 hr		265		575	Dudley and Neal 1942
3	Rabbit	4 hr		135		260	Dudley and Neal 1942
4	Dog	4 hr		30		65 <sup>a</sup>	Dudley and Neal 1942
5	Cat	4 hr		275		600	Dudley and Neal 1942
6	Monkey	4 hr		90			Dudley and Neal 1942
Systemic							
7	Human	20-45 min (occup)	Derm/Oc		16 (skin irritation)		Wilson et al. 1948
8	Rat	5 d 8hr/d	Renal	129			Gut et al. 1984
9	Rat	5 d 8hr/d	Hepatic	125			Gut et al. 1985
10	Rat	12 hr	Hepatic	26			Gut et al. 1984
11	Rats	4 hr	Renal	100	200 (glucosuria, proteinuria)		Rouisse et al. 1986
12	Rat	5 d 8hr/d	Hepatic		129 (lower liver wt.)		Gut et al. 1984
13	Rat	5 d 8hr/d	Resp	129			Gut et al. 1984

TABLE 2-1 (Continued)

Figure Key	Species	Exposure Frequency/ Duration	Effect	NOAEL (ppm)	LOAEL (Effect)		Reference
					Less Serious (ppm)	Serious (ppm)	
14	Rat	5 d 8hr/d	Other			125 (unhealthy appearance)	Gut et al. 1985
15	Rat	4 hr	Derm/Oc		315 (skin redness)		Dudley and Neal 1942
16	Gn Pig	4 hr	Derm/Oc	100	575 (irritation)		Dudley and Neal 1942
17	Rabbit	4 hr	Derm/Oc		100 (skin redness)		Dudley and Neal 1942
18	Monkey	4 hr	Derm/Oc	65	90 <sup>a</sup> (skin redness)		Dudley and Neal 1942
Neurological							
19	Human	20-45 min (occup)			16 (irritability)		Wilson et al. 1948
20	Human	8 hr		4.6 <sup>b</sup>			Jakubowski et al. 1987
21	Dog	4 hr			30 <sup>a</sup> (salivation)	100 (paralysis)	Dudley and Neal 1942
22	Cat	4 hr			100 (salivation)	275 (pain)	Dudley and Neal 1942
23	Monkey	4 hr		65	90 (weakness)		Dudley and Neal 1942
Developmental							
24	Rat	10 d Gd6-15 6hr/d		40		80 <sup>a</sup> (malformations)	Murray et al. 1978
CHRONIC EXPOSURE							
Death							
25	Rat	2 yr 5d/wk 6hr/d				20 <sup>a</sup> (early deaths)	Quast et al 1980a


TABLE 2-1 (Continued)

Figure Key	Species	Exposure Frequency/ Duration	Effect	NOAEL (ppm)	LOAEL (Effect)		Reference
					Less Serious (ppm)	Serious (ppm)	
Systemic							
26	Rat	2 yr 5d/wk 6hr/d	Hemato	80			Quast et al 1980a
27	Rat	2 yr 5d/wk 6hr/d	Resp	20	80 (irritation nasal mucosa)		Quast et al 1980a
28	Rat	2 yr 5d/wk 6hr/d	Hepatic	80			Quast et al 1980a
29	Rat	2 yr 5d/wk 6hr/d	Renal	80			Quast et al 1980a
Neurological							
30	Rat	2 yr 5d/wk 6hr/d				80 <sup>a</sup> (focal gliosis)	Quast et al 1980a
Cancer							
31	Rat	104 wk 5d/wk 7hr/d				60 CEL (multiple tumors)	Maltoni et al. 1988
32	Rat	2 yr 5d/wk 6hr/d				20 CEL (multiple tumors)	Quast et al 1980a

<sup>a</sup>Presented in Table 1-2.

<sup>b</sup>Used to derive acute inhalation MRL; dose adjusted for intermittent exposure and divided by an uncertainty factor of 10 (for human variability), resulting in an MRL of 0.1 ppm. This MRL is presented in Table 1-1.

LOAEL = lowest-observed-adverse-effect level; NOAEL = no-observed-adverse-effect level; ppm = parts per million; hr = hour; derm/oc = dermal/ocular; min = minute; occup = occupational; d = day; wt = weight; Resp = respiratory; Gd = gestation day; yr = year; hemato = hematological; CEL = cancer effect level.



**FIGURE 2-1. Levels of Significant Exposure to Acrylonitrile – Inhalation**




FIGURE 2-1 (Continued)

## 2. HEALTH EFFECTS

Section 2.2.1.2) while in the other species convulsions and coma occurred (see Section 2.2.1.4) (Dudley and Neal 1942). It should be noted that this study is based on nominal concentrations without analytical verification.

Chronic exposure to acrylonitrile has been reported to result in early deaths in male and female rats (Quast et al. 1980a). A statistically significant increase in mortality was observed within the first year of a study in which animals were exposed to 80 ppm of acrylonitrile. At 20 ppm, increased deaths were noted in females during the last 10 weeks of the study. The cause of death was not specifically identified.

The highest NOAEL values and all reliable LOAEL values for death in each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

### 2.2.1.2 Systemic Effects

**Respiratory Effects.** In workers exposed to acrylonitrile at concentrations of 16 to 100 ppm for periods of 20 to 45 minutes, irritation of the nose and throat and a feeling of fullness in the chest was reported by Wilson et al. (1948).

Acute effects on the respiratory tract of animals demonstrate species differences. In guinea pigs exposed to 575 ppm for 4 hours, marked irritation of the respiratory tract was evidenced by coughing and nasal exudate, with delayed death from lung edema (Dudley and Neal 1942). In other species (rats, rabbits, dogs and monkeys), death occurred at lower doses than in guinea pigs, but was not related to respiratory effects. In these animals, mild irritation of the respiratory tract and effects resembling cyanide poisoning were noted. Respiration was initially stimulated, but then followed by rapid shallow breathing (Dudley and Neal 1942).

Reported exposure at lower doses shows no acrylonitrile-related effects on the respiratory system. For example, rats repeatedly exposed for 5 days to 129 ppm of acrylonitrile showed no gross or histological changes in lung tissue (Gut et al. 1984). However, chronic exposure of rats to acrylonitrile appears to cause irritation of the nasal passages. In a 2-year study, histopathological evaluation of tissues of the respiratory system revealed degenerative and inflammatory changes of the nasal turbinates at 80 ppm (Quast et al. 1980a). These effects were not noted at 20 ppm.

## 2. HEALTH EFFECTS

The highest NOAEL values and all reliable LOAEL values for respiratory effects in each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

**Cardiovascular Effects.** In humans, tachycardia was among the symptoms described in a child (age 3) who was exposed by sleeping in a room that had been fumigated with acrylonitrile. The child died as a result of the exposure (see Section 2.2.1.1) (Grunske 1949). No studies were located regarding cardiovascular effects in animals following inhalation exposure to acrylonitrile.

**Gastrointestinal Effects.** Though tumors of the small intestine were noted in a comprehensive and well-conducted study by Quast et al. (1980a) (see Section 2.2.1.8), there were no nonneoplastic histopathologic changes observed in any portion of the gastrointestinal tract of rats at doses up to 80 ppm for 2 years.

**Hematological Effects.** Humans exposed to acrylonitrile at concentrations where nausea, vomiting and weakness occurred (16 to 100 ppm for 20 to 45 minutes), were also reported to have low grade anemia. However, complete recovery after cessation of exposure was reported (Wilson 1944; Wilson et al. 1948). No adverse hematological effects were detected in Japanese workers exposed to acrylonitrile for 10 to 13 years at exposure levels averaging 2.1 to 14.1 ppm (Sakurai et al. 1978).

In a chronic study in rats (Quast et al. 1980a), some changes in the blood parameters measured were observed at various intervals during the study, but the findings did not occur consistently and were not dose-related. Therefore, the authors concluded that these findings were not direct effects of exposure to acrylonitrile, but rather were a secondary response to other effects such as weight loss, tumor formation, or inflammatory reactions.

The highest NOAEL values and all reliable LOAEL values for hematological effects in each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

**Musculoskeletal Effects.** No studies were located regarding musculoskeletal effects in humans or animals following inhalation exposure to acrylonitrile.

**Hepatic Effects.** Acrylonitrile is metabolized in the liver to potentially toxic metabolites (see Section 2.3). There are limited indications that the liver is a target organ for acrylonitrile toxicity.

## 2. HEALTH EFFECTS

In humans, mild jaundice lasting several days to 4 weeks has been observed after acute occupational exposure to acrylonitrile vapors at high concentrations (Wilson 1944); however, the concentrations of acrylonitrile to which workers were exposed were not reported. The effects were fully reversible. In factory workers exposed to acrylonitrile for 10 years or more, Sakurai et al. (1978) reported an increase in palpable livers of workers. However, the authors considered these results to be inconclusive because the increase was not statistically significant and subjective judgments were involved. Also, blood chemistry evaluations did not indicate liver damage.

In animals, acrylonitrile does not appear to cause damage to the liver following acute or chronic inhalation exposure, though some biochemical changes are noted. Exposure of rats for 5 days to 129 ppm of acrylonitrile resulted in slightly lower liver weight, but no histopathological changes were found (Gut et al. 1984). Depletion of glutathione and effects on carbohydrate and lipid metabolism were observed (Gut et al. 1984, 1985). These effects would be expected considering the affinity of acrylonitrile for sulfhydryl groups on proteins and the involvement of glutathione in the metabolism of acrylonitrile (see Section 2.6). A 2-year chronic study in rats showed no liver injury as evaluated by serum enzyme activity and histopathological evaluation of the tissue (Quast et al. 1980a).

The highest NOAEL values and all reliable LOAEL values for hepatic effects in each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

**Renal Effects.** Most studies indicate that inhalation exposure to acrylonitrile does not result in significant kidney injury. For example, physical examination of workers exposed to acrylonitrile vapors in the workplace for 10 or more years provided no indication of renal effects (Sakurai et al. 1978). In animals, no histological or biochemical signs of renal injury were seen following exposure of rats to 129 ppm of acrylonitrile for 5 days (Gut et al. 1984), or to 80 ppm for 2 years (Quast et al. 1980a). Small increases in urinary levels of glucose, gamma-glutamyl transpeptidase, and N-acetyl-glucosaminidase were observed in rats exposed to 200 ppm of acrylonitrile for 4 hours (Rouisse et al. 1986), but this was not accompanied by any significant effect on urinary creatinine or blood urea nitrogen. No significant effects were noted at 100 ppm for this duration of exposure.

The highest NOAEL values and all reliable LOAEL values for renal effects in each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.



## 2. HEALTH EFFECTS

**Dermal/Ocular Effects.** In humans, direct skin irritation resulting from exposure to acrylonitrile vapors has been observed (see Section 2.2.3).

A skin redness reported in experimental animals (rats, rabbits, cats and monkeys) after inhalation exposure to acrylonitrile may be due to a vasodilatory effect, rather than a direct irritant action (Ahmed and Patel 1981). Guinea pigs, which do not exhibit the cyanide-type effects of acrylonitrile poisoning (see Section 2.2.1.4), were observed to have nose and eye irritation from the acrylonitrile vapors (Dudley and Neal 1942).

The highest NOAEL values and all reliable LOAEL values for dermal/ocular effects in each species and duration category are recorded in Table 2-1 and plotted in Figure 2-1.

### 2.2.1.3 Immunological Effects

No studies were located regarding immunologic effects in humans or animals following inhalation exposure to acrylonitrile.

### 2.2.1.4 Neurological Effects

In humans, acute exposure to acrylonitrile results in characteristics of cyanide-type toxicity. Symptoms in humans associated with acrylonitrile poisoning include limb weakness, labored and irregular breathing, dizziness and impaired judgment, cyanosis, nausea, collapse, and convulsions (Baxter 1979). However, the doses that produce these effects were not clearly defined. Workers exposed to 16 to 100 ppm for 20 to 45 minutes complained of headaches and nausea, apprehension and nervous irritation (Wilson et al. 1948). The workers exposed to acrylonitrile vapors fully recovered. In a study with human volunteers exposed to acrylonitrile at doses of 2.3 and 4.6 ppm, no symptoms attributable to effects on the nervous system were observed (Jakubowski et al. 1987). The dose of 4.6 ppm was used to calculate the acute inhalation MRL of 0.1 ppm, as described in the footnote in Table 2-1.

Animals exposed to acrylonitrile usually display acute neurological symptoms. Dogs appear to be particularly sensitive. Excessive salivation in dogs occurred at concentrations of 30 ppm of acrylonitrile, and paralysis of the hind limbs occurred at 100 ppm. In cats, excessive salivation occurred at 100 ppm; at 275 ppm animals were observed to paw their heads and stomachs and howl as if in pain. In monkeys, "weakness" was reported at 90 ppm, but higher doses were not tested. Guinea pigs showed no measurable signs of neurological effects from acute exposure to acrylonitrile at a dose that caused death

## 2. HEALTH EFFECTS

(575 ppm) (Dudley and Neal 1942). It should be noted that this study is based on nominal concentrations with no analytical verification. Therefore, the overall usefulness of the study is limited.

Chronic exposure of rats to 80 ppm acrylonitrile resulted in focal gliosis and perivascular cuffing in the brain (Quast et al. 1980a). The gliosis appeared to be a pre-malignant lesion related to the formation of brain tumors, and is discussed in more detail in Section 2.2.1.8 (below).

The highest NOAEL values and all reliable LOAEL values for neurological effects are recorded in Table 2-1 and plotted Figure 2-1.

### **2.2.1.5 Developmental Effects**

Inhalation exposure to acrylonitrile results in teratogenic effects in rats. Inhalation of 80 ppm acrylonitrile during days 6 to 15 of gestation (the critical period of organogenesis) resulted in a significant increase in fetal malformations. These malformations included short tail, missing vertebrae, short trunk, omphalocele and hemivertebra. In this well-conducted study, mean number of implantations, live fetus and resorptions were not significantly altered by exposure to 40 or 80 ppm of acrylonitrile. No effects on fetal body size were evident. Maternal toxicity was observed at both dose levels tested (40 and 80 ppm), as evidenced by decreased weight gain (Murray et al. 1978).

### **2.2.1.6 Reproductive Effects**

No studies were located regarding reproductive effects in humans or animals following inhalation exposure to acrylonitrile.

### **2.2.1.7 Genotoxic Effects**

Factory workers exposed for an average of 15 years to acrylonitrile vapors showed no increase in chromosomal aberrations in the peripheral lymphocytes (Thiess and Fleig 1978). As in most human studies, the actual concentration of acrylonitrile to which these workers were exposed was not reported. However, monitoring data indicated that the average exposure concentration for the workers was 5 ppm for the majority of the exposure period (approximately 10 years); at the time the study was conducted, acrylonitrile levels in the workplace had been reduced to 1.5 ppm.

No studies were located regarding genotoxic effects in animals following inhalation exposure to acrylonitrile.

## 2. HEALTH EFFECTS

### 2.2.1.8 Cancer

A number of epidemiological studies have been conducted to evaluate the association between lung cancer and occupational exposure to acrylonitrile (Collins et al. 1989; Delzell and Monson 1982; Kiesselbach et al. 1979; O'Berg 1980; O'Berg et al. 1985). However, many of the studies suffer from deficiencies such as an insufficient quantification of exposure, short follow-up, small and relatively youthful cohorts or lack of consideration of the effects of smoking, and the results of the studies are often inconsistent.

The most reliable of the epidemiology studies on acrylonitrile is that conducted by O'Berg (1980) in which 1,345 male workers in a textile factory were followed over a period of 25 years. Exposures to acrylonitrile were divided into three groups (low, medium and high), although quantitative estimates of exposure levels were not assigned. O'Berg (1980) concluded that there may be an association between human lung cancer and acrylonitrile inhalation exposure. The incidence of cancer did not increase significantly in exposed workers over the unexposed population. The data from this on-going study were updated by O'Berg et al. (1985) to include an additional 7 years of follow-up. Though an increased incidence of lung cancer in workers exposed to acrylonitrile was observed, the effect was not as pronounced as it had been in the previous report. The authors concluded that continued follow-up was needed.

Statistically significant cancer excesses have been observed in other studies; however, due to limitations in the studies firm conclusions cannot be made. In an analysis of mortality among 327 rubber workers, nine deaths from lung cancer were observed compared to 4.7 to 5.9 expected deaths among the general population and other rubber workers (Delzell and Monson 1982). The observed and expected numbers of death were small and the study did not evaluate cigarette smoking as a confounding factor contributing to excess death.

Another well-designed epidemiology study (Collins et al. 1989) did not find an increased risk of lung cancer from acrylonitrile exposure. In this study, 2,671 males (1,774 with known acrylonitrile exposure) were followed for 32 years. Workers in acrylic fiber manufacturing plants were categorized into four dose ranges (expressed in ppm/year). Smoking histories were taken into consideration. No significant relationship between lung cancer and acrylonitrile exposure was established, although the authors found some evidence in this study of confounding association between smoking status and cumulative exposure to acrylonitrile. In this study, as in the previously discussed studies (O'Berg 1980; O'Berg et al. 1985), the subjects were relatively young,

## 2. HEALTH EFFECTS

and only 9% of the cohort had died. Additional follow-up will be necessary to determine whether death can be associated with cancers resulting from acrylonitrile exposure.

Epidemiology studies have also suggested a possible association between acrylonitrile exposure and prostate cancer (Chen et al. 1987; O'Berg et al. 1985), but the data are too limited to warrant any firm conclusion. Collins et al. (1989) found no increase in prostate cancer in acrylonitrile-exposed workers, but again the data are too limited to be meaningful.

Deaths from cancer of the stomach were statistically elevated in one study of six manufacturing plants in the United Kingdom involved in the polymerization of acrylonitrile and the spinning of acrylic fibers (Werner and Carter 1981). No quantitative data on the exposure levels to acrylonitrile were given and the numbers of expected deaths was too small to provide confidence in the results.

In animals, chronic studies provide convincing evidence that acrylonitrile is carcinogenic when administered by the inhalation route of exposure. Tumors in the brain, Zymbal gland and mammary glands in rats exposed to acrylonitrile for 2 years were reported by both Quast et al. (1980a) and Maltoni et al. (1988). Quast et al. (1980a) reported two types of lesions in the central nervous system: astrocytomas and focal or multifocal glial cell proliferation. The glial cell proliferation was considered to be a microscopic change suggestive of early tumorigenesis. Brain tumors were observed in males exposed to 80 ppm of acrylonitrile, and in females at 20 and 80 ppm. In addition, Quast et al. (1980a) reported tumors of the tongue and small intestine in males (80 ppm). Maltoni et al. (1988) noted tumors in the liver (60 ppm).

### 2.2.2 Oral Exposure

No studies were located regarding health effects in humans associated with the oral ingestion of acrylonitrile. Table 2-2 and Figure 2-2 summarize the health effects observed in experimental animals following oral exposure to acrylonitrile. These effects are discussed below.

#### 2.2.2.1 Death

The acute oral LD<sub>50</sub> has been estimated to be 93 mg acrylonitrile/kg body weight (mg/kg) in rats (Back et al. 1972) and 27 mg/kg for mice (Ahmed and Patel 1981). Longer term intake (6 months) in dogs at 16 mg/kg/day resulted in early deaths. Animals lost weight and were described as depressed, lethargic, weak and emaciated prior to death.

TABLE 2-2. Levels of Significant Exposure to Acrylonitrile - Oral

Figure Key	Species	Route	Exposure Frequency/ Duration	Effect	NOAEL (mg/kg/day)	LOAEL (Effect)		Reference
						Less Serious (mg/kg/day)	Serious (mg/kg/day)	
ACUTE EXPOSURE								
Death								
1	Rat		ND				93 (LD50)	Back et al. 1972
2	Mouse	(G)	ND				27 <sup>a</sup> (LD50)	Ahmed and Patel 1981
Systemic								
3	Rat	(G)	1x	Gastro		50 (GI bleeding)		Ghanayem and Ahmed 1983
4	Rat	(G)	10 d Gd6-15 1x/d	Hepatic	25	65 (incr. liver wt.)		Murray et al. 1978
5	Rat	(G)	1x	Gastro		46.5 (covalent binding to proteins)		Farooqui and Ahmed 1983
6	Rat	(G)	10 d Gd6-15 1x/d	Gastro	10	25 (stomach effects)		Murray et al. 1978
7	Rat	(G)	1x	Hemato		80 (RBC enzyme alteration)		Farooqui and Ahmed 1983
Neurological								
8	Rat	(G)	1x				93 (CNS effects)	Ahmed and Patel 1981
9	Mouse	(G)	1x				27 (CNS effects)	Ahmed and Patel 1981
Developmental								
10	Rat	(G)	10 d Gd6-15 1x/d		10 <sup>b</sup>		25 <sup>c</sup> (malformations)	Murray et al. 1978

TABLE 2-2 (Continued)

Figure Key	Species	Route	Exposure Frequency/ Duration	Effect	NOAEL (mg/kg/day)	LOAEL (Effect)		Reference
						Less Serious (mg/kg/day)	Serious (mg/kg/day)	
INTERMEDIATE EXPOSURE								
Death								
11	Dog	(W)	6 mo		10		16	Quast et al. 1975
Systemic								
12	Rat	(W)	50 wk	Gastro	70			Beliles et al. 1980
13	Rat	(W)	3 wk	Other	14	70 (adrenal atrophy)		Szabo et al. 1984
14	Rat	(W)	21 d	Hepatic	14	70 (incr. GSH)		Szabo et al. 1977
15	Dog	(W)	6 mo	Gastro	10		16 <sup>d</sup> (esophageal ulcerations)	Quast et al. 1975
16	Dog	(W)	6 mo	Renal	18			Quast et al. 1975
17	Dog	(W)	6 mo	Hemato	10	16 (decr. RBC)		Quast et al. 1975
Neurological								
18	Rat	(W)	50 wk		70			Beliles et al. 1980
Reproductive								
19	Dog	(W)	6 mo		10		16 (depression, lethargy)	Quast et al. 1975
20	Rat	(W)	220 d		14		70 <sup>e</sup> (decr. reproduct. indices)	Beliles et al. 1980
21	Mouse	(G)	60 d 1x/d				10 <sup>f</sup> (decr. sperm count, tubular degeneration)	Tandon et al. 1988

TABLE 2-2 (Continued)

Figure Key	Species	Route	Exposure Frequency/ Duration	Effect	NOAEL (mg/kg/day)	LOAEL (Effect)		Reference
						Less Serious (mg/kg/day)	Serious (mg/kg/day)	
Cancer								
22	Rat	(W)	50 wk				14 CEL (multiple tumors)	Beliles et al. 1980
CHRONIC EXPOSURE								
Death								
23	Rat	(W)	M:26 mo F:23 mo		4.2		14 (early deaths)	Bio/dynamics 1980b
24	Rat	(G)	20 mo 7d/wk 1x/d		0.1		10 (early deaths)	Bio/dynamics 1980c
25	Rat	(W)	M:22 mo F:19 mo		0.14		14 (early deaths)	Bio/dynamics 1980a
26	Rat	(W)	2 yr				4.4 <sup>o</sup> (early deaths)	Quast et al. 1980b
Systemic								
27	Rat	(W)	M:22 mo F:19 mo	Hemato	0.14	14 (decr. red cells)		Bio/dynamics 1980a
28	Rat	(W)	M:26 mo F:23 mo	Hepatic	4.2	14 (incr. liver wt.)		Bio/dynamics 1980b
29	Rat	(W)	M:22 mo F:19 mo	Renal		14 (incr. kidney wt.)		Bio/dynamics 1980a
30	Rat	(W)	M:22 mo F:19 mo	Cardio	14			Bio/dynamics 1980a
31	Rat	(G)	20 mo 7d/wk 1x/d	Hemato		10 (decr. red cells)		Bio/dynamics 1980c

TABLE 2-2 (Continued)

Figure Key	Species	Route	Exposure Frequency/ Duration	Effect	NOAEL (mg/kg/day)	LOAEL (Effect)		Reference
						Less Serious (mg/kg/day)	Serious (mg/kg/day)	
32	Rat	(G)	20 mo 7d/wk 1x/d	Hepatic	0.1	10 (incr. liver wt.)		Bio/dynamics 1980c
33	Rat	(G)	20 mo 7d/wk 1x/d	Renal	0.1	10 (incr. kidney wt.)		Bio/dynamics 1980c
34	Rat	(W)	2 yr	Hemato	25			Quast et al. 1980b
35	Rat	(W)	M:26 mo F:23 mo	Renal	4.2	14 (incr. kidney wt.)		Bio/dynamics 1980b
36	Rat	(W)	M:22 mo F:19 mo	Hepatic		14 (incr. liver wt.)		Bio/dynamics 1980a
37	Rat	(W)	2 yr	Renal	25			Quast et al. 1980b
38	Rat	(W)	M:26 mo F:23 mo	Hemato	4.2 <sup>b</sup>	14 <sup>i</sup> (decr. red cells)		Bio/dynamics 1980b
Neurological								
39	Rat	(W)	18 mo			70 (decr. activity)		Bigner et al. 1986
Reproductive								
40	Rat	(W)	M:22 mo F:19 mo		0.14	14 (incr. testes wt.)		Bio/dynamics 1980a
Cancer								
41	Rat	(W)	95 wk				70 CEL (tumors)	Beliles et al. 1980
42	Rat	(W)	2 yr				28 CEL (zymbal gland)	Gallagher et al. 1988



TABLE 2-2 (Continued)

Figure Key	Species	Route	Exposure Frequency/ Duration	Effect	NOAEL (mg/kg/day)	LOAEL (Effect)		Reference
						Less Serious (mg/kg/day)	Serious (mg/kg/day)	
43	Rat	(G)	20 mo 7d/wk 1x/d				10 CEL (multiple tumors)	Bio/dynamics 1980c
44	Rat	(W)	M:26 mo F:23 mo				1.4 CEL (multiple tumors)	Bio/dynamics 1980b
45	Rat	(W)	M:22 mo F:19 mo				14 CEL (multiple tumors)	Bio/dynamics 1980a
46	Rat	(W)	18 mo				70 CEL (brain tumor)	Bigner et al. 1986
47	Rat	(W)	2 yr				3.4 CEL (multiple tumors)	Quast et al. 1980b

<sup>a</sup>Converted to an equivalent concentration of 142 ppm in water for presentation in Table 1-4.

<sup>b</sup>Used to derive acute oral MRL; dose divided by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability), resulting in a minimal risk level of 0.1 mg/kg/day. This MRL has been converted to an equivalent concentration in water (3.4 ppm) for presentation in Table 1-3.

<sup>c</sup>Converted to an equivalent concentration of 180 ppm in water for presentation in Table 1-4.

<sup>d</sup>Converted to an equivalent concentration of 200 ppm in water for presentation in Table 1-4.

<sup>e</sup>Converted to an equivalent concentration of 500 ppm in water for presentation in Table 1-4.


<sup>f</sup>Converted to an equivalent concentration of 52 ppm in water for presentation in Table 1-4. Value also used to derive intermediate oral MRL; dose divided by an uncertainty factor of 1,000 (10 for extrapolation from animals to humans and 10 for human variability), resulting in a minimal risk level of 0.01 mg/kg/day.

<sup>g</sup>Converted to an equivalent concentration of 35 ppm in water for presentation in Table 1-4.

<sup>h</sup>Used to derive chronic oral MRL; dose divided by an uncertainty factor of 100 (10 for extrapolation from animals to humans and 10 for human variability) resulting in a minimal risk level of 0.04 mg/kg/day. This MRL has been converted to an equivalent concentration in water (1.4 ppm) for presentation in Table 1-3.


<sup>i</sup>Converted to an equivalent concentration of 100 ppm in water for presentation in Table 1-4.

LOAEL = lowest-observed-adverse-effect level; NOAEL = no-observed-adverse-effect level; mg/kg/day = milligram/kilogram/day; ND = no data; (G) = gavage; LD50 = lethal dose, mortality 50%; d = day; Gd = gestation day; 1x = one time; hemato = hematological; gastro = gastrointestinal; RBC = red blood cell; CNS = central nervous system; (W) = water; mo = month; wk = week; cardio = cardiovascular; incr. = increased; GSH = glutathione; decr = decreased; reproduct. = reproductive; histo = histological; CEL = cancer effect level; M = males; F = females; wt. = weight.




**FIGURE 2-2. Levels of Significant Exposure to Acrylonitrile – Oral**

**INTERMEDIATE  
(15-364 Days)**



**FIGURE 2-2 (Continued)**



**FIGURE 2-2 (Continued)**

## 2. HEALTH EFFECTS

Chronic studies in rats indicate that lifetime exposure to doses of 4.4 mg acrylonitrile/kg body weight/day (mg/kg/day) or higher may result in premature death (Bio/dynamics 1980a, 1980b, 1980c; Gallagher et al. 1988; Quast et al. 1980b). The cause of death was not specifically identified.

The highest NOAEL values and all reliable LOAEL values for death in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

### 2.2.2.2 Systemic Effects

**Respiratory Effects.** Chronic oral exposure of animals to low levels of acrylonitrile has not been found to result in damage to the lungs. Histopathological evaluation of lung tissues from rats showed no lung injury at doses up to 25 mg/kg/day for 2 years (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1980b).

**Cardiovascular Effects.** No injury to the cardiovascular system has been detected in studies in animals. No effects were seen in dogs exposed to 18 mg/kg/day in drinking water for 6 months (Quast et al. 1975). In chronic studies in rats exposed to 14 mg/kg/day in drinking water (Bio/dynamics 1980b) or by gavage (Bio/dynamics 1980c), an increased heart-to-body-weight ratio was observed. However, the effects were not seen in other studies where acrylonitrile was given to rats in drinking water (Bio/dynamics 1980a; Quast et al. 1980b). No histopathological changes were reported in heart tissues in any of the chronic studies (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1980b).

The highest NOAEL values and all reliable LOAEL values for cardiovascular effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

**Gastrointestinal Effects.** Studies in animals suggest that oral exposure to acrylonitrile is irritating to the esophagus and stomach. An acute oral dose of acrylonitrile resulted in hemorrhagic necrosis of the forestomach of rats (Szabo et al. 1984). In dogs given acrylonitrile in drinking water for 6 months at 16 mg/kg/day, gross and microscopic evaluation showed focal erosions and ulcerations in the esophagus (Quast et al. 1975). The authors considered these lesions to be due to irritation to the mucous membranes. No esophageal lesions were present at 10 mg/kg/day. Rats given acrylonitrile in drinking water for 50 weeks at doses up to 70 mg/kg/day showed no effects on the gastrointestinal tract (Beliles et al. 1980), indicating that rats are less sensitive to this effect than dogs. Rats exposed to acrylonitrile in drinking water for 2 years developed papillomatous proliferations of the epithelium of the forestomach at a dose of 28 mg/kg/day, but not at

## 2. HEALTH EFFECTS

7.1 mg/kg/day (Gallagher et al. 1988). This effect is discussed under Section 2.2.2.8 below. Nonproliferative effects on the stomach were not reported.

The method by which an oral dose is administered may also determine whether gastrointestinal effects occur. While no effects were observed in rats given acrylonitrile in drinking water at doses up to 70 mg/kg/day for 50 weeks (Beliles et al. 1980), administration by gavage of 25 mg/kg/day acrylonitrile to rats for 10 days resulted in thickening of the nonglandular portion of the stomach (Murray et al. 1978). A single gavage dose of 50 mg/kg to rats was reported to cause gastrointestinal bleeding, as measured by increased heme content in the gastrointestinal tract (Ghanayem and Ahmed 1983).

The highest NOAEL values and all reliable LOAEL values for gastrointestinal effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

**Hematological Effects.** Decreased red blood cell counts, hematocrit, and hemoglobin content have been reported following acute, intermediate and chronic oral studies in animals (Bio/dynamics 1980a, 1980b, 1980c; Farooqui and Ahmed 1983a; Quast et al. 1975). Although the mechanism of these hemotoxic effects is not clear, Farooqui and Ahmed (1983) found that acrylonitrile bound covalently both to red blood cell membranes and to hemoglobin. In dogs administered acrylonitrile at doses up to 18 mg/kg/day for 6 months, decreased red cell counts, hematocrit, and hemoglobin content were seen only in animals that died (Quast et al. 1975). In chronic studies in rats, lower red cell parameters were seen at 14 mg/kg/day, but these effects were not specifically identified with animals that died during the study (Bio/dynamics 1980a, 1980b, 1980c). In another chronic study in rats at doses up to 25 mg/kg/day for 2 years, no effects on red cell parameters were observed, though early deaths were observed in the study (Quast et al. 1980b). Based on lower red cell parameters (Bio/dynamics 1982a), a chronic oral MRL of 0.04 mg/kg/day was calculated as described in the footnote in Table 2-2.

The highest NOAEL values and all reliable LOAEL values for hematological effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

**Musculoskeletal Effects.** No studies were located regarding musculoskeletal effects in humans or animals following oral exposure to acrylonitrile.

## 2. HEALTH EFFECTS

**Hepatic Effects.** Although acrylonitrile is metabolized by the liver to potentially toxic metabolites (see Section 2.3.3), the liver does not appear to be an important target organ for acrylonitrile toxicity (Silver et al. 1982). In chronic studies in rats, microscopic examination of liver tissue showed no damage after 2 years of exposure to acrylonitrile at doses up to 25 mg/kg/day (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1980b).

Some effects in the liver have been reported, but they may be adaptive changes related to increased metabolic activity. Increased glutathione levels were reported at doses of 70 mg/kg/day for 21 days (Szabo et al. 1977). Because the metabolism of acrylonitrile includes pathways that utilize glutathione (see Section 2.3.3), the higher glutathione levels in the liver may be due to increased demand for glutathione for the metabolism of acrylonitrile.

A consistent observation in rats exposed to acrylonitrile was increased liver weight, both in acute studies at 65 mg/kg/day (Murray et al. 1978) and chronic studies at 10 mg/kg/day (Bio/dynamics 1980a, 1980b, 1980c). Again, this may be an adaptive change related to increased metabolic activity by the liver due to the presence of acrylonitrile in the body.

The highest NOAEL values and all reliable LOAEL values for hepatic effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

**Renal Effects.** No adverse effects on the renal system have been reported in animals administered acrylonitrile via the oral route. In chronic studies in rats, increased kidney weights relative to body weight were observed (Bio/dynamics 1980a, 1980b, 1980c). However, the significance of this observation, if any, is not known, because no histopathological, blood chemistry, or urinalysis findings suggestive of kidney injury were observed in intermediate and chronic studies in rats or dogs (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1975, 1980b).

The highest NOAEL values and all reliable LOAEL values for renal effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

**Dermal/Ocular Effects.** No studies were located regarding dermal or ocular effects in humans or animals following oral exposure to acrylonitrile.

## 2. HEALTH EFFECTS

**Other Systemic Effects.** Decreased adrenal weight, adrenal atrophy, and decreased aldosterone and corticosterone levels in plasma were observed in rats administered acrylonitrile in drinking water for 3 weeks at 70 mg/kg/day (Szabo et al. 1984). Similar effects were observed when the dosing period was extended to 60 days (Szabo et al. 1984). However, no effects on adrenal weight or histopathology of the adrenals were observed in several chronic studies in which doses were similar to the 21- and 60-day studies by Szabo et al. (1984) (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1980b).

The highest NOAEL values and all reliable LOAEL values for other systemic effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

### 2.2.2.3 Immunological Effects

No studies were located regarding immunologic effects in humans or animals following oral exposure to acrylonitrile.

### 2.2.2.4 Neurological Effects

Exposure to acrylonitrile caused effects similar to those of cyanide poisoning. Intermediate and chronic exposure of animals to acrylonitrile resulted in decreased activity and depression (Beliles et al. 1980; Bigner et al. 1986; Quast et al. 1975). Species differences are apparent, dogs being more susceptible than rats. In dogs administered acrylonitrile for 6 months at 16 mg/kg/day, depression and lethargy were reported, with the majority of the experimental groups animals dying before the end of the study period (Quast et al. 1975). In rats, acute exposure results in a transient cholinomimetic stimulatory phase (salivation, diarrhea, lachrymation, vasodilation), followed by a central nervous system depression (Ahmed and Farooqui 1982; Ahmed and Patel 1981). In rats exposed for 1 to 2 years at 70 mg/kg/day, decreased activity, abnormal gait, and prostration were noted but no deaths were reported (Beliles et al. 1980; Bigner et al. 1986). No histopathological changes in the nervous system were reported except those related to tumor formation (see Section 2.2.2.8) (Beliles et al. 1980).

The highest NOAEL values and all reliable LOAEL values for neurological effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.



## 2. HEALTH EFFECTS

### 2.2.2.5 Developmental Effects

Oral exposure to acrylonitrile has been shown to result in teratogenic effects in rats (Murray et al. 1978). Oral administration of 25 mg/kg/day of acrylonitrile during days 6 to 15 of gestation resulted in an increased incidence of malformations. Although the incidence of malformations at 25 mg/kg/day was not statistically significant, the authors considered them related to the administration of acrylonitrile. At a higher dose (65 mg/kg/day), the increase in malformations was statistically significant, and the same malformations were seen at both doses. Statistically significant lowered fetal body weight and shorter crown-rump length also were observed at 65 mg/kg/day. The number of live pups and resorption per litter were not affected by the administration of acrylonitrile. In this study a NOAEL of 10 mg/kg/day was established. Based on this value an acute oral MRL of 0.1 mg/kg/day was calculated as described in the footnote in Table 2-2.

The highest NOAEL values and all reliable LOAEL values for developmental effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

### 2.2.2.6 Reproductive Effects

In a three-generation reproduction study in rats, Beliles et al. (1980) found that exposure of animals to acrylonitrile in drinking water at 70 mg/kg/day resulted in reduced viability and lactation indices in all generations. The authors considered the reduced pup indices to be the result of maternal toxicity, possibly related to reduced milk production due to decreased water intake by the dams. Fostering of pups on untreated dams lessened pup mortality. Measurement of acrylonitrile in the pups was not performed.

Animal studies suggest possible effects of acrylonitrile on the male reproductive system. Tandon et al. (1988) observed histological and biochemical evidence of degenerative changes in testicular tubules of mice exposed to 10 mg/kg/day of acrylonitrile for 60 days. These changes were accompanied by a 45% decrease in sperm count. Whether the tubular changes or the decreased sperm count have an effect on fertility is not known because no studies have been conducted to evaluate the effects of acrylonitrile on the reproductive capability of treated mice. Based on the reduced sperm count and the degenerative changes in testicular tubules, an intermediate oral MRL of 0.01 mg/kg/day was calculated as described in the footnote in Table 2-2.

There appear to be significant inter-species differences with respect to histopathological effects of acrylonitrile on the male reproductive system. Histological examination did not reveal testicular degeneration in rats or dogs given acrylonitrile for 2 years (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1975, 1980b), although

## 2. HEALTH EFFECTS

an increase in testicular weight was observed in rats given acrylonitrile in drinking water at doses of 14 mg/kg/day for 2 years (Bio/dynamics 1980a). The biological significance of this effect is not clear, because this was observed in only one chronic drinking water study in rats, and was not observed in a gavage study in the same strain of rat (Spartan) or at comparable doses in Fisher 344 rats (Bio/dynamics 1980b, 1980c).

The highest NOAEL values and all reliable LOAEL values for reproductive effects in each species and duration category are recorded in Table 2-2 and plotted in Figure 2-2.

### 2.2.2.7 Genotoxic Effects

In vivo studies in animals suggest that oral exposure to acrylonitrile has limited genotoxic potential. Acrylonitrile produced no significant increase in chromosomal aberrations in bone marrow cells from rats administered up to 21 mg/kg/day (Rabello-Gay and Ahmed 1980), and no significant effect in a dominant lethal assay in rats at doses up to 60 mg/kg/day (Working et al. 1987). Studies on unscheduled DNA synthesis in liver and brain of rats exposed to doses of 50 mg/kg showed increased DNA synthesis in the liver but not in the brain, suggesting limited potential for acrylonitrile to be genotoxic (Hogy and Guengerich 1986). However, unscheduled DNA synthesis was not observed in an in vivo spermatocyte assay in rats exposed to 60 mg/kg/day (Hurtt et al. 1987).

### 2.2.2.8 Cancer

No studies were located regarding cancer in humans following oral exposure to acrylonitrile.

Chronic studies in rats provide convincing evidence that acrylonitrile is carcinogenic to animals when administered orally (Beliles et al. 1980; Bigner et al. 1986; Bio/dynamics 1980a, 1980b, 1980c; Gallagher et al. 1988; Quast et al. 1980a). Tumors of the central nervous system and Zymbal gland were identified in most studies at doses of 1.4 mg/kg/day and higher. Tumors of the gastrointestinal tract (forestomach, intestines) and mammary gland (in females) were also common at doses of 3.4 mg/kg/day or higher (Bigner et al. 1986; Bio/dynamics 1980a, 1980c; Maltoni et al. 1977; Quast et al. 1980b). Gallagher et al. (1988) noted a dose-dependent decrease in the number of rats with pituitary tumors (mainly prolactinomas) following chronic oral exposure to acrylonitrile in drinking water.

## 2. HEALTH EFFECTS

### 2.2.3 Dermal Exposure

#### 2.2.3.1 Death

The death of a 10-year-old girl following dermal exposure to acrylonitrile was reported by Lorz (1950). An acrylonitrile preparation had been applied to the scalp of the child as a treatment for head lice. The child experienced nausea, headache and dizziness. Death occurred 4 hours after application. The concentration was not specified in this case report.

In guinea pigs, the dermal LD<sub>50</sub> was reported to be 370 mg/kg (Roudabush et al. 1965). Values of 226 to 250 mg/kg have been noted in rabbits (Back et al. 1972; Roudabush et al. 1965).

The highest NOAEL values and all reliable LOAEL values for death in each species and duration category are recorded in Table 2-3.

#### 2.2.3.2 Systemic Effects

No studies were located regarding systemic effects in animals following dermal exposure to acrylonitrile.

In humans, one study of a worker accidentally sprayed with acrylonitrile indicated that transient injury to liver and muscle may have occurred, but the data are too limited to draw any firm conclusions (Vogel and Kirkendall 1984).

Workers exposed to acrylonitrile vapors at 16 to 100 ppm for 20 to 45 minutes complained of intolerable itching of the skin, but no dermatitis was observed (Wilson et al. 1948). This is presumably a direct irritant effect of acrylonitrile on the skin.

In a study of Japanese workers exposed to acrylonitrile, irritation of the conjunctiva and upper respiratory tract was reported. Workers who may have been exposed to particularly high concentrations (inside polymerization tanks) experienced transient irritation of the scrotal skin. No concentrations of acrylonitrile were specified (Sakurai et al. 1978).

#### 2.2.3.3 Immunological Effects

No studies were located regarding immunological effects in humans or animals following dermal exposure to acrylonitrile.

TABLE 2-3. Levels of Significant Exposure to Acrylonitrile - Dermal

Figure Key	Species	Exposure Frequency/ Duration	Effect	NOAEL (mg/kg/day)	LOAEL (Effect)		Reference
					Less Serious (mg/kg/day)	Serious (mg/kg/day)	
ACUTE EXPOSURE							
Death							
1	Gn Pig	ND				370 (LD50)	Roudabush et al. 1965
2	Rabbit	ND				250 (LD50)	Back et al. 1972
3	Rabbit	ND				226 (LD50)	Roudabush et al. 1965

LOAEL = lowest-observed-adverse-effect level; NOAEL = no-observed-adverse-effect level; mg/kg/day = milligram/kilogram/day;  
 ND = no data; LD50 = lethal dose, mortality 50%.

## 2. HEALTH EFFECTS

### 2.2.3.4 Neurological Effects

Signs of cyanide poisoning were exhibited by a man accidentally sprayed with acrylonitrile. Dizziness, redness, nausea, vomiting and hallucinations were reported (Vogel and Kirkendall 1984). The symptoms persisted for 3 days.

No studies were located regarding the following health effects in humans or animals after dermal exposure to acrylonitrile.

### 2.2.3.5 Developmental Effects

### 2.2.3.6 Reproductive Effects

### 2.2.3.7 Genotoxic Effects

### 2.2.3.8 Cancer

## 2.3 TOXICOKINETICS

### 2.3.1 Absorption

#### 2.3.1.1 Inhalation Exposure

In a well-controlled and conducted study with human volunteers, Jakubowski et al. (1987) reported that an average of 52% of the inhaled dose of acrylonitrile (5 or 10 mg/m<sup>3</sup>) is absorbed by the lungs. Similar results were reported by Rogaczewska and Piotrowski (1968), who found that 46% of inhaled acrylonitrile is retained by the lungs of humans.

Pilon et al. (1988b) demonstrated in rats exposed to 4 mg/kg acrylonitrile (2,3-<sup>14</sup>C) in a closed-circuit inhalation chamber that the absorption of acrylonitrile was biphasic, characterized by a rapid dose-dependent phase that was followed by a slower dose-independent phase.

#### 2.3.1.2 Oral Exposure

No studies were located regarding absorption in humans following oral exposure to acrylonitrile.

Results of studies in laboratory animals with <sup>14</sup>C-acrylonitrile indicate acrylonitrile is rapidly and extensively absorbed by the oral route. Radiolabeled acrylonitrile is detected in blood within 30 minutes after administration of an oral dose and peak plasma concentrations are reached 6 hours after administration (Farooqui and Ahmed 1982). Extensive absorption is indicated by the fact that only 2 to 10% of administered radioactivity is recovered in the feces (Ahmed et al. 1982, 1983; Farooqui and Ahmed 1982; Young et al. 1977).

## 2. HEALTH EFFECTS

### 2.3.1.3 Dermal Exposure

In studies in human volunteers conducted by Rogaczewska and Piotrowski (1968), absorption by skin was estimated to be 0.6 mg/cm<sup>2</sup>/hr. Though no quantitative estimates of dermal absorption could be made, absorption of acrylonitrile via the dermal route by humans was demonstrated in a case study by Vogel and Kirkendall (1984). Accidental spraying of a man with acrylonitrile resulted in marked symptoms of acrylonitrile toxicity, indicating that significant amounts of acrylonitrile had been absorbed, primarily through the skin.

No studies were located regarding absorption in animals following dermal exposure to acrylonitrile.

### 2.3.2 Distribution

#### 2.3.2.1 Inhalation Exposure

No studies were located regarding distribution of acrylonitrile in humans following inhalation exposure to acrylonitrile.

Acrylonitrile is rapidly distributed throughout the body after inhalation exposure. Measurable amounts of acrylonitrile derived radiolabel were present in the brain, stomach, liver, kidney, lung and blood within 1 hour of initiation of exposure (Pilon et al. 1988b).

#### 2.3.2.2 Oral Exposure

Tissue distribution of radioactivity in rats after a single oral dose of <sup>14</sup>C-acrylonitrile indicates that acrylonitrile and its metabolites are rapidly distributed to all tissues (Ahmed et al. 1982, 1983; Silver et al. 1987; Young et al. 1977). Species differences are apparent. In mice, cyanide levels in the blood peaked at 1 hour, while in rats peak levels were not reached until 3 hours after administration (Ahmed and Patel 1981). The highest levels of radioactivity were recovered in the gastrointestinal tract, in particular in the stomach. The retention of acrylonitrile and its metabolites in the stomach appears to be due, at least in part, to covalent binding (Ahmed et al. 1982; Silver et al. 1987). Young et al. (1977) noted that label accumulated in the stomach following intravenous as well as oral exposure to <sup>14</sup>C-acrylonitrile, suggesting that enterogastric circulation may be important in the preferential retention in the stomach.

Distribution studies by whole-body autoradiography in rats and monkeys revealed accumulation of radiolabel in the liver, kidney, lung, adrenal cortex and stomach. In fetuses exposed in utero, only the eye lens accumulated radiolabel at a higher concentration than that observed in maternal blood (Sandberg and Slanina 1980).

## 2. HEALTH EFFECTS

### 2.3.2.3 Dermal Exposure

No studies were located regarding distribution in humans or animals following dermal exposure to acrylonitrile.

### 2.3.3 Metabolism


Proposed pathways for the metabolism of acrylonitrile are presented in Figure 2-3 (Ahmed et al. 1983; EPA 1980a; Langvardt et al. 1980; Linhart et al. 1988; Muller et al. 1987; Pilon et al. 1988a; Roberts et al. 1989). Studies indicate that the metabolism of acrylonitrile in animals proceeds by the same pathways whether exposure is by the oral (Ahmed et al. 1983; Langvardt et al. 1980; Pilon et al. 1988a) or the inhalation route (Gut et al. 1985; Muller et al. 1987; Tardif et al. 1987). No data were located regarding the metabolism of acrylonitrile following dermal exposure.

Both enzymatic and nonenzymatic biotransformation of acrylonitrile occurs. Acrylonitrile is capable of covalently binding to proteins and other macromolecules such as lipids or nucleic acids, or acrylonitrile can also be directly conjugated to glutathione and excreted in urine as cyanoethylmercapturic acid.

Alternatively, acrylonitrile is metabolized to 2-cyanoethylene oxide by the microsomal enzyme system. 2-Cyanoethylene oxide can react directly with tissue macromolecules or it can be further metabolized to oxidation products that release cyanide. Cyanide is converted to thiocyanate and excreted in the urine. 2-Cyanoethylene oxide is also conjugated with glutathione and metabolized to 2-hydroxyethylmercapturic acid which is excreted in the urine.

Acrylonitrile is also metabolized to CO<sub>2</sub> which is eliminated through the lungs. Carbon dioxide is produced when acrylonitrile is metabolized to ethylene oxide and degraded to oxidation products and cyanide via the epoxide hydratase pathways (Farooqui and Ahmed 1982; Young et al. 1977).

Studies indicate that acrylonitrile conjugation with glutathione is the preferred pathway for metabolism (Ghanayem and Ahmed 1982; Miller and Villaume 1978; Pilon et al. 1988a). However, if glutathione is depleted or the pathway is overloaded (as may be the case at high doses), metabolism to the thiocyanate via 2-cyanoethylene oxide is increased. Increased thiocyanate excretion with glutathione depletion or increased dose was demonstrated by Pilon et al. (1988a). Glutathione



**FIGURE 2-3. Proposed Metabolic Scheme for Acrylonitrile**

Adapted from Ahmed et al. 1983, Linhart et al. 1988, Muller et al. 1987, Roberts et al. 1989



## 2. HEALTH EFFECTS

depleted rats excreted 58% of an orally administered dose as thiocyanate, while normal rats given the same dose (4 mg/kg) of acrylonitrile excreted only 23% as thiocyanate. Normal rats given acrylonitrile at 4 or 10 mg/kg excreted 16% and 23% of the dose as thiocyanate, respectively.

The increased metabolism of acrylonitrile to 2-cyanoethylene oxide has significant implications in acrylonitrile toxicity. 2-Cyanoethylene oxide has been shown to react with cell macromolecules (including nucleic acids) both in vivo and in vitro (Guengerich et al. 1981; Hogy and Guengerich 1986). This metabolite may be responsible for the carcinogenic effects of acrylonitrile.

Urinary excretion patterns of thiocyanate suggest that there are quantitative species differences in acrylonitrile metabolism (Ahmed and Patel 1981). Thiocyanate was identified as a metabolite in rats, mice, rabbits and Chinese hamsters. About 20 to 23% of the administered dose was excreted as thiocyanate in rats, rabbits and Chinese hamsters, while 35% was excreted as thiocyanate in mice (Gut et al. 1975). It has also been observed that mice metabolize acrylonitrile more rapidly than rats (Ahmed and Patel 1981; Gut et al. 1975). Maximum blood cyanide concentrations were observed 1 hour after dosing in mice, but 3 hours after dosing in rats (Ahmed and Patel 1981). In mice, thiocyanate was present in the urine within 4 hours of dosing, while in rats, thiocyanate was present in urine only at time intervals longer than 4 hours (Gut et al, 1975).

In humans, metabolites of acrylonitrile have been identified in urine following occupational exposure (assumed to be by the inhalation route), and also in controlled exposure studies. Metabolites identified in humans were the same as those in animals (Jakubowski et al. 1987; Sakurai et al. 1978). Acrylonitrile and thiocyanate were quantified in urine of workers exposed to acrylonitrile. Dose-related increases in thiocyanate were observed, indicating that cyanide is liberated with the metabolism of acrylonitrile. In a study with human volunteers under controlled conditions, 2-cyanoethylmercapturic acid (CMA) was monitored in urine as an indication of exposure. On average, 22% of the absorbed acrylonitrile was metabolized to CMA; however, considerable individual variability was observed. The CMA excretion ranged from 13% to 39% of the absorbed dose (Jakubowski et al. 1987).

In a case study of a human male accidentally sprayed with acrylonitrile, recurring signs of cyanide poisoning were seen over a 3 day period (Vogel and Kirkendall 1984). This indicates that acrylonitrile is also metabolized to cyanide following predominantly dermal exposure.

## 2. HEALTH EFFECTS

### 2.3.4 Excretion

#### 2.3.4.1 Inhalation Exposure

Studies on workers in an occupational setting showed a dose-response relationship between the concentration of acrylonitrile of inspired air and the recovery of metabolites in the urine (Houthuijs et al. 1982; Sakurai et al. 1978). In a controlled study using human volunteers, urinary metabolite data suggested that the elimination of acrylonitrile followed first-order kinetics, with a half-life of seven to eight hours (Jakubowski et al. 1987).

The predominant route of excretion in rats is via urine (Gut et al. 1985; Tardif et al. 1987; Young et al. 1977). In rats exposed to 5 ppm of 1-<sup>14</sup>C-acrylonitrile for 6 hours, 68% of the absorbed radioactivity was excreted in the urine within 220 hours, with 3.9% in the feces, 6.1% in expired air as <sup>14</sup>C<sub>2</sub>O, and 18% of the radioactivity being retained in the body tissues. Following exposure to a higher concentration (100 ppm), a larger fraction of the dose was recovered in urine (82%) and a smaller fraction (2.6%) was retained in the body (Young et al. 1977), indicating that urinary excretion is dose-dependent. Percent fecal excretion was similar at both doses.

#### 2.3.4.2 Oral Exposure

Following oral exposure, the major route of excretion of acrylonitrile in rats is via the urine, either as thiocyanate or as other products of conjugation. Within the first 24 hours of a single oral dose, 40% to 60% was recovered in the urine (Ahmed et al. 1983). Farooqui and Ahmed (1982) reported that 10 days after the administration of a single dose, 61% of the dose had been accounted for in the urine, 3% in feces and 13% in the expired air. Approximately 25% was retained in the body covalently bound to tissues (see Section 2.3.3).

A study by Young et al. (1977) showed that retention and excretion of acrylonitrile are not directly proportional to dose. The data suggest a saturation process, perhaps due to covalent binding to tissue macromolecules. Seventy-two hours after administration of single oral doses of either 0.1 or 10 mg/kg, the proportion of the dose retained in the carcass was 37% at the low dose (0.1 mg/kg) and 27% at the high dose (10 mg/kg).

#### 2.3.4.3 Dermal Exposure

No studies were located regarding excretion in humans or animals following dermal exposure to acrylonitrile.

## 2. HEALTH EFFECTS

### 2.4 RELEVANCE TO PUBLIC HEALTH

Acrylonitrile is a common industrial chemical, and humans may be exposed to acrylonitrile around factories where it is made or used, or near chemical waste sites where it has been improperly stored or disposed of. The two most likely exposure pathways are breathing acrylonitrile that has evaporated into air (it is readily volatile), or drinking water that has been contaminated (acrylonitrile is highly soluble and stable in water). However, data describing acrylonitrile levels in drinking water are currently lacking. The low odor threshold of acrylonitrile in water (19 ppm) could limit exposure to this substance through contaminated drinking water.

Short exposures at high concentrations show cyanide-type toxicity. In humans, symptoms include headaches, feelings of nausea, irritability, and apprehension. These symptoms are reversible. However, prolonged exposure can result in death. Limited data suggest that children may be more susceptible to acrylonitrile toxicity than adults. However, current data are not significant to draw firm conclusions. In animals, acrylonitrile has been shown to result in teratogenic effects when administered by either the inhalation or oral routes of exposure. Impaired reproductive capability has also been demonstrated in mice. Exposures high enough to cause neurological, developmental, or reproductive effects in humans are rather unlikely to occur except in cases of spills or accidents.

Epidemiologic studies have associated acrylonitrile exposure in the workplace with increased incidence of lung cancer and possibly prostate cancer. Because only six cases of prostate cancer were observed and all were found in workers at ages when the incidence rate increases, firm conclusions cannot be made about the potential for acrylonitrile to cause prostate cancer. In animals, acrylonitrile has been demonstrated to cause tumors at multiple sites, including the nervous system, gastrointestinal tract, Zymbal gland and mammary gland. Because of this clear carcinogenic potential, chronic low-level exposure to acrylonitrile is of special concern.

**Death.** Several cases of death in children following inhalation or dermal exposure to acrylonitrile has been reported (Grunske 1949; Lorz 1950). Children seem to be more susceptible to the lethal effect of acrylonitrile than adults. However, current data are not sufficient to draw firm conclusions regarding risk. In the case studies where children had died after sleeping in rooms fumigated with acrylonitrile, the adults that shared the quarters reported few, if any, effects.

## 2. HEALTH EFFECTS

In animals, deaths from acrylonitrile have been reported in several species following inhalation, oral or dermal exposure. In most species, death appears to be related to cyanide poisoning. That the cyanide moiety is involved in human toxicity of acrylonitrile has been reported in a case study in which a human male was sprayed with acrylonitrile when a valve burst (Vogel and Kirkendall 1984). This individual suffered symptoms characteristic of cyanide poisoning, and treatments designed to reduce cyanide levels in the blood were required in order to save his life.

**Systemic Effects.** Humans exposed to high levels of acrylonitrile in the workplace have complained of nasal irritation and an oppressive feeling in the upper respiratory passages (Wilson 1944; Wilson et al. 1948). Acute exposure of guinea pigs to acrylonitrile has resulted in severe pulmonary irritation and lung edema. In other species, only mild pulmonary irritation has been reported (Dudley and Neal 1942). Chronic exposure to acrylonitrile via inhalation has resulted in degenerative and inflammatory changes in the nasal passages in rats (Quast et al. 1980a). The concentrations at which these effects have been seen in animals are irritating to the respiratory tract of humans.

The gastrointestinal tract does not appear to be a target organ of acrylonitrile toxicity in humans. In animals, noncarcinogenic lesions in the esophagus and stomach suggest an irritational effect on the gastrointestinal tract (Murray et al. 1978; Quast et al. 1975). However, several investigators (Cote et al. 1984; Ghanayem and Ahmed 1983; Ghanayem et al. 1985; Szabo et al. 1983) have demonstrated that gastrointestinal bleeding and duodenal ulcers may occur even when rats were given acrylonitrile by routes other than oral, suggesting that the effects on the gastrointestinal tract may not be due exclusively to irritation. More study is needed in this area before any conclusions about the relevance to human health can be drawn.

In humans, severe cases of acrylonitrile poisoning have resulted in low grade anemia (Wilson 1944; Wilson et al. 1948), but complete recovery was reported. Chronic occupational exposure to low levels of acrylonitrile has not resulted in detectable effects on the hematological system (Sakurai et al. 1978). In intermediate and chronic studies in animals, decreased red cell count, hemoglobin concentration and hematocyte were observed (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1975; 1980a). However, it is not certain if these effects are due to acrylonitrile binding to red blood cell membranes and proteins (Farooqui and Ahmed 1982), or are secondary effects related to the poor physical condition which develops in the exposed animals.

## 2. HEALTH EFFECTS

Some liver damage has been reported in humans after acute exposure to high doses. In workers exposed to high levels of acrylonitrile vapors, mild jaundice was diagnosed (Wilson 1944). In a case of an accidental dermal exposure of a man, enzyme levels in the blood suggestive of liver injury were reported for several days (Vogel and Kirkendall 1984). These effects appeared to be fully reversible.

The liver has been reported to be the primary site of acrylonitrile metabolism. Metabolism studies indicate that a toxic metabolite (2-cyanoethylene oxide) is formed that covalently binds to liver tissue. However, the liver has not been indicated as a target organ for intermediate or chronic exposure in humans or animals. In factory workers exposed to acrylonitrile for 10 or more years, no liver damage has been observed (Sakurai et al. 1978). In animal studies, microscopic examination of liver tissue has revealed no hepatic damage after 90 days to 2 years of exposure by the inhalation or oral routes (Bio/dynamics 1980a, 1980b, 1980c; Humiston et al. 1975; Quast 1975, 1980a, 1980b).

Some effects seen in the livers of animals may suggest that adaptive changes occur as a response to increased metabolic activity. These include increased glutathione levels, possibly due to an increased demand for glutathione for the metabolism of acrylonitrile. Increased liver weights have also been observed in both acute and chronic oral studies (Bio/dynamics 1980a, 1980b, 1980c; Murray et al. 1978).

Skin irritation results from direct contact with liquid or gaseous acrylonitrile. Workers exposed to high levels of acrylonitrile in air have complained of itching, but no dermatitis was observed (Wilson et al. 1948). Also, transient irritation of scrotal skin has been noted by workers after entering areas with high ambient acrylonitrile concentrations. Direct contact of acrylonitrile with the skin has resulted in erythema, desquamation and slow healing (Dudley and Neal 1942). In both humans and animals, skin redness has been reported subsequent to acute exposures (Dudley and Neal 1942; Vogel and Kirkendall 1984).

**Immunological Effects.** Immunological effects of acrylonitrile have not been studied in humans or animals.

**Neurological Effects.** Acute effects have been described as being a cyanide-type poisoning (Dudley and Neal 1942; Vogel and Kirkendall 1984). These observations are consistent with results from studies in animals where it has been shown that the major metabolic pathways include cyanide and thiocyanate as products of acrylonitrile metabolism (Ahmed et al. 1983; EPA 1980a; Langvardt et al. 1980; Pilon et al. 1988a). The formation of cyanide following oral exposure is greater than inhalation or subcutaneous exposure (Gut et al. 1975), indicating

## 2. HEALTH EFFECTS

that the amount of acrylonitrile reaching the liver in free form (i.e., not covalently bound) may determine the amount of cyanide formed. Symptoms which have been associated with acrylonitrile poisoning in humans include limb weakness, labored and irregular breathing, dizziness and impaired judgment, cyanosis and nausea, collapse, and convulsions (Baxter 1979). Case studies of occupational exposure to acrylonitrile suggest that the acute nonlethal effects in humans may be fully reversible (Vogel and Kirkendall 1984; Wilson 1944; Wilson et al. 1948).

Acute exposure of several species of animals has been reported to lead to convulsions and coma (Dudley and Neal 1942). Exposure of animals for 6 months or more at 16 mg of acrylonitrile per kilogram body weight per day (mg/kg/day) results in neurological effects characterized by decreased activity and depression (Bigner et al. 1986; Beliles et al. 1980; Quast et al. 1975). However, the only histopathological changes seen in the nervous system have been those associated with the formation of tumors (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1975, 1980a, 1980b).

**Developmental Effects.** There are no data available on developmental effects of acrylonitrile in humans; however, two well-conducted studies in rats have shown that acrylonitrile is teratogenic in animals by both inhalation and oral exposure (Murray et al. 1978). Fetal malformations occurred in a dose-related manner. When administered orally, malformations were present even at doses in which no maternal or fetal toxicity was apparent.

**Reproductive Effects.** There are no data available on the reproductive effects in humans. However, a three-generation reproduction study in rats showed that acrylonitrile has an effect on reproduction. In all three generations, viability and lactation indices were affected by 'the presence of acrylonitrile in the drinking water (Beliles et al. 1980). Species differences were observed with respect to the effects of acrylonitrile on male reproductive organs. In mice, adverse effects on male reproductive organs were demonstrated at 10 mg/kg/day in a 60-day study (Tandon et al. 1988). In rats, no effects on male reproductive organs were seen even when exposure was at 14 mg/kg/day for 2 years (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1980a, 1980b).

**Genotoxic Effects.** The genotoxicity of acrylonitrile in vitro has been extensively studied (Table 2-4). Although the results are not entirely consistent, positive results have been observed for all end points evaluated: gene mutations, chromosomal aberrations, DNA damage, and cell transformation. However, in vivo genotoxicity studies have generally been negative in mammalian systems, including humans (Table 2-5). Tests in Drosophila melanogaster suggest that

TABLE 2-4. Genotoxicity of Acrylonitrile In Vitro

End Point	Species (Test System)	Results		Reference
		With Activation	Without Activation	
Prokaryotic organisms:				
Gene mutation	<u>Salmonella typhimurium</u> (plate incorporation)	+	+	Khudoley et al. 1987
	<u>S. typhimurium</u> (plate incorporation)	+	-	Lijinsky and Andrews 1980
	<u>S. typhimurium</u> (plate incorporation)	-	+	Baker and Bonin 1985
	<u>S. typhimurium</u> (liquid preincubation)	+	+	Zeiger and Haworth 1985
	<u>S. typhimurium</u> (liquid preincubation)	-	-	Matsushima et al. 1985
	<u>S. typhimurium</u> (gas exposure)	+	-	DeMeester et al. 1978
	<u>Escherichia coli</u>	No data	+	Venitt et al. 1977
Eukaryotic organisms:				
Fungi:				
Gene conversion	<u>Saccharomyces cerevisia</u> D7	-	+	Arni 1985
	<u>S. cerevisiae</u> JD1	+	-	Brooks et al. 1985
Mammalian cells:				
Gene mutation	Mouse lymphoma L5178Y thymidine kinase locus	No data	+	Myhr et al. 1985
	Mouse lymphoma L5178Y thymidine kinase locus	+	+	Amacher and Turner 1985
	Mouse lymphoma L5178Y thymidine kinase locus	+	+	Lee and Webber 1985
	Mouse lymphoma L5178Y ouabain resistance	-	-	Garner and Campbell 1985
	Mouse lymphoma L5178Y 6-thioguanine resistance	+	+	Garner and Campbell 1985

TABLE 2-4 (Continued)

End Point	Species (Test System)	Results		Reference
		With Activation	Without Activation	
Sister chromatid exchange	Chinese hamster V79/HGPT	-	-	Lee and Webber 1985
	Mouse lymphoma P388F thymidine kinase locus	+	-	Anderson and Cross 1985
	Human lymphoblasts AHH-1 TK6	+	-	Crespi et al. 1985
	Human lymphoblastoid TK6	+	-	Recio and Skopek 1988
	Human lymphoblasts	NA	+	Crespi et al. 1985
	Rat liver RL4	NA	-	Priston and Dean 1985
	Human lymphocytes	-	-	Obe et al. 1985
	Human lymphocytes	+	-	Perocco et al. 1982
	Chinese hamster ovary	+	-	Brat and Williams 1982
	DNA Synthesis	Hepatocyte primary cultures	NA	+
Hepatocyte primary cultures		NA	+	Glauert et al. 1985
Hepatocyte primary cultures		NA	-	Probst and Hill 1985
Cell transformation	Syrian hamster embryo cells	NA	+	Sanner and Rivedal 1985; Parent and Casto 1979
	Balb/C-3T3	+	-	Matthews et al. 1985
	C3H/10T1/2	+	-	Lawrence and McGregor 1985
	C3H/10T1/2	NA	+	Banerjee and Segal 1986
	NIH/3T3	NA	+	Banerjee and Segal 1986

+ = positive result; - = negative result; NA = not applicable.



TABLE 2-5. Genotoxicity of Acrylonitrile In Vivo

End Point	Species (Test System)	Results	Reference
Mammalian systems:			
Chromosomal aberrations	Mouse bone marrow	-	Leonard et al. 1981
	Mouse bone marrow	-	Sharief et al. 1986
	Mouse bone marrow	-	Rabello-Gay 1980; Ahmed 1980
Micronuclei	Mouse bone marrow	-	Leonard et al. 1981
Dominant lethals	Mouse	-	Leonard et al. 1981
Host mediated assays:			
Gene mutations	<u>S. typhimuriam</u> (mouse host mediated)	-	Lambotte-Vandepaer et al. 1980
	<u>S. typhimuriam</u> (rat host mediated)	-	Lambotte-Vandepaer et al. 1980, 1981
Non-mammalian systems:			
Gene mutations	<u>Drosophila melanogaster</u>	+	Fujikawa et al. 1985; Vogel 1985; Wurgler et al. 1985

- = negative result; + = positive result.

## 2. HEALTH EFFECTS

acrylonitrile may have potential for genotoxicity in vivo. Acrylonitrile and its metabolites have been shown to bind to nucleic acids both in vivo and in vitro suggesting that acrylonitrile does have potential for genetic damage.

**Cancer.** Studies in rats by inhalation and oral exposure have all demonstrated that acrylonitrile is carcinogenic. Multiple tumor sites have been identified (Bio/dynamics 1980a, 1980b, 1980c; Quast et al. 1980a, 1980b). There is limited evidence that acrylonitrile is a human carcinogen. Some epidemiology studies indicate that occupational exposure to acrylonitrile results in an increased incidence of lung cancer (O'Berg 1980; O'Berg et al. 1985). However, other studies have not reached this conclusion (Collins et al. 1989; Kiesselbach et al. 1979). There is also a suggestion that acrylonitrile exposure may result in an increased incidence of prostate cancer (Chen et al. 1987; O'Berg et al. 1985). Limitations of these studies do not allow firm conclusions about the potential for acrylonitrile to cause prostate cancer. EPA has classified acrylonitrile as a probable human carcinogen (IRIS 1988). There is sufficient evidence in animals that acrylonitrile is carcinogenic, but the human evidence is limited.

### 2.5 BIOMARKERS OF EXPOSURE AND EFFECT

Biomarkers are broadly defined as indicators signaling events in biologic systems or samples. They have been classified as biomarkers of exposure, biomarkers of effect, and biomarkers of susceptibility (NAS/NRC 1989).

A biomarker of exposure is a xenobiotic substance or its metabolite(s) or the product of an interaction between a xenobiotic agent and some target molecule or cell that is measured within a compartment of an organism (NAS/NRC 1989). The preferred biomarkers of exposure are generally the substance itself or substance-specific metabolites in readily obtainable body fluid or excreta. However, several factors can confound the use and interpretation of biomarkers of exposure. The body burden of a substance may be the result of exposures from more than one source. The substance being measured may be a metabolite of another xenobiotic (e.g., high urinary levels of phenol can result from exposure to several different aromatic compounds). Depending on the properties of the substance (e.g., biologic half-life) and environmental conditions (e.g., duration and route of exposure), the substance and all of its metabolites may have left the body by the time biologic samples can be taken. It may be difficult to identify individuals exposed to hazardous substances that are commonly found in body tissues and fluids (e.g., essential mineral nutrients such as copper, zinc and selenium). Biomarkers of exposure to acrylonitrile are discussed in Section 2.5.1.

## 2. HEALTH EFFECTS

Biomarkers of effect are defined as any measurable biochemical, physiologic, or other alteration within an organism that, depending on magnitude, can be recognized as an established or potential health impairment or disease (NAS/NRC 1989). This definition encompasses biochemical or cellular signals of tissue dysfunction (e.g., increased liver enzyme activity or pathologic changes in female genital epithelium cells), as well as physiologic signs of dysfunction such as increased blood pressure or decreased lung capacity. Note that these markers are often not substance specific. They also may not be directly adverse, but can indicate potential health impairment (e.g., DNA adducts). Biomarkers of effects caused by acrylonitrile are discussed in Section 2.5.2.

A biomarker of susceptibility is an indicator of an inherent or acquired limitation of an organism's ability to respond to the challenge of exposure to a specific xenobiotic. It can be an intrinsic genetic or other characteristic or a preexisting disease that results in an increase in absorbed dose, biologically effective dose, or target tissue response. If biomarkers of susceptibility exist, they are discussed in Section 2.7, "POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE."

### **2.5.1 Biomarkers Used to Identify and/or Quantify Exposure to Acrylonitrile**

Parent acrylonitrile molecule and its metabolites have been measured in blood and urine, but, except for measurement of thiocyanate, methods have not been developed for routine monitoring of exposed humans.

Factory workers exposed to an average of 0.1, 0.5 and 4.2 ppm of acrylonitrile in the air during an 8-hour work day averaged 3.9, 19.7, and 360 µg/L acrylonitrile in the urine, respectively, and 4.5, 5.78, and 11.4 mg/L thiocyanate in the urine, respectively (Sakurai et al. 1978). No acrylonitrile was detected in the urine of a control group, but an average of 4.00 mg/L of thiocyanate was found in the urine. The presence of thiocyanate in the urine of workers not exposed to acrylonitrile has been related to cigarette smoking (Houthuijs et al. 1982; Sakuria et al. 1978). Houthuijs et al. (1982) reported post-shift acrylonitrile values of 39 µg/L when the mean acrylonitrile concentration in the air was 0.13 ppm.

### **2.5.2 Biomarkers Used to Characterize Effects Caused by Acrylonitrile**

A variety of effects have been demonstrated following acrylonitrile exposure in humans and animals. These effects show a close similarity to an underlying cyanide effect, particularly for acute exposures. Effects can be detected in groups of exposed individuals by monitoring

## 2. HEALTH EFFECTS

signs and symptoms such as increased salivation, dizziness, and labored and irregular breathing. In some cases convulsions and coma may occur. Because the release of cyanide for producing toxic effects is common for other compounds, measuring these effects is not specific for acrylonitrile exposure. These effects do identify potential health impairment. It should be noted that the toxicity of acrylonitrile resides not only in the cyanide radical but in the entire molecule. The latter structure explains various chronic effects such as cancer that result from acrylonitrile, as opposed to cyanide whose effects are more relevant for acute toxicity. Studies that identify subtle physiological changes that can be used to detect or predict risk of disease following long-term exposure to acrylonitrile are not available.

### 2.6 INTERACTIONS WITH OTHER CHEMICALS

The interaction between acrylonitrile and other chemicals has not been thoroughly studied. O'Berg (1980) noted that out of 8 workers exposed to acrylonitrile who developed lung cancer, 7 were smokers (smoking history was not available for the eighth individual). This suggests that smoking might increase lung cancer risk from acrylonitrile, but the data are too limited to draw any firm conclusions on this point.

Radimer et al. (1974) described four cases of severe epidermal necrolysis in individuals who had been exposed to the residual fumes of a mixture of acrylonitrile and carbon tetrachloride used to fumigate their homes. Three of the people died. The authors thought this was most likely due to the effects of acrylonitrile, but noted that an interaction between carbon tetrachloride and acrylonitrile was possible.

In animals, the hemorrhagic effects of acrylonitrile exposure on the adrenals may be reduced by prior exposure of the animals to adrenergic blockers or chemicals that deplete the adrenal cortex of catecholamines (Silver et al. 1987; Szabo et al. 1980). It is difficult to judge whether adrenergic antagonists would have a similar protective effect in humans, because effects of acrylonitrile on the adrenal have not been described in humans.

Acrylonitrile alone has little tendency to produce duodenal ulcers in animals, but pretreatment with phenobarbital or Aroclor results in a marked increase in the incidence of such ulcers (Szabo et al. 1983, 1984). Although the mechanism of the ulcerogenic effect is not obvious, these data indicate that agents which enhanced mixed-function oxidase activity may also increase the toxicity of acrylonitrile.

## 2. HEALTH EFFECTS

### 2.7 POPULATIONS THAT ARE UNUSUALLY SUSCEPTIBLE

Case studies of acrylonitrile poisoning in humans following fumigation of living quarters in post-World War II Germany suggest that children are more susceptible to acrylonitrile than adults (Grunske 1949). Children died after sleeping in rooms recently fumigated with acrylonitrile for lice and bed bugs, while adults sharing the same quarters reported few, if any, effects (skin or eye irritation).

Studies in animals indicate that acrylonitrile can produce teratogenic effects at doses that have little maternal toxicity, suggesting that pregnant women may also be susceptible. It also seems likely that individuals in poor health or with respiratory problems might be particularly susceptible to acrylonitrile, but there are no data on this point.

### 2.8 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of acrylonitrile is available. Where adequate information is not available, ATSDR, in conjunction with the National Toxicology Program (NTP), is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of acrylonitrile.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that, if met would reduce or eliminate the uncertainties of human health assessment. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

#### 2.8.1 Existing Information on the Health Effects of Acrylonitrile

Figure 2-4 summarizes areas concerning the health effects of acrylonitrile where studies have and have not been performed. There are some data available on the effects in humans following acute or chronic exposure to acrylonitrile via the inhalation route of exposure. The target organ for acute toxicity is the nervous system. Chronic exposure to acrylonitrile has been associated with cancer. However, many of the available reports lack quantitative information on exposure levels. In humans, there are no data on oral exposure to acrylonitrile and a very limited amount of acute data on dermal exposure. In animals, there are data available on inhalation and oral exposure and limited dermal exposure data.

2. HEALTH EFFECTS

	Death	SYSTEMIC			Immunologic	Neurologic	Developmental	Reproductive	Genotoxic	Cancer
		Acute	Intermed.	Chronic						
Inhalation	●	●		●		●			●	●
Oral										
Dermal	●	●				●				

**HUMAN**

	Death	SYSTEMIC			Immunologic	Neurologic	Developmental	Reproductive	Genotoxic	Cancer
		Acute	Intermed.	Chronic						
Inhalation	●	●	●	●		●	●			●
Oral	●	●	●	●		●	●	●	●	●
Dermal	●									

**ANIMAL**

● Existing Studies

**FIGURE 2-4. Existing Information on Health Effects of Acrylonitrile**

## 2. HEALTH EFFECTS

### 2.8.2 Identification of Data Needs

**Acute-Duration Exposure.** Information is available regarding the effects of acute-duration inhalation exposure of humans to acrylonitrile and the effects are characteristic of cyanide-type toxicity. Quantitative data are limited but are sufficient to derive an acute inhalation MRL. Further studies of humans exposed to low levels of acrylonitrile in the workplace would increase the confidence of the acute MRL. Studies in animals support and confirm these findings. No studies are available on the effects of acute-duration oral exposure in humans; however, exposure to acrylonitrile reveals neurological disturbances characteristic of cyanide-type toxicity and lethal effects in rats and mice. Rats also develop birth defects. Animal data are sufficient to derive an acute oral MRL. Additional studies employing other species and various dose levels would be useful in confirming target tissues and determining thresholds for these effects. In humans, acrylonitrile causes irritation of the skin and eyes. No data are available on acute dermal exposures in animals.

**Intermediate-Duration Exposure.** No information is available on the effects of intermediate-duration inhalation, oral, or dermal exposure in humans or inhalation and dermal exposures in animals. There is information on intermediate-duration oral exposure in animals. Studies revealed decreased reproductive indices, decreased sperm count and tubular degeneration in rats and mice. Blood cell counts were low in rats and there was evidence of ulcerogenic effects in dogs. Acrylonitrile was also lethal in dogs. Data in animals were sufficient to derive an intermediate-duration oral MRL. Further studies in animals would be useful in defining threshold for these effects.

**Chronic-Duration Exposure and Cancer.** No studies were located evaluating chronic effects associated with acrylonitrile exposure in humans by any route of exposure. There is evidence of hematological effects and alterations in organ weights (kidney, liver) in rats following oral exposure. Additional chronic-duration studies in other species by inhalation and oral exposures would be useful in determining other target organs and thresholds for these effects. There are studies of humans chronically exposed to acrylonitrile in workplace air. These studies link acrylonitrile exposure to lung cancer and it has been suggested that the chemical may have the potential to cause prostate cancer. Due to limitations of these studies, firm conclusions cannot be made. Additional studies providing quantitative exposure data and evaluating confounding associations would be useful in clarifying potential risk.

## 2. HEALTH EFFECTS

**Genotoxicity.** Microbiol and mammalian assays are available on the genotoxic effects of acrylonitrile. Results of in vitro tests evaluating gene mutation in prokaryotic and eukaryotic cell types are variable. Similarly, results were mixed for in vitro mammalian assays evaluating gene mutation, chromosomal aberration and DNA repair and damage. In vivo tests evaluating chromosomal aberrations were negative. Additional investigations evaluating structural and numerical chromosomal aberrations in humans exposed in the workplace would be useful in determining the significance of effects found in animal assays.

**Reproductive Toxicity.** No studies have been conducted in humans regarding reproductive toxicity of acrylonitrile after inhalation, oral, or dermal exposure or inhalation and dermal exposures in animals. Studies in male mice have shown that exposure to acrylonitrile in drinking water affects sperm count and results in tubular degeneration. No effects on male reproductive organs have been reported in rats. Studies to further evaluate the significance of the testicular effects on reproductive capability in mice and other species would be very valuable. Studies on reproduction via the inhalation route would be valuable in determining whether the effects seen in oral studies are unique to that route of exposure.

**Developmental Toxicity.** No information is available on developmental effects of acrylonitrile in humans by any route of exposure. Acrylonitrile is teratogenic and embryotoxic in rats both by the oral and inhalation routes of exposure. Developmental studies on other animal species have not been conducted. Because species differences for acute acrylonitrile toxicity and metabolism have been demonstrated, additional developmental studies in other species using various dose levels would be valuable in evaluating the potential for acrylonitrile to cause developmental effects in humans. Because the available oral study was conducted by gavage, additional studies are needed to determine if these effects will occur following ingestion of drinking water or food.

**Immunotoxicity.** No information was found on the immunological effects of acrylonitrile in humans or animals by any route of exposure. Because no immunopathological effects have been reported in subchronic and chronic studies involving multiple species, additional studies employing a more specific testing battery are not warranted at this time.

**Neurotoxicity.** Clinical signs indicative of disturbances of the nervous system in exposed humans have been well documented in short-term studies at high doses and appear to be reversible. These effects are characteristic of cyanide toxicity. Animal studies confirm findings in



## 2. HEALTH EFFECTS

humans. In longer-term studies, effects on the nervous system have also been reported, but it is not certain if these effects are permanent or reversible following termination of acrylonitrile exposure.

**Epidemiological and Human Dosimetry Studies.** There are studies on the adverse effects of acrylonitrile in humans. These studies link acrylonitrile exposure and lung cancer. It has also been suggested that acrylonitrile may have the potential to cause prostate cancer. Many of the studies have major limitations including insufficient quantification of exposure, short follow-up, small study population, and inadequate evaluation of confounding associations. Additional studies would be useful in clarifying the cancer risk and estimating the exposure levels that lead to these effects.

**Biomarkers of Exposure and Effect.** The presence of thiocyanate and 2-cyanoethyl mercapturic acid have been monitored in urine as indicators of acrylonitrile exposure.

Effects produced by exposure to acrylonitrile, particularly after acute exposures, are characteristic of cyanide toxicity. These effects can be detected in people exposed by evaluating signs and symptoms such as limb weakness, labored and irregular breathing, dizziness and impaired judgement, cyanosis and convulsions. While tests are not specific for acrylonitrile-induced toxicity, they do identify potential health impairment. Studies to develop more specific biomarkers of acrylonitrile-induced effects would be useful in assessing the potential health risk of acrylonitrile near hazardous waste sites.

**Absorption, Distribution, Metabolism, and Excretion.** Metabolism and excretion in animals exposed to acrylonitrile by the inhalation and oral routes have been studied extensively. However, only limited data on absorption and distribution are available. Some data on humans exposed by inhalation are available. No data are available on the toxicokinetics of acrylonitrile when the exposure route is dermal. More extensive information on absorption and distribution of acrylonitrile would be valuable to fully understand the toxicokinetics of acrylonitrile. Some data on the toxicokinetics of acrylonitrile by the dermal route would be valuable in order to determine if metabolism of acrylonitrile differs by route of exposure. Development of physiologically-based toxicokinetic models would also be valuable in extrapolation of animal data to humans.

**Comparative Toxicokinetics.** The absorption, distribution, metabolism, and excretion of acrylonitrile in rats has been studied. Limited work in other species suggests that important species

## 2. HEALTH EFFECTS

differences do exist. Further evaluation of these differences, and comparison of metabolic patterns in humans with those of animals would assist in determining the most appropriate animal species for evaluating the hazard and risk of human exposure to acrylonitrile.

### **2.8.3 On-going Studies**

A number of research projects in progress are investigating the mechanism of toxicity and tumor formation of acrylonitrile. The projects are summarized in Table 2-6.

A number of on-going epidemiology studies are evaluating the effects of occupational exposure to acrylonitrile. Independent investigations are being conducted by Dr. R. D. Jones of the Employment Medical Advisory Service of the United Kingdom; Dr. L.J. Lucas of American Cyanamid Company; and Dr. L. I. Glass of SRA Technologies, Inc.

## 2. HEALTH EFFECTS

TABLE 2-6. On-going Studies on Acrylonitrile

Investigator	Affiliation	Research Description	Sponsor
D.D. Bigner	Duke University	Immunological and histological studies on brain tumors following acrylonitrile exposure	NCI
F.P. Guengerich	Vanderbilt University	Bioactivation and covalent binding	NIEHS
A. Segal	New York University	Tumor induction by acrylonitrile and its epoxide	NIEHS
-	CIIT	Comparative studies of the metabolism and pharmacokinetics of acrylonitrile and cyanoethylene oxide in mice and rats	BP
T.R. Fennell, J.P. MacNeela, M.J. Turner, et al.	CIIT	Hemoglobin adducts formed on administration of acrylonitrile (AN) to rats	BP
-	CIIT	Molecular studies of the mutagenic effects of acrylonitrile metabolites and investigation of the formation of DNA adducts	BP
T.R. Fennell, S.C. Sumner, S.D. Held, et al.	CIIT	Detection of eight urinary metabolites of [1,2,3- <sup>13</sup> C] acrylonitrile in the rat and mouse using <sup>13</sup> C nuclear magnetic resonance spectroscopy	-
G.L. Kedderis S.D. Held, R. Batru, et al.	CIIT	Dose-dependent urinary excretion of four acrylonitrile (ACN) metabolites in F-344 rats and B6C3F1 mice	-

## 2. HEALTH EFFECTS

Table 2-6 (Continued)

Investigator	Affiliation	Research Description	Sponsor
D.E. Rickert, A.E. Roberts, D. Pilon, et al.	CIIT	Distribution of acrylonitrile (ACN) in tissues of control and glutathione (GSH) depleted B6C3F1 mice	-
M.L. Gargas, G.L. Kedderis, T.R. Fennell, et al.	CIIT	A physiologically-based pharmacokinetic (PB-PK) model for acrylonitrile (ACN) in the rat	-
G.M.H. Swaen	University of Limburg, Netherlands	Mortality study with quantitative exposure assessment of 3,000 Dutch workers exposed to acrylonitrile until 1987 and 4,000 employed controls	-
-	CIIT	Mechanisms of acrylonitrile carcinogenicity in experimental test systems	BP
D.D. Bigner	Duke University	Tumor transplantation to determine whether acrylonitrile is genotoxic in the rat	-
C.H. Tamburro	University of Louisville	Health surveillance systems and molecular dosimetry	-
-	CIIT	Studies on the binding of acrylonitrile and its metabolites to hemoglobin for use as a biomarker	BP
-	CIIT	Development of a physiologically-based dosimetry model for acrylonitrile and cyanoethylene oxide	-

## 2. HEALTH EFFECTS

Table 2-6 (Continued)

Investigator	Affiliation	Research Description	Sponsor
C.H. Tamburro	University of Louisville	Development and implementation of prospective medical surveillance system for cancer control involving 1,200 active employees, 5,500 previous employees and 3,500 family members of employees from 1974	-
M. Dosemeci	National Cancer Institute	Case control epidemiology study concerning occupational exposure and cancer risk in Turkey in workers exposed between 1978 and 1984	-
L.M. Pottern	National Cancer Institute	Mortality study of 25,316 workers potentially exposed to acrylonitrile in eight U.S. facilities between 1952 and 1965	NCI/NIOSH
P.A. Stewart	National Cancer Institute	Assessment of historic exposures of acrylonitrile in eight U.S. facilities between 1952 and 1965	NCI/NIOSH

BP = British Petroleum America; CIIT = Chemical Industry Institute of Toxicology; NCI = National Cancer Institute; NIEHS = National Institute of Environmental Health Sciences; NIOSH = National Institute for Occupational Safety and Health.



### 3. CHEMICAL AND PHYSICAL INFORMATION

#### 3.1 CHEMICAL IDENTITY

Table 3-1 lists common synonyms, trade names and other pertinent identification information for acrylonitrile.

#### 3.2 PHYSICAL AND CHEMICAL PROPERTIES

Table 3-2 lists important physical and chemical properties of acrylonitrile.

## 3. CHEMICAL AND PHYSICAL INFORMATION

TABLE 3-1. Chemical Identity of Acrylonitrile

	Value	Reference
Chemical name	Acrylonitrile	NLM 1988
Synonyms	Cyanoethylene; 2-propenenitrile; vinyl cyanide	NLM 1988
Trade names	Acritet, Caswell No. 010, ENT 54, Fumigrain, Ventox	HSDB 1988; Windholz 1983
Chemical formula	C <sub>3</sub> H <sub>3</sub> N	NLM 1988
Chemical structure		
Identification numbers:		
CAS registry	107-13-1	NLM 1988
NIOSH RTECS	AT5250000	NIOSH 1981, 1982
EPA Hazardous Waste	U009	NLM 1988
OHM/TADS	7216574	HSDB 1988
DOT/UN/NA/IMCO	UN1093	NLM 1988
Shipping	IMCO 3.2	HSDB 1988
HSDB	176	NLM 1988
NCI	C50215	NLM 1988

CAS = Chemical Abstracts Service; NIOSH = National Institute for Occupational Safety and Health; RTECS = Registry of Toxic Effects of Chemical Substances; EPA = Environmental Protection Agency; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; DOT/UN/NA/IMCO = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute.



## 3. CHEMICAL AND PHYSICAL INFORMATION

TABLE 3-2. Physical and Chemical Properties of Acrylonitrile

Property	Value	Reference
Molecular weight	53.06	Weast 1985
Color	Colorless	Verschuieren 1983
Physical state	Liquid	Verschuieren 1983
Melting point	-83°C	Verschuieren 1983
Boiling point	77.4°C	Verschuieren 1983
Density at 20°C	0.8060	Weast 1985
Odor	Pungent (onion, garlic)	Verschuieren 1983
Odor threshold:		
Water	18.6 mg/L	Verschuieren 1983
Air	47 mg/m <sup>3</sup>	Verschuieren 1983
Solubility:		
Water at 20°C	79,000 mg/L	Klein et al. 1957
Organic solvents	Soluble in all common organic solvents	Sax and Lewis 1987
Partition coefficients:		
Log octanol/water	-0.92	Verschuieren 1983
Log K <sub>oc</sub>	-0.07	Mabey et al. 1982
Vapor pressure at 22.8°C	100 mmHg	Mabey et al. 1982
Henry's law constant:	8.8x10 <sup>-5</sup> atm-m <sup>3</sup> /mol	Mabey et al. 1982
Autoignition temperature	481°C	Sax 1984
Flashpoint (closed cup)	-1°C	Sax 1984
Flammability limits	3% to 17%	Sax and Lewis 1987
Conversion factors	1 ppm = 2.203 mg/m <sup>3</sup> 1 mg/m <sup>3</sup> =0.454 ppm	Verschuieren 1983



#### 4. PRODUCTION, IMPORT, USE, AND DISPOSAL

##### 4.1 PRODUCTION

Acrylonitrile is produced commercially by the process of propylene ammoxidation, in which propylene, ammonia and air are reacted in a fluidized bed in the presence of a catalyst (EPA 1984, 1985a). Production in the United States has increased gradually over the past 20 years from 304,300 kkg<sup>a</sup> in 1967 (Cogswell 1984) to 1,112,754 kkg in 1987 (USITC 1988).

Acrylonitrile is currently produced by five manufacturers at six locations in the United States: American Cyanamid Company, Avondale, Louisiana; BP America, Inc., Green Lake, Texas and Lima, Ohio; E.I. duPont de Nemours and Company, Inc., Beaumont, Texas; Monsanto Company, Chocolate Bayou, Texas; and Sterling Chemicals, Inc., Texas City, Texas (SRI 1988).

##### 4.2 IMPORT

Imports of acrylonitrile have been relatively small. In recent years, imports of acrylonitrile have decreased from a high of 7,000 kkg in 1974 to negligible level in 1981 and thereafter.

A substantial fraction of the acrylonitrile produced in the United States is exported. Exports rose from 23,500 kkg in 1972 to 364,000 kkg (39% of United States production) in 1982 (Cogswell 1984).

##### 4.3 USE

The primary use of acrylonitrile is as the raw material for the manufacture of acrylic and modacrylic fibers. Other major uses include the production of plastics (acrylonitrile-butadiene-styrene (ABS) and styrene-acrylonitrile (SAN)), nitrile rubbers, nitrile barrier resins, adiponitrile and acrylamide (EPA 1984).

Acrylonitrile has been used, in a mixture with carbon tetrachloride, as a fumigant for flour milling and bakery food processing equipment and for stored tobacco. However, most pesticide products containing acrylonitrile have been voluntarily withdrawn by the manufacturers (IARC 1979). Currently, acrylonitrile in combination with carbon tetrachloride is registered as a restricted-use pesticide.

---

<sup>a</sup>1 kkg = 1,000 kg (1 metric ton, equivalent to 2,200 pounds)

#### 4. PRODUCTION, IMPORT, USE, AND DISPOSAL

In 1982, 51% of the United States consumption of acrylonitrile was used for acrylic fibers, 18% for ABS and SAN resins, 14% for adiponitrile, 5% for acrylamide and 3% for nitrile elastomers. The remaining 9% was for miscellaneous uses (Cogswell 1984).

##### **4.4 DISPOSAL**

Because acrylonitrile is listed as a hazardous substance, disposal of waste acrylonitrile is controlled by a number of federal regulations (see Chapter 7). Rotary kiln, fluidized bed and liquid injection incineration are acceptable methods of acrylonitrile disposal (HSDB 1988). Underground injection is another disposal method. The most recent quantitative information on amount of acrylonitrile disposed in waste sites is for 1987. Emissions were 0.9 metric tons in surface water, 152 metric tons disposed through Publically Owned Treatment Works (POTW), 92 metric tons disposed of on land and 1,912 metric tons by underground injection (TRI 1988). Because acrylonitrile is relatively volatile and is also readily soluble in water, release to the environment from waste sites is of concern.

## 5. POTENTIAL FOR HUMAN EXPOSURE

### 5.1 OVERVIEW

Humans may be exposed to acrylonitrile through air, water and food. Acrylonitrile has been identified in 3 of 1,177 NPL sites. The frequency of these sites within the United States can be seen in Figure 5-1.

Acrylonitrile is readily volatile, and significant quantities escape into air during manufacture and use. Volatilization may also occur from hazardous waste sites. In air, acrylonitrile is degraded primarily by reaction with hydroxyl radicals, with an estimated half-life of 5 to 50 hours. Acrylonitrile has been detected in air in the vicinity of various industrial sources at concentrations from 0.1 to 325 ppb, but has not been detected in typical ambient air.

Acrylonitrile is also readily soluble in water, and current total discharges to water via industrial effluents are low. Water contamination may also occur following a spill or near a chemical site. In water, acrylonitrile has little tendency to adsorb to sediment, but is subject to biodegradation by microorganisms. The rate and extent of degradation depend upon conditions and upon the time for Microbiol acclimation. Degradation may approach 100% under favorable circumstances, but may be inhibited by high concentrations of acrylonitrile.

For members of the general public who do not live near an industrial source or a chemical waste site, exposure to very low levels of acrylonitrile may occur through contact with consumer products such as acrylic carpeting or by ingestion of food stored in acrylic plastic containers. Acrylonitrile may enter human food materials by leaching from plastic food containers. For people who do live near such a source, inhalation of acrylonitrile in air is likely to be the main route of exposure, although intake through water could also be of concern. Inhalation of acrylonitrile in air may be much higher for workers in industries that produce or use acrylonitrile. Acrylonitrile has been detected in surface water and groundwater near some industrial sources and hazardous waste sites, usually at concentrations between 20 and 4,700 ppb.

### 5.2 RELEASES TO THE ENVIRONMENT

#### 5.2.1 Air

Because acrylonitrile is readily volatile, significant releases to air may occur during acrylonitrile production and use (Hughes and Horn 1977; Miller and Villaume 1978). Kayser et al. (1982) estimated that 11,790 kkg/yr (metric tons per year) of acrylonitrile was released from these sources, accounting for 87% of all acrylonitrile released to the




FIGURE 5-1. FREQUENCY OF SITES WITH ACRYLONITRILE CONTAMINATION

## 5. POTENTIAL FOR HUMAN EXPOSURE

environment. Quantitative data on air releases from other sources were not located, but volatilization might be significant following a spill or in the vicinity of a chemical waste site containing acrylonitrile.

### 5.2.2 Water

Acrylonitrile may also be released to water during production and use. According to data collected under SARA 313, total reported releases to water during 1987 were 0.9 metric tons (TRI 1988). No data were located on acrylonitrile releases to water from other sources, but because acrylonitrile is readily soluble and is not strongly adsorbed to soil or sediment (see Section 5.3.1 below), large accidental spills or leaks from chemical waste sites could lead to significant water contamination. While acrylonitrile is rapidly biodegraded in water, high concentrations may inhibit degradation. Several examples of groundwater contamination following spills have been reported (Miller and Villaume 1978). Acrylonitrile may also be released to water by leaks or emissions from hazardous waste sites. Acrylonitrile has been detected in surface water at 2 sites and in groundwater at 5 sites of 862 hazardous waste sites (including NPL and other sites) being investigated under Superfund (CLPSD 1988).

### 5.2.3 Soil

Direct release of acrylonitrile to soil during acrylonitrile production and use is believed to be minimal (less than 1 kkg/yr) (Kayser et al. 1982). Accidental spills or leaks from hazardous waste sites could lead to local areas of soil contamination, and acrylonitrile has been detected in soil at 3 chemical waste sites (NPL and other sites) being investigated under Superfund (CLPSD 1988).

## 5.3 ENVIRONMENTAL FATE

### 5.3.1 Transport and Partitioning

Acrylonitrile is both readily volatile in air (0.13 atm at 23°C) (Mabey et al. 1982) and highly soluble in water (79,000 mg/L) (Klein et al. 1957). These characteristics dominate the behavior of acrylonitrile in the environment. While present in air, acrylonitrile has little tendency to adsorb to particulate matter (Cupitt 1980), so air transport of volatilized material is determined mainly by wind speed and direction. Similarly, acrylonitrile dissolved in water has only a low tendency to adsorb to suspended soils or sediments (Roy and Griffin 1985), so surface transport is determined by water flow parameters. Based on its relatively high water solubility, acrylonitrile is expected to be highly mobile in moist soils. In addition, acrylonitrile may

## 5. POTENTIAL FOR HUMAN EXPOSURE

penetrate into groundwater from surface spills or from contaminated surface water. The high vapor pressure indicates that evaporation from dry soil samples is expected to occur rapidly (EPA 1987).

The tendency of acrylonitrile to partition between air and water is described by Henry's law constant (H). The value of H for acrylonitrile has not been determined experimentally, but has been calculated to be  $8.8 \times 10^{-5}$  atm-m<sup>3</sup>/mole (Mabey et al. 1982). This value indicates that acrylonitrile will occur in both air and water, tending to transfer between air and water phases only slowly. Cupitt (1980) estimated the half-time of acrylonitrile clearance from air in wet precipitation to be greater than 10 months.


Based on the relatively low value of the octanol/water partition coefficient ( $K_{ow}$ ) for acrylonitrile ( $\log K_{ow} = -0.92$ ) (Verschueren 1983), it would not be expected that acrylonitrile will strongly bioaccumulate in the tissues of aquatic organisms (Kenaga 1980; Neely et al. 1974). However, data in aquatic organisms exposed to water containing acrylonitrile show that it does accumulate in fat tissue. Barrows et al. (1978) measured a steady-state bioconcentration factor (BCF) of 48 in bluegill sunfish. Based on the relative proportion of fat in sunfish and other aquatic organisms, EPA (1980a) estimated an average BCF of about 30 for the edible portions of freshwater and marine species.

### 5.3.2 Transformation and Degradation

#### 5.3.2.1 Air

The principal pathway leading to degradation of acrylonitrile in air is believed to be photooxidation, mainly by reaction with hydroxyl radicals (OH). The rate constant for acrylonitrile reaction with OH has been measured as  $4.1 \times 10^{-12}$  cm<sup>3</sup>/molecule/second (Harris et al. 1981). This would correspond to an atmospheric half-life of about 5 to 50 hours. This is consistent with a value of 9 to 10 hours measured in a smog chamber (Suta 1979).

The photooxidation of acrylonitrile by hydroxyl radicals in the presence of nitric oxide has been observed to yield formaldehyde (HCHO) and formyl cyanide (HCOCN) (Hashimoto et al. 1984). From these results, the following reaction was proposed:



Data given by Hashimoto et al. (1984) suggest that the half-life of acrylonitrile in the atmosphere may be on the order of 12 hours.



## 5. POTENTIAL FOR HUMAN EXPOSURE

Acrylonitrile may also be oxidized by other atmospheric components such as ozone and oxygen, but the rates of these reactions are much lower than for OH. and are not considered to be an important degradative pathway (Harris et al. 1981).

**5.3.2.2 Water**

Very little is known about nonbiologically mediated transformations of acrylonitrile in water. There are no data to suggest that acrylonitrile hydrolyzes under ambient conditions. While it is known that acrylonitrile photooxidizes in air, no reliable information was found on photochemical reactions in water. There were also no data on the oxidation of acrylonitrile in water. Acrylonitrile is susceptible to oxidation by strong oxidants such as chlorine used to disinfect drinking water.

Acrylonitrile is readily degraded by aerobic microorganisms in water, especially if there is time for acclimation (Cherry et al. 1956; Mills and Stack 1953, 1955; Stover and Kincannon 1983). After 27 days of acclimation, about 70% of the acrylonitrile initially present in river water was degraded under laboratory conditions, yielding acrylic acid and ammonia. Complete degradation occurred under ideal conditions where nutrients were added to promote Microbiol growth (Cherry et al. 1956).

A bacterium classified as Nocardia rhodochrous LL 100-2 has been reported to be able to degrade acrylonitrile (DiGeronimo and Antoine 1976). An aerobic bacterium classified as Arthrobacter in an acclimated sludge decompletely degraded acrylonitrile after 48 hours yielding acrylic acid (Yamada et al. 1979). It was proposed that acrylonitrile was biodegraded by the following reaction:



It has been shown that low concentrations of acrylonitrile in solution (10 mg/L or less) can be completely degraded in a laboratory, static-culture batch experiment where domestic sewage water was the source of the Microbiol inoculum (Tabak et al. 1981). A solution of acrylonitrile (152 mg/L) was degraded to less than 0.05 mg/L in a continuous flow activated sludge system under laboratory conditions (Kincannon et al. 1983).

Studies performed using sewage sludge indicate that acrylonitrile may also be degraded by methanogenic bacteria under anaerobic conditions, although concentrations of 50 to 1,000 mg/L lead to moderate inhibition of bacterial fermentation (Miller and Villaume 1978). This

## 5. POTENTIAL FOR HUMAN EXPOSURE

suggests that Microbiol degradation of acrylonitrile in anaerobic groundwater may not proceed efficiently if acrylonitrile levels were high, as might occur after a spill.

### 5.3.2.3 Soil

No studies were located regarding the biodegradation of acrylonitrile in soil. However, it seems likely that acrylonitrile in moist soil would be subject to biodegradation similar to that observed in aerobic water, although degradation rates might differ.

## 5.4 LEVELS MONITORED OR ESTIMATED IN THE ENVIRONMENT

### 5.4.1 Air

Acrylonitrile has not been found to occur at measurable concentrations in ambient air (Brodzinsky and Singh 1983). Measurable levels of atmospheric acrylonitrile are associated with industrial sources.

Air samples collected in one acrylonitrile-fiber plant ranged from 3 to 20 mg/m<sup>3</sup> (EPA 1980a). Mean 24-hour acrylonitrile concentrations in atmospheric samples collected within 5 km of 11 factories producing or using acrylonitrile ranged from less than 0.1 to 325 µg/m<sup>3</sup> (Suta 1979). The occurrence of acrylonitrile was correlated to wind patterns; the highest concentrations were downwind of and in close proximity to the plant. The median concentration of acrylonitrile for 43 measurements in "source-dominated areas" (i.e., near chemical plants) was 2.1 µg/m<sup>3</sup> (Brodzinsky and Singh 1983). There were no data available on the concentration of acrylonitrile in air near chemical waste sites, but because acrylonitrile is easily volatilized, this is an exposure pathway of concern.

### 5.4.2 Water

Acrylonitrile is not a common contaminant of typical surface water or groundwater. In a state-wide survey of over 1,700 wells in Wisconsin, acrylonitrile was not detected in any sample (Krill and Sonzogni 1986). Acrylonitrile was detected in 46 of 914 samples of surface water and groundwater taken across the United States (Staples et al. 1985), generally at levels less than 10 ppb.

The most likely source of acrylonitrile in water is industrial discharges. Levels of acrylonitrile measured in the effluents from a variety of industrial sites (iron and steel factories, textile mills, chemical plants) have ranged from 20 to 4,700 ppb, resulting in

## 5. POTENTIAL FOR HUMAN EXPOSURE

concentrations in nearby rivers ranging from below detection limits to 4,300 ppb (EPA 1983c). Data collected under SARA indicated total discharges to water during 1987 were 0.9 metric tons (TRI 1988).

Another source of acrylonitrile in water is leachate from chemical waste sites. Preliminary data from the Contract Laboratory Program (CLP) Statistical Database indicates that acrylonitrile has been detected in surface water samples collected at two of 862 hazardous-waste sites (including NPL and other sites) being investigated under Superfund. The median concentration of the positive samples was 100  $\mu\text{g/L}$  (CLPSD 1988). Acrylonitrile was detected in 12 groundwater samples collected at 5 sites, also at a median concentration of 100  $\mu\text{g/L}$ .

### 5.4.3 Soil

Staples et al. (1985) reported that acrylonitrile was not present at detectable concentrations in 351 sediment samples collected from lake and river bottoms across the United States. Preliminary data from the Contract Laboratory Program (CLP) Statistical Database (CLPSD 1988) indicated that acrylonitrile was detected in soils at 3 of 862 hazardous waste sites (including NPL and other sites) being investigated under Superfund. The median concentration of five samples was 120  $\mu\text{g/kg}$ .

### 5.4.4 Other Media

Foods may become contaminated with acrylonitrile as a result of the migration of the monomer from chemical containers made of acrylonitrile polymers. Acrylonitrile has been found to desorb from polyacrylonitrile resins and partition into cooking oil (Gilbert et al. 1980). Other foods which may be contaminated by acrylonitrile from their containers include luncheon meat, peanut butter, margarine, fruit juice, and vegetable oil (EPA 1980a, 1983c; FDA 1984). There are few data on the extent of food-related acrylonitrile exposure. The FDA reported typical acrylonitrile concentrations in margarine of 25  $\mu\text{g/kg}$  (FDA 1984), and the Commission of European Communities (1983) reported that the levels of acrylonitrile in contaminated foods are generally about 1  $\mu\text{g/kg}$ . While past data suggest potential exposure, there is little current migration of the monomer from current packaging materials because food is packaged in vastly different resins that have been drastically improved (AN Group 1990).

Acrylonitrile was detected in the smoke of cigarettes made in the United States in the 1960s and 1970s, usually at levels of 1 to 2 mg per cigarette (IARC 1979). At that time, acrylonitrile was used as a fumigant for stored tobacco. Most pesticide registrations for acrylonitrile were cancelled in 1978, and the use of acrylonitrile as a

## 5. POTENTIAL FOR HUMAN EXPOSURE

fumigant has been discontinued. Although no data were located, cigarette smoking is probably no longer a major source of human exposure to acrylonitrile.

Residual acrylonitrile monomer may also occur in commercially-made polymeric materials used in rugs and other products. Estimated levels include acrylic and modacrylic fibers (less than 1 mg acrylonitrile/kg polymeric material), acrylonitrile-based resins (15 to 50 mg/kg), and nitrile rubber and latex (0 to 750 mg/kg) (IARC 1979; Miller and Villaume 1978). It is possible that acrylonitrile may evaporate into air or leach into water from these products, but no data on this topic were located.

### 5.5 GENERAL POPULATION AND OCCUPATIONAL EXPOSURE

As shown in Table 5-1, only people living near chemical factories or work sites are likely to be exposed to measurable amounts of acrylonitrile in air and water. Dispersion modeling studies have indicated that approximately 2.6 million people living within 30 km of emission sources may be exposed to atmospheric acrylonitrile (Suta 1979). Members of the general population may also be potentially exposed to acrylonitrile through the consumption of acrylonitrile-contaminated food. However, it should be recalled that only foods in direct contact with acrylonitrile-based plastics are subject to contamination, and then only at very low levels.

Occupational exposures via inhalation of acrylonitrile vapor at the work place are likely to be considerably greater than exposures outside the workplace (Table 5-1). Exposure levels may be highest for workers in plants where the chemical is synthesized (EPA 1984).

### 5.6 POPULATIONS WITH POTENTIALLY HIGH EXPOSURES

It has been estimated that over 100,000 workers are potentially exposed to acrylonitrile during production and use (NIOSH 1977, 1988). Occupational exposures include plastic and polymer manufacturers, polymer molders, polymer combustion workers, furniture makers, and manufacturers of fibers and synthetic rubber (EPA 1980a). Other populations who could have elevated exposure to acrylonitrile are residents in the vicinity of industrial sources or chemical waste sites.

### 5.7 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of acrylonitrile is available. Where adequate information is not available, ATSDR, in conjunction with the NTP, is

## 5. POTENTIAL FOR HUMAN EXPOSURE

**TABLE 5-1. Estimated Levels of Human Exposure to Acrylonitrile for Nonoccupational and Occupational Exposure**

Population Type	Medium	Typical Concentration in Medium	Assumed Rate of Intake of Medium	Assumed Absorption Fraction	Estimated Dose ( $\mu\text{g}/\text{kg}/\text{day}$ )
General <sup>a</sup> (70-kg Adult)	Air	0.0 $\mu\text{g}/\text{m}^3$	20 $\text{m}^3/\text{day}$	0.9	0
	Water	0.0 $\mu\text{g}/\text{L}$	2 L/day	0.9	0
	Food	1 $\mu\text{g}/\text{kg}$	2 kg/day	0.5	0.01
Population living within 5 km of a chemical factory or waste site	Air	2 to 12 $\mu\text{g}/\text{m}^3$	20 $\text{m}^3/\text{day}$	0.9	0.5 to 3.0
	Water <sup>b</sup>	0.1 $\mu\text{g}/\text{L}$	2 L/day	0.9	0.003
Workers in an acrylonitrile factory	Air	0.1 to 4 $\text{mg}/\text{m}^3$	10 $\text{m}^3/\text{day}$	0.9	12.9 to 514

<sup>a</sup>Potential exposures from chemical spills and acrylic clothing were not considered.

<sup>b</sup>Untreated well water assuming waste effluent or leachate initially containing 10  $\mu\text{g}/\text{L}$  is reduced by a factor of 100 by groundwater dilution and biodegradation before it reaches the well.

## 5. POTENTIAL FOR HUMAN EXPOSURE

required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of acrylonitrile.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that, if met would reduce or eliminate the uncertainties of human health assessment. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

### 5.7.1 Identification of Data Needs

**Physical and Chemical Properties.** Most of the important physical-chemical properties of acrylonitrile have been determined (see Chapter 3). However, the partitioning of acrylonitrile between the air and water has been evaluated by using an estimated value for a Henry's law constant. This general approach assumes that the concentration of the chemical in water is low. Because acrylonitrile is relatively soluble in water, this approach may not be accurate. Experimental measurement of the partition coefficient for acrylonitrile at water-air interfaces would be useful in refining models on the behavior of acrylonitrile in the environment.

**Production, Use, Release and Disposal.** Substantial data exist on past production, use and emissions of acrylonitrile in the United States. Data on current releases to the environment and disposal practices are collected under SARA 313. Additional studies are not needed at this time because these data are readily available.

According to the Emergency Planning and Community Right to Know Act of 1986 (EPCRTKA), (§313), (Pub. L. 99-499, Title III, (§313), industries are required to submit release information to the EPA. The Toxic Release Inventory (TRI), which contains release information for 1987 and 1988, became available in May of 1989. This database will be updated yearly and should provide a more reliable estimate of industrial production and emission.

**Environmental Fate.** Laboratory studies indicate that acrylonitrile is biodegraded in aqueous systems promoting microbial growth, but typical degradation rates in lakes or rivers have not been studied in detail. Similarly, it is not known if acrylonitrile will biodegrade significantly in soil. Data on the chemical oxidation, photodegradation, and biodegradation of acrylonitrile in surface and groundwater would be helpful.

## 5. POTENTIAL FOR HUMAN EXPOSURE

**Bioavailability from Environmental Media.** There are limited data on the bioavailability of acrylonitrile in different environmental media. Data on the bioavailability of acrylonitrile would be valuable.

**Food Chain Bioaccumulation.** There are no data on the bioaccumulation of acrylonitrile in the food chain. The lack of data may not be a major limitation, because limited data suggest that acrylonitrile has a relatively low tendency to be bioconcentrated by lower trophic levels.

**Exposure Levels in Environmental Media.** Existing studies have not provided data on acrylonitrile levels in typical ambient air. Studies using analytical methods with lower detection limits would be helpful in determining if ambient air is an exposure medium of concern. Because higher levels of exposure are most likely near industrial sources or chemical waste sites, additional data on the occurrence of acrylonitrile in the atmosphere, surface water, and groundwater near such sites would be useful.

**Exposure Levels in Humans.** Human exposure levels to acrylonitrile can only be estimated based on average concentrations in air, food and water. Direct studies of personal exposure levels for individuals with exposures judged to be average and above average (e.g., people living near industrial sources or hazardous waste sites) would be helpful in improving total dose estimates, and in identifying exposure pathways of concern.

**Exposure Registries.** No exposure registries for acrylonitrile were located. This compound is not currently one of the compounds for which a subregistry has been established in the National Exposure Registry. The compound will be considered in the future when chemical selection is made for subregistries to be established. The information that is amassed in the National Exposure Registry facilitates the epidemiological research needed to assess adverse health outcomes that may be related to the exposure to this compound.

### 5.7.2 On-going Studies

The Acrylonitrile Group, Inc. is currently performing an acrylonitrile degradation study in groundwater and surrounding soil to obtain kinetic data for hydrolysis, biodegradation, primary metabolites and rates of mineralization. Radiolabeled material at three concentrations over three orders of magnitude will be used. The data will also establish the effect of accumulation on biodegradation kinetics.





## 6. ANALYTICAL METHODS

The purpose of this chapter is to describe the analytical methods that are available for detecting and/or measuring and monitoring acrylonitrile in environmental media and in biological samples. The intent is not to provide an exhaustive list of analytical methods that could be used to detect and quantify acrylonitrile. Rather, the intention is to identify well-established methods that are used as the standard methods of analysis. Many of the analytical methods used to detect acrylonitrile in environmental samples are the methods approved by federal agencies such as EPA and the National Institute for Occupational Safety and Health (NIOSH). Other methods presented in this chapter are those that are approved by a trade association such as the Association of Official Analytical Chemists (AOAC) and the American Public Health Association (APHA). Additionally, analytical methods are included that refine previously used methods to obtain lower detection limits, and/or to improve accuracy and precision.

### 6.1 BIOLOGICAL MATERIALS

Acrylonitrile in both biological and environmental samples is most commonly determined by gas chromatography with a nitrogen-phosphorus detector (GC/NPD) (Page 1985), gas chromatography/flame ionization detection (GC/FID) (EPA 1982a), or gas chromatography/mass spectroscopy (GC/MS) (Anderson and Harland 1980). Infrared spectroscopy (Jacobs and Syrjala 1978) may also be used. In the handling of samples and standards, acrylonitrile should be treated with precautions appropriate for a probable human carcinogen.

Comparatively little information is available in the literature on the determination of acrylonitrile in biological materials. Methods have been published for the determination of acrylonitrile in blood (Anderson and Harland 1980; Freshour and Melcher 1983) and urine (Houthuijs et al. 1982; Sakurai et al. 1978). Acrylonitrile was widely used in the past as a fumigant, and it is possible that methods used for the determination of other fumigants in biological materials (Walters 1986) could be adapted to the determination of acrylonitrile in such samples.

Methods for detection of acrylonitrile in biological materials are summarized in Table 6-1.

### 6.2 ENVIRONMENTAL SAMPLES

The methods of choice for the determination of acrylonitrile in environmental samples are GC/NPD (Page 1985), GC/FID (EPA 1982a), and GC/MS (EPA 1982b). Multiple internal reflectance infrared spectrometry (Jacobs and Syrjala 1978) is useful for monitoring low levels of acrylonitrile in air.

TABLE 6-1. Analytical Methods for Determining Acrylonitrile in Biological Materials

Sample Matrix	Sample Preparation	Analytical Method	Sample Detection Limit	Accuracy (% Recovery)	Reference
Blood	Perfusion of whole blood with nitrogen at 95°C in a perfusion flask, collection on Tenax	GC/MS	1 ng/mL	50-80	Anderson and Harland 1980
Plasma	Sample injected onto precolumn, water evaporated, analyte flushed from precolumn by heating	GC/NPD	2 ng/mL	No data	Freshour and Melcher 1983
Urine	Headspace gas	GC/NPD	2 ng/mL	3.3% <sup>a</sup>	Houthuijs et al. 1982
Urine	Azeotropic distillation	GC	5 ng/mL	No data	Sakurai et al. 1982

<sup>a</sup>Coefficient of variation

GC = gas chromatography; MS = mass spectrometry; NPD = nitrogen-phosphorus detector.

## 6. ANALYTICAL METHODS

The determination of acrylonitrile in air may be accomplished by collection on a solid sorbet, such as activated charcoal, followed by elution and gas chromatographic measurement (NIOSH 1984).

EPA has published a method of analysis specific for acrylonitrile in water (EPA 1982a) and a method applicable to its determination in water along with other purgeable organics (EPA 1982b). Other standard EPA methods are adaptable for the determination of acrylonitrile in wastes (EPA 1986b).

Methods for the determination of acrylonitrile in environmental samples are summarized in Table 6-2.

### 6.3 ADEQUACY OF THE DATABASE

Section 104(i)(5) of CERCLA, directs the Administrator of ATSDR (in consultation with the Administrator of EPA and agencies and programs of the Public Health Service) to assess whether adequate information on the health effects of acrylonitrile is available. Where adequate information is not available, ATSDR, in conjunction with the NTP, is required to assure the initiation of a program of research designed to determine the health effects (and techniques for developing methods to determine such health effects) of acrylonitrile.

The following categories of possible data needs have been identified by a joint team of scientists from ATSDR, NTP, and EPA. They are defined as substance-specific informational needs that, if met would reduce or eliminate the uncertainties of human health assessment. In the future, the identified data needs will be evaluated and prioritized, and a substance-specific research agenda will be proposed.

#### 6.3.1 Identification of Data Needs

**Method for Determining Biomarkers of Exposure and Effect.** Exposure to acrylonitrile can be determined by measuring parent acrylonitrile, acrylonitrile metabolites and adducts. Good methods exist for detecting and quantifying acrylonitrile in biological materials. These methods are based on capillary gas chromatography, where a special problem is introduction of analytes quantitatively and in a narrow band onto the capillary GC. One of the most promising approaches to analyte transfer is the use of cyrofocusing, in which analytes are condensed in a narrow band on a trap or on the GC column itself. Improvements in cryofocusing of volatile organic analytes for capillary GC determination (Washall and Wampler 1988) should improve sensitivity for the determination of acrylonitrile.

TABLE 6-2. Analytical Methods for Determining Acrylonitrile in Environmental Samples

Sample Matrix	Sample Preparation	Analytical Method	Sample Detection Limit	Accuracy	Reference
Air	Direct monitoring in ambient air	IR	0.2 ppm (volume)	±5% at 20 ppm	Jacobs and Syrjala 1978
Air	Collect on charcoal, elute to GC	GC/FID	0.015 mg/sample	Not evaluated	NIOSH 1984
Air	Charcoal adsorption, elute with acetone, inject acetone solution	GC/NPD	10 ppb (volume, estimated)	70-82% recovery	Marano et al. 1978
Water	Purge, trap on soil sorbent trap, heat and desorb, backflush to GC	GC/IDMS	50 µg/L	Not evaluated	EPA 1980b
Water	Purge at 85°C, trap on Tenax sorbent trap, heat and desorb, backflush to GC	GC/FID	0.5 µg/L	107 ± 5.6% recovery	EPA 1982a
Water	Purge, trap on Tenax and silica gel, heat and desorb, backflush to GC	GC/MS	No data	Not evaluated	EPA 1982b
Soil, sediment, solid waste	Isolation of volatile acrylonitrile from sample followed by GC separation	GC/FID	0.5 µg/L	84 - 104% recovery	EPA 1986b

IR = infrared spectrometry; GC = gas chromatography; FID = flame ionization detector; NPD = nitrogen-phosphorous detector; IDMS = isotope dilution mass spectrometry; MS = mass spectrometry.

## 6. ANALYTICAL METHODS

Acrylonitrile metabolites have been measured in blood and urine, but, except for measurement of thiocyanate, these methods have not been developed for routine monitoring of exposed humans. Supercritical fluid extraction/chromatography and immunoassay analysis are two areas of intense current activity from which substantial advances in the determination of acrylonitrile and its metabolites in biological samples can be anticipated. The two techniques are complementary because supercritical fluid extraction is especially promising for the removal of analytes from sample material and immunoassay is very analyte-selective and sensitive (Vanderlaan et al. 1988).

Studies using radioactivity-labeled acrylonitrile indicate that acrylonitrile or its metabolites form covalent adducts with cellular macromolecules in most tissues. Studies to develop chemical or immunological methods for measuring these adducts would be especially valuable in detecting and perhaps even quantifying human exposure to acrylonitrile. Adverse health effects demonstrated following exposure to acrylonitrile, particularly acute exposures, were characteristic of cyanide toxicity. Because these effects are also indicative of exposure to many other toxicants, additional methods are needed for more specific biomarkers of effects of acrylonitrile exposure.

**Methods for Determining Parent Compounds and Degradation Products in Environmental Media.** Methods for determining acrylonitrile in environmental samples are quite good. It may be assumed that the normal incentives for both research and the development of commercial methods of analysis will result in new analytical methods for acrylonitrile that have improved sensitivity and selectivity. Degradation products of acrylonitrile in environmental media are difficult to determine. This difficulty is not as much an analytical problem as it is a problem of knowing the fundamental environmental chemistry of these compounds in water, soil, air and biological systems.

### 6.3.2 On-going Studies

There exists an ongoing effort to develop a "Master Analytical Scheme" for organic compounds in water (Michael et al. 1988). The overall goal is the development of technology capable of detecting and quantitatively measuring organic compounds at 0.1 µg/L in effluent waters. Analytes are to include numerous semivolatile compounds and some compounds that are only "semisoluble" in water, as well as volatile compounds (boiling point, less than 150°C).

Studies designed to improve the determination of environmental contaminants will continue to provide refinements and improvements in the determination of acrylonitrile. The current high level of activity in supercritical fluid extraction of solid and semisolid samples should

## 6. ANALYTICAL METHODS

yield improved recoveries and sensitivities for the determination of acrylonitrile in solid wastes, and the compound should be amenable to supercritical fluid chromatographic analysis. Immunoassay analysis is another area of intense current activity from which substantial advances in the determination of acrylonitrile in environmental samples can be anticipated (Vanderlaan et al. 1988).

## 7. REGULATIONS AND ADVISORIES

Because of its potential to cause adverse health effects in exposed people, a number of regulations and guidelines have been established for acrylonitrile by various national and state agencies. These values are summarized in Table 7-1.

## 7. REGULATIONS AND ADVISORIES

TABLE 7-1. Regulations and Guidelines Applicable to Acrylonitrile

Agency	Description	Value <sup>a</sup>	Reference
IARC	Carcinogenic classification	Group 2A <sup>b</sup>	IARC 1982
<b><u>National</u></b>			
Regulations:			
a. Air:			
OSHA	PEL TWA Ceiling	2 ppm 10 ppm	OSHA 1978a, 1978b 29 CFR 1910.1045
b. Water:			
EPA OWRS	General permits under the NPDES	NA	40 CFR 122 Appendix D Table II
	Criteria and Standards for the NPDES	NA	40 CFR 125
	General pretreatment regulations for existing and new sources of pollution	NA	40 CFR 403
	Hazardous substance Reportable quantity	NA 100 lb	EPA 1985b, (40 CFR 116) 40 CFR 117.3
c. Nonspecific media:			
EPA OERR	Reportable quantity	100 lb	EPA 1989c, (40 CFR 302.4)
	Extremely hazardous substance list threshold planning quantity	10,000 lb	EPA 1987d, (40 CFR 355)
EPA OPP	Restricted use pesticide	NA	40 CFR 162.31
EPA OSW	Hazardous waste constituent (Appendix VIII)	NA	EPA 1980c, (40 CFR 261)
	Groundwater monitoring list (Appendix IX)	NA	EPA 1987e, (40 CFR 264)
	Restriction on land disposal, proposed treatment standards	NA	EPA 1989b, (40 CFR 268.10)
EPA OTS	Toxic chemical release reporting	NA	EPA 1988, (40 CFR 372)
FDA	Use of copolymers as components of food packaging materials	NA	21 CFR 177.1030, 177.1040
Guidelines:			
a. Air:			
ACGIH	TLV TWA	2 ppm (4.5 mg/m <sup>3</sup> )	ACGIH 1986
NIOSH	Recommended exposure limit guidelines for occupational exposure		NIOSH 1985, 1988b
	TWA	1 ppm	
	Ceiling (15 minutes)	10 ppm	
	IDLH	4000 ppm	
b. Water:			
EPA OWRS	Ambient water quality criteria		
	Ingesting water and organisms	5.8x10 <sup>-5</sup> mg/L	EPA 1980d
	Ingesting organisms only	6.5x10 <sup>-4</sup> mg/L	EPA 1980d







**8. REFERENCES**

- \* ACGIH. 1986. Documentation of the threshold limit values and biological exposure indices. 5th ed. American Conference of Governmental Industrial Hygienists. Cincinnati, OH.
- Adams M, Collins M. 1988. Sensitive portable gas chromatograph with data retrieval and communications capability for remote surveillance of toxic gases and vapors in plant. Anal Proc 25:190-191.
- Ahmed AE, Abreu ME. 1982. Microsomal metabolism of acrylonitrile in liver and brain. Adv Exp Med Biol 136B:1229-1238.
- Ahmed AE, Farooqui, MYH. 1982. Comparative toxicities of aliphatic nitriles. Toxicol Lett 12:157-163.
- \* Ahmed AE, Patel K. 1981. Acrylonitrile: In vivo metabolism in rats and mice. Drug Metab Dispos 9:219-222.
- \* Ahmed AE, Farooqui MY, Upreti RK, et al. 1982. Distribution and covalent interactions of [1-<sup>14</sup>C]acrylonitrile in the rat. Toxicology 23:159-175.
- \* Ahmed AE, Farooqui MYH, Upreti RK, et al. 1983. Comparative toxicokinetics of 2,3-<sup>14</sup>C- and 1-<sup>14</sup>C-acrylonitrile in the rat. J Appl Toxicol 3:39-47.
- Alridge WN. 1944. A new method for the estimation of micro quantities of cyanide and thiocyanate. Analyst 69:262-265.
- \* Amacher DE, Turner GN. 1985. Tests for gene mutational activity in L5178Y/TK assay system. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 487-496.
- Amoore JE, Hautala E. 1983. Odor as an aid to chemical safety: Odor thresholds compared with threshold limit values and volatilities for 214 industrial chemicals in air and water dilution. J Appl Toxicol 3:272-290.
- \* AN Group. 1990. Public comments on scientific and technical issues on the Draft Toxicological Profile for Acrylonitrile. Submitted to Agency for Toxic Substances and Disease Registry (ATSDR) February 15.

---

\* - Cited in text.

## 8. REFERENCES

- \* Anderson D, Cross MF. 1985. Suitability of the P388F mouse lymphoma system for detecting potential carcinogens and mutagens. *Food Chem Toxicol* 23:115-118.
- \* Anderson RA, Harland WA. 1980. The analysis of volatiles in blood from fire fatalities. In: Oliver JS, ed. *Forensic Toxicology. Proc. Eur. Meet. Int. Assoc. Forensic Toxicol.* Baltimore, MD: University Park Press, 279-292.
- Appel KE, Peter H, Bolt M, et al. 1981a. Effect of potential antidotes on the acute toxicity of acrylonitrile. *Int Arch Occup Environ Health* 49:157-163.
- Appel KE, Peter H, Bolt M, et al. 1981b. Interaction of acrylonitrile with hepatic microsomes of rats and men. *Toxicol Lett* 7:335-339.
- \* Arni P. 1985. Induction of various genetic effects in the yeast Saccharomyces cerevisiae strain D7. In: Ashby J, de Serres FJ, et al., eds. *Progress in mutation research, Vol. 5. Evaluation of short-term tests for carcinogens.* Amsterdam, The Netherlands: Elsevier Science Publishers, 217-224.
- Atkinson R, Carter WPL, Aschmann SM, et al. 1985. Atmospheric fates of organic chemicals: Prediction of ozone and hydroxyl radical reaction rates and mechanisms. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/3-85/063. NTIS No. PB85-241529.
- \* Back KC, Thomas AA, MacEwen JD. 1972. Reclassification of materials listed as transportation health hazards. Washington DC: Department of Transportation, Office of Hazardous Materials. NTIS No. PB-214270.
- \* Baker RS U, Bonin AM. 1985. Tests with the Salmonella plateincorporation assay. In: Ashby J, de Serres FJ, et al., eds. *Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens.* Amsterdam, The Netherlands: Elsevier Science Publishers, 177-180.
- \* Banerjee S, Segal A. 1986. In vitro transformation of C3H/10T1/2 and NIH/3T3 cells by acrylonitrile and acrylamide. *Cancer Lett* 32:293-304.
- \* Barnes D, Bellin J, DeRosa C, et al. 1987. Reference dose (RfD): Description and use in health risk assessments. Vol. I. Appendix A: Integrated risk information system supportive documentation. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. EPA/600/8-86/032a.

## 8. REFERENCES

- Barrett WJ, Dillon HK, James RH. 1974. Sampling and analysis of four organic compounds using solid sorbents, No. 1. Cincinnati, OH: National Institute for Occupational Safety and Health. NTIS No. PB82-151861.
- \* Barrows ME, Petrocelli SR, Macek KJ, et al. 1978. Bioconcentration and elimination of selected water pollutants by bluegill sunfish. Proceedings - American Chemical Society, Division of Environmental Chemistry. 18:345-346.
- \* Baxter RA. 1979. Evaluation and control of industrial exposure to acrylonitrile. Ann Occup Hyg 22:429-435.
- \* Beliles RP, Paulin HJ, Makris NG, et al. 1980. Three-generation reproduction study of rats receiving acrylonitrile in drinking water. Chemical Manufacturers Association. Washington, DC. LB1 Project No. 2660, February.
- Berg GL, ed. 1984. Farm chemicals handbook 1984. Willoughby, OH: Meister Publishing Co., C5, C44.
- Betso SR, McLean JD. 1976. Determination of acrylamide monomer by differential pulse polarography. Anal Chem 48:766-770.
- \* Bigner DD, Bigner SH, Burger PC, et al. 1986. Primary brain tumors in Fischer 344 rats chronically exposed to acrylonitrile in their drinking water. Food Chem Toxicol 24:129-137.
- \* Bio/dynamics. 1980a. A twenty-four month oral toxicity/carcinogenicity study of acrylonitrile administered to Spartan rats in the drinking water. Biodynamics, Inc., Division of Biology and Safety Evaluation, East Millstone, NJ. Project No. BDN-77-28.
- \* Bio/dynamics. 1980b. A twenty-four month oral toxicity/carcinogenicity study of acrylonitrile administered in the drinking water to Fischer 344 rats. Biodynamics, Inc., Division of Biology and Safety Evaluation, East Millstone, NJ. Project No. BDN-77-27.
- \* Bio/dynamics. 1980c. A twenty-four month oral toxicity/carcinogenicity study of acrylonitrile administered by intubation to Spartan rats. Biodynamics, Inc., Division of Biology and Safety Evaluation, East Millstone, NJ. Project No. BDN-77-29.
- Bishop CS, Dye A. 1982. Microwave heating enhances the migration of plasticizers out of plastics. J Environ Health 44:231-235.

## 8. REFERENCES

- Bocek K. 1976. Relationships among activity coefficients, partition coefficients and solubilities. In: Tichy M, ed. Quantitative structure-activity relationships. Basel and Stuttgart: Birkhauser Verlag, 231-240.
- Bolt HM, Laib RJ, Peter, H et al. 1986. DNA adducts of halogenated hydrocarbons. J Cancer Res Clin Oncol 112:92-96.
- Borders RA, Gluck SJ, Sowle WF, et al. 1986. Development and validation of personal monitoring methods for low levels of acrylonitrile in workplace atmosphere: II. Thermal desorption and field validation. Am Ind Hyg Assoc J 47:158-163.
- Bradley MO. 1985. Measurement of DNA single-strand breaks by alkaline elution in rat hepatocytes. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 353-357.
- \* Brat SV, Williams GM. 1982. Hepatocyte-mediated production of sister chromatid exchange in co-cultured cells by acrylonitrile: Evidence for extra cellular transport of a stable reactive intermediate. Cancer Lett 17:213-216.
- Brieger H, Rieders F, Hodes WA. 1951. Acrylonitrile: spectrophotometric determination, acute toxicity and mechanism of action. Arch Ind Health Occup Toxicol 6:128-140.
- \* Brodinzky R, Singh HB. 1983. Volatile organic chemicals in the atmosphere: An assessment of available data. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA-600/3-83-027(A).
- \* Brooks TM, Gonzalez LP, Calvert R, et al. 1985. The induction of mitotic gene conversion in the yeast Saccharomyces cerevisiae strain JD1. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 225-228.
- Buchter A, Peter H. 1984. Clinical toxicology of acrylonitrile. G Ital Med Law 6:83-86.
- Cadena F, Eiceman GA, Vandiver VJ. 1984. Removal of volatile organic pollutants from rapid streams. J Water Pollut Control Fed 5:460-463.

## 8. REFERENCES

- CCTTE. 1988. Computerized Listing of Chemicals Being Tested for Toxic Effects. United Nations Environment Programme, International Programme on Chemical Safety, International Register of Potentially Toxic Chemicals, Geneva, Switzerland.
- \* Chen JL, Walrath J, O'Berg MT, et al. 1987. Cancer incidence and mortality among workers exposed to acrylonitrile. *Am J Ind Med* 11:157-163.
- \* Cherry AB, Gabaccia AJ, Senn HW. 1956. The assimilation behavior of certain toxic organic compounds in natural water. *Sewage and Industrial Wastes*. 28:1137-1146.
- Clamsky KB, ed. 1986. Chemical guide to the OSHA hazard communication standard. Burlingame, CA: Roytech Publications, Inc., 68-E-3.
- CLC. 1989. Coordinated List of Chemicals. U.S. Environmental Protection Agency, Office of Research and Development, Washington, DC.
- \* CLPSD. 1988. Contract Laboratory Program Statistical Database. Viar and Company, Management Services Division, Alexandria, VA. December 1988.
- \* Cogswell SA. 1984. CEH marketing research report: acrylonitrile. In: Chemical economics handbook. Menlo Park, CA: SRI International.
- \* Collins JJ, Page LC, Caporossi JC, et al. 1989. Mortality patterns among employees exposed to acrylonitrile. *J Occup Med* 31:368-371.
- \* Commission of the European Communities. 1983. Reports of the scientific committee for food (thirteenth series). Luxembourg. NTIS No. PB83-172759.
- Connor TH, Forti GC, Sitra P, et al. 1979. Bile as a source of mutagenic metabolites produced in vivo and detected by Salmonella typhimurium. *Environ Mutagen* 1:269-276.
- Cooke M, Degner KB, Osborne MC, et al. 1984. Candidate sampling and analysis methods for twenty-one suspect carcinogens in combustion emissions. Proceedings - Air Pollution Control Association Annual Meeting, 77th 84-12.4.
- \* Cote IL, Bowers A, Jaeger RJ. 1983. Induced tolerance to acrylonitrile toxicity by prior acrylonitrile exposure. *Res Commun Chem Pathol Pharmacol* 42:169-172.

## 8. REFERENCES

- \* Cote IL, Bowers A, Jaeger RJ. 1984. Effects of acrylonitrile on tissue glutathione concentrations in rat, mouse, and hamster. *Res Commun Chem Pathol Pharmacol* 43:507-510.
- Crane CR, Sanders DC, Endecott BR, et al. 1986. Inhalation toxicology: VI. Evaluation of the relative toxicity of thermal decomposition products from nine aircraft panel materials. Washington, DC: Federal Aviation Administration, Office of Aviation Medicine. DOT/FAA/AM-86/3.
- \* Crespi CL, Ryan CG, Seixas GM, et al. 1985. Tests for mutagenic activity using mutation assays at two loci in the human lymphoblast cell lines TK6 and AHH-1. In: Ashby J, de Serres FJ, et al., eds. *Progress in mutation research*. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 497-516.
- \* Cupitt LT. 1980. Fate of toxic and hazardous materials in the air environment. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA-600/3-80-084. NTIS No. PB80-221948.
- \* De Meester C, Poncelet F, Roberfroid, et al. 1978. Mutagenicity of acrylonitrile. *Toxicology* 11:19-27.
- De Meester C, Duverger-Van Bogaert M, Lambotte-Vandepaer M, et al. 1979. Liver extract mediated mutagenicity of acrylonitrile. *Toxicology* 13:7-15.
- \* Delzell E, Monson RR. 1982. Mortality among rubber workers: VI. Men with potential exposure to acrylonitrile. *J Occup Med* 24:767-769.
- \* DiGeronimo MJ, Antoine AD. 1976. Metabolism of acetonitrile and propionitrile by *Nocardia rhodochrous* LL100-21. *Appl Environ Microbiol* 31:900-906.
- Dorigan J, Fuller B, Duffy R. 1976. Preliminary scoring of organic air pollutants: Chemistry, production and toxicity of selected synthetic organic chemicals (chemicals A-C) Appendix I. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. EPA-450/3-77-008b. NTIS No. PB-264443.
- Dossing M, Ranek L. 1984. Isolated liver damage in chemical workers. *Br J Ind Med* 41:142-144.
- \* Dudley HC, Neal PA. 1942. Toxicology of acrylonitrile (vinyl cyanide): I. A study of the acute toxicity. *Journal of Industrial Hygiene and Toxicology* 24:27-36.



## 8. REFERENCES

- Duverger M, Lambotte M, Malvoisin E, et al. 1981a. Metabolic activation and mutagenicity of 4 vinylic monomers (vinyl chloride, styrene, acrylonitrile, butadiene). *Toxicol Eur Res* 3:131-140.
- Duverger-Van Bogaert M, Lambotte-Vandepaer M, de Meester C, et al. 1981b. Effect of several factors on the liver extract mediated mutagenicity of acrylonitrile and identification of four new in vitro metabolites. *Toxicol Lett* 7:311-319.
- Duverger-van Bogaert M, Lambotte-Vandepaer M, Mercier M, et al. 1982a. In vitro covalent binding of acrylonitrile to rat liver proteins. *Toxicol Lett* 13:203-209.
- Duverger-van Bogaert M, Lambotte-Vandepaer M, de Meester C, et al. 1982b. Role of glutathione in liver-mediated mutagenicity of acrylonitrile. *Toxicol Lett* 11:305-311.
- Duverger-van Bogaert M, Lambotte-Vandepaer M, de Meester C. 1982c. Vinyl chloride and acrylonitrile: Activation mechanism and mutagenicity. *Toxicol Eur Res* 4:35-37.
- Duverger M, Lambotte M, Mercier, et al. 1983. Role of glutathione in the in vivo and in vitro mutagenicity of acrylonitrile. *Mutat Res* 113:247-248.
- Edney E, Mitchell S, Bufalini JJ. 1982. Atmospheric chemistry of several toxic compounds. Research Triangle Park, NC: U.S.
- Environmental Protection Agency, Office of Research and Development. EPA-600/3-82-092. NTIS No. PB83-146340.
- \* EPA. 1980a. Ambient water quality criteria for acrylonitrile. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards. EPA 440/5-80-017. NTIS No. PB81-111285.
- \* EPA. 1980b. Volatile organic compounds by purge-and-trap isotope dilution GC-MS. Method 1624. Washington, DC: U.S. Environmental Protection Agency.
- \* EPA. i9aoc. U.S. Environmental Protection Agency. Federal Register. 45:33084-33133.
- \* EPA. 1980d. U.S. Environmental Protection Agency. Part V. Federal Register. 45:79318-79379.

## 8. REFERENCES

- \* EPA. 1982a. Acrolein and acrylonitrile - method 603. In: Longbottom JE, Lichtenberg JJ, eds. Methods for organic chemical analysis of municipal and industrial wastewater. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory. EPA-600/4-82-057.
- \* EPA. 1982b. Purgeables - method 624. In: Longbottom JE, Lichtenberg JJ, eds. Methods for organic chemical analysis of municipal and industrial wastewater. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory. EPA-600/4-82-057.
- \* EPA. 1983a. Treatability manual: Volume I. Treatability data. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA-600/2-82-001a.
- \* EPA. 1983b. Reportable quantity document for acrylonitrile. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Criteria and Assessment Office. External Review Draft ECAO-CIN-R005.  
  
EPA. 1983c. Health assessment document for acrylonitrile. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA-600/8-82-007F.
- \* EPA. 1984. Locating and estimating air emissions from sources of acrylonitrile. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. EPA-450/4-84-007a.
- \* EPA. 1985a. Health and environmental effects profile for acrylonitrile. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/X-85/372. NTIS No. PB88-170832.
- \* EPA. 1985b. U.S. Environmental Protection Agency. Part II. Federal Register. 50:13456-13522.  
  
EPA. 1985c. U.S. Environmental Protection Agency. Federal Register 50:24319-24321.
- \* EPA. 1986a. Reference values for risk assessment. Final draft. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Solid Waste. ECAO-CIN-477.
- \* EPA. 1986b. Gas chromatographic analysis of acrolein, acrylonitrile, and acetonitrile. Method 8030. In: Test methods for evaluating solid wastes. SW-846. 3rd ed. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.

## 8. REFERENCES

- \* EPA. 1987a. Health effects assessment for acrylonitrile. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/8-88/014. NTIS No. PB88-179411.
  
- EPA. 1987b. Toxic air pollutant/source crosswalk: A screening tool for locating possible sources emitting toxic air pollutants. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. EPA-450/4-87-023a.
  
- EPA. 1987c. U.S. Environmental Protection Agency. Part II Federal Register. 52:8140.
  
- \* EPA. 1987d. U.S. Environmental Protection Agency. Part II. Federal Register. 52:13378-13410.
  
- \* EPA. 1987e. U.S. Environmental Protection Agency. Part II. Federal Register. 52:25942-25953.
  
- \* EPA. 1988. U.S. Environmental Protection Agency. Part II. Federal Register. 53:4500-4539.
  
- \* EPA. 1989a. Interim Methods for Development of Inhalation Reference Doses. U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. Washington, DC. EPA 600/8-88/066F.
  
- \* EPA. 1989b. U.S. Environmental Protection Agency. Part II. Federal Register. 54:1056-1120.
  
- \* EPA, 1989c. U.S. Environmental Protection Agency. Part II. Federal Register. 54:33425-33431.
  
- EXICHEM Database. 1989. Organization for Economic Cooperation and Development.
  
- Fanini D, Trieff NM, Sadagopa Ramanujam VM, et al. Effect of acute acrylonitrile exposure on metrazol induced seizures in the rat. *Neurotoxicology* 6:29-34.
  
- \* Farooqui MYH, Ahmed AE. 1982. Molecular interaction of acrylonitrile and potassium cyanide with rat blood. *Chem Biol Interact* 38:145-159.
  
- \* Farooqui MYH, Ahmed AE. 1983a. The effects of acrylonitrile on hemoglobin and red cell metabolism. *J Toxicol Environ Health* 12:695-707.
  
- Farooqui MYH, Ahmed AE. 1983b. In vivo interactions of acrylonitrile with macromolecules in rats. *Chem Biol Interact* 47:363-371.

## 8. REFERENCES

- \* FDA. 1984. Food and Drug Administration. Federal Register 49:36635-36644.
- Fennell TR, MacNeela JP, Turman MN, et al. 1989. Hemoglobin adducts formed on administration of acrylonitrile (AN) to rats. CIIT, Research Triangle Park, NC.
- Fennell TR, Sumner SC, Held SD, et al. 1990. Detection of urinary metabolites of [1,2,3-<sup>13</sup>C] acrylonitrile in the rat and mouse detected by <sup>13</sup>C nuclear magnetic resonance spectroscopy. CIIT, Research Triangle Park, NC.
- Fossa AK, Riano MM, Lavin PJ, et al. 1984. New York State techniques and experience in regulating toxic air contaminants. Proceedings - Air Pollution Control Association Annual Meeting 77. 84-6.2.
- Freed DJ, Majsce AM. 1977. In situ generation of standards for gas chromatographic analysis. Anal Chem 49:139-141.
- \* Freshour NL, Melcher RG. 1983. Analytical method for the determination of acrylonitrile in rat plasma at the nanograms per milliliter level. J Anal Toxicol 7:103-105.
- \* FSTRAC. 1988. Summary of state and federal drinking water standards and guidelines. Washington, DC: Federal-State Toxicology and Regulatory Alliance Committee, Chemical Communication Subcommittee.
- \* Fujikawa K, Ryo H, Kondo S. 1985. The Drosophila reversion assay using the unstable zeste-white somatic eye color system. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 319-324.
- Fujisawa S, Masuhara E. 1981. Determination of partition coefficients of acrylates, methacrylates, and vinyl monomers using high performance liquid chromatography (HPLC). J Biomed Mater Res 15:787-793.
- \* Gallagher GT, Maul EA, Kovacs K, et al. 1988. Neoplasms in rats ingesting acrylonitrile for two years. J Am Coll Toxicol 7:603-615.
- \* Garner RC, Campbell J. 1985. Tests for the induction of mutations to ouabain or 6-thioguanine resistance in mouse lymphoma L5178Y cells. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 525-529.

## 8. REFERENCES

- Garrett NE, Stack HF, Gross MR, et al. 1984. An analysis of the spectra of genetic activity produced by known or suspected human carcinogens. *Mutat Res* 134:89-111.
- Geiger LE, Hogy LL, Guengerich FP. 1983. Metabolism of acrylonitrile by isolated rat hepatocytes. *Cancer Res* 43:3080-3087.
- Gerin M, Tardif R, Brodeur J. 1988. Determination of specific urinary thioethers derived from acrylonitrile and ethylene oxide. *IARC Sci Pub* 275-278.
- \* Ghanayem BI, Ahmed AE. 1982. In vivo biotransformation and biliary excretion of 1-<sup>14</sup>C-acrylonitrile in rats. *Arch Toxicol* 50:175-185.
- \* Ghanayem BI, Ahmed AE. 1983. Acrylonitrile-induced gastrointestinal hemorrhage and the effects of metabolism modulation in rats. *Toxicol Appl Pharmacol* 68:290-296.
- Ghanayem BI, Ahmed AE. 1986. Prevention of acrylonitrile-induced gastrointestinal bleeding by sulfhydryl compounds, atropine and cimetidine. *Res Commun Chem Pathol Pharmacol* 53:141-144.
- \* Ghanayem BI, Boor PJ, Ahmed AE. 1985. Acrylonitrile-induced gastric mucosal necrosis: Role of gastric glutathione. *J Pharmacol Exp Ther* 232:570-577.
- \* Gilbert SG, Miltz .I, Giacini JR. 1980. Transport considerations of potential migrants from food packaging materials. *Journal of Food Processing and Preservation* 4:27-49.
- \* Glauert HP, Kennan WS, Sattler GL, et al. 1985. Assays to measure the induction of unscheduled DNA synthesis in cultured hepatocytes. In: Ashby J, de Serres FJ, et al., eds. *Progress in mutation research*. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 371-373.
- Going JE, Kuykendahl P, Long S, et al. 1979. Environmental monitoring near industrial sites: acrylonitrile. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA-560/6-79-003. NTIS No. PB-295928.
- Goncharova LN, Chirkov VV, Rudakova LE. 1984. [Effect of the manufacture of acrylonitrile on endocrine function.] *Gig Tr Prof Zabol* 23-25. (Russian)
- Gortseva IV, Tarasova NA, Shutova TV, et al. 1987. [Determination of acrylonitrile in polymer materials and aqueous and oily extracts of them by the gas-chromatography method.] *Gig Sanit* 61-62. (Russian)

## 8. REFERENCES

- Gosselin RE, Smith RP, Hodge HC, et al. 1984. Clinical toxicology of commercial products. 5th ed. Baltimore, MD: Williams and Wilkins, 11-215.
- \* Grunske F. 1949. Health care and occupational medicine. Ventox and Ventox intoxication. Dtsch Med Wochenschr 74:1081-1083.
- \* Guengerich FP, Geiger LE, Hogg LL, et al. 1981. In vitro metabolism of acrylonitrile to 2-cyanoethylene oxide, reaction with glutathione, and irreversible binding to proteins and nucleic acids. Cancer Res 41:4925-4933.
- Guengerich FP, Hogg LL, Inskeep PB, et al. 1986. Metabolism and covalent binding of  $\alpha$ -dihaloalkanes, vinyl halides and acrylonitrile. IARC Sci Publ 70:255-260.
- Guicherit R, Schulting FL. 1985. The occurrence of organic chemicals in the atmosphere of The Netherlands. Sci Total Environ 43:193-219.
- \* Gut I, Nerudova J, Kopecky J, et al. 1975. Acrylonitrile biotransformation in rats, mice, and Chinese hamsters as influenced by the route of administration and by phenobarbital, SKF 525-A, cysteine, dimercaprol, or thiosulfate. Arch Toxicol 33:151-161.
- Gut I, Kopecky J, Filip J. 1981. Acrylonitrile- $^{14}\text{C}$  metabolism in rats: Effect of the route of administration on the elimination of thiocyanate and other radioactive metabolites in urine and feces. J Hyg Epidemiol Microbiol Immunol 25:12-16.
- \* Gut I, Nerudova J, Frantik E, et al. 1984. Acrylonitrile inhalation in rats: I. Effect on intermediary metabolism. J Hyg Epidemiol Microbiol Immunol 28:369-376.
- \* Gut I, Nerudova J, Stiborova A, et al. 1985. Acrylonitrile inhalation in rats: II. Excretion of thioethers and thiocyanate in urine. J Hyg Epidemiol Microbiol Immunol 29:9-13.
- Hachiya N, Sato M, Takizawa Y. 1984. Detection of DNA damage in mutagen-treated mammalian tissues by alkaline elution assay. Mutat Res 130:363.
- Haemisegger E, Jones A, Steigerwald B, et al. 1985. The air toxics problem in the United States: An analysis of cancer risks for selected pollutants. Washington, DC: U.S. Environmental Protection Agency, Office of Air and Radiation (ANR-443). EPA-450/1-85-001. NTIS No. PB85-225175.

## 8. REFERENCES

- \* Harris GW, Kleindienst TE, Pitts JN Jr. 1981. Rate constants for the reaction of OH radicals with CH<sub>3</sub>CN, C<sub>2</sub>H<sub>5</sub>CN and CH<sub>2</sub> = CH-CN in the temperature range 298-424 K. *Chemical Physics Letters* 80:479-483.
- \* Hashimoto S, Bandow H, Akimoto H, et al. 1984. Products and mechanism for the OH radical initiated oxidation of acrylonitrile, methacrylonitrile and allylcyanoide in the presence of NO. *International Journal of Chemical Kinetics* 16:1385-1399.
- Haskovec C, Gut I, Volmerova D, et al. 1988. Acrylonitrile depletes glutathione without changing calcium sequestration in hepatic microsomes and mitochondria. *Toxicology* 48:87-92.
- \* Hassett JJ, Banwart WL, Griffin RA. 1983. Correlation of compound properties with sorption characteristics of nonpolar compounds by soils and sediments: concepts and limitations. In: Francis CW, Auerbach SI, Jacobs VA, eds. *Environment and solid wastes: characterization, treatment, and disposal*. Boston, MA: Butterworths, 161-176.
- \* Hogy LL, Guengerich FP. 1986. In vivo interaction of acrylonitrile and 2-cyanoethylene oxide with DNA in rats. *Cancer Res* 46:3932-3938.
- Holecek V, Kopecky J. 1983. Conjugations of acrylonitrile and glycidonitrile with glutathione--a contribution to problems of metabolism of acrylonitrile. *Czech Med* 6:116-121.
- \* Houthuijs D, Remijn B, Willems H, et al. 1982. Biological monitoring of acrylonitrile exposure. *Am J Ind Med* 3:313-320.
- Hradec J, Spiegelhalder B, Preussmann R. 1988. The initiator tRNA acceptance assay as a short-term test for carcinogens. 2. Results with ten compounds selected by the International Programme on Chemical Safety for the evaluation of short-term tests for carcinogens. *Carcinogenesis* 9:843-846.
- \* HSDB. 1988. Hazardous Substances Data Bank. National Library of Medicine, National Toxicology Information Program, Bethesda, MD. December 1988.
- \* Hughes TW, Horn DA. 1977. Source assessment: Acrylonitrile manufacture (air emissions). Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA-60V2-77-107j. NTIS No. PB-271969.
- \* Humiston CG, Frauson LO, Quast JF, et al. 1975. A go-day oral toxicity study incorporating acrylonitrile in the drinking water of rats. Dow Chemical Company, Health and Environmental Research, Toxicology Research Laboratory, Midland, MI.

## 8. REFERENCES

- \* Hurtt ME, Bentley KS, Working PK. 1987. Effects of acrylamide and acrylonitrile on unscheduled DNA synthesis (UDS) in rat spermatocytes. *Environ Mutagen* 9:49-50.
- IAPA. 1987. Occupational health bulletin: acrylonitrile. Industrial Accident Prevention Association. Toronto, Ontario.
- \* IARC. 1982. IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans. Chemicals, industrial processes and industries associated with cancer in humans. IARC monographs, Volumes 1 to 29, supplement 4. International Agency for Research on Cancer, Lyon, France.
- \* IARC. 1979. Acrylonitrile, acrylic and modacrylic fibres, and acrylonitrile-butadiene-styrene and styrene-acrylonitrile copolymers. IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans: Some monomers, plastics and synthetic elastomers, and acrolein. International Agency for Research on Cancer, Lyon, France 19:73-113.
- IARC. 1988. International Agency for Cancer Research, Lyon, France.
- \* IRIS. 1988. Integrated Risk Information System. U.S. Environmental Protection Agency, Washington, DC. December 1988.
- IRPTC. 1989. International Register of Potentially Toxic Chemicals. United Nations Environment Programme, Geneva, Switzerland. January, 1989.
- \* Jacobs HW, Syrjala RH. 1978. The use of infrared analyzers for monitoring acrylonitrile. *Am Ind Hyg Assoc J* 39:161-165.
- Jaeger RJ, Cote IL. 1982. Effect of hypoxia on the acute toxicity of acrylonitrile. *Res Commun Chem Pathol Pharmacol* 36:345-348.
- Jaeger RJ, Cote IL, Rogers AE, et al. 1984. Acute toxicity of acrylonitrile: Effect of diet on tissue nonprotein sulfhydryl content and distribution of 1- <sup>14</sup>C-acrylonitrile or its metabolites. *J Am Coll Toxicol* 3:93-102.
- \* Jakubowski M, Linhart I, Pielas G, et al. 1987. 2-Cyanoethylmercapturic acid (CEMA) in the urine as a possible indicator of exposure to acrylonitrile. *Br J Ind Med* 44:834-840.
- Kamijo K, Kovacs K, Szabo S, et al. 1986. Effect of acrylonitrile on the rat pituitary: enlargement of Golgi region in prolactin cells, crinophagy in prolactin cells and growth hormone cells. *Br J Exp Pathol* 67:439-451.



## 8. REFERENCES

- Kankaanpaa J, Elovaara E, Hemminiki K, et al. 1979. Embryotoxicity of acrolein, acrylonitrile and acrylamide in developing chick embryos. *Toxicol Lett* 4:93-96.
- \* Kayser R, Sterling D, Viviani D. 1982. Intermedia priority pollutant guidance documents. Washington, DC: U.S. Environmental Protection Agency, Office of Toxics Integration, Office of Pesticides and Toxic Substances.
- Kedderis GL, Held SD, Batin R, et al. 1989. Dose-dependent urinary excretion of acrylonitrile (ACN) metabolites in F-344 rats and B6C3F1 mice. CIIT, Research Triangle Park, NC.
- \* Kenaga EE. 1980. Predicted bioconcentration factors and soil sorption coefficients of pesticides and other chemicals. *Ecotoxicol Environ Safety* 4:26-38.
- \* Khudoley VV, Mizgirev I, Pliss GB. 1987. The study of mutagenic activity of carcinogens and other chemical agents with *Salmonella typhimurium* assays: Testing of 126 compounds. *Arch Geschwulstforsch* 57:453-462.
- \* Kiesselbach N, Korallus U, Lange HJ, et al. 1979. [Acrylonitrile--epidemiological study--Bayer 1977: a report on a prospective epidemiological study with a past beginning of coworkers at the Leverkusen plant of Bayer AG with acrylonitrile (ACN) exposure.] *Zentralbl Arbeitsmed Arbeitsschutz Prophyl* 10:256-259. (German)
- Kim NK, Stone DW. (NA) Organic chemicals and drinking water. New York State Department of Health, pp. 132.
- \* Kincannon DF, Stover EL, Nichols V, et al. 1983. Removal mechanisms for toxic priority pollutants. *J Water Pollut Control Fed* 55:157-163.
- \* King L, Sherbin G. 1986. Point sources of toxic organics to the upper St. Clair River. *Water Poll Res J Canada* 21:433-446.
- \* Klein E, Weaver JW, Webre BG. 1957. Solubility of acrylonitrile in aqueous bases and alkali salts. *Industrial and Engineering Chemistry* 2:72-75.
- Koerselman W, van der Graaf M. 1984. Acrylonitrile: A suspected human carcinogen. *Int Arch &cup Environ Health* 54:317-324.
- Kopecky J, Gut I, Nerudova J, et al. 1980. Acrylonitrile metabolism in the rat. *Arch Toxicol* 4(Suppl):322-324.

## 8. REFERENCES

- Kopecky J, Gut I, Nerudova J, et al. 1980. Two routes of acrylonitrile metabolism. *J Hyg Epidemiol Microbiol Immunol* 24:356-362.
- \* Krill RM, Sonzogni WC. 1986. Chemical monitoring of Wisconsin's groundwater. *JAWWA* 78:70-75.
- \* Lambotte-Vandepaer M, Duverger-van Bogaert M. 1984. Genotoxic properties of acrylonitrile. *Mutat Res* 134:49-59.
- \* Lambotte-Vandepaer M, Duverger-van Bogaert M, de Meester C, et al. 1980. Mutagenicity of urine from rats and mice treated with acrylonitrile. *Toxicology* 16:67-71.
- \* Lambotte-Vandepaer M, Duverger-van Bogaert M, de Meester C, et al. 1981. Identification of two urinary metabolites of rats treated with acrylonitrile; influence of several inhibitors on the mutagenicity of those urines. *Toxicol Lett* 7:321-327.
- Lambotte-Vandepaer M, Duverger-van Bogaert M, Rollmann B. 1985. Metabolism and mutagenicity of acrylonitrile: An in vivo study. *Environ Mutagen* 7:655-662.
- \* Langvardt PW, Putzig CL, Braun WH, et al. 1980. Identification of the major urinary metabolites of acrylonitrile in the rat. *J Toxicol Environ Health* 6:273-282.
- \* Lawrence N, McGregor DB. 1985. Assays for the induction of morphological transformation in C3H/10T1/2 cells in culture with and without S9-mediated metabolic activation. In: Ashby J, de Serres FJ, et al., eds. *Progress in mutation research*. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 651-658.
- \* Lee CG, Webber TD. 1985. The induction of gene mutations in the mouse lymphoma L5178Y/K<sup>+</sup> assay and the Chinese hamster V79/HGPRT assay. In: Ashby J, de Serres FJ, et al, eds. *Progress in mutation research*. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 547-554.
- \* Leonard A, Garny V, Poncelet F, et al. 1981. Mutagenicity of acrylonitrile in mouse. *Toxicol Lett* 7:329-334.
- Levins P, Adams J, Brenner P, et al. 1979. Sources of toxic pollutants found in influents to sewage treatment plants. VI. Integrated interpresentation. Washington, DC: U.S. Environmental Protection Agency, Office of Water Planning and Standards. EPA 440/4-81-008. NTIS No. PB81-219685.

## 8. REFERENCES

- \* Lijinsky W, Andrews AW. 1980. Mutagenicity of vinyl compounds in *Salmonella typhimurium*. *Teratogenesis Carcinog Mutagen* 1:259-267.
- \* Linhart I, Smejkal J, Novak J. 1988. N-acetyl-S-(1-cyano-2-hydroxyethyl)-L-cysteine, a new urinary metabolite of acrylonitrile and oxiranecarbonitrile. *Arch Toxicol* 61:484-488.

Litvak S, Sarih L, Fournier M, et al. 1983. Involvement of tRNA in retrovirus expression: biological implications of reverse transcriptase-primer tRNA interactions. *Recent Results Cancer Res* 84:184-190.

- \* Lorz H. 1950. Percutaneous poisoning with acrylonitrile *Dtsch Med Wochenschr* 75:1087-1088.
- \* Ludzack FJ, Schaffer RB, Bloomhuff RN. 1961. Experimental treatment of organic cyanides by conventional processes. *J Water Pollut Control Fed* 33:492-505.
- \* Mabey WR, Smith JH, Pod011 RT, et al. 1982. Aquatic fate process data for organic priority pollutants. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards. EPA 440/4-81-014.
- \* Maltoni C, Ciliberti A, Di Maio V. 1977. Carcinogenicity bioassays on rats of acrylonitrile administered by inhalation and by ingestion. *Med Lav* 68:401-411.

Maltoni C, Ciliberti A, Carretti D. 1982. Experimental contributions in identifying brain potential carcinogens in the petrochemical industry. *Ann NY Acad Sci* 381:216-249.

- \* Maltoni C, Ciliberti A, Cotti G, et al. 1988. Long-term carcinogenicity bioassays on acrylonitrile administered by inhalation and by ingestion to Sprague-Dawley rats. *Ann NY Acad Sci* 534:179-202.
- \* Marano RS, Levine SP, Harvey TM. 1978. Trace determination of subnanogram amounts of acrylonitrile in complex matrices by gas chromatography with a nitrogen-selective detector. *Anal Chem* 50:1948-1950.

MATHTECH. 1982. Draft: Level I economic evaluation: Acrylonitrile. Report to U.S. Environmental Protection Agency, Office of Pesticides and Toxic Substances, Washington, DC, by MATHTECH, Inc., Arlington, VA.

## 8. REFERENCES

- \* Matsushima T, Muramatsu M, Haresaku M. 1985. Mutation tests on *Salmonella typhimurium* by the preincubation method. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 181-186.
- \* Matthews RD. 1980. Estimated permissible levels, ambient concentrations, and adverse effects of the nitrogenous products of combustion: the cyanides, nitroolefins, and nitroparaffins. *Journal of Combustion Toxicology* 7:157-172.
- \* Matthews EJ, DelBalzo T, Rundell JO. 1985. Assays for morphological transformation and mutation to ouabain resistance of Balb/c-3T3 cells in culture. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam the Netherlands: Elsevier Science Publishers, 639-650.
- McMahon LW. 1983. Organic priority pollutants in wastewater. Proceedings of the 1982 UCC-ND/GAT Environmental Protection Seminar. Oak Ridge, TN: Oak Ridge National Laboratory.
- \* McNally ME, Wheeler JR. 1988. Supercritical fluid extraction coupled with supercritical fluid chromatography for the separation of sulfonylurea herbicides and their metabolites from complex matrices. *J Chromatogr* 435:63-71.
- McNeal T, Brumley WC, Breder C, et al. 1979. Gas-solid chromatographic-mass spectrometric confirmation of low levels of acrylonitrile after distillation from food-simulating solvents. *J Assoc Off Anal Chem* 62:41-46.
- Melcher RG, Borders RA, Coyne LB. 1986. Development and validation of personal monitoring methods for low levels of acrylonitrile in workplace atmosphere: I. Test atmosphere generation and solvent desorption methods. *Am Ind Hyg Assoc J* 47:152-157.
- \* Michael-LC, Pellizzari ED, Wiseman RW. 1988. Development and evaluation of a procedure for determining volatile organics in water. *Environ Sci Technol* 22:565-570.
- Michel F, Mercklein L, Crastes de Paulet A, et al. 1984. The effect of various acrylonitriles and related compounds on prostaglandin biosynthesis. *Prostaglandins* 27:69-84.
- \* Miller LM, Villaume JE. 1978. Investigation of selected potential environmental contaminants: acrylonitrile. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA 560/2-78-003. NTIS No. PB-285881.

## 8. REFERENCES

- \* Mills EJ Jr, Stack VT Jr. 1953. Biological oxidation of synthetic organic chemicals. *Engineering Bulletin, Proceedings of the Eighth Industrial Waste Conference, Series No. 83*, 492-517.
  - \* Mills EJ Jr, Stack VT Jr. 1955. Acclimation of microorganisms for the oxidation of pure organic chemicals. *Proceedings of the Ninth Industrial Waste Conference, Series No. 87*, 449-464.
  - \* Muller G, Verkoyen C, Soton N, et al. 1987. Urinary excretion of acrylonitrile and its metabolites in rats. *Arch Toxicol* 60:464-466.
  - \* Murray FJ, Schwetz BA, Nitschke KD, et al. 1978. Teratogenicity of acrylonitrile given to rats by gavage or by inhalation. *Food Cosmet Toxicol* 16:547-551.
  - \* Myhr B, Bowers L, Caspary WJ. 1985. Assays for the induction of gene mutations at the thymidine kinase locus in L5178Y mouse lymphoma cells in culture. In: Ashby J, de Serres FJ, et al., eds. *Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens*. Amsterdam, The Netherlands: Elsevier Science Publishers, 555-568.
  - \* NAS. 1980. *Drinking water and health: Volume 3*. Washington, DC: National Academy Press, 71-76.
  - \* NAS/NRC. 1989. *Biological markers in reproductive toxicology*. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press.
  - \* NATICH. 1988. NATICH data base report on state, local and EPA air toxics activities. July, 1988. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards, National Air Toxics Information Clearinghouse.
  - \* Neely WB, Branson DR, Blau GE. 1974. Partition coefficient to measure bioconcentration potential of organic chemicals in fish. *Environ Sci Technol* 8:1113-1115.
- Nerudova J, Gut I, Savolainen H. 1988. Consequences of acrylonitrile metabolism in rat hepatocytes: Effects on lipid peroxidation and viability of the cells. *Environ Res* 46:133-141.
- \* NIOSH. 1977. *Current intelligence bulletin 18: acrylonitrile*. Rockville, MD: U.S. Department of Health, Education, and Welfare, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health.

## 8. REFERENCES

- NIOSH. 1980. Health hazard evaluation determination report no. 79-36-656, Bell Helmets, Inc., Norwalk, California. Cincinnati, OH: U.S. Department of Health, Education, and Welfare, National Institute for Occupational Safety and Health. NTIS No. PB80-163181.
- \* NIOSH. 1981-82. Registry of Toxic Effects of Chemical Substances. Acrylonitrile. National Institute of Occupational Safety and Health.
- \* NIOSH. 1984. Acrylonitrile - Method 1604. In: NIOSH manual of analytical methods. 3rd ed. Cincinnati, OH: National Institute of Occupational Safety and Health.
- \* NIOSH. 1985. Pocket guide to chemical hazards. Washington, DC: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health. DHHS (NIOSH) Publication NO. 85-114.
- \* NIOSH. 1988a. National occupational exposure survey. Cincinnati, OH: National Institute for Occupational Safety and Health.
- \* NIOSH. 1988b. NIOSH recommendations for occupational safety and health standards. Morbidity and mortality weekly report [supplement] Vol. 37:s-7. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health.
- \* NLM. 1988. National Library of Medicine - Chemline database printout for acrylonitrile. August, 1988.
- NTP. 1988. National Toxicology Program. Review of current DHHS, DOE, and EPA research related to toxicology. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service.
- Obayashi AW, Gorgan JM. 1983. Management of industrial pollutants by anaerobic processes. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Research and Development. EPA-600/2-83-119. NTIS No. PB84-133024.
- \* Obe G, Hille A, Jonas R, et al. 1985. Tests for the induction of sister-chromatid exchanges in human peripheral lymphocytes in culture. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 439-442.
- \* O'Berg MT. 1980. Epidemiologic study of workers exposed to acrylonitrile. J Occup Med 22:245-252.
- \* O'Berg MT, Chen JL, Burke CA, et al. 1985. Epidemiologic study of workers exposed to acrylonitrile: An update. J Occup Med 27:835-40.

## 8. REFERENCES

- \* OSHA. 1978a. U.S. Occupational Safety and Health Administration. Part IV. Federal Register. 43:2586-2607.
- \* OSHA. 1978b. U.S. Occupational Safety and Health Administration. Part VI. Federal Register. 43:45762-45819.
- OSHA. 1985. U.S. Department of Labor. Occupational Safety and Health Administration. Code of Federal Regulations. 29 CFR 1910.1045.
- \* Page BD. 1985. Determination of acrylonitrile in foods by headspace-gas chromatography with nitrogen-sensitive detection: Collaborative study. *J Assoc Off Anal Chem* 68:776-782.
- \* Parent RA, Casto BC. 1979. Effect of acrylonitrile on primary Syrian golden hamster embryo cells in culture: Transformation and DNA fragmentation. *J Natl Cancer Inst* 62:1025-1029.
- Parkin DM, Wahrendorf J, Demaret E, et al. 1987. Directory of ongoing research in cancer epidemiology. New York, NY: Oxford University Press, 267, 332-333, 344, 366, 599.
- Parry JM, Danford N, Parry EM. 1984. In vitro techniques for the detection of chemicals capable of inducing mitotic chromosome aneuploidy. *Altern Lab Anim* 11:117-128.
- \* Perocco P, Pane G, Bolognesi S, et al. 1982. Increase of sister chromatid exchange and unscheduled synthesis of deoxyribonucleic acid by acrylonitrile in human lymphocytes in vitro. *Scand J Work Environ Health* 8:290-293.
- Peter H, Bolt HM. 1984. Experimental pharmacokinetics and toxicology of acrylonitrile. *G Ital Med Lav* 6:77-81.
- Peter H, Appel KE, Berg R, et al. 1983. Irreversible binding of acrylonitrile to nucleic acids. *Xenobiotica* 13:19-25.
- PHRED. 1988. Public Health Risk Evaluation Database. U.S. Environmental Protection Agency, Washington, DC. March 1988.
- \* Pilon D, Roberts AE, Rickert DE. 1988a. Effect of glutathione depletion on the irreversible association of acrylonitrile with tissue macromolecules after oral administration to rats. *Toxicol Appl Pharmacol* 95:311-320.
- \* Pilon D, Roberts AE, Rickert DE. 1988b. Effect of glutathione depletion on the uptake of acrylonitrile vapors and on its irreversible association with tissue macromolecules. *Toxicol Appl Pharmacol* 95:265-278.

## 8. REFERENCES

- \* Priston RA J, Dean BJ. 1985. Tests for the induction of chromosome aberrations, polyploidy and sister chromatid exchanges in rat liver (RL,) cells. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 387-395.
  
- \* Probst GS, Hill LE. 1985. Tests for the induction of DNA repair synthesis in primary cultures of adult rat hepatocytes. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands. Elsevier Science Publishers, 381-386.
  
- Prokopczyk B, Bertinato P, Hoffmann D. 1988. Synthesis and kinetics of decomposition of 7-(2-cyanoethyl)guanine and 6-(2-cyanoethyl)guanine, markers for reaction of acrylonitrile and 3-(methylnitrosamino)propionitrile with DNA. *Carcinogenesis* 9:2125-2128.
  
- \* Quast JF, Humiston CG, Schwetz BA, et al. 1975. A six-month oral toxicity study incorporating acrylonitrile in the drinking water of purebred beagle dogs. Dow Chemical Co., Toxicology Research Laboratory, Midland, MI.
  
- \* Quast JF, Schuetz DJ, Balmer MF, et al. 1980a. A two-year toxicity and oncogenicity study with acrylonitrile following inhalation exposure of rats. Dow Chemical Co., Toxicology Research Laboratory, Midland MI.
  
- \* Quast JF, Wade CE, Humiston CG, et al. 1980b. A two-year toxicity and oncogenicity study with acrylonitrile incorporated in the drinking water of rats. Dow Chemical Co., Toxicology Research Laboratory, Midland, MI.
  
- \* Rabello-Gay MN, Ahmed AE. 1980. Acrylonitrile: In vivo cytogenetic studies in mice and rats. *Mutat Res* 79:249-255.
  
- \* Radimer GF, Davis JH, Ackerman AB. 1974. Fumigant-induced toxic epidermal necrolysis. *Arch Dermatol* 110:103-104.
  
- Rajendran- S, Muthu M. 1980. Nutritional and histopathological studies on rats fed on energy food fumigated with acrylonitrile. *Nutrition Reports International* 22:677-685.
  
- \* Recio L, Skopek TR. 1988. Mutagenicity of acrylonitrile and its metabolite 2-cyanoethylene oxide in human lymphoblasts in vitro. *Mutat Res* 206:297-305.
  
- \* Rickert DE, Roberts AE, Pilon D. 1988. Distribution of acrylonitrile (ACN) in tissues of control and glutathione (GSH) depleted B6C3F1 mice. CIIT, Research Triangle Park, NC.



## 8. REFERENCES

- Riddell RJ, Panacer DS, Wilde SM, et al. 1986. The importance of exposure period and cell type in in vitro cytotoxicity tests. *Altern Lab Anim* 14:86-92.
- Rinkus SJ, Legator MS. 1985. Fluorometric assay using high-pressure liquid chromatography for the microsomal metabolism of certain substituted aliphatics to 1,N<sup>6</sup>-ethenoadenine-forming metabolites. *Anal Biochem* 150:379-393.
- \* Roberts AE, Lacy SA, Pilon D, et al. 1989. Metabolism of acrylonitrile to 2-cyanoethylene oxide in F-344 rat liver microsomes, lung microsomes, and lung cells. *Drug Metab Dispos* 17:481-486.
- \* Rogaczewska T, Piotrowski J. 1968. Experimental evaluation of the absorption routes of acrylonitrile in man. *Med Pr* 19:349-354.
- \* Roudabush RL, Terhaar CJ, Fassett DW, et al. 1965. Comparative acute effects of some chemicals on the skin of rabbits and guinea pigs. *Toxicol Appl Pharmacol* 7:559-565.
- \* Rouisse L, Chakrabarti S, Tuchweber B. 1986. Acute nephrotoxic potential of acrylonitrile in Fischer-344 rats. *Res Commun Chem Pathol Pharmacol* 53:347-360.
- \* Roy WR, Griffin RA. 1985. Mobility of organic solvents in watersaturated soil materials. *Environ Geol Water Sci* 7:241-247.
- \* Sakurai H, Kusumoto M. 1972. An epidemiological study of health impairment among acrylonitrile workers. *J Sci Labor* 48:273-282.
- \* Sakurai H, Onodera M, Utsunomiya T, et al. 1978. Health effects of acrylonitrile in acrylic fibre factories. *Br J Ind Med* 35:219-225.
- \* Sandberg EC, Slanina P. 1980. Distribution of [1-<sup>14</sup>C] acrylonitrile in rat and monkey. *Toxicol Lett* 6:187-191.
- \* Sanner T, Rivedal E. 1985. Tests with the Syrian hamster embryo (SHE) cell transformation assay. In: Ashby J, de Serres FJ, et al., eds. *Progress in mutation research. Vol. 5. -Evaluation of short-term tests for carcinogens.* Amsterdam, The Netherlands: Elsevier Science Publishers, 665-671.
- Sato M, Ishizu S, Momotuni H. 1975. [Determination of acrylonitrile, cyanide and thiocyanate in blood and urine.] *Jap J Ind Health* 17:99-105. (Japanese).

## 8. REFERENCES

- Sato M. 1978. [Studies on the toxic effect of acrylonitrile: Its metabolism, absorption, and excretion.] *Nippon Eiseigaku Zasshi* 33:497-505. (Japanese)
- \* Sax NI. 1984. *Dangerous properties of industrial materials*. 6th ed. New York, NY: Van Nostrand Reinhold Company, 132-133.
- \* Sax NI, Lewis RJ Sr. 1987. *Hawley's condensed chemical dictionary*. 11th ed. New York, NY: Van Nostrand Reinhold Company, 19.
- \* Sharief Y, Brown AM, Backer LC, et al. 1986. Sister chromatid exchange and chromosome aberration analyses in mice after *in vivo* exposure to acrylonitrile, styrene, or butadiene monoxide. *Environ Mutagen* 8:439-448.
- Silver EH, Szabo S. 1982. Possible role of lipid peroxidation in the actions of acrylonitrile on the adrenals, liver and gastrointestinal tract. *Res Commun Chem Pathol Pharmacol* 36:33-43.
- \* Silver EH, McComb DJ, Kovacs K, et al. 1982. Limited hepatotoxic potential of acrylonitrile in rats. *Toxicol Appl Pharmacol* 64:131-139.
- \* Silver EH, Szabo S, Cahill M, et al. 1987. Time-course studies of the distribution of [ $1-^{14}\text{C}$ ]Acrylonitrile in rats after intravenous administration. *J Appl Toxicol* 7:303-306.
- Sipos-Meszlenyi M, Adamis T, Hollo A, et al. 1986. Some applications of infrared and ultraviolet spectroscopy for the toxicological examination of medical plastic products. *Periodica Polytechnica Chemical Engineering* 30:165-170.
- Sittig M. 1985. *Handbook of toxic and hazardous chemicals and carcinogens*. 2nd ed. Park Ridge, NJ: Noyes Publications, 43-45.
- Solomon JJ, Fedky J, Mukai F, et al. 1985. Direct alkylation of 2'-deoxynucleosides and DNA following *in vitro* reaction with acrylamide. *Cancer Res* 45:3465-3470.
- Solomon JJ, Segal A. 1985. Direct alkylation of calf thymus DNA by acrylonitrile. Isolation of cyanoethyl adducts of guanine and thymine and carboxyethyl adducts of adenine and cytosine. *Environ Health Perspect* 62:227-230.
- Spicer CW, Riggin RM, Holdren MW, et al. 1985. Atmospheric reaction products from hazardous air pollutant degradation. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA/600/3-85/028. NTIS No. PB85-185841.

## 8. REFERENCES

- SRI. 1985. Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- SRI. 1986. Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- SRI. 1987a. Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- SRI. 1987b. Directory of chemical producers: United States of America: Supplement II. Menlo Park, CA: SRI International.
- \* SRI. 1988a. Directory of chemical producers: United States of America. Menlo Park, CA: SRI International.
- SRI. 1988b. Directory of chemical producers: United States of America: Supplement I. Menlo Park, CA: SRI International.
- \* Staples CA, Werner AF, Hoogheem TJ. 1985. Assessment of priority pollutant concentrations in the United States using STORET database. Environ Toxicol Chem 4:131-142.
- Steichen RJ. 1976. Modified solution approach for the gas chromatographic determination of residual monomers by head-space analysis. Anal Chem 48:1398-1402.
- \* Stover EL, Kincannon DF. 1983. Biological treatability of specific organic compounds found in chemical industry wastewaters. J Water Pollut Control Fed 55:97-109.
- Strother DE, Mast RW, Kraska RC, et al. 1988. Acrylonitrile as a carcinogen: Research needs for better risk assessment. Ann NY Acad Sci 534:169-178.
- \* Suta BE. 1979. Human exposure to atmospheric concentrations of selected chemicals. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards. NTIS No. PB81-193278.
- Szabo S, Gallagher GT. 1984. Effects of alkyl nitriles on the gastrointestinal tract. Surv Synth Path Res 3:11-30.
- \* Szabo S, Bailey KA, Boor PJ, et al. 1977. Acrylonitrile and tissue glutathione: Differential effect of acute and chronic interactions. Biochem Biophys Res Commun 79:32-37.

## 8. REFERENCES

- \* Szabo S, Huttner I, Kovacs K, et al. 1980. Pathogenesis of experimental adrenal hemorrhagic necrosis ("apoplexy"): Ultrastructural, biochemical, neuropharmacologic, and blood coagulation studies with acrylonitrile in the rat. *Lab Invest* 42:533-546.
- Szabo S, Reynolds ES, Unger SH. 1982. Structure-activity relations between alkyl nucleophilic chemicals causing duodenal ulcer and adrenocortical necrosis. *J Pharmacol Exp Ther* 223:68-76.
- \* Szabo S, Silver EH, Gallagher GT, et al. 1983. Potentiation of duodenal ulcerogenic action of acrylonitrile by PCB or phenobarbital in the rat. *Toxicol Appl Pharmacol* 71:451-454.
- \* Szabo S, Gallagher GT, Silver EH, et al. 1984. Subacute and chronic action of acrylonitrile on adrenals and gastrointestinal tract: Biochemical, functional and ultrastructural studies in the rat. *J Appl Toxicol* 4:131-140.
- \* Tabak HH, Quave SA, Mashni CI, et al. 1981. Biodegradability studies with organic priority pollutant compounds. *J Water Pollut Control Fed* 53:1503-1518.
- \* Tandon R, Saxena DK, Chandra SV, et al. 1988. Testicular effects of acrylonitrile in mice. *Toxicol Lett* 42:55-63.
- Tanii H, Hashimoto K. 1984a. Structure-toxicity relationship of aliphatic nitriles. *Toxicol Lett* 22:267-272.
- Tanii H, Hashimoto K. 1984b. Studies on the mechanism of acute toxicity of nitriles in mice. *Arch Toxicol* 55:47-54.
- Tanii H, Hashimoto K. 1986. Influence of ethanol on the in vivo and in vitro metabolism of nitriles in mice. *Arch Toxicol* 58:171-176.
- \* Tardif R, Talbot D, Gerin M, et al. 1987. Urinary excretion of mercapturic acids and thiocyanate in rats exposed to acrylonitrile: Influence of dose and route of administration. *Toxicol Lett* 39:255-261.
- \* Thiess AM, Fleig I. 1978. Analysis of chromosomes of workers exposed to acrylonitrile. *Arch Toxicol* 41:149-152.
- Thuroff E, Kaufer NF, Lochmann ER. 1986. Effect of acrylonitrile on the transcription of specific genes in Saccharomyces cerevisiae. *Mol Gen Genet* 202:336-337.
- TPCDB. 1988. Testing Priority Committee Data Base. U.S. Environmental Protection Agency, Office of Toxic Substances, Washington, DC. November 18, 1988.

## 8. REFERENCES

- \* TRI. 1988. Toxic Chemical Release Inventory. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.
- \* USITC. 1988. Synthetic organic chemicals: United States production and sales, 1987. Washington, DC: U.S. International Trade Commission. USITC Publication 2118.
- Van Bladeren PJ, Delbressine LP, Hoogeterp JJ, et al. 1981. Formation of mercapturic acids from acrylonitrile, crotononitrile, and cinnamonnitrile by direct conjugation and via an intermediate oxidation process. Drug Metab Dispos 9:246-249.
- \* Vanderlaan M, Watkins BE, Stanker L. 1988. Environmental monitoring by immunoassay. Environ Sci Technol 22:247-254.
- \* Venitt S, Bushell CT, Osborne M. 1977. Mutagenicity of acrylonitrile (cyanoethylene) in Escherichia coli. Mutat Res 45:283-288.
- \* Verschueren K. 1983. Handbook of environmental data on organic chemicals. 2nd ed. New York, NY: Van Nostrand Reinhold Company, 162-165.
- \* VIEW Database. 1989. Agency for Toxic Substances and Disease Registry (ATSDR), Office of External Affairs, Exposure and Disease Registry Branch, Atlanta GA. June 20, 1989. (Map based on VIEW Database, June 12, 1989).
- Vilim V, Nerudova J, Frantik E, et al. 1988. Acrylonitrile potentiation of oxygen toxicity in rats. Biomed Biochim Acta 47:206-209.
- \* Vogel EW. 1985. The Drosophila somatic recombination and mutation assay (SRM) using the white-coral somatic eye color system. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 313-317.
- \* Vogel RA, Kirkendall WM. 1984. Acrylonitrile (vinyl cyanide) poisoning: A case report. Tex Med 80:48-51.
- \* Walters SM. 1986. Cleanup of samples. In: Zweig G, Sherma J, eds. Analytical methods for pesticides and plant growth regulators. Vol. 15. New York, NY: Academic Press, 67-110.
- \* Washall JW, Wampler TP. 1988. Purge and trap analysis of aqueous samples with cryofocusing. American Laboratory (July): 70-74.

## 8. REFERENCES

- \* Weast RC, ed. 1985. CRC handbook of chemistry and physics. Boca Raton, FL: CRC Press, Inc., C-58.
- Werner JB, Carter JT. 1981. Mortality of United Kingdom acrylonitrile polymerisation workers. Br J Ind Med 38:247-253.
- \* West CR. 1989. Letter with accompanying data, from Carol Rowan West, Director, Office of Research and Standards, Department of Environmental Quality Engineering, Commonwealth of Massachusetts, to Barry L. Johnson, ATSDR, dated May 8, 1989.
- \* WHO. 1983. Environmental health criteria 28: acrylonitrile. World Health Organization. Geneva, Switzerland.
- \* Williams GM, Tong C, Brat SV. 1985. Tests with the rat hepatocyte primary culture/DNA-repair test. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 341-345.
- \* Wilson RH. 1944, Health hazards encountered in the manufacture of synthetic rubber. JAMA 124:701-703.
- \* Wilson RH, Hough GV, McCormick WE. 1948. Medical problems encountered in the manufacture of American-made rubber. Ind Med 17:199-207.
- Windholz M, ed. 1983. The Merck index: An encyclopedia of chemicals, drugs, and biologicals. 10th ed. Rahway, NJ: Merck and Co., Inc., 20.
- \* Working PK, Bentley KS, Hurtt ME, et al. 1987. Comparison of the dominant lethal effects of acrylonitrile and acrylamide in male Fischer 344 rats. Mutagenesis 2:215-220.
- \* Wurgler FE, Graff U, Frei H. 1985. Somatic mutation and recombination test in wings of Drosophila melanogaster. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 325-340.
- \* Yamada H, Asano Y, Hino T, et al. 1979. Microbiol utilization of acrylonitrile. J Ferment Technol 57:8-14.
- \* Yamanaka S, Kanamaru F, Koizumi M. 1974. Role of interlayer cations in the formation of acrylonitrile-montmorillonite complexes. J Phys Chem 78:42-44.

## 8. REFERENCES

- \* Young JD, Slauter RW, Karbowski RJ. 1977. The pharmacokinetic and metabolic profile of r4C-acrylonitrile given to rats by three routes. Dow Chemical Co., Toxicology Research Laboratory, Midland, MI,
- Young RH, Ryckman DW, Buzzell JC Jr. 1968. An improved tool for measuring biodegradability. J Water Pollut Control Fed 40(Pt 2):R354-R368.
- Zeiger E, Haworth S. 1985. Tests with a preincubation modification of the Salmonella/microsome assay. In: Ashby J, de Serres FJ, et al., eds. Progress in mutation research. Vol. 5. Evaluation of short-term tests for carcinogens. Amsterdam, The Netherlands: Elsevier Science Publishers, 187-199.





## 9. GLOSSARY

**Acute Exposure** -- Exposure to a chemical for a duration of 14 days or less, as specified in the Toxicological Profiles.

**Adsorption Coefficient ( $K_{oc}$ )** -- The ratio of the amount of a chemical adsorbed per unit weight of organic carbon in the soil or sediment to the concentration of the chemical in solution at equilibrium.

**Adsorption Ratio ( $K_d$ )** -- The amount of a chemical adsorbed by a sediment or soil (i.e., the solid phase) divided by the amount of chemical in the solution phase, which is in equilibrium with the solid phase, at a fixed solid/solution ratio. It is generally expressed in micrograms of chemical sorbed per gram of soil or sediment.

**Bioconcentration Factor (BCF)** -- The quotient of the concentration of a chemical in aquatic organisms at a specific time or during a discrete time period of exposure divided by the concentration in the surrounding water at the same time or during the same time period.

**Cancer Effect Level (CEL)** -- The lowest dose of chemical in a study or group of studies which produces significant increases in incidence of cancer (or tumors) between the exposed population and its appropriate control.

**Carcinogen** -- A chemical capable of inducing cancer.

**Ceiling value (CL)** -- A concentration of a substance that should not be exceeded, even instantaneously.

**Chronic Exposure** -- Exposure to a chemical for 365 days or more, as specified in the Toxicological Profiles.

**Developmental Toxicity** -- The occurrence of adverse effects on the developing organism that may result from exposure to a chemical prior to conception (either parent), during prenatal development, or postnatally to the time of sexual maturation. Adverse developmental effects may be detected at any point in the life span of the organism.

**Embryotoxicity and Fetotoxicity** -- Any toxic effect on the conceptus as a result of prenatal exposure to a chemical; the distinguishing feature between the two terms is the stage of development during which the insult occurred. The terms, as used here, include malformations and variations, altered growth, and in utero death.

**EPA Health Advisory** -- An estimate of acceptable drinking water levels for a chemical substance based on health effects information. A health advisory is not a legally enforceable federal standard, but serves as technical guidance to assist federal, state, and local officials.

## 9. GLOSSARY

**Immediately Dangerous to Life or Health (IDLH)** -- The maximum environmental concentration of a contaminant from which one could escape within 30 min without any escape-impairing symptoms or irreversible health effects.

**Intermediate Exposure** -- Exposure to a chemical for a duration of 15-364 days as specified in the Toxicological Profiles.

**Immunologic Toxicity** -- The occurrence of adverse effects on the immune system that may result from exposure to environmental agents such as chemicals.

**In Vitro** -- Isolated from the living organism and artificially maintained, as in a test tube.

**In Vivo** -- Occurring within the living organism.

**Lethal Concentration<sub>(L0)</sub> (LC<sub>L0</sub>)** -- The lowest concentration of a chemical in air which has been reported to have caused death in humans or animals.

**Lethal Concentration<sub>(50)</sub> (LC<sub>50</sub>)** -- A calculated concentration of a chemical in air to which exposure for a specific length of time is expected to cause death in 50% of a defined experimental animal population.

**Lethal Dose<sub>(L0)</sub> (LD<sub>L0</sub>)** -- The lowest dose of a chemical introduced by a route other than inhalation that is expected to have caused death in humans or animals.

**Lethal Dose<sub>(50)</sub> (LD<sub>50</sub>)** -- The dose of a chemical which has been calculated to cause death in 50% of a defined experimental animal population.

**Lethal Time<sub>(50)</sub> (LT<sub>50</sub>)** -- A calculated period of time within which a specific concentration of a chemical is expected to cause death in 50% of a defined experimental animal population.

**Lowest-Observed-Adverse-Effect Level (LOAEL)** -- The lowest dose of chemical in a study or group of studies which produces statistically or biologically significant increases in frequency or severity of adverse effects between the exposed population and its appropriate control.

**Malformations** -- Permanent structural changes that may adversely affect survival, development, or function.

## 9. GLOSSARY

**Minimal Risk Level (MRL)** -- An estimate of daily human exposure to a chemical that is likely to be without an appreciable risk of deleterious effects (noncancerous) over a specified duration of exposure.

**Mutagen** -- A substance that causes mutations. A mutation is a change in the genetic material in a body cell. Mutations can lead to birth defects, miscarriages, or cancer.

**Neurotoxicity** -- The occurrence of adverse effects on the nervous system following exposure to a chemical.

**No-Observed-Adverse-Effect Level (NOAEL)** -- That dose of chemical at which there are no statistically or biologically significant increases in frequency or severity of adverse effects seen between the exposed population and its appropriate control. Effects may be produced at this dose, but they are not considered to be adverse.

**Octanol-Water Partition Coefficient ( $K_{ow}$ )** -- The equilibrium ratio of the concentrations of a chemical in n-octanol and water, in dilute solution.

**Permissible Exposure Limit (PEL)** -- An allowable exposure level in workplace air averaged over an 8-hour shift.

**$q_1^*$**  -- The upper-bound estimate of the low-dose slope of the dose-Response curve as determined by the multistage procedure. The  $q_1^*$  can be used to calculate an estimate of carcinogenic potency, the incremental excess cancer risk per unit of exposure (usually  $\mu\text{g/L}$  for water,  $\text{mg/kg/day}$  for food, and  $\mu\text{g/m}^3$  for air).

**Reference Dose (RfD)** -- An estimate (with uncertainty spanning perhaps an order of magnitude) of the daily exposure of the human population to a potential hazard that is likely to be without risk of deleterious effects during a lifetime. The RfD is operationally derived from the NOAEL (from animal and human studies) by a consistent application of uncertainty factors that reflect various types of data used to estimate RfDs and an additional modifying factor, which is based on a professional judgment of the entire database on the chemical. The RfDs are not applicable to nonthreshold effects such as cancer.

**Reportable Quantity (RQ)** -- The quantity of a hazardous substance that is considered reportable under CERCLA. Reportable quantities are: (1) 1 lb or greater or (2) for selected substances, an amount established by regulation either under CERCLA or under Sect. 311 of the Clean Water Act. Quantities are measured over a 24-hour period.

## 9. GLOSSARY

**Reproductive Toxicity** -- The occurrence of adverse effects on the reproductive system that may result from exposure to a chemical. The toxicity may be directed to the reproductive organs and/or the related endocrine system. The manifestation of such toxicity may be noted as alterations in sexual behavior, fertility, pregnancy outcomes, or modifications in other functions that are dependent on the integrity of this system.

**Short-Term Exposure Limit (STEL)** -- The maximum concentration to which workers can be exposed for up to 15 min continually. No more than four excursions are allowed per day, and there must be at least 60 min between exposure periods. The daily TLV-TWA may not be exceeded.

**Target Organ Toxicity** -- This term covers a broad range of adverse effects on target organs or physiological systems (e.g., renal, cardiovascular) extending from those arising through a single limited exposure to those assumed over a lifetime of exposure to a chemical.

**Teratogen** -- A chemical that causes structural defects that affect the development of an organism.

**Threshold Limit Value (TLV)** -- A concentration of a substance to which most workers can be exposed without adverse effect. The TLV may be expressed as a TWA, as a STEL, or as a CL.

**Time-weighted Average (TWA)** -- An allowable exposure concentration averaged over a normal 8-hour workday or 40-hour workweek.

**Toxic Dose (TD<sub>50</sub>)** -- A calculated dose of a chemical, introduced by a route other than inhalation, which is expected to cause a specific toxic effect in 50% of a defined experimental animal population.

**Uncertainty Factor (UF)** -- A factor used in operationally deriving the RfD from experimental data. UFs are intended to account for (1) the variation in sensitivity among the members of the human population, (2) the uncertainty in extrapolating animal data to the case of humans, (3) the uncertainty in extrapolating from data obtained in a study that is of less than lifetime exposure, and (4) the uncertainty in using LOAEL data rather than NOAEL data. Usually each of these factors is set equal to 10.

## APPENDIX

## PEER REVIEW

A peer review panel was assembled for acrylonitrile. The panel consisted of the following members: Dr. Ahmed Ahmed, Professor of Environmental Pathology and Toxicology, Department of Pathology, University of Texas Medical Branch, Galveston; Dr. Norman M. Trieff, Professor of Environmental Toxicology, University of Texas Medical Branch, Galveston; Dr. Sandor Szabo, Associate Professor of Pathology, Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston. These experts collectively have knowledge of acrylonitrile's physical and chemical properties, toxicokinetics, key health end points, mechanisms of action, human and animal exposure, and quantification of risk to humans. All reviewers were selected in conformity with the conditions for peer review specified in the Superfund Amendments and Reauthorization Act of 1986, Section 110.

A joint panel of scientists from ATSDR and EPA has reviewed the peer reviewers' comments and determined which comments will be included in the profile. A listing of the peer reviewers' comments not incorporated in the profile, with a brief explanation of the rationale for their exclusion, exists as part of the administrative record for this compound. A list of databases reviewed and a list of unpublished documents cited are also included in the administrative record.

The citation of the peer review panel should not be understood to imply their approval of the profile's final content. The responsibility for the content of this profile lies with the Agency for Toxic Substances and Disease Registry.