

Staff Paper Series

Targeting Agricultural Drainage to Reduce Nitrogen Losses in a Minnesota Watershed

by

Daniel R. Petrolia, Prasanna H. Gowda,
and David J. Mulla

**DEPARTMENT OF APPLIED ECONOMICS
COLLEGE OF AGRICULTURAL, FOOD, AND ENVIRONMENTAL SCIENCES
UNIVERSITY OF MINNESOTA**

Targeting Agricultural Drainage to
Reduce Nitrogen Losses in a Minnesota Watershed

Daniel R. Petrolia, Prasanna H. Gowda,
and David J. Mulla

The analyses and views reported in this paper are those of the authors. They are not necessarily endorsed by the Department of Applied Economics or by the University of Minnesota.

The University of Minnesota is committed to the policy that all persons shall have equal access to its programs, facilities, and employment without regard to race, color, creed, religion, national origin, sex, age, marital status, disability, public assistance status, veteran status, or sexual orientation.

Copies of this publication are available at <http://agecon.lib.umn.edu/>. Information on other titles in this series may be obtained from: Waite Library, University of Minnesota, Department of Applied Economics, 232 Classroom Office Building, 1994 Buford Avenue, St. Paul, MN 55108, U.S.A.

Copyright (c) (2005) by Daniel R. Petrolia, Prasanna H. Gowda, and David J. Mulla. All rights reserved. Readers may make copies of this document for non-commercial purposes by any means, provided that this copyright notice appears on all such copies.

Targeting Agricultural Drainage to Reduce Nitrogen Losses in a Minnesota Watershed¹

Daniel R. Petrolia, Prasanna H. Gowda, and David J. Mulla²

Abstract

Agricultural nitrogen losses are the major contributor to nitrogen loads in the Mississippi River, and consequently, to the existence of a hypoxic, or “dead”, zone in the Gulf of Mexico. Focusing on two small agricultural watersheds in southeast Minnesota, simulation results from the Agricultural Drainage And Pesticide Management (ADAPT) model were combined with a linear-optimization model to evaluate the environmental and economic impact of alternative land-use policies for reducing nitrogen losses. Of particular importance was the study’s explicit focus on agricultural subsurface (tile) drainage, which has been identified as the major pathway for agricultural nitrogen losses in the upper Midwest, and the use of drainage-focused abatement policies. Results indicate that tile-drained land plays a key role in nitrogen abatement, and that a combined policy of nutrient management on tile-drained land and retirement of non-drained land is a cost-effective means of achieving a 20- or 30-percent nitrogen-abatement goal. Results also indicate that although it is cost-effective to abate on tile-drained land, it is not cost-effective to undertake policies that plug or remove tile drains from the landscape, regardless of whether the land would be retired or kept in production. Therefore, results imply that although tile-drained land is a major source of nitrogen lost to waterways, it is not cost-effective to remove the land from production or to remove the drainage from the land. Because of its value to agricultural production, it is better to keep tile-drained land in production under nutrient management and focus retirement policies on relatively less-productive, non-drained acres.

Introduction

The Mississippi-Atchafalaya River Basin drains 41 percent of the continental United States and accounts for about 80 percent of the United States’ total freshwater input to the Gulf of Mexico. The basin discharges an estimated 1.6 million metric tons of nitrogen each year into the Gulf, with 61 percent of that as nitrates (the mobile form of

¹ The authors wish to thank Ford Runge and Steven Taff for reviewing this manuscript.

² Authors are, respectively, Ph.D. candidate, Department of Applied Economics, University of Minnesota; Agricultural Engineer, Conservation and Production Research Laboratory, USDA-ARS, Bushland, TX; Professor, Department of Soil, Water, and Climate, University of Minnesota

nitrogen) (Goolsby, 1999; Rabalais, 1999), which leads to the formation of a hypoxic (<2 mg oxygen/L) zone along the northern Gulf coast (Rabalais et al., 2002). This “dead” zone is a significant environmental and economic problem, given that the Gulf accounted for almost one-fifth of the nation’s 2003 commercial fish landings, and just over one-fifth of the \$3 billion total value of these landings. Furthermore, the state most affected by Gulf hypoxia, Louisiana, accounts for over 10 percent of the nation’s recreational fish landings alone (Pritchard, 2004).

The birthplace of the Mississippi River, Minnesota is also home to the Minnesota River, a tributary of the Mississippi. The 17,000 square miles of land in the Minnesota River Basin comprise about one percent of the total area drained by the Mississippi River, but account for an estimated 3.3 percent of total nitrate loads. The Minnesota River is similar to many basins in the Upper Mississippi River Basin in that it is dominated by corn and soybean production, receives heavy applications of nitrogen fertilizer, has soils high in organic matter content, experiences a significant excess of precipitation over evapotranspiration, and is extensively managed with artificial drainage systems (Davis et al., 2000).

Tile (or subsurface) drainage accelerates removal of excess surface and subsurface water from fields, which, in turn, promotes well-aerated roots, enhancing plant uptake of nutrients. Such drainage also allows for timely field operations, promotes earlier plant growth, and improves yields (Zucker and Brown, 1998). In addition to these positive on-farm characteristics, tile drainage has been shown to reduce the loss of phosphorus, organic nitrogen, and other pollutants, such as certain pesticides, to

waterways (Skaggs, 1994). Because the primary method of transport of nitrate-nitrogen is at the subsurface level, however, tile drainage can significantly increase nutrient loads in waterways.³


Further, the combination of tile drainage with certain crops can significantly affect nitrate losses.⁴ The results of Randall and Mulla (2001) are, perhaps, most telling: high concentrations of nitrate-nitrogen can be lost through tile drainage from high organic matter soils even if little or no nitrogen is applied, especially in wet years that follow very dry years. In short, significant levels of nitrates can be lost on tilled land regardless of the nutrient management techniques adopted. This result has grave implications for policies that promote certain production methods, such as Best Management Practices (BMPs).

This study addresses the issue of nitrogen loading and agricultural drainage at the watershed scale by focusing on two minor watersheds, the Highwater Creek/Dutch Charlie Creek (HDCC) and Sleepy Eye Creek (SEC) watersheds, comprising 133,058 and 174,180 acres, respectively, of the Cottonwood River Watershed (USGS Cataloging Unit 07020008) in southwestern Minnesota (see Figure 1). The Cottonwood River watershed is dominated by agriculture, with 88 percent of land devoted to row crops and accounts for an estimated 9 percent of nitrate-nitrogen loads to the Minnesota River (Mulla and Mallawatantri, 1997).

³ Jackson et al. (1973), found that during a three-year study period, subsurface tile drainage accounted for 99.1 percent of the nitrate loss (annual average of 0.299 and 34.34 kg/ha for surface runoff and subsurface tile drainage, respectively). Logan et al. (1994), found that during their four-year study period, nitrate losses from tile drainage had a range of 0.01-86.0 kg/ha annually, while that of surface runoff was 0.01-2.3 kg/ha.

⁴ In a four-year study by Randall et al. (1997), average annual nitrate loss via tile drainage was 54.5, 50.75, 1.8, and 1.125 kg/ha, for continuous corn, corn-soybean/soybean-corn, alfalfa, and CRP plots, respectively.

Figure 1. Minnesota River Basin and Cottonwood River Watershed



Left: Minnesota River Basin with Cottonwood River watershed highlighted. **Above:** Cottonwood, with Sleepy Eye, Highwater, and Dutch Charlie Creek subwatersheds highlighted (clockwise from top, respectively).

Methods

Land-use⁵, climate, and soil data were collected from satellite imagery and an agricultural survey (Strock et al., 2005), the Minnesota Climatology Working Group (2004), and the State Soil Geographic (STATSGO) database (NRCS, 1994), respectively, to conduct simulations over the years 1974-2003, using the Agricultural Drainage And Pesticide Transport (ADAPT) model. ADAPT (Chung *et al.*, 1992) is a daily time step field-scale water table management simulation model that was developed by integrating GLEAMS (Groundwater Loading Effects of Agricultural Management Systems; Leonard *et al.*, 1987), a root zone water quality model, with subsurface drainage algorithms from

⁵ Land units were developed using a two-part process consisting of the development of Hydrologic Response Units (HRUs), and HRU aggregation into Transformed Hydrologic Response Units (THRUs). In HRU formation, spatial data layers of land cover, soils, and slope (STATSGO map-unit average) were overlain with ARC/INFO GIS software, resulting in a GIS layer consisting of many polygons that each contains hydrologic characteristics that are unique from those around it. Polygons which are similar in every aspect except location were then aggregated into THRUs, the functional modeling unit.

DRAINMOD (Skaggs, 1978), a subsurface drainage model. It has been calibrated at the field scale for a variety of Midwestern conditions. Results from ADAPT were used as input parameters to conduct economic analysis. Economic data were taken from Lazarus and Selley (2003), NASS (2003), and the University of Minnesota's FINBIN database (CFFM, 2004). Physical and economic data were then used as inputs to a linear constrained-optimization model solved using the Generalized Algebraic Modeling System (GAMS, 2004).

The optimization model was used to test the effectiveness of four specific land-use policies for achieving 20- and 30-percent nitrogen-load abatement. These two abatement levels have been recommended by the Mississippi River-Gulf of Mexico Watershed Nutrient Task Force as necessary for significant reductions in Gulf hypoxia (2001). The first policy was nutrient (fertilizer) management, which called for the adoption of a spring-applied 112 lb. /acre rate of nitrogen fertilizer. Spring application has been established as a Best Management Practice by the University of Minnesota (Randall and Schmitt, 1993), and the rate mentioned was the lowest of the three most commonly used application rates within the study region. The second policy was land retirement, where the current row crop was no longer grown, replaced by pasture. The third policy, labeled "plug and crop" called for land having artificial drainage to be plugged, but for the current crop to be retained. It was assumed that a loss of drainage would reduce crop yields by 20 percent. The last policy, labeled "plug and retire", called for tile-drained land to have its lines plugged in addition to being put to pasture. Because

nutrient management has been cited as one the most critical methods of nitrogen abatement, it was first tested alone. All four policies were then tested simultaneously.

Each policy was implemented under three “rules” which governed the way land was chosen for abatement. The first rule required uniform abatement, such that the required percentage of abatement was achieved on each unique land unit, and required that abatement was achieved at minimum cost. In other words, each farmer had to meet the constraint individually, and did it the cheapest way possible. The second rule placed the abatement constraint at the watershed level, and chose land for abatement based on pounds of nitrogen abated per acre. The third rule, also at the watershed level, chose land based on cost per pound of abatement. Thus, the latter two rules represent the decision of a hypothetical watershed manager to abate either using the fewest number of acres or at the lowest cost. For ease of reference, these rules are referred to as the “uniform” rule, the “per-acre” rule, and the “per-dollar” rule, respectively.

Cost, here, refers to foregone net returns to agricultural producers. Net returns on a given acre are defined as crop price per bushel multiplied by yield per acre, minus enterprise cost per acre. Because the role of drainage in nitrogen abatement was of central concern, acreage and abatement levels attributed to tile-drained land were accounted for under each scenario. It was estimated that 10 percent of the land (13,024 acres) in the HDCC watershed contains subsurface drainage, whereas 30 percent (52,255 acres) is tiled in the SEC watershed. Although crop yields were unique to each land unit, the average yields for HDCC were 150 and 43 bushels per acre for corn and soybeans, respectively, and 155 and 43 bushels per acres for SEC. Average enterprise cost was

\$270 per acre for corn and \$174 per acre for soybeans. Value per bushel of corn and beans was \$2.19 and \$6.04, respectively. The cost of plugging an acre of tile-drained land was estimated to be \$25, and an incentive payment necessary for a producer to alter fertilizer practices was estimated to be \$30 per acre.

Results

Under the base case, agricultural net returns were estimated at \$8.3 million in the HDCC and \$12.6 million in the SEC watershed. Base-case nitrogen loads were estimated at 1.2 million pounds for HDCC and 2.8 million pounds for SEC. Fertilizer management alone was not sufficient to achieve a 30-percent reduction in nitrogen loads in either of the watersheds, and was able to achieve 20-percent abatement only in the SEC watershed. Consequently, comparison of the nutrient-management policy under the three rules was impossible. The policy was able to achieve a 7-percent reduction in the HDCC watershed and a 23-percent reduction in the SEC watershed when all relevant acres were nutrient-managed, with an associated \$3.3 million and \$4.2 million price tag, respectively (see Table 1). The role of tile-drained land was critical under this policy. Although tile-drained acreage comprised only 10 and 30 percent, respectively, of all land in the HDCC and SEC watersheds, it accounted for 99 and 100 percent of abatement, respectively. Therefore, implementing nutrient management on tile-drained land alone resulted in roughly the same level of abatement at a cost of \$358,967 and \$1.4 million, respectively. Extending the policy to non-drained crop land boosted abatement in HDCC by an additional 1-percent at an additional cost of \$2.9 million, and actually reduced abatement in SEC for an additional \$2.8 million. (This reduction in abatement was due, in part, to

Table 1. Nutrient-management policy results for the HDCC and SEC watershed.

Nutrient-Management Policy			
HDCC			
Policy	Net Return	N Load (lbs.)	% Abatement on Tile-drained Land
Base Scenario	\$8,291,310	1,233,080	0%
All Land N-Managed	\$5,007,233	1,147,522	99%
<i>% N Abated</i>		<i>(7%)</i>	
Tile-drained Land N-Managed	\$7,932,343	1,148,383	100%
<i>% N Abated</i>		<i>(7%)</i>	
SEC			
Policy	Net Return	N Load (lbs.)	% Abatement on Tile-drained Land
Base Scenario	\$12,627,309	2,777,748	0%
All Land N-Managed	\$8,412,698	2,134,301	100%
<i>% N Abated</i>		<i>(23%)</i>	
Tile-drained Land N-Managed	\$11,219,355	2,134,185	100%
<i>% N Abated</i>		<i>(23%)</i>	

the atypical result of some soils whose nitrogen-losses actually increased when fertilizer application was switched from fall to spring.)

Tables 2 and 3 contain the results of implementing the four abatement policies under the three rules in the HDCC and SEC watersheds to achieve 20- and 30-percent abatement, respectively. Because the “plug and crop” policy was not part of any solution, it was omitted from the tables. The per-dollar rule resulted in the highest net returns under 20-percent abatement, saving \$1.4 million in the HDCC relative to the uniform-abatement rule, and \$2.3 million in the SEC. Differences in net returns were similar under 30-percent abatement. Net returns under the per-acre abatement rule fell between those of per-dollar and uniform abatement. In addition to cost differences, land-use choices were substantially different under the three abatement rules. The uniform-abatement rule required the greatest number of retired acres to satisfy a given constraint,

Table 2. Policy results for the HDCC and SEC watersheds with 20% abatement.

<u>20% N-LOAD ABATEMENT</u>					
<u>HDCC</u>					
Policy Rule	Net Return	Acres Retired	Acres Plugged & Retired	Acres Nutrient-Managed	% Abatement on Tile-Drained Land
Base Scenario	\$8,291,310	0	0	0	0%
Per-Dollar	\$7,716,599	22,140	0	5,247	43%
Per-Acre	\$6,575,254	7,834	10,693	0	76%
Uniform	\$6,317,563	33,657	0	4,760	17%

<u>SEC</u>					
Policy Rule	Net Return	Acres Retired	Acres Plugged & Retired	Acres Nutrient-Managed	% Abatement on Tile-Drained Land
Base Scenario	\$12,627,309	0	0	0	0%
Per-Dollar	\$11,780,977	3,928	0	31,979	95%
Per-Acre	\$11,054,926	18,596	0	0	100%
Uniform	\$9,483,958	33,476	0	21,907	50%

and relied least on tile-drained acreage. When the per-acre rule was followed to achieve 20-percent abatement, there was a four-and-a-half-fold increase in the dependence on tile-drained land relative to uniform abatement in the HDCC watershed. In the SEC, it increased abatement from half to all on tile-drained land. Furthermore, the number of retired acres necessary to achieve abatement fell by half in both watersheds.

Additionally, abatement by nutrient management was relatively inefficient under the per-acre rule, and no land followed that policy under that rule. Based on these results, it can be concluded that targeting for tile-drained land was more effective in terms of abatement per acre. It remains to be seen whether it was cost-effective. Thus, hereafter, the discussion will focus on differences between abatement under the per-acre and per-dollar rules.

When the per-dollar rule was followed to achieve 20-percent abatement, the

Table 3. Policy results for the HDCC and SEC watersheds with 30% abatement.

30% N-LOAD ABATEMENT					
<u>HDCC</u>					
Policy Rule	Net Return	Acres Retired	Acres Plugged & Retired	Acres Nutrient-Managed	% Abatement on Tile-Drained Land
Base Scenario	\$8,291,310	0	0	0	0%
Per-Dollar	\$6,861,092	39,080	0	4,692	37%
Per-Acre	\$5,915,503	23,289	11,383	0	52%
Uniform	\$5,226,218	47,719	0	6,723	18%

<u>SEC</u>					
Policy Rule	Net Return	Acres Retired	Acres Plugged & Retired	Acres Nutrient-Managed	% Abatement on Tile-Drained Land
Base Scenario	\$12,627,309	0	0	0	0%
Per-Dollar	\$10,617,160	25,714	0	40,214	76%
Per-Acre	\$9,915,311	27,510	3,855	0	100%
Uniform	\$7,891,215	49,839	0	31,945	50%

Proportion attributed to tile-drained acreage fell from 76 to 43 percent relative to the per-acre rule in the HDCC, and from 100 to 76 percent in the SEC. Also, nutrient management was part of the solution under the per-dollar rule whereas the “plug and retire” policy was not, indicating that nutrient management was cost-effective and the “plug and retire” policy was not. In the SEC watershed in particular, following the per-dollar rule reduced the number of retired acres under 20-percent abatement from 18,596 to 3,928. Thus, nutrient management was relatively ineffective on an abatement-per-acre basis, with no acreage being subject to it under the per-acre rule, but was effective on a per-dollar basis, with 4,700-5,200 acres in the HDCC and 32,000-40,000 acres in the SEC being nutrient-managed. Further, between 93-100 percent of all acres selected for nutrient-management were tile-drained acres.

With regard to the level of abatement, cost differences were of the same magnitude under 20- and 30-percent abatement. Land-uses changes, however, were substantial when abatement increased from 20 to 30 percent. Following the per-acre rule, increasing abatement from 20 to 30 percent increased the number of retired and plug-and-retired acres by 53 percent in the HDCC and by 68 percent in the SEC. Similar results were found under the per-dollar rule, with the increase in abatement requiring an additional 77 percent of retired acreage in the HDCC. In the SEC watershed, the most significant change in land use when abatement was increased was found under the per-dollar rule, where retired acres increased 554 percent.

Discussion

The results of this study indicate that choice of policy and the rule by which that policy is implemented can have a significant impact on the income and land use in a given watershed. Further, the degree to which these differences occur depends, largely, on the characteristics of the watershed in question. These results indicate, as expected, that a watershed with relatively more drainage, such as the SEC watershed, will rely more heavily on tile-drained land to achieve a given level of nitrogen abatement. Results also indicate that abatement is more effective on tile-drained land, and hence focusing attention on land that is tile-drained is a more efficient means of achieving abatement objectives. This conclusion was especially true under the nutrient-management policy. Therefore, the results indicate that close attention should be paid to nutrient practices on tile-drained land, but not on non-drained land.

Furthermore, results indicate that policies that are effective on an abatement-per-acre basis are not necessarily cost-effective. This was true of a policy of plugging and retiring an acre of tile-drained land. Although it was cost-effective to retire tile-drained land from production, it was not at all cost-effective to remove the drainage on that land as well. In other words, it is more cost-effective to address what activity is undertaken on tile-drained land rather than to address the drainage itself. This conclusion is bolstered by the fact that under no policy was it effective to remove drainage on an acre of land but keep it in production. Finally, meeting an abatement constraint of 20 percent required relatively few acres to be retired or managed; however, increasing the abatement constraint, by just 10 percent in this case, required an almost two-fold increase in retired acreage.

The implications of this study are summarized in Table 4, which indicates, based on the results, whether a policy was cost-effective in achieving nitrogen-load abatement on tile-drained and non-drained land, respectively. As the table shows, a combination of nutrient management on row-cropped tile-drained acres combined with land retirement (mostly on non-drained acres) is the most cost-effective means of achieving nitrogen-load abatement, given the tested policy choices. Further, these results indicate that because crop yields are relatively higher on tile-drained land, it is more cost-effective to

Table 4. Cost-effectiveness of each policy on tile-drained and non-drained land to achieve nitrogen-load abatement.

Is the Policy Cost Effective on this Land Type?				
Land Type	Retire	Plug & Retire	Plug & Crop	Nutrient Manage
Tile-drained Land	NO*	NO	NO	YES
Non-drained Land	YES	-	-	NO

*Marked as “NO” because tile-drained land never comprised more than 15% of retired acreage under any scenario following the per-dollar rule.

keep tile-drained acres in production under nutrient management, and to focus most of the land retirement on non-drained acres (tile-drained acreage never comprised more than 15 percent of retired acreage under any scenario following the per-dollar rule).

Results also imply that gains from abatement trading may be possible if watersheds abate cooperatively. Under the per-dollar rule policies, the marginal cost per pound of abatement in the SEC watershed was half that of the HDCC watershed. Subsequent research on agricultural nitrogen loading would do well to investigate the benefits of cooperative abatement and abatement trading.

Literature Cited

- Chung, S.O., A.D. Ward, and C.W. Shalk. "Evaluation of the hydrologic component of the ADAPT water table management model." *Trans. ASAE* 35(1992):571-579.
- Davis, D.M., P.H. Gowda, D.J. Mulla, and G.W. Randall. "Modeling Nitrate Nitrogen Leaching in Response to Nitrogen Fertilizer Rate and Tile Drain Depth or Spacing for Southern Minnesota." *JEQ*, 29(2000): 1568-1581.
- Center for Farm Financial Management (CFFM). "FINBIN" online database, <http://www.finbin.umn.edu/>, Dept. of Applied Economics, University of Minnesota, St. Paul, 2004.
- GAMS Development Corporation. *Generalized Algebraic Modeling System*, version 2.5, 2004.
- Goolsby, D.A., W.A. Battaglin, G.B. Lawrence, R.S. Artz, B.T. Aulenbach, R.P. Hooper, D.R. Kenney, and G.J. Stensland. "Flux and Sources of Nutrients in the Mississippi-Atchafalaya River Basin: Topic 3 Report for the Integrated Assessment of Hypoxia in the Gulf of Mexico." NOAA Coastal Ocean Program Decision Analysis Series No. 17. NOAA Coastal Ocean Program, Silver Spring, MD, May 1999.
- Jackson, W.A., L.E. Asmussen, E.W. Hauser, and A.W. White. "Nitrate in Surface and Subsurface Flow from a Small Agricultural Watershed." *JEQ*, 2(1973): 480-482.
- Lazarus, William F. and Roger Selley. "Minnesota farm machinery economic cost estimates for 2004". FO-6696, University of Minnesota Extension Service, 2003.
- Leonard, R.A., W.G. Knisel, and D.A. Still. "GLEAMS: groundwater loading effects of agricultural management systems." *Trans. ASAE* 30(1987): 1403-1418.
- Logan, T.J., D.J. Eckert, and D.G. Beak. "Tillage, crop and climatic effects on runoff and tile drainage losses of nitrate and four herbicides." *Soil and Tillage Res.*, 30(1994): 75-103.
- Minnesota Climatology Working Group. "Historical Climate Data", <http://www.climate.umn.edu/doc/historical.htm>, 2004.
- Mississippi River/Gulf of Mexico Watershed Nutrient Task Force. *Action Plan for Reducing, Mitigating, and Controlling Hypoxia in the Northern Gulf of Mexico*. Washington, D.C., 2001.
- Mulla, David J. and Prasanna H. Gowda. Unpublished manuscript, 2003.
- Mulla, David J. and A.P. Mallawatantri. "Minnesota River Basin Water Quality Overview." Minnesota Extension Service Pub. No. F-7079-E, University of Minnesota, St. Paul, 1997.
- National Agricultural Statistics Service. *Agricultural Statistics*, various issues, USDA, 2004.
- Natural Resource Conservation Service. *State Soil Geographic Database (STATSGO)*, 1994.
- Pritchard, Elizabeth S, ed. *Fisheries of the United States 2003*. National Marine Fisheries Service, Office of Science and Technology, Fishery Statistic and Economics Division, Silver Spring, MD, October 2004.
- Rabalais, N.N., R.E. Turner, D. Justic, Q. Dortch, and W.J. Wiseman, Jr.

- “Characterization of Hypoxia: Topic 1 Report for the Integrated Assessment of Hypoxia in the Gulf of Mexico.” NOAA Coastal Ocean Program Decision Analysis Series No. 15. NOAA Coastal Ocean Program, Silver Spring, MD, May 1999.
- Rabalais, N.N., R.E. Turner, and D. Scavia. “Beyond Science into Policy: Gulf of Mexico Hypoxia and the Mississippi River.” *BioScience*, February 2002, 52(2): 129-142.
- Randall, G.W., D.R. Huggins, M.P. Russelle, D.J. Fuchs, W.W. Nelson, and J.L. Anderson. “Nitrate Losses through Subsurface Tile Drainage in Conservation Reserve Program, Alfalfa, and Row Crop Systems.” *JEQ*, 26(1997): 1240-1247.
- Randall, G.W. and M.S. Schmitt. “Best Management Practices for Nitrogen Use Statewide in Minnesota.” Extension Publication FO-06125-GO, University of Minnesota, 1993.
- Skaggs, R.W. “A water management model for shallow water table soils.” Rep. No. 134. Water Research Institute of North Carolina, Raleigh, N.C, 1978.
- Strock, J.S., D. Bruening, J.D. Apland, and D.J. Mulla. “Farm nutrient management practices in two geographically diverse watersheds in the Cottonwood River Watershed of Minnesota”, submitted for review, 2005.
- Zucker, Leslie A. and Larry C. Brown (Eds.). *Agricultural Drainage: Water Quality Impacts and Subsurface Drainage Studies in the Midwest*. OSU Extension Bulletin 871. The Ohio State University, 1998.