



United States
Department of
Agriculture

Marketing and
Regulatory
Programs

Animal and
Plant Health
Inspection
Service



Proposed Program for the Control of the Woodwasp *Sirex noctilio* F. (Hymenoptera: Siricidae) in the Northeastern United States

Environmental Assessment,
May 2008

Proposed Program for the Control of the Woodwasp *Sirex Noctilio* F. (Hymenoptera: Siricidae) in the Northeastern United States

Environmental Assessment May 2008

Agency Contact:

Lynn Evans-Goldner
Emergency and Domestic Programs
Plant Protection and Quarantine
Animal and Plant Health Inspection Service
U.S. Department of Agriculture
4700 River Road, Unit 137
Riverdale, MD 20737

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 1400 Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

Mention of companies or commercial products in this report does not imply recommendation or endorsement by the U.S. Department of Agriculture over others not mentioned. USDA neither guarantees nor warrants the standard of any product mentioned. Product names are mentioned solely to report factually on available data and to provide specific information.

This publication reports research involving pesticides. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.

Table of Contents

I. Background and Introduction	1
A. Description and Biology of <i>S. noctilio</i>	2
B. History of Invasiveness of <i>S. noctilio</i>	3
C. <i>Sirex noctilio</i> in North America	3
II. Purpose and Need for the Proposed Action	4
A. Decisions to be Made	5
B. Relevant Issues	5
III. Alternatives	5
A. No Action	6
B. Quarantine Only	6
C. Biological Control Only	11
D. Quarantine and Biological Control	13
IV. Affected Environment	14
A. North American Siricid Complex, Their Fungal Food Sources, and the Nematodes That Infect Them	14
B. Pine Resources of North America	16
V. Environmental Consequences	17
A. No Action	17
B. Quarantine Only	19
C. Biological Control Only	24
D. Quarantine and Biological Control	23
E. Cumulative Effects	24
F. Endangered Species Act	25
VI. Other Issues	26
VII. Agencies, Organizations, and Individuals Consulted	27
VIII. References	28
Appendices	
A. Organism Pest Risk Assessment	
B. Dicamba and Triclopyr Toxicity Data	

I. Background and Introduction

The United States Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS), is proposing a program to control a nonnative, invasive woodwasp, *Sirex noctilio* (Hymenoptera: Siricidae). This program is necessary to control *S. noctilio* and reduce the potential for damage from this major pest of pine trees. This environmental assessment¹ (EA) has been prepared, consistent with USDA, APHIS' National Environmental Policy Act of 1969 (NEPA) implementing procedures (Title 7 of the Code of Federal Regulations (CFR), part 372). It examines the potential effects on the quality of the human environment that may be associated with a proposal to control infestations of *S. noctilio* within the currently infested States of New York, Pennsylvania, Michigan, and Vermont, as well as States where *S. noctilio* would likely naturally spread within the next 10 years (based on a 25-mile per year rate of spread), including Connecticut, Delaware, Indiana, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, Ohio, Rhode Island, Virginia, and West Virginia (USDA, APHIS, 2007a (refer to appendix A)). This EA considers the potential effects of the proposed action and its alternatives, including no action.

APHIS prepared an EA and finding of no significant impact (FONSI) in March 2007, and both are incorporated here by reference (USDA, APHIS, 2007b). In the EA, APHIS considered a program to control *S. noctilio* in New York and Pennsylvania, the only States where *S. noctilio* had been detected at that time. The EA evaluated four alternatives: no action, a quarantine program, a biological control program, and a combination of quarantine and biological control (preferred action) in New York and Pennsylvania. A FONSI was signed on June 21, 2007, and determined that the proposed program (including quarantine and biological control) would not have a significant impact on the quality of the human environment in those States. Since that time, *S. noctilio* has been detected in additional States, and APHIS would like to implement a control program in those States. In addition, if *S. noctilio* were detected in other States, APHIS would want to implement a control program in those States, as well. Therefore, this EA considers the potential effects on the quality of the human environment from implementing a program for control of *S. noctilio* in the currently infested States and surrounding States where *S. noctilio* would most likely be detected.

¹ Regulations implementing the National Environmental Policy Act of 1969 (42 United States Code 4321 et seq.) provide that an EA "[shall include brief discussions of the need for the proposal, of alternatives as required by section 102(2)(E), of the environmental impacts of the proposed action and alternatives, and a listing of agencies and persons consulted." 40 CFR § 1508.9.

A. Description and Biology of *S. noctilio*

Sirex noctilio is a large insect, usually 1.0 to 1.5 inches long, with a metallic blue-black body coloration. The adult female is metallic blue except for legs which are reddish-brown. Pointing backwards from the underside of the abdomen is a spike-like projection that protects the ovipositor (egg-laying tube) when it is not in use. In the adult male, the middle segments of the abdomen are orange-yellow and the hind legs are thickened and almost wholly black. Larvae are creamy white, legless, and have a distinctive dark spine at the rear of the abdomen. Unlike some other wasps in the insect wasp family (Hymenoptera), siricid wasps (like *Sirex* spp.) do not sting animals.

The life cycle of *S. noctilio* is typically 1 year, but it can be completed in as little as 3 months in the laboratory or may last up to 2 years in cool climates. Adult *S. noctilio* wasps emerge during the summer through mid-autumn. Males emerge before the females and may outnumber the females by 20:1. Females that do not find mates lay eggs that produce all males. Mated females produce offspring of either sex. Females can begin to lay eggs 1 day after adult emergence. Adults do not feed but instead live on fat stored in their bodies. The adult life span can last up to 12 days; however, a female who has deposited all her eggs may live for as little as 3 or 4 days. Throughout their short life, and depending on their size, females deposit between 20 and 500 eggs.

The female inserts her ovipositor into the wood of a pine tree and drills as many as three to four holes. In the first and second drill holes, the female injects mucus that is toxic to pine trees and spores of a fungus (*Amylostereum areolatum* (*A. areolatum*)). The mucus disables the tree's transport system for water and nutrients, creating ideal conditions for the spread of the fungus. The fungus kills the tree and causes the wood to dry out, creating a more favorable environment for *S. noctilio* larval development and stimulating egg hatch. The fungus is later fed on by the *S. noctilio* larvae. Larvae hatch from the eggs after a minimum of 9 days, but the eggs may remain dormant for several months, overwintering in cooler climates. Larval tunnels in the wood are tightly packed with very fine sawdust. The pupal stage lasts between 16 and 21 days. When a female wasp emerges from her pupal skin, she takes up fungus spores from the tree from which she emerges and stores them in a special organ in her abdomen. The adult wasp bores its way out of the tree leaving a characteristic round exit-hole which varies in diameter, according to the size of the wasp.

Sirex noctilio can attack living, healthy pines, while woodwasps native to the United States attack only dead and dying trees. At low populations, *S. noctilio* selects suppressed, stressed, and injured trees for laying eggs,

however, at high densities will select healthy trees. Generally, foliage of infested trees initially wilts, and then changes color from dark green to light green, to yellow, and finally to red during the 3 to 6 months following attack. Infested trees may have resin beads or dribbles at the egg-laying sites.

B. History of Invasiveness of *S. noctilio*

Sirex noctilio is native to Europe, Asia, and northern Africa, where it is generally considered to be a secondary pest. In its native range, it attacks pines almost exclusively, for example, Scotch (*Pinus sylvestris*), Austrian (*P. nigra*), and maritime (*P. pinaster*) pines. It was introduced inadvertently into Argentina, Australia, Brazil, Chile, New Zealand, South Africa, and Uruguay. In these Southern Hemisphere countries, *S. noctilio* attacks exotic pine plantations where it has caused up to 80 percent tree mortality. Most of the plantations are planted with North American pine species, especially Monterey pine (*P. radiata*) and loblolly pine (*P. taeda*). Pines are the main hosts of *S. noctilio*, and it can complete its life-cycle on multiple species within the pine genus (*Pinus*). The following *Pinus* species are known to be hosts of *S. noctilio*: *P. radiata*, *P. nigra*, *P. calabrica*, *P. nigra austriaca*, *P. ponderosa*, *P. elliotii*, *P. patula*, *P. contorta*, *P. caribaea*, *P. pinaster*, *P. attenuata*, *P. muricata*, *P. banksiana*, *P. canariensis*, *P. densiflora*, *P. echinata*, *P. halepensis*, *P. jeffreyi*, *P. palustris*, *P. pinaster*, *P. pinea*, *P. brutia*, *P. sylvestris*, and *P. taeda*.

Thus far, in North America, jack pine (*P. banksiana*), *P. resinosa* (red pine), *P. strobus* (white pine), *P. sylvestris* (Scotch pine) and have served as hosts for *S. noctilio*. In addition, *S. noctilio* has been recorded on other conifers such as larch, Douglas fir, spruce, and fir.

C. *Sirex noctilio* in North America

In February, 2005, a single female *S. noctilio* woodwasp was identified in a sample collected as part of the New York State Cooperative Agricultural Pest Survey for Exotic Wood Borers and Bark Beetles. It had been collected on September 7, 2004, from a trap placed among mixed hardwoods and pine, just inside a forest edge near the city of Fulton, New York, in Oswego County. From this initial site, positive findings from subsequent surveys have also been identified in a number of New York counties including: Allegany, Broome, Cattaraugus, Cayuga, Chautauqua, Chemung, Cortland, Erie, Genesee, Hamilton, Herkimer, Jefferson, Livingston, Madison, Monroe, Niagara, Oneida, Onondaga, Ontario, Orleans, Oswego, Schuyler, Seneca, Steuben, St. Lawrence, Tomkins, Wayne, Wyoming, and Yates.

The Pennsylvania Department of Agriculture announced, on July 27, 2006, that *S. noctilio* was found in that State for the first time. A single adult female was found in a trap in Hills Creek State Park in Tioga County. Subsequently, there have been finds in Bradford, Elk, Erie, McKean, Monroe, and Tioga Counties. Agriculture and forestry officials are conducting an intensive trapping survey from the infested site in the northern Pennsylvania counties to determine the range of the wasp.

On September 15, 2007, APHIS confirmed the detection of a single adult *S. noctilio* female in a trap in Lamoille County, Vermont. Although the survey has continued in that State, no additional detections have occurred.

In Michigan, *S. noctilio* was first detected in Macomb County on July 6, 2007. Since then, two *S. noctilio* adult females were trapped in Sanilac County on July 7 and July 25, 2007.

In addition to these recent infestations, *S. noctilio* has also been identified and occurs in several locations in Ontario, Canada.

A prediction model for potential distribution of *S. noctilio* in North America found that many areas of the United States, Canada, and Mexico were suitable for establishment (Carnegie et al., 2006). APHIS expects that distribution of pines will probably limit the spreading of *S. noctilio* rather than climate. Thus, it is believed that *S. noctilio* could likely establish wherever pines are found in North America.

If not controlled, *S. noctilio* poses a serious threat to pine forests and plantations in the United States. It attacks many *Pinus* species, including those of economic, environmental, and aesthetic importance and, unlike other siricid (belonging to the insect family Siricidae) species, it attacks and kills healthy, living trees. Uncontrolled *S. noctilio* populations can build up to outbreak levels rapidly. It is a serious pest of pine plantations in Australia, New Zealand, South America, and South Africa.

II. Purpose and Need for the Proposed Action

APHIS is proposing a program for the purpose of managing the natural and artificial spread of *S. noctilio* with the release of a parasitic nematode, *Beddingia siricidicola* (*B. siricidicola*), and the implementation of a quarantine wherever *S. noctilio* is detected in the United States. This program is not expected to eradicate *S. noctilio*; it is not feasible to do so (USDA, APHIS, 2007a). However, there is a need to control *S. noctilio*, because it has the potential to be a serious threat to pine forests, plantations, and landscape plantings in the United States if not managed.

APHIS has responsibility for taking actions to exclude, eradicate, and/or control plant pests, including *S. noctilio*, under the Plant Protection Act (7 United States Code (U.S.C.) 7701 et seq.). APHIS has been delegated the authority to administer these statutes, and has promulgated quarantines and regulations (7 CFR 319) which regulate the importation of commodities and means of conveyance to help protect against the introduction and spread of harmful pests. The underlying strategy of the proposed program is to contain *S. noctilio* by reducing the population density and spread of the woodwasp from currently infested areas to other counties and States.

A. Decisions to Be Made

A decision that APHIS must make is whether or not to expand the proposed program beyond New York and Pennsylvania to control *S. noctilio* in the United States. Depending on the first decision, a second decision that APHIS must make is what control measures will be included (quarantine, biological control, or a combination of both) in the expanded program to protect *Pinus* species while avoiding adverse impacts on the human environment.

B. Relevant Issues

This EA will deal in detail with the issues of—

Biological Resources: The protection of *Pinus* species (and the species that depend on them) in the United States and impact of program activities on native siricids and other nontarget species.

Threatened and Endangered Species: Section 7 of the Endangered Species Act requires that APHIS determine that the actions it authorizes, funds, or carries out do not jeopardize the existence of threatened or endangered species or critical habitat.

III. Alternatives

APHIS considered four alternatives in response to the need to manage *S. noctilio* and contain infestations: (A) no action, (B) quarantine only, (C) biological control only, and (D) the combination of quarantine and biological control (preferred alternative). Each alternative is described briefly in this section, and the potential impacts of each are considered in the following section. Eradication and chemical control are alternatives that are not considered further because they are not feasible. (A summary of issues and impacts associated with each of the alternatives is found in table 1.)

A. No Action

Under the no action alternative, APHIS would not implement any measures to manage *S. noctilio* infestations. Some control or management measures could be taken by other Federal or non-Federal entities; those actions would not be under APHIS' control nor funded by APHIS. Local business owners and area residents could attempt to limit damage from *S. noctilio* infestations by removing the infested trees from their properties. The lack of measures to prevent the spread of *S. noctilio* from infested areas (occurring via natural dispersal of the insect or artificial spread from movement of infested pine products) could lead to an increase in *S. noctilio* populations and increase its range of distribution within the United States.

B. Quarantine Only

Under this alternative, APHIS would implement a quarantine area within and around *S. noctilio*-infested States or portions of States. APHIS would work cooperatively with the appropriate State regulatory officials to establish a quarantine area in an effort to limit the artificial spread of *S. noctilio*. Actions will be taken, as needed, to increase the quarantine areas as the areas of infestation expand. This alternative is expected to prevent the artificial spread of *S. noctilio* by limiting the movement of *Pinus* spp. wood products from *S. noctilio*-infested areas into uninfested areas; however, it is not expected to prevent the natural dispersal of the insect. To reduce the risk of artificial movement of *S. noctilio* from infested areas to uninfested areas, the following conditions would apply:

1. Regulated Articles

Certain articles present a significant risk of spreading *S. noctilio* if the articles are moved from quarantined areas without restriction; these are called "regulated articles." The following are designated as regulated articles—

- Pine wood packaging materials, pine pallets, loose wood packing material made of pine, and nonmanufactured pine wood articles used in the repair and remanufacture of wood packaging material;
- Except for mugo pine (*Pinus mugo*) nursery stock, all pine nursery stock with trunks 4 inches or larger in diameter, as measured at the soil line;
- Dug, balled, and burlapped pine Christmas trees with trunks 4 inches or larger in diameter, as measured at the soil line;
- Cut pine Christmas trees that are collected from the wild or taken from an established landscape planting;

- Pine firewood;
- Any of the following pine items with components greater than 1 inch in thickness in any two dimensions: Wood shavings, wood chips, mulch, wood millings and fines, wood pellet components, wood residue, wood scraps, wood slabs, and wood waste;
- Pine logs, log home components, kit homes, lumber, log furniture, decorative items, sawn wood, round wood, and nonmanufactured wood products; and
- Pine pulpwood with components greater than 1 inch in thickness in any two dimensions.

In addition, any other article, product, or means of conveyance may be designated as a regulated article if an inspector determines that it presents a risk of spreading *S. noctilio* and notifies the person in possession of the article, product, or means of conveyance that it is subject to the restrictions in the regulations. This allows an inspector who discovers evidence of *S. noctilio* in an article, product, or means of conveyance to take immediate action after informing the person in possession of it that it is being regulated.

2. Quarantined Areas

Each State or portion of a State will be quarantined where *S. noctilio* has been found by an inspector, where APHIS has reason to believe that *S. noctilio* is present, or which APHIS deems necessary to regulate because of its inseparability for quarantine purposes from localities where *S. noctilio* has been found. Less than an entire State will be designated as a quarantined area only under certain conditions. Such a designation may be made if APHIS determines that: (1) the State has adopted and is enforcing restrictions on the intrastate movement of regulated articles listed above that are equivalent to the interstate movement restrictions imposed by APHIS; and (2) the designation of less than an entire State will be adequate to prevent the artificial spread of *S. noctilio*.

In accordance with these criteria and because of *S. noctilio*, APHIS is quarantining 8 counties in Michigan, 1 county in New Jersey, 60 counties in New York, 1 county in Ohio, 23 counties in Pennsylvania, and 6 counties in Vermont.

3. Interstate Movement of Regulated Articles

Regulated articles may be moved interstate from a quarantined area into or through an area that is not quarantined if they are accompanied by a certificate or limited permit.

A regulated article may be moved interstate without a certificate or limited permit by USDA for experimental or scientific purposes or if the regulated article originates outside the quarantined area. Articles originating outside the quarantined area that are moved interstate through a quarantined area must be moved under the following conditions: (1) the points of origin and destination are indicated on a waybill accompanying the regulated article; (2) the regulated article, if moved between June 1 and October 31, is moved in an enclosed vehicle or is completely covered to prevent access by *S. noctilio*; (3) the regulated article is moved directly through the quarantined area without stopping (except for refueling or for traffic conditions such as traffic lights or stop signs); and (4) the article has not been combined or commingled with other articles so as to lose its individual identity. These requirements are to prevent contamination of articles moving from uninfested areas through quarantined areas.

Articles may be moved with a limited permit issued by APHIS if the regulated articles are moved by USDA for experimental or scientific purposes. These articles must be moved under conditions specified on the permit, which APHIS has found to be adequate to prevent the spread of *S. noctilio*. A tag or label bearing the number of the permit issued for the regulated article must be attached to the outside of the container of the regulated article, or attached to the regulated article itself if the regulated article is not in a container.

4. Certificates and Limited Permits

An inspector or person operating under a compliance agreement (discussed below in section III.B.5) may issue a certificate for the interstate movement of a regulated article if he or she determines that the regulated article:

- Has been treated (discussed below in section III.B.6.) within the quarantined area under the direction of an inspector and, if moved between June 1 and October 31, is moved in an enclosed vehicle or is completely covered to prevent access by *S. noctilio*, is moved directly through the quarantine area without stopping (except for refueling or for traffic conditions, such as traffic lights or stop signs), and has not been combined or comingled with other articles so as to lose its individual identity; or
- Is pine nursery stock (other than mugo pine nursery stock) 4 inches or larger in diameter as measured at the soil line, including balled and burlapped Christmas trees, and it has been harvested from a nursery and not collected from the wild or taken from an

established landscape planting, has been inspected and found free of signs and symptoms of *S. noctilio* attack, and will be moved only between March 15 and June 15; and

- Is to be moved in compliance with any additional emergency conditions that APHIS may impose under section 414 of the Plant Protection Act (7 U.S.C. 7714) in order to prevent the artificial spread of *S. noctilio*; and
- Is eligible for unrestricted movement under all other Federal domestic plant quarantines and regulations applicable to the regulated articles.

A limited permit (rather than a certificate) can be issued by an inspector or person operating under a compliance agreement for movement of a regulated article if he or she determines that the regulated article:

- Is to be moved between November 1 and May 31 to a facility operating under a compliance agreement with APHIS and treated at the destination within 30 days of arrival at the destination, or by June 1, whichever is sooner; or
- Is to be moved interstate between November 1 and May 31 to a specified destination for specific processing, handling, or utilization (the origin, destination, and other conditions to be listed on the limited permit), and this interstate movement will not result in the spread of *S. noctilio* because *S. noctilio* will be destroyed or the risk otherwise mitigated by the specific processing, handling, or utilization; and
- Is to be moved in compliance with any additional emergency conditions that APHIS may impose under section 414 of the Plant Protection Act (7 U.S.C. 7714) in order to prevent the spread of *S. noctilio*; and
- Is eligible for unrestricted movement under all other Federal domestic plant quarantines and regulations applicable to the regulated article.

5. Compliance Agreements

Compliance agreements are provided for the convenience of persons who are involved (1) in the growing, handling, or movement of regulated articles from quarantined areas, or (2) the movement of articles that originate outside the regulated area and are moved through a regulated area. A person may enter into a compliance agreement when an inspector has determined that the person requesting the compliance agreement has been made aware of the requirements of the regulations, and the person

has agreed to comply with the requirements of the regulations and the provisions of the compliance agreement.

Treatments to destroy *S. noctilio* listed in 7 CFR part 305, “Phytosanitary Treatments,” (discussed in section III.B.6.) are authorized for use on certain regulated articles if those articles are to travel under a certificate. If the regulated articles are moved between June 1 and October 31, they must be treated before movement and the treatment must be listed on certificate. If the regulated articles are moved between November 1 and May 31, they may be treated either in the quarantined area or moved to a facility operating under a compliance agreement and treated at the facility within 30 days of arrival or by June 1, whichever is sooner.

Pulpwood, loose wood packing material, wood waste, millings, wood pellet components, wood scraps, and other residue greater than 1 inch thickness in any two dimensions may also be treated by chipping to 1 inch or less in any two dimensions.

6. Treatments

The regulations in 7 CFR part 305, “Phytosanitary Treatments,” provide treatment schedules and other requirements related to approved treatments. Paragraph (v) of § 305.2 lists the treatment schedules approved for certain wood articles, including containers, oak logs and lumber, and Christmas trees. APHIS will amend these regulations to add entries for several of the regulated pine wood articles. Specifically, APHIS will add entries for: nonmanufactured pine wood articles less than 3 inches in diameter; nonmanufactured pine wood articles 3 inches or larger in diameter; pine pulpwood, loose wood packing material, wood waste, millings, wood pellet components, wood scraps, and other residue greater than 1-inch thickness in two dimensions; and pine logs, log home components, log home kits, lumber sawn wood, round wood, and log furniture. For each entry, APHIS will indicate whether the article may be treated with heat or methyl bromide fumigation.

APHIS will add schedule T404-e-1 (methyl bromide treatment) to § 305.6 for use on pine wood packing materials, including pine components used to repair and remanufacture pallets. APHIS will also add schedule T404-e-2 (heat treatment) to § 305.25 to provide an alternative treatment for logs, nonmanufactured wood articles, pulpwood, loose wood packing material, solid wood packaging material, wood waste, and certain other articles. This treatment schedule calls for these articles to be heated to a minimum core temperature of 132.8 °F (56 °C) and held at that temperature for a minimum time period of 30 minutes. Research indicates that these treatments will be effective to destroy *S. noctilio* in regulated articles.

C. Biological Control Only

Under this alternative, APHIS would work cooperatively with the USDA Forest Service and the State Departments of Agriculture of infested States to implement a biological control program. Implementation of this alternative would reduce, but not eradicate, *S. noctilio* in the United States. *Sirex noctilio* has been successfully managed in Australia, New Zealand, and South America by using *B. siricidicola* as a biological control agent. This parasitic nematode infects *S. noctilio* larvae and, ultimately, sterilizes the adult females. Female *S. noctilio* woodwasps infected by the nematode *B. siricidicola* emerge and lay infertile eggs that are filled with nematodes. This facilitates the spread of the nematode population. The goal is to reduce and maintain *S. noctilio* below damaging levels and reduce the natural spread of *S. noctilio*.

1. Description and Biology of *Beddingia siricidicola*

A nematode is a very small (approximately 0.4 millimeters (mm)-1.5 mm), unsegmented worm with an elongated, rounded body that is pointed at both ends. Like other nematodes, *B. siricidicola* has an egg stage, several juvenile stages, and an adult stage. It reproduces sexually. Unlike other nematodes, it has two very different potential forms, dependent upon the microenvironment—a mycetophagous type (fungus-feeding) and entomophagous type (insect-feeding or parasitic) (Bedding, 1972). Mycetophagous nematodes grow, develop, and reproduce entirely by consuming the fungus that *S. noctilio* carries, *A. areolatum*. These nematodes can do so indefinitely, producing many generations in a season.

If nematode juveniles develop near *S. noctilio* larva, however, they develop into entomophagous adults. The sexes mate and the females bore into *S. noctilio* larvae. Shortly before pupation of the *S. noctilio* larva, a female nematode produces many infective juveniles which migrate to the host gonads (reproductive glands, such as ovaries or testes). The infective stages do not sterilize males but only enlarge the testes; however, they move into the ovaries of *S. noctilio* females and enter the eggs, rendering them sterile.

After the *S. noctilio* female emerges, she then lays eggs containing only nematode juveniles. Inside the tree, the juvenile nematodes develop as the mycetophagous form; however, after the first generation, they can develop as either form (mycetophagous or entomophagous), dependent upon microenvironment, and then the cycle begins again. In the absence of human-assisted movement, the dispersal capacity of *B. siricidicola* is completely dependent upon the dispersal and egg-laying behavior of its *S. noctilio* host. The presence of nematodes apparently does not affect the flight of *S. noctilio* females appreciably.

The primary damage inflicted by *B. siricidicola* is the sterilization of *S. noctilio* females. As *S. noctilio* establishes in new areas, this nematode can be easily mass-reared in the laboratory and introduced by inoculation into *S. noctilio*-infested trees. The strain of nematode that has been proposed for release in the United States is known as the Kamona strain. It has been selected for high effectiveness against *S. noctilio*. When these nematodes are mass-reared, they are maintained on their fungus food source, *A. areolatum*. The nematodes that will be released in the United States are reared on a strain of *A. areolatum* that was introduced to North America with *S. noctilio*.

2. Application Method Using *Beddingia siricidicola*

Sirex noctilio is attracted to trees that are stressed. Trap trees can be created by artificially stressing a tree to attract *S. noctilio*. Trap trees are an integral part of all successful *S. noctilio* biological control programs and can also be used for early detection. Trap trees should be located in susceptible, unthinned stands of pine, in sites subjected to drought/moisture stress, or in stands that have been damaged by other pests and disease.

Trap trees are prepared by trimming the lower branches to make the tree base accessible. Herbicide (dicamba or triclopyr) is delivered slowly into the sapwood near the base of the tree approximately 1/2 to 1 meter from the ground by using an injector, drilling holes, or by frilling.² The recommended application rate for program herbicides is 2 milliliters of undiluted herbicide applied every 5 centimeters (cm) around the tree. The herbicide will be applied to the tree in early summer so that the tree is stressed at the time of adult *S. noctilio* emergence (USDA, APHIS, 2006b).

At the end of the summer or early fall, after the trap tree has been infested with *S. noctilio*, these trees will be felled and trimmed of branches on the top side to facilitate the ease of inoculation with *B. siricidicola*. The felled tree should be raised off the ground to prevent attack by wood-decay fungi. The cut ends should be coated with a timber sealant to prevent moisture loss and invasion of fungi.

Inoculation holes approximately 10-mm deep are made using a rebound hammer fitted with a wood punch. The holes are regularly spaced along the length of the log in the following pattern, which is adjusted for the size of the log:

- Log diameter greater than 15 cm: 2 rows of staggered holes spaced 30 cm apart along the log approximately in the “10 o’clock to 2 o’clock” position.

² Frilling inflicts a single line of overlapping downward axe cuts around the bole (stem) leaving a frill.

- Log diameter less than 15 cm: 1 row of holes 30 cm apart along the top of the log down to about 5-cm diameter.
- Deep fissured bark at the base of the tree: not inoculated since *S. noctilio* is less likely to inhabit this part of the tree.

A nematode suspension, formulated in polyacrylamide gel, is poured into a gel dispenser which can simply consist of a “plastic mustard bottle” with a long, tapered nozzle. Within this suspension, a small amount of the fungus, *A. areolatum*, may be present as a result of the nematode rearing culture. However, because a strain of the fungus isolated from *S. noctilio*-infested logs in New York is used for mass-rearing of *B. siricidicola*, there is no concern of introducing a strain of *A. areolatum* that is not already present. The tip of the gel dispenser is inserted into the base of the inoculation hole and withdrawn as the nematode suspension is squeezed into the hole. Once filled, the gel is pressed into the hole with a finger or thumb. This latter step is critical as it ensures close contact between the nematode suspension and the wood fibers inside the hole. This allows the nematode to move into the wood easily. Inoculation during hot or rainy days should be avoided. Warm temperatures may dry out the gel and the nematodes may then dry out before moving into the wood. Rain may wash the gel out of the inoculation holes.

The use of biological control to combat *S. noctilio* began in Australia when the nematode *Deladenus siricidicola* Bedding (Nematoda: Neotylenchidae) and hymenopterous (belonging to the insect order Hymenoptera) parasitoids were imported and introduced. *Deladenus siricidicola*, which was later placed in the new genus *Beddingia*, is the biological control agent of choice against *S. noctilio* because of its effectiveness, biological characteristics, and ease of production in the laboratory. There are other biological control agents available for the management of *S. noctilio*, including several species of parasitic wasps, including *Ibalia leucospoides*, *Megarhyssa nortoni*, and *Rhyssa persuasoria*; however, the most widespread and efficacious is *B. siricidicola* (Bedding and Iede, 2005). Currently, *B. siricidicola* is utilized in the management of *S. noctilio* in Argentina, Australia, Brazil, Chile, New Zealand, South Africa, and Uruguay, saving millions of dollars in pine timber (Bedding and Iede, 2005).

D. Quarantine and Biological Control (Preferred Alternative)

Under this alternative, APHIS would implement a combination of alternatives B (quarantine) and C (biological control) to contain the artificial and natural spread of *S. noctilio* to other States. Combining these two alternatives addresses the artificial spread of *S. noctilio* by means of a

regulatory quarantine and reduces the natural spread of populations of *S. noctilio* by releasing parasitic nematodes (*B. siricidicola*).

Table 1. Summary of Issues and Impacts for the Four Alternatives.

Issue	No Action	Quarantine Only	Biological Control Only	Quarantine and Biological Control
Pine Resources	Will not protect <i>Pinus</i> species in the United States.	Will slow the human-assisted (artificial) spread of <i>S. noctilio</i> , offering some protection to <i>Pinus</i> species but does not control natural spread of <i>S. noctilio</i> .	Will slow the natural spread of <i>S. noctilio</i> , offering some protection to <i>Pinus</i> species but does not slow human-assisted (artificial) spread of <i>S. noctilio</i> .	Will slow natural and human-assisted spread of <i>S. noctilio</i> , thus offering the most protection to <i>Pinus</i> species in the United States.
Native Siricid and Nontarget Species	Would not affect native siricid species but could adversely affect plant and animal species, dependent on pine habitats.	Will have no effect on native siricids. Methyl bromide would have minimal environmental effect.	Biological control agent may attack some native siricid species. Nontarget exposure to herbicides is minimal.	Quarantine will have no effect on native siricids. Methyl bromide would have minimal environmental effect. Biological control agent may attack some native siricid species. Nontarget exposure to herbicides is minimal.
Threatened and Endangered (T&E) Species	May result in adverse effects to Kirtland's warbler and other T&E species, dependent on <i>Pinus</i> species.	Will have no effect on T&E species.	Biological control is not likely to adversely affect T&E species.	Combining quarantine and biological control is not likely to adversely affect T&E species.

IV. Affected Environment

A. The North American Siricid Complex, Their Fungal Food Sources, and the Nematodes That Infect Them

Native siricid woodwasps occur in the eastern and western United States (table 2). Three genera make up a group of 18 known species that attack coniferous (cone-bearing) trees. These genera are *Sirex*, *Urocerus*, and *Xeris*. Two of the genera, *Sirex* and *Urocerus*, have a symbiotic relationship (a mutually beneficial relationship) with the fungal species *Amylostereum chailletii* (*A. chailletii*). The genus *Xeris* has no known fungal symbiont.

Table 2. Native Species of Siricidae Reported from North America and Their Fungal Symbionts

Genus	Species	Subspecies	Region of N. America	Host Type	Attacks <i>Pinus</i> ?	Fungal Symbiont
<i>Sirex</i>	<i>areolatus</i>		East, West	C ^a	Yes	<i>A. chailletii</i>
<i>Sirex</i>	<i>behrensii</i>		West	C	Yes	???
<i>Sirex</i>	<i>cyaneus</i>		East, West	C	Yes	<i>A. chailletii</i>
<i>Sirex</i>	<i>edwardsii</i>		East	C	Yes	<i>A. chailletii</i>
<i>Sirex</i>	<i>juvencus</i>	<i>californicus</i>	West	C	Yes	<i>A. chailletii</i>
<i>Sirex</i>	<i>juvencus</i>	<i>juvencus</i>	East, West	C	Yes	<i>A. chailletii</i>
<i>Sirex</i>	<i>longicauda</i>		West	C	Yes	<i>A. chailletii</i>
<i>Sirex</i>	<i>nigricornis</i>		East	C	Yes	<i>A. chailletii</i>
<i>Urocerus</i>	<i>albicornis</i>		East, West	C	Yes	<i>A. chailletii</i>
<i>Urocerus</i>	<i>californicus</i>		West	C	Yes	<i>A. chailletii</i>
<i>Urocerus</i>	<i>cressoni</i>		East	C	Yes	???
<i>Urocerus</i>	<i>gigas</i>	<i>flavicornis</i>	East, West	C	Yes	<i>A. chailletii</i>
<i>Urocerus</i>	<i>taxodii</i>		East	C	No	???
<i>Xeris</i>	<i>morrisonii</i>	<i>indecisus</i>	West	C	Yes	none
<i>Xeris</i>	<i>morrisonii</i>	<i>morrisonii</i>	West	C	No	none
<i>Xeris</i>	<i>spectrum</i>	<i>spectrum</i>	East, West	C	Yes	none
<i>Xeris</i>	<i>spectrum</i>	<i>townesi</i>	West	C	Yes	none
<i>Xeris</i>	<i>tarsalis</i>		West	C	No	none
<i>Eriotremex</i>	<i>formosanus</i>		SE U.S.	D	No	???
<i>Tremex</i>	<i>columba</i>		East, West	D	No	<i>Cerrena unicolor</i>

^aC=coniferous ^bD=deciduous

Source: Morgan, 1968; Baker 1972; Furniss and Carolin, 1977; Talbot, 1977; Bedding and Akhurst, 1978; Krombein et al., 1979; Smith and Schiff, 2002; Slippers et al., 2003. (From Williams, 2007).

Other *Amylostereum* species have not been identified in the United States (although *A. areolatum* has now been introduced with *S. noctilio*). The widespread distribution of *A. chailletii* in North America is related to the species' ability to be spread by native woodwasps, as well as by windborne spores (Slippers et al., 2003). *Amylostereum areolatum* is dispersed exclusively by siricid woodwasps and may be present in areas of the United States where *S. noctilio* has been identified.

The proposed biological control organism, *B. siricidicola*, belongs to a group of nematodes within the genus *Deladenus* which has approximately 23 species. Seven of the species were reclassified into the new genus *Beddingia* because they occur in two forms, a mycetophagous form and an entomophagous form. Of the seven *Beddingia* species, five are known to occur with siricid woodwasps in North America (Chitambar, 1991; Bedding and Akhurst, 1978). Of these five nematode species, *B. nevexii* can occur in both *Sirex* and *Urocerus* species, while *B. canii* and *B. proximus* have been identified in only *Sirex* woodwasps. *Beddingia wilsonii* and *B. rudyi*, have been isolated from woodwasps from the genus *Urocerus* (Bedding and Akhurst, 1978). While the nematodes do not

appear to be specific to an insect host, they are specific in regard to the *Amylostereum* fungus species on which they rely for food. *Beddingia wilsonii* is the only nematode that will feed on both *A. chailletii* and *A. areolatum*, while *B. siricidicola* feeds exclusively on *A. areolatum*. The remaining *Beddingia* species feed on *A. chailletii* (Bedding and Akhurst, 1978).

B. Pine Resources of North America

Pine ecosystems are diverse and widespread throughout the United States. These ecosystems provide a substantial economic value and provide numerous environmental benefits. Softwood production in the United States is a multibillion dollar industry that provides numerous commodities. In the southern States where pine production typically occurs in large, even-aged, managed stands, the combined value of logs and bolts, lumber, veneer, and pulpwood production is greater than \$8 billion per year (USDA, APHIS, 2007a). The value of the same commodities in the western United States is greater than \$10 billion. Other commodities, such as Christmas tree production, result in a revenue of approximately \$2 million in the northeastern and north-central United States (USDA, APHIS, 2007a).

In addition to the large economic benefits of pine resources, they also provide valuable and unique habitat to a variety of flora and fauna throughout the United States. Multiple diverse pine habitats exist across the United States with the result being numerous plant and animal species that depend on these types of ecosystems.

As an example in the southeastern United States, longleaf pine (*P. palustris*) ecosystems occupy nine States where stands may occur on dry sites as savannahs, or in wet areas as flatwoods. These habitats occupy less than 3 percent of their previous extent, and are considered one of the rarest ecosystems in the United States (Brockaway et al., 2005). These habitats have reported high biodiversity with plant diversities reaching 140 vascular plant species in a 1,000-square meter area. The high biodiversity and rare occurrence of this type of habitat results in many species that are considered rare and, in some cases, are listed as threatened or endangered. This is exemplified by the number and diversity of threatened and endangered species that currently utilize longleaf habitat. Some of the listed species that are dependent on the longleaf pine ecosystem are the red cockaded woodpecker (*Picoides borealis*), gopher tortoise (*Gopherus polyphemus*), and flatwoods salamander (*Ambystoma cingulatum*).

In the southwestern United States, ponderosa pine (*P. ponderosa*) stands provide habitat for over 250 vertebrate species, including the federally

listed Mexican spotted owl (*Strix occidentalis lucida*) (Allen et al., 2002). The high diversity in vertebrate species is related to the high relative plant diversity that has been identified in ponderosa pine habitats when compared to other adjacent habitats (Stohlgren et al., 2001).

On the California coast, three relict stands of Monterey pine (*P. radiata*) provide a unique ecological habitat for a variety of plants and wildlife, including the federally protected Yadon's piperia (*Piperia yadonii*) (USFWS, 2004) and the Santa Cruz long-toed salamander (*Ambystoma macrodactylum croceum*). In addition, these trees serve as a source of genetic material for *P. radiata* plantations in the Southern Hemisphere (Ciesla, 2003).

In the north-central and northeastern United States, pine stands intermixed with hardwoods and native grasses form barrens that provide unique plant diversity and habitat for wildlife, including rare and endangered species. In Michigan, pine barrens of jack pine (*P. banksiana*) provide habitat for the endangered Kirtland's warbler (*Dendroica kirtlandii*), and in Wisconsin these areas provide habitat for the endangered Karner blue butterfly (*Lycaeides melissa samuelis*) (Houseman and Anderson, 2002; USFWS, 2003a). Rare and uncommon flora and fauna also exist in the pine barren habitats of New England with State-listed species, and in important areas, such as the barrens in New Jersey, that are considered globally imperiled. While barren habitats typically occur on poor soils, the plants that grow in these areas tend to be unique and support uncommon and rare wildlife.

The above mentioned habitats are not inclusive of all pine habitats in the United States but represent ecosystems that contain pine species known to be susceptible to *S. noctilio*. In addition, these ecosystems occur in plant hardiness zones that are known to support *S. noctilio*, with current estimates suggesting that survivorship for the woodwasp is likely within all pine-growing regions in the United States (Carnegie et al., 2006; USDA, APHIS, 2007a).

V. Environmental Consequences

A. No Action

Based on its ability to colonize North American pines, as well as other factors, *S. noctilio* has been rated as a high-risk species for North American forests (USDA, APHIS, 2007a). Environmental impacts related to the implementation of the no action alternative could be widespread throughout the United States. Carnegie et. al. (2006) used a climatic model to assess the potential areas of survivorship for *S. noctilio* in North America, and was able to determine a 75-percent likelihood of survival in

all areas where *Pinus* occurs east of the Mississippi River, from Canada to the southeastern United States. West of the Mississippi River (in Arkansas, Missouri, and the Pacific Northwest) *Pinus* forests also have a 75-percent likelihood of *S. noctilio* survivorship. Other *Pinus* areas west of the Mississippi have a 50-percent likelihood of survival. These predictions are consistent with other information that *S. noctilio* currently occurs in multiple USDA plant hardiness zones that are found in the United States (USDA, APHIS, 2007a).

The rate of spread of *S. noctilio* is difficult to quantify since native and nonnative *Pinus* in the United States occur as intermixed species in some habitats where the spread may be slower, versus large *Pinus* plantations in the southeastern United States where the spread could be much quicker. Additionally, *S. noctilio* may be introduced into areas by means other than natural dispersion, which further confounds the possible rate of spread. In plantation environments in Africa and Australia, *S. noctilio* has been shown to spread at a rate of 19 to 30 miles per year (Tribe and Cillie, 2004; Carnegie et al., 2006). For the natural rate of spread in the United States, estimates have ranged from 18 to 25 miles per year if no control program is implemented (USFS, 2006; USDA, APHIS, 2007a).

In the Southern Hemisphere, where *S. noctilio* has been demonstrated to be a significant pest of pine, *S. noctilio* is an invasive insect attacking monocultures of nonnative pines. In areas where it has been inadvertently introduced, it has caused up to 80 percent tree mortality in plantations planted with North American pine species. These high mortality rates suggest that *S. noctilio* has the potential to cause extensive damage and mortality in commercially mature timber and future-growing stock timber in the United States. However, North American forests may vary in their susceptibility to *S. noctilio* because many noncommercial forests that contain pine are more species-diverse and have established populations of predators that will likely attack *S. noctilio* (Dodds et al., 2007).

In January 2006, the U.S. Forest Service issued an economic analysis for the potential impact of *S. noctilio* in the United States. This economic analysis assumed that *S. noctilio* would spread from the current New York infestation at 25 miles per year and take 55 years to infest the entire southeastern United States. At a 10 percent mortality threshold, in the South there would be 244-million square feet lost, valued at \$1.9 billion dollars; in the total infested area, 360-million square feet and \$2.9 billion would be lost. If mortality reaches 50 percent, losses would reach 1.4-billion square feet and \$11 billion lost in the South, and 2.1-billion square feet and \$17 billion in the infested area (USFS, 2006). The continued spread of *S. noctilio* could cause widespread mortality of *Pinus* species resulting in negative economic and environmental impacts.

Environmental impacts related to the no action alternative relate to the loss of *Pinus*-associated habitats. While these habitat types vary across the United States, effects to pine-dependent nontarget species would be expected, including threatened and endangered species. For example, listed bird species, such as the red cockaded woodpecker (*P. borealis*), that depend on mature longleaf pine stands in the southern United States, and Kirtland's warbler (*D. kirtlandii*), that depends on jack pine in Michigan, are two bird species that would be directly impacted by the loss of pine habitat (USFWS, 2003b). In the western United States, rare native Monterey pine (*P. radiata*) habitat and dependent flora and fauna could also be impacted if *S. noctilio* is able to establish reproducing populations. In addition to federally listed species, numerous other pine-dependent species would be at risk due to the loss of pine habitat. The loss of *Pinus* species in these habitats would alter the plant succession process in pine habitats, and would allow for other undesirable plant species to become established, altering plant diversity.

B. Quarantine Only

The quarantine only alternative is designed to stop the spread of *S. noctilio* through human actions. Quarantine is widely accepted as an effective method to prevent the artificial or human-assisted spread of pests; however, it does not address the natural spread of *S. noctilio*. Quarantine requirements would prohibit the transport of any wood or wood product that can harbor the pest unless it has been treated in such a way as to eliminate any living *S. noctilio*. The environmental impacts of any required treatments (e.g., heat treatment) are expected to be minimal as the treatments are generally considered to be safe and routine measures when conducted according to general practices. Insecticides, other than methyl bromide, would not be used as they would be ineffective when used against larval *S. noctilio*.

One of the treatments that could be used under the quarantine alternative is fumigation with methyl bromide. Methyl bromide is a substance classified by the U.S. Environmental Protection Agency under the Clean Air Act as a Class I ozone-depleting chemical. Ozone depletion is the primary environmental concern related to fumigations with methyl bromide.

The contribution of the worldwide usage of methyl bromide in 1996 (63,960 metric tons) to atmospheric ozone depletion was less than 1 percent (UNEP, 1998). Global methyl bromide use was approximately 72,000 metric tons or 143,000,000 pounds in 2006 (EPA, 2006). The amount of methyl bromide required to treat articles from *S. noctilio* quarantine zones would be applied at a maximum dosage rate of 4 pounds per 1,000 cubic feet. Based on the maximum application rate for

S. noctilio treatment and the total quantity of methyl bromide used in 2006, approximately 400-million board feet would need to be treated in a year to reach just 1/10 of a percent of the global total stratospheric ozone depletion. This level of treatment is impractical and treatment with methyl bromide is expected to be minimal for this program; however, it demonstrates the very minor contribution of program use of methyl bromide to its total use. Additionally, the fumigation of certain commodities, such as wood packing products, releases 11 to 21 percent less methyl bromide than other fumigations, further reducing the atmospheric contribution of methyl bromide (USDA, APHIS, 2007c).

The expected use of methyl bromide in the fumigation of articles from pine trees for the control of *S. noctilio* would be minimal and is well below levels that could contribute substantially to ozone depletion. This conclusion is consistent with a previous environmental impact statement (EIS) regarding the importation of unmanufactured wood articles from Mexico (USDA, APHIS, 2002). The EIS found that the impact of methyl bromide from routine commodity treatments on ozone depletion was not significant (USDA, APHIS, 2002). Any other potential environmental impacts would not be expected due to the carefully controlled manner in which methyl bromide is used, and the environmental fate which reduces exposure and risk to human health and the environment.

C. Biological Control Only

1. Nontarget Effects

Nontarget effects related to the release of *B. siricidicola* and *A. areolatum* are not expected due to lack of toxicity to nontarget organisms and low exposure for most of them. The method of application and natural dispersal for the nematode/fungal complex eliminates potential exposure to all organisms that do not utilize trees that have been infected with *B. siricidicola*. In cases where exposure could occur, effects data for other entomopathogenic (causing diseases to insects) nematodes have demonstrated low toxicity in laboratory and field experiments to birds, mammals, and amphibians (Georgis et al., 1991; Bathon, 1996; Ehlers, 1996). Significant impacts to nontarget arthropods are not expected with the introduction of *B. siricidicola*. Impacts to soil-borne arthropods are not expected due to application and natural dispersal methods of the nematode. Nematode-related impacts to trees would also not be expected since the nematode feeds exclusively on *A. areolatum*.

The primary concern regarding release of *B. siricidicola* into the environment is its impact on native siricid species. The primary factor affecting exposure of a siricid species to *B. siricidicola* is its fungal symbiont. The two most common siricid symbionts worldwide are *A. areolatum*, which is native to Europe and Japan, and *A. chailletii*, which is native to North America, Europe, and Japan. *Beddingia siricidicola*

feeds only on *A. areolatum* (Bedding and Akhurst, 1978). Thus, species that feed on *A. chailletii*, including most of those in North America, will not encounter *B. siricidicola* (Williams, 2007).

Only a few North American siricid species may potentially feed on wood colonized by *A. areolatum*. In particular, *Xeris* species are not symbiotic with any fungus species but, instead, lay eggs in trees already attacked by another siricid species and inoculated with its fungus. The subspecies *X. morrisoni indecisus*, *X. spectrum spectrum*, and *X. spectrum townesi* would be exposed to *B. siricidicola* as *A. areolatum* spreads across North America with the dispersal of *S. noctilio*, but only if these woodwasps attack trees already infested with *S. noctilio* (Williams, 2007). It is unknown whether native *Xeris* species will be able to use *A. areolatum* or will be attracted to it (Williams, 2007).

It is difficult to predict the impact of *B. siricidicola* parasitism on *Xeris* species in North America. Those populations developing on *A. chailletii* will not be affected. The cosmopolitan subspecies *X. spectrum spectrum* is parasitized by *B. siricidicola* in Eurasia, yet it apparently remains a common insect there. However, the natural control situation in Eurasia and a classical biological control program in North America may not be entirely comparable. The Kamona strain, (collected from the Kamona forest in Tasmania, Australia) of *B. siricidicola* has been selected for high pathogenicity on *S. noctilio*. Its impact on nontarget siricid species may be greater than that of naturally occurring *B. siricidicola* strains in Eurasia. However, even the most successful biological control programs have not driven their target to extinction. Thus, it is unlikely that impacts on *Xeris* species would be great (Williams, 2007).

In addition, two *Sirex* species, *S. juvencus juvencus* and *S. cyaneus*, may be attacked by *B. siricidicola*. Because these species are capable of symbiosis with both *A. chailletii* and *A. areolatum*, it is possible that they may adapt in North America to using the introduced *A. areolatum* as a resource, exposing them to attack by *B. siricidicola* (Williams, 2007).

Once *B. siricidicola* is released into the environment and if other insect species were to be attacked by it, the resulting effects could be environmental impacts that may not be easily reversed. Biological control agents, such as *B. siricidicola*, generally spread even without the agency of man. In principle, therefore, release of these nematodes, even at one site, must be considered equivalent to release over the entire area in which potential hosts occur and in which the climate is suitable for reproduction and survival. Post-release evaluations of *B. siricidicola* populations and their effects on *S. noctilio* will be conducted for several years by APHIS researchers after initial release.

2. Herbicide Effects

Successful implementation of the *S. noctilio* control program is based on the ability to detect insect damage and establish populations of nematodes in areas where they have the greatest chance of parasitizing *S. noctilio*. This can be accomplished through the establishment of trap trees which are chemically stressed trees used to attract *S. noctilio*. An herbicide is used to stress individual trees at optimal adult emergence times for *S. noctilio*, and to minimize attacks from bark beetles which may reduce suitability for *S. noctilio* colonization (USDA, APHIS, 2006a). The herbicides recommended for use in the *S. noctilio* control program are triclopyr and dicamba. Both herbicides have multiple formulations that are used in a variety of agricultural and nonagricultural applications. Due to the low toxicity to nontarget organisms and the application method being implemented in the *S. noctilio* program, the risk to nontarget organisms is considered minimal for both herbicides.

a. Triclopyr and Dicamba Toxicity

Triclopyr is a broadleaf and woody plant herbicide used in a variety of applications and sold in formulations as a triethylamine salt (TEA) or a butoxyethyl ester (BEE). The formulation proposed for use in this program contains the triethylamine salt. Dicamba is the other herbicide that may be used. It also has a wide variety of uses with the formulated material containing the dimethylamine salt (DMA). In the environment, DMA degrades to dicamba acid and the TEA salt degrades to form triclopyr acid. Toxicity for both herbicides varies depending on the test organism, as well as the formulation being tested (EPA, 2005; EPA, 1998). Both herbicides have low acute and chronic toxicity to nontarget organisms (appendix B).

b. Triclopyr and Dicamba Exposure and Risk

Exposure from either herbicide, triclopyr or dicamba, will be extremely low for all nontarget organisms under labeled use due to the method of application in the *S. noctilio* control program. Application techniques for either herbicide involve direct injection into the sapwood of the tree, 20 cm above the ground, as a solution of undiluted formulated material every 5 cm around the trunk (USDA, APHIS, 2006b). Because the material is being injected, the material will not enter aquatic environments via drift or runoff. The fact that herbicide residues will not enter aquatic systems under labeled use mitigates aquatic exposure and subsequent risk to aquatic organisms.

While the material will be applied in terrestrial environments, no drift to off-site plant material is expected and, therefore, exposure and risk is minimal to nontarget organisms such as mammals, birds, terrestrial invertebrates, and plants. There is the potential for exposure to

invertebrates that may colonize treated trees; however, the number of trap trees in any given area will be considerably lower than the available trees per acre, and the application rates are low, thus, further reducing exposure. In cases where colonizing insects may be exposed to herbicide residues in the tree trunk, risk is expected to be minimal based on the available terrestrial invertebrate toxicity data, and the lack of reports that suggest either herbicide has insecticidal properties.

Residues are not expected to accumulate in any terrestrial invertebrates that may inhabit trap trees due to the environmental fate of both herbicides. Invertebrates from injected trees that could serve as prey items would pose low risk to nontarget organisms since neither herbicide accumulates in tissue, both are applied at low concentrations, and both have dietary toxicity values to birds and mammals that demonstrate low acute and chronic toxicity. Logging debris from injected trees could contain low levels of both herbicides and associated metabolites; however, due to the environmental fate and low numbers of trees being injected, the risk to nontarget organisms would be minimal.

3. Human Health Impacts

No human health impacts are expected from the release of *B. siricidicola* and *A. areolatum* to control *S. noctilio*. In areas of the world where *B. siricidicola* and *A. areolatum* have been introduced, no known human health related effects have been observed. The low risk to human health is related to a lack of toxicity and low exposure due to the method of application. While the vertebrate toxicity data regarding *Beddingia/Deladenus* is unknown, other nematode species used as entomopathogenic biological control agents have been shown to have no toxicity or risk to vertebrates (Bathon, 1996; Boemare, 1996; Ehlers and Hokannen, 1996). The lack of toxicity to vertebrates is a function of host specificity for entomopathogenic nematodes and the inability of nematodes to survive in vertebrate hosts.

In addition to the lack of toxicity for the nematode, exposure to *B. siricidicola* and *A. areolatum* is not expected due to the method of application. Since the nematode and fungus are inoculated into each tree during the preparation of trap trees by applicators, or by the wasp through natural dispersion, exposure would be highest for applicators and extremely low for other individuals. Standard protective equipment used during applications and the lack of toxicity would preclude any risk to applicators.

D. Quarantine and Biological Control

Environmental impacts from this alternative will consist of a combination of the impacts anticipated from alternatives B and C. The impacts are expected to be additive in nature. These impacts have been discussed

above. This alternative is the one most likely to result in successful control and containment of *S. noctilio* because it combines population control methodologies that should limit the natural spread of the pest, and quarantine methodologies that are designed to eliminate the artificial spread of the pest due to human factors.

E. Cumulative Effects

“Cumulative impacts are defined as the impact on the environment which results from the incremental impact of the action when added to other past, present and reasonably foreseeable future actions regardless of what agencies or person undertakes such other actions” (40 CFR 1508.7).

Direct cumulative impacts attributed to the management of *S. noctilio* could be related to the use of the herbicides triclopyr or dicamba, release of *B. siricidicola*, the creation of trap trees, or the possible use of methyl bromide. The use of an herbicide (triclopyr or dicamba) in the program will have no negative cumulative impacts in areas where the program is implemented due to the unique application method which minimizes exposure and risk to human health and the environment. The use of *B. siricidicola* is unique to *S. noctilio* woodwasp control, and the nematode is not used to control other forestry or agronomic pests. Risks to human health and the environment from the release of *B. siricidicola* are low due to the method of application and the lack of toxicity that has been identified. The establishment of trap trees for inoculation will create additional stressed pine trees in areas where stressed trees already may exist; however, the number of trees is small relative to the total number of trees per acre. If methyl bromide is used as part of a *S. noctilio* management program, the direct cumulative impacts to ozone depletion are expected to be inconsequential due to its extremely minor contribution, as described previously in this EA. The limited quantities of methyl bromide needed to treat the pine tree articles under quarantine are within the negligible treatments anticipated under previous cumulative analysis (USDA, APHIS, 2002).

Without successful implementation of a control and containment program, many more trees would likely be killed by *S. noctilio* infestation. These additional, newly-stressed trees could also lead to secondary forest pest outbreaks which could result in negative cumulative impacts on pine resources. Loss of additional pine acreage over time, due to *S. noctilio* infestation, would result in negative economic impacts and alter native pine ecosystems.

F. Endangered Species Act

Section 7 of the Endangered Species Act (ESA) and its implementing regulations require Federal agencies to ensure that their actions are not likely to jeopardize the continued existence of federally listed threatened and endangered species, or result in the destruction or adverse modification of critical habitat.

APHIS prepared a biological assessment (BA) that considered the impact of the release of *B. siricidicola* on federally listed insects in the United States. The host specificity of the nematode precludes any impact on federally listed threatened or endangered insects. The only known hosts of *B. siricidicola*, in addition to *S. noctilio*, are certain siricid species. There are no listed siricid species in the United States. Therefore, there will be no effect on any listed or proposed insect species and no adverse modification of designated critical habitat in the continental United States. APHIS prepared a BA that documented this no effect determination.

APHIS also prepared BAs that analyzed impacts of the creation of trap trees and inoculation of those trees with *B. siricidicola* on threatened and endangered species and designated critical habitat in New York, Pennsylvania, Vermont, and Michigan. APHIS determined that this activity would have no effect on threatened and endangered species or critical habitat in Vermont. In New York, Pennsylvania, and Michigan, APHIS determined that these activities may affect, but are not likely to adversely affect, listed species in those States with the implementation of certain protection measures. APHIS submitted the BAs to U.S. Fish and Wildlife Service (FWS) field offices in New York, Pennsylvania, and Michigan, requesting concurrence with these determinations; FWS has concurred. If *S. noctilio* spreads into new States and *B. siricidicola* releases are planned, APHIS will consult with the appropriate FWS field office within the State before proceeding with program activities.

Implementation of a quarantine will have no effect on federally listed species or their habitats because heat and methyl bromide treatments are conducted in facilities that do not occur in habitats of listed species. APHIS is not requiring insecticide treatments of nursery stock or other *Pinus* species items that could potentially expose listed species or habitat to insecticides.

Positive impacts are expected from the control program (both quarantine and biological control) for those listed species that depend on native pine habitats. A majority of the sensitive pine habitats in the United States contain pine species that have been shown to be susceptible to *S. noctilio* (Carnegie et al., 2006). These habitats contain a large number of federally

listed species, as well as other rare and uncommon species that, in the future, could require Federal protection if impacted by *S. noctilio*.

VI. Other Issues

Consistent with Executive Order (EO) 12898, “Federal Actions to Address Environmental Justice in Minority Populations and Low-income Populations,” APHIS considered the potential for disproportionately high and adverse human health and environmental effects on any minority populations and low-income populations. The environmental and human health effects from all of the alternatives are minimal and are not expected to have disproportionate adverse effects to any minority or low-income populations.

Consistent with EO 13045, “Protection of Children from Environmental Health Risks and Safety Risks,” APHIS considered the potential for disproportionately high and adverse environmental health and safety risks to children. No circumstances that would trigger the need for special environmental reviews are involved in implementing any of the alternatives. The program applicators will ensure that the general public is not in or around areas being treated and, therefore, no exposure will occur from the application of any herbicides. Hence, it is expected that no disproportionate effects on children are anticipated as a consequence of implementing any of the alternatives evaluated above.

VII. Agencies, Organizations, and Individuals Consulted

This EA was prepared and reviewed by APHIS. The addresses of participating APHIS units, cooperators, and consultants (as applicable) follow.

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine
Program Support
4700 River Road, Unit 134
Riverdale, MD 20737-1236

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Policy and Program Development
Environmental Services
4700 River Road, Unit 149
Riverdale, MD 20737-1238

U.S. Department of Agriculture
Animal and Plant Health Inspection Service
Plant Protection and Quarantine
4700 River Road, Unit 150
Riverdale, MD 20737-1229

U.S. Department of Interior
U. S. Fish & Wildlife Service
3817 Luker Road
Cortland, NY 13045

U.S. Department of Interior
U.S. Fish & Wildlife Service
315 South Allen Street, Suite 322
State College, PA 16801-4850

U.S. Department of Interior
U.S. Fish & Wildlife Service
East Lansing Field Office
2651 Coolidge Road, Suite 101
East Lansing, MI 48823-6316

VIII. References

- Allen, C.D., Savage, M., Falk, D.A., Suckling, K.F., Swetnam, T.W., Schulke, T., Stacey, P.B., Morgan, P., Hoffman, M., and Klingel, J.T., 2002. Ecological restoration of southwestern ponderosa pine ecosystems: A broad perspective. *Ecological Applications*. 12(5): 1418–1433.
- Baker, W.L., 1972. *Eastern Forest Insects*. USDA Forest Service, Miscellaneous Publication No. 1175.
- Bathon, H., 1996. Impact of entomopathogenic nematodes on non-target hosts. *Biocontrol Science and Technology*. 6:421–434.
- Bedding, R.A., 1972. Biology of *Deladenus siricidicola*, an entomophagous-mycetophagous nematode parasitic in siricid woodwasps. *Nematologica*. 18: 482–493.
- Bedding, R.A., and Akhurst, R.J., 1978. Geographical distribution and host preferences of *Deladenus* species (Nematoda: Neotylenchidae) parasitic in siricid woodwasps and associated hymenopterous parasitoids. *Nematologica*. 24:286–294.
- Bedding, R.A., and Iede, E.T., 2005. Application of *Beddingia siricidicola* for Sirex Woodwasp Control. *Nematodes as Biocontrol Agents*. CPC Publishing. Cambridge, pp 385–399.
- Boemare, N., Laumond, C., and Mauleon, H., 1996. The entomopathogenic nematode-bacterium complex: biology, life cycle and vertebrate safety. *Biocontrol Science and Technology*. 6:333–345.
- Brockway, D.G., Outcalt, K.W., Tomczak, D.J., and Johnson, E.E., 2005. Restoration of Longleaf Pine Ecosystems. Gen. Tech. Rep. SRS-83. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 34 p.
- Carnegie, A.J., Matsuki M., Haugen, D.A, Hurley, B.P., Ahumada, R., Kklasmer, P., Sun, J. and Iede, ET., 2006. Predicting the potential distribution of *Sirex noctilio* (Hymenoptera: Siricidae), a significant exotic pest of *Pinus* plantations. *Annals of Forestry Science*. 63:119–128.
- Chitambar, J., 1991. On the genus *Deladenus* Thorne, 1941 (Nemata: Allantonematidae). Review of the mycetophagous stage. *Revue Nematol*. 14(3): 427–444.
- Ciesla, W.M., 2003. European woodwasp: a potential threat to North America's conifer forests. *Journal of Forestry*. 101(2):18–23.

Dodds, K.J., Cooke, R.R., and Gilmore, D.W., 2007. Silvicultural options to reduce pine susceptibility to attack by a newly detected invasive species, *Sirex noctilio*. Northern Journal of Applied Forestry. 24: 165–167.

Ehlers, R.U., 1996. Current and future use of nematodes in biocontrol: practice and commercial aspects with regard to regulatory policy issues. Biocontrol Science and Technology. 6:303–316.

Ehlers, R.U., and Hokkanen, H.M.T., 1996. Insect biocontrol with non-endemic entomopathogenic nematodes (*Steinernema* and *Heterorhabditis* spp.): conclusions and recommendations of a combined OECD and COST workshop on scientific and regulatory policy issues. Biocontrol Science and Technology. 6:295–302.

EPA—See U.S. Environmental Protection Agency

Furniss, R.L., and Carolin, V.M., 1977. Western Forest Insects. USDA, Forest Service, Miscellaneous Publication No. 1339.

Georgis, R., Kaya, H.K., and Gaugler, R., 1991. Effect of steinernematid and heterorhabditid nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) on nontarget arthropods. Environmental Entomology. 20(3): 815–822.

Houseman, G.R., and Anderson, R.C., 2002. Effects of jack pine plantation management on barrens flora and potential Kirtland's warbler nest habitat. Restoration Ecology. 10(1): 27–36.

Krombein, K.V., Hurd, P.D., Smith, D.R., and Burks, B.D., 1979. Catalog of Hymenoptera North of Mexico, Volume 1. Smithsonian Institution Press, Washington, DC.

Morgan, F.D., 1968. Bionomics of Siricidae. Annual Review of Entomology. 13: 239–256.

Slippers, B., Coutinho, T.A., Wingfield, B.D., and Wingfield, M.J., 2003. A review of the genus *Amylostereum* and its association with woodwasps. South African Journal of Science. 99: 70–74.

Smith, D.R. and Schiff, N.M., 2002. A review of siricid woodwasps and their ibaliid parasitoids in the eastern United States, with emphasis on the mid-Atlantic region. Proceedings of the Entomological Society of Washington. 104: 174–194.

Stohlgren, T.J., Otsuki, Y., Villa, C.A., Lee, M., and Belnap, J., 2001. Patterns of plant invasions: a case example in native species hotspots and rare habitats. *Biological Invasions*. 3:37–50.

Talbot, P.H.B., 1977. The *Sirex-Amylostereum-Pinus* association. *Annual Review of Phytopathology*. 15: 41-54.

Tribe, G.D., and Cillie, J.J., 2004. The spread of *Sirex noctilio* Fabricius (Hymenoptera: Siricidae) in South African pine plantations and the introduction and establishment of its biological control agents. *African Entomology*. 12: 9–17.

UNEP—See United Nations Environment Programme.

United Nations Environment Programme, 1998. Environmental effects of ozone depletion: 1998 assessment. Nov. 1998.

USDA—See U.S. Department of Agriculture

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2007a. Organism Pest Risk Analysis: Risks to the Conterminus United States Associated with the Woodwasp, *Sirex noctilio* Fabricius, and the Symbiotic Fungus, *Amylostereum areolatum* (Fries: Fries) Boidin (Revision 1). 40 pp.

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2007b. Proposed Program for Management of the Woodwasp *Sirex noctilio* Fabricius (Hymenoptera: Siricidae), Environmental Assessment, June 2007. [April 16, 2008.] Available: http://www.aphis.usda.gov/plant_health/ea/sirex.shtml

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2007c. Importation of Solid Wood Packing Material. Draft Supplemental to the Final Environmental Impact Statement—February 2007. Riverdale, MD.

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2006a. *Sirex* Science Advisory Panel Report, January 9-10, 2006. Annapolis, MD. 65 pp.

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2006b. Section 6 Control Procedures. Draft Version 2.0. 18 pp.

U.S. Department of Agriculture, Animal and Plant Health Inspection Service, 2002. Rule for the Importation of Unmanufactured Wood Articles From Mexico, With Consideration for Cumulative Impact of Methyl Bromide Use. Final Environmental Impact Statement, September, 2002. USDA, APHIS, Riverdale, MD.

U.S. Environmental Protection Agency, 2006. Methyl Bromide Questions and Answers. March 14, 2008. [April 15, 2008.] Available: <http://www.epa.gov/ozone/mbr/qa.html>.

U.S. Environmental Protection Agency, Office of Pesticide Programs, 2005. Environmental fate and ecological risk assessment for the reregistration of dicamba and dicamba sodium, potassium, diglycoamine, dimethylamine and isopropylamine salts. 220 pp.

U.S. Environmental Protection Agency, Prevention, Pesticides and Toxic Substances, 1998. Reregistration Eligibility Decision (RED): Triclopyr. Washington, DC. 269 pp. [April 15, 2008.] Available: <http://www.epa.gov/oppsrrd1/REDs/2710red.pdf>.

U.S. Fish and Wildlife Service, 2004. Recovery plan for five plants from Monterey County, California. U.S. Fish and Wildlife Service, Portland, Oregon. 159 pp.

U.S. Fish and Wildlife Service, 2003a. Final recovery plan for the Karner Blue Butterfly (*Lycaeides melissa samuelis*). U.S. Fish and Wildlife Service, Fort Snelling, Minnesota. 273 pp.

U.S. Fish and Wildlife Service, 2003b. Recovery plan for the red-cockaded woodpecker (*Picoides borealis*): second revision. U.S. Fish and Wildlife Service, Atlanta, GA. 296 pp.

U.S. Forest Service—Forest Health Protection, 2006. Economic analysis of the potential impact of *Sirex noctilio*, with emphasis on pines in the Southeastern United States. 5 pp.

USFS—See U.S. Forest Service

USFWS—See U.S. Fish and Wildlife Service

Williams, D.W., 2007. Potential non-target effects of *Beddingia siricidicola* when used as a biological control agent of *Sirex noctilio* in North America. White Paper. In: *Sirex* Science Panel Report, December 13-14, 2006, Indianapolis, Indiana. [April 10, 2008.] Available: http://www.aphis.usda.gov/plant_health/plant_pest_info/sirex/downloads/ap12-14-06.pdf

Appendix A. Organism Pest Risk Assessment

**Organism Pest Risk Analysis:
Risks to the Conterminous United States Associated with the
Woodwasp, *Sirex noctilio* Fabricius, and the Symbiotic
Fungus, *Amylostereum areolatum* (Fries: Fries) Boidin.**

May 2007



Paula Klasmer
Instituto Nacional de Tecnología Agropecuaria
www.forestryimages.org

Agency Contact:

Daniel Borchert
USDA-APHIS-PPQ-CPHST-PERAL
1730 Varsity Drive, Suite 300
Raleigh, NC 27606-5202
Phone: 919-855-7545
Fax: 919-855-7599
daniel.m.borchert@aphis.usda.gov

Executive Summary

This pest risk analysis (PRA), was conducted at the request of the United State Department of Agriculture (USDA), Animal Plant Health Inspection Service (APHIS), Plant Protection and Quarantine (PPQ), Emergency and Domestic Programs (EDP) unit, to examine the risks associated with the woodwasp, *Sirex noctilio* Fabricius, and the symbiotic fungus, *Amylostereum areolatum* (Fries: Fries) Boidin.

This risk analysis evaluates the overall risk *S. noctilio* and *A. areolatum* pose to the United States using an APHIS-PPQ organism pest risk assessment template. Because of the obligate relationship between *S. noctilio* and *A. areolatum*, the organisms were analyzed together, with regard to potential risk; however, *S. noctilio* is the primary focus.

Sirex noctilio scored High with regard to Cumulative Risk, Habitat Suitability, Dispersal Potential and Economic Impact. Its Environmental Impact was Medium for the United States, based on the wasp's potential to impact the endangered *Pinus radiata*, and introduction of a non-native nematode species, *Beddingia siricidicola*, for use as a biological control agent on *S. noctilio*. *Sirex noctilio* scored Low with regard to Host Range because it prefers species in the genus *Pinus*. These scores indicate that *S. noctilio* could pose a serious potential economic threat to the U.S. forestry industry.

Climate will most likely not limit the distribution of *S. noctilio* in the United States. Consequently, its projected area of U.S. colonization will depend on the distribution of pines. Pines are found throughout the United States, with the highest concentrations in the south, west, northeast, and north central states, respectively.

The Southern United States has a concentrated and uniform distribution of pine forests, while those in the West, North Central and Northeast are more dispersed. *Pinus taeda* L. is the major planted pine species in the South; this species is a suitable host plant for *S. noctilio*. The estimated annual value of southern softwood logs, pulpwood, timber and veneer is greater than \$8 billion. By using the U.S. Forest Service regional breakdown, the South is the world's largest softwood timber producer, and its output is projected to increase. Moreover, the South's tendency to encounter tropical storms and hurricanes has left many forests damaged, producing host material for *S. noctilio*. With southern pine resources affected by these storms, the economic vitality of this commodity is at risk, as is the tendency for *S. noctilio* introduction is increased.

However, there is uncertainty regarding the rate at which spread will be accomplished, and the degree of impact that *S. noctilio* will have on U.S. pines resources. A source of uncertainty with regard to the spread of *S. noctilio*, relates to the potential competition with native bark beetles, native Siricids and stand vigor. Given the potential consequences of *S. noctilio* introduction in the United States, we recommend that pine resources be protected by regulatory means until more is known about its ability to impact U.S. pines.

Central questions to be addressed by future research on *Sirex noctilio* include:

- Can *S. noctilio* displace indigenous bark beetles and Siricids?

- Will *S. noctilio* cause minimal forest damage in the United States, as it does in its native range of Europe and North Africa, or will it become a major forest pest as observed in Southern Hemisphere countries?
- Will native parasitoids effectively use *S. noctilio* as a host?
- Will *S. noctilio* become like *S. cyaneus* F., e.g. spread by commerce, be nominally invasive, and currently a minor pest (Arnett, 1985)?

Sirex noctilio can spread through natural or artificial means. Natural spread mechanisms include flight and wind dispersal. Artificial, (*i.e.* human mediated), pathways of *S. noctilio* movement include pine logs and lumber products; large pine branches; and pine stumps. *Sirex cyaneus* is a similar species of woodwasp, with a native range corresponding to that of *S. noctilio*; this species' establishment in the United States was due to commerce and may indicate how *S. noctilio* could spread without proper safeguards (Smith, 2004).

The development of control strategies for *S. noctilio* originated in Australia, and is being utilized by various countries in the Southern Hemisphere. Control strategies include the restricted movement of infested materials; population monitoring through survey and trap trees; good silvicultural management practices; and the utilization of biological control agents, primarily the parasitic nematode, *Beddingia siricidicola*.

We recommend that similar strategies be implemented for the management of *S. noctilio* in the United States. These strategies will prevent serious economic damage, which would be the result of pest population outbreaks. To implement these strategies, we must determine the extent and the level of *S. noctilio* infestation; prevent the artificial spread of *S. noctilio* in potentially infested materials through treatment and regulated movement of the materials; promote good silvicultural management practices; and utilize the parasitic nematode, *B. siricidicola*, as a biological control agent when, and if, the nematode is approved for release in the United States.

Table of Contents

Executive Summary	ii
I. Introduction	1
II. Life History.....	1
2.1 Biology and Ecology.....	1
III. Geographic Distribution.....	4
3.1 Worldwide Distribution	4
3.2 Current Distribution in the United States and Canada, and History of Introduction	4
IV. Consequences of Introduction.....	5
4.1 Risk Element #1. Habitat Suitability.....	5
4.2 Risk Element #2: Host Range	8
4.3 Risk Element #3: Dispersal Potential.....	9
4.4 Risk Element #4: Economic Impact	10
4.5 Risk Element #5: Environmental Impact	17
4.6 Cumulative Risk Element Score	18
V. Pathways of Introduction	18
5.1 Natural Spread	18
5.2 Artificial Spread.....	21
VI. Control Options	27
VII. Risk Mitigation Options	29
VIII. Conclusions	30
IX. Recommendations.....	31
X. Future Research Needs.....	31

XI. Authors & Reviewers 32
XII. Acknowledgements 32
XIII. References Cited..... 33

I. Introduction

Sirex noctilio Fabricius (Hymenoptera: Siricidae) is a member of the horntail wasp family, Siricidae. This species belongs to the subfamily Siricinae and attacks gymnosperms. *Sirex noctilio* is the only species of the woodwasp family that can kill relatively healthy pine trees (Spradbery, 1973; Haugen *et al.*, 1990; Smith and Schiff, 2002). The native range of *S. noctilio* includes Europe, Asia, and North Africa, where it is a minor pest (CPC, 2005; Cielsa, 2003). *Sirex noctilio* is a serious pest of Monterey pine, *Pinus radiata*, plantations in New Zealand, Australia, South Africa, Argentina, Uruguay, and Chile. In Brazil, loblolly pine, *Pinus taeda*, is the primary host of *S. noctilio*.

In September 2004, a Lindgren funnel trap in Fulton, New York (NY) captured an adult *S. noctilio* (Hoebeke *et al.*, 2005). The wasp was positively identified as *S. noctilio* by Richard Hoebeke of Cornell University. Additional delimiting trapping, aerial and ground surveys for potentially infested trees were conducted around Fulton and Oswego NY. As a result, there were additional finds of *S. noctilio*. Because positive trap detections continued to increase in the survey areas, the survey radius expanded 30 to 70 miles from Oswego during the summer and early fall of 2005. The most distant positive find occurred approximately 50 miles southwest of Oswego. Sixty-one of 576 traps were positive for *S. noctilio* in 2005, yielding 85 female specimens. Currently, Oswego is the only location where *S. noctilio* larvae have been collected from infested trees; it is a focal point of the *S. noctilio* infestation in the United States.

The purpose of this pest risk analysis is to examine the risk associated with the woodwasp, *Sirex noctilio* Fabricius and the symbiotic fungus *Amylostereum areolatum* (Fries:Fries) Boidin to the conterminous United States. *Sirex noctilio* and *Amylostereum areolatum* exist in an obligate relationship, with the fungus exhibiting a very rare or absent sexual stage. *Amylostereum areolatum* relies on *S. noctilio* for dispersal and inoculation into the tree and the wasp relies on the fungus for wood breakdown and food (Slippers *et al.*, 2003). Due to this obligate relationship, we did not separate the risks associated with the two organisms, but considered them together in the analysis. Topic areas addressing specific issues were added as necessary. We identified areas of uncertainty and made recommendations regarding regulations and future research needs. This information can be used to facilitate the implementation of regulatory practices regarding the wasp with the goal of protecting U.S. pine resources in an efficacious and economically expedient manner.

II. Life History

2.1 Biology and Ecology

Sirex noctilio is a woodwasp that attacks coniferous trees. The adult wasps are large, ranging from 1 to 1.5 inches (2.5 to 4 centimeters) in length, with a metallic (iridescent) blue-black coloration and a stout upturned spine (cornus) at the end of the abdomen. The body of the female wasp is uniform in color with a prominent robust ovipositor located beneath the spine; and the legs are reddish-yellow, with black feet (tarsi). The male wasp has orange middle abdominal segments (segments 3 to 7) and the hind legs are black in color (Smith and Schiff,

2002; Haugen and Hoebeke, 2005). Detailed identification keys for *Sirex noctilio*, and other Siricid species, can be found in Smith and Schiff (2002).

Adult *S. noctilio* wasps emerge during the summer through mid-autumn (Madden, 1988) through circular holes (3 to 8 millimeters, depending on the size of the adult). The wasps are sexually mature upon emergence, with male wasps emerging first and in greater numbers than the females (Neumann and Minko, 1981). A post-emergence rest period of 15 minutes or more is followed by short bursts of flight. Males fly above and around the tree tops on sunny days, and can form small swarms when present in sufficient numbers. *Sirex noctilio* females have an initial flight of 100 meters or more after emergence, followed by shorter noisy flights similar to the male wasps (Morgan and Stewart, 1966). The females will then move up the trunk of the tree and begin inserting the ovipositor; females prefer stressed trees, which may be the result of drought, nutrition deficiency, unsuitable site selection, damage from storms, and climate, organisms, or suppression (Neumann *et al.*, 1987). *Sirex noctilio* attack both plantation and wild pine trees under stressed conditions. The volatiles released from attractive trees are monoterpene hydrocarbons and ketones (Simpson and McQuilkin, 1976). The exploratory insertion of the ovipositor assesses the suitability of the host tree. If the osmotic pressure of the tree is high, greater than 1210 kPa, the female will deposit only phytotoxic mucus (Madden, 1974); the mucus will cause a disruption in the functioning of the tree's needles and respiration (Coutts, 1969b), weakening the tree, and making it susceptible to future *Sirex* infestations. The mucus secretion is a blend of polysaccharides enzymes, oxidases and proteolases (King, 1966; Fong and Crowden, 1976) produced by glands and stored in a reservoir (Spradbery, 1973). *Sirex noctilio* has a larger mucus gland compared to other members of the Siricidae family (Spradbery, 1977).

Sirex noctilio is arrhenotokous, meaning that unmated females produce only male offspring, while mated females produce either male or female offspring (Taylor, 1981). In suitable trees, (*i.e.*, trees with low osmotic pressure 200-810 kPa), the female wasp will make multiple drills and deposit egg(s), spores of the fungi *Amylostereum areolatum*, and mucus (Neumann *et al.*, 1987). *Sirex noctilio* will oviposit in wood with a moisture content ranging from 20-200 percent oven-dry weight (Morgan and Stewart, 1966). *Amylostereum areolatum* is a saprotrophic decay fungi that causes extensive rot within infected trees over time, spreading up to 2.8 meters in ten years (Vasiliauskas *et al.*, 1998). *Amylostereum areolatum* and *S. noctilio* have a symbiotic relationship; the fungi is deposited into the tree by the wasp, and then dries and breaks down the wood into digestible hyphal fragments that the larvae feed and develop on (Neumann *et al.*, 1987). Slippers *et al.* (2003) completed a review of the symbiotic association between *Amylostereum* and woodwasps. Adult *S. noctilio* do not feed and have a short life span ranging from a few days to a few weeks. The females can sometimes be found dead on trees with the ovipositor still inserted (Taylor, 1981). The number of eggs per female *S. noctilio* is dependent on the size of the wasp, *e.g.*, 212 eggs were found per average size female in Victoria, Australia (Neumann *et al.*, 1987). The mucus and *A. areolatum* work in conjunction to dry the wood of moisture to create a more suitable environment for eggs to hatch (30 to 70 percent less than saturated) (Neumann and Minko, 1981).

The eggs of *S. noctilio* are elongate-oval (1.0-1.5 mm x 0.2-0.3 mm), white, and soft and smooth in appearance (Morgan, 1968). They are deposited into the xylem (or outer sapwood of the tree) within drill shafts bored by the ovipositor (Madden, 1988). The oviposition depth is reported to

be 10-20 millimeters into the sapwood, but can be influenced by the width of the growth rings in the tree (Neumann and Minko, 1981; Madden 1988). In Australia, eclosion of the larvae from the eggs typically occurs in 14 days, but the length of time can be delayed and affected by environmental conditions (Neumann *et al.*, 1987).

Larvae are creamy-white with deep segmentations, s-shaped and uniform in diameter (Neumann and Minko, 1981). The average body length and thoracic width of the *S. noctilio* larvae is 1.06 by 0.26, and 27.17 by 6.23 millimeters for first and sixth instars, respectively (Neumann and Minko, 1981). *Sirex noctilio* typically passes through six to seven larval instars before pupating, but the number of instars can range from 5-12 (Neumann and Minko, 1981). The first instar larvae move up or down from the egg gallery along the wood tracheid line 8 to 12 millimeters from the bark (Morgan and Stewart, 1966). The second instar female larvae acquire mycelium of *A. areolatum* from the infected wood and store it in hypopleural organs (Neumann and Minko, 1981). At the end of the third instar larval stage, *S. noctilio* will have progressed 15 to 20 millimeters in the wood from the initial egg chamber (Morgan and Stewart, 1966). The fourth and fifth instar larvae generally turn inward toward the heartwood (center) about 60 mm, creating a meandering mine that turns up and out toward the surface of the wood (Morgan and Stewart, 1966). The total length of the larval mine is reported to be 120-150 mm long, expanding in diameter as it progresses, with the pupation chamber constructed 50 mm from the surface of the tree (Morgan and Stewart, 1966). The larval tunnel is filled with a coarse frass consisting of chewed wood and excrement. A pre-pupal period of 20 to 28 days is typical (Morgan and Stewart, 1966), but the length is influenced by environmental conditions. The pupae and newly molted (teneral) adult retain the cast off skin (exuviae) from pre-pupal and pupal molts; this skin acts as a cap over the distal third of the abdomen until the adult wasp begins emergence. The cap helps the transfer of the fungus to the female wasp (Morgan and Stewart, 1966).



Figure 1. Wood Infested with *Sirex noctilio* Larvae. Paula Klasmer, Instituto Nacional de Tecnologia Agropecuaria, www.forestryimages.org

III. Geographic Distribution

3.1 Worldwide Distribution

Sirex noctilio occurs in the following European countries where it is considered a minor pest: Austria, Belgium, Cyprus, Czechoslovakia (former), Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Italy, Norway, Poland, Portugal, Azores, Romania, Russian Federation (localized), Serbia, Spain (Canary Islands), and the United Kingdom (CPC, 2005). In Asia and Africa, *S. noctilio* occurs in Mongolia and South Africa (CPC, 2005). In South America, *S. noctilio* is a major pest in Argentina, Brazil (Parana, Rio Grande Do Sul, and Santa Catarina), Uruguay, and Chile (CPC, 2005). The infestation of woodwasp in Chile is confined and under quarantine controls (CPC, 2005). In Oceania, *S. noctilio* occurs in Australia and New Zealand (CPC, 2005). The worldwide geographic distribution of *Sirex noctilio*, represented by country or region, is illustrated in Figure 2.




Figure 2. Reported Worldwide Distribution of *S. noctilio* (CPC, 2005).

3.2 Current Distribution in the United States and Canada, and History of Introduction

To date, *Sirex noctilio* has been detected in 25 New York counties and 2 Pennsylvania counties. It has been found in the Canadian province of Ontario in multiple counties and municipalities. Figure 3 displays the United States and Canadian counties in which *S. noctilio* has been detected.




Figure 3. Counties in North America where *S. noctilio* has been detected (CFIA 2006; APHIS, 2006a).

IV. Consequences of Introduction

4.1 Risk Element #1. Habitat Suitability

Rating	Numerical Score	Explanation
High	3	Attacks and survives on hosts in 4 or more plant hardiness zones
Medium	2	Attacks and survives on hosts in 2 or 3 plant hardiness zones
Low	1	Attacks and survives on hosts in a single plant hardiness zone

Rank: High

Sirex noctilio occurs in USDA Plant Hardiness Zones 3 to 10 in Europe (Spradbery and Kirk, 1978; CPC, 2005; Kelley, 1998), 8 to 10 in Australia (Neumann *et al.*, 1987; Dawson, 1991), New Zealand (Liddle wonder, 2002; CPC 2005), 6 to 10 in South America (CPC, 2005; Plant Ideas, 2006), and 7 to 9 in South Africa (CPC, 2005; Backyard Gardener, 2006).

A degree day model for *S. noctilio* (based on the work of Madden (1981)) was developed using the NAPPFAST system (Borchert and Magarey, 2005). The model demonstrated that *S. noctilio* could complete one generation (life-cycle) of development per year in much of the United States, with generation development taking potentially two years at higher latitudes (Figure 4). A CLIMEX prediction model for *S. noctilio* potential distribution found that many areas of the United States, Canada and Mexico were suitable for establishment (Carnegie *et al.*, 2006). Consequently, host distribution will probably limit *S. noctilio*'s distribution, for example, the distribution of pine hosts in the United States indicates that *S. noctilio* can establish in USDA Plant Hardiness Zones 3 to 9 (Figures 4, 5 and 6). This confers a rank of **High** to *S. noctilio* with regard to Habitat Suitability in the United States (USDA-ARS, 1990).

Below is a list of the regions and their pine host distribution, as per the USDA Plant Hardiness Zones: Northeastern and North Central States, pine host distribution Zones 3 to 6 (Figures 5, 6 and 7) (USDA-ARS, 1990); the Southern States, pine host distribution Zones 6 to 9; and the Western States, pine host distribution Zones 3 to 8. This confers a rank of **High** to *S. noctilio* with regard to Habitat Suitability in each of these regions.




Figure 4. Generation Potential Estimated for *Sirex noctilio* in the United States. (NAPPFAST degree day models with a base developmental temperature of 6.8 C and 2500 DD estimate for the completion of one generation.)




Figure 5. United States *S. noctilio* Pine Hosts Distribution on Timberland (CPC, 2005; USDA-USFS, 1991).




Figure 6. *Sirex noctilio* Pine Hosts Density in the United States (CPC, 2005; USDA-USFS, 1991). Pine hosts density is reported in percent forest cover in timberland (USDA-USFS, 1991).




Figure 7. Major Pine Forest Types in the Conterminous United States (USDA-USFS, 1991).

4.2 Risk Element #2: Host Range

Rating	Numerical Score	Explanation
High	3	Insect attacks multiple species within multiple host families
Medium	2	Insect attacks multiple species within a single host family
Low	1	Insect attacks only a single species or multiple species within a single genus

Rank: Low

Pines are the principal hosts of *S. noctilio*, and the wasp can complete its life-cycle on multiple species within this genus (Chrystal, 1928; CPC, 2005; Spradberry and Kirk, 1978, Morgan and Stewart, 1966). The following *Pinus* species are hosts of *S. noctilio*: *P. radiata*, *P. nigra calabrica*, *P. nigra austriaca*, *P. ponderosa*, *P. elliotii*, *P. patula*, *P. contorta*, *P. caribaea*, *P. pinaster*, *P. attenuata*, *P. muricata*, *P. banksiana*, *P. canariensis*, *P. densiflora*, *P. echinata*, *P. halepensis*, *P. jeffreyi*, *P. palustris*, *P. pinaster*, *P. pinea*, *P. brutia*, *P. sylvestris* and *P. taeda* (Morgan and Stewart, 1966; CPC, 2005; Madden 1988; Spradberry and Kirk, 1978). *Sirex noctilio* has been collected from *P. strobis* (white pine) and *P. resinosa* (red pine) in Oswego, New York. *Sirex noctilio* has also been recorded on other genera of conifers, such as *Larix* (Larch), *Pseudotsuga* (Douglas Firs), *Picea* (spruce) and *Abies* (firs) (Morgan and Stewart, 1966; Madden, 1988, Spradberry and Kirk, 1978; Chrystal, 1928). *Sirex noctilio* attacked *Larix* and *Pseudotsuga* in New Zealand when moribund (Morgan and Stewart, 1966), while Chrystal

(1928) reported that it “sometimes attacked silver firs and very occasionally spruce.” In an extensive survey by Spradberry and Kirk (1978), 99 percent of the 8,265 *S. noctilio* wasps collected were from *Pinus* spp.; the remaining 0.8 percent and 0.05 percent were found on *Picea* and *Abies* spp., respectively. Based on this information, we consider pines to be the only viable hosts with regard to risk scoring. We confer a rating of **Low** to *S. noctilio* with regard to Host Range; however, the genus *Pinus* is a large, diverse and important genus of plants, and *S. noctilio* has been reported to attack numerous species within this genus.

4.3 Risk Element #3: Dispersal Potential

	Dispersal Considerations	Source
X	Consistent and prolific reproduction	Madden, 1988
	Rapid growth to reproductive maturity	
	Wide range of hosts	
X	Tolerant to temperature extremes	Spradberry and Kirk, 1978
	Phoresy, <i>i.e.</i> dispersal by utilizing another organism	
	Ability to utilize different host during different life stages	
	Social behavior	
	Migratory behavior/swarming	
X	Alteration of generations/parthenogenesis/phase polymorphism	Taylor, 1981
X	Can reside within host	Neumann <i>et al.</i> , 1987
	Diapause/overwintering	
X	Stress tolerance, including ability to resist insecticides and/or adverse weather conditions	Fisher, 1955
	Lack of natural control agents	
X	Natural dispersal	Morgan and Stewart, 1966 Neumann <i>et al.</i> , 1987
X	Wind dispersal	Morgan and Stewart, 1966
	Water dispersal	
	Machinery dispersal	
	Animal dispersal	
X	Human dispersal	USDA, 1992

Rating	Numerical Score	Explanation
High	3	Insect has high reproductive potential (<i>e.g.</i> , prolific egg production, high survival rate), AND highly mobile life stages (<i>i.e.</i> , capable of moving long distances aided by wind, water or vectors)
Medium	2	Insect has either high reproductive potential OR highly mobile life stages
Low	1	Insect has neither high reproductive potential nor highly mobile life stages

Rank: High

Sirex noctilio females lay an average of 212 eggs (Neumann *et al.*, 1987); a single female can produce 53 female offspring (Neumann and Minko, 1981). *Sirex noctilio* are arrhenotokous parthenogenic, meaning that unmated females produce only male offspring, while mated females can produce either male or female offspring (Taylor, 1981).

The immature stages of *S. noctilio* remain within the wood during development; survival during larval development is affected by the ability of the tree to compartmentalize the fungus and larvae; parasitization of the wasp by natural enemies; moisture content of the wood; and the amount of resin. In newly invaded areas, the survival rate is generally high (Neumann and Minko, 1981; Taylor, 1981).

The maximum yearly dispersal rate of *Sirex noctilio* observed in Australia is 30 to 40 km (18 to 25 miles) and 48 kilometers (29.8 miles) per year in South Africa (Carnegie *et al.*, 2006). In flight tunnel experiments, large females have been reported to fly up to 160 kilometers (100 miles) (Taylor, 1981). Observation shows that when female wasps emerge later in the season, or in areas where resources are limited, they disperse farther than early emerging females. Male wasps remain near the initial emergence area unless spread by wind (Morgan and Stewart, 1966).

Sirex noctilio can be spread by humans through the movement of infested material, *e.g.* logs. *Sirex noctilio* has been observed ovipositing on freshly cut logs on logging trucks in New Zealand, often continuing to attack the logs during transit (Morgan and Stewart, 1966). The immature stages develop within the wood and are difficult to detect, even in rough sawn lumber (USDA, 1992; Fisher, 1955). The wood wasp *Sirex cyaneus*, already well-established in the United States, is a likely example of an invasive species that followed a similar pathway, and has been reported to be moved in commerce (Arnett, 1985).

4.4 Risk Element #4: Economic Impact

Impact Categories include reduced commodity yield (*e.g.*, feeding, disease vector); lower commodity value (*e.g.*, by increasing costs of production, lowering the market price, or a combination or, if not an agricultural insect, by increasing costs of control); loss of markets (foreign or domestic) due to presence of a new quarantine pest.

Rating	Numerical Score	Explanation
High	3	Insect causes all three of the above impacts, or causes any one impact over a wide range of economic plants, plant products or animals (over five types)
Medium	2	Insect causes any two of the above impacts, or causes any one impact to three or four types of economic plants, plant products, or animals
Low	1	Insect causes any one of the above impacts to one or two types of economic plants, plant products, or animals
Nil	0	Insect causes none of the above impacts

Rank: High

Table 1. Annual U.S. Christmas Tree Production in the Northeastern and North Central United States (Helmsing, 2004; Michigan Ag Connection, 2005; Olsen pers. comm., 2005; USDA-ERS, 1990; USDA-NASS, 2002).

State	Sales	Proportion of Trees Sold (Pines)	Estimated Annual Pine Christmas Tree Sales
Connecticut	3,407,000	0.10	340,700
Delaware	401,000	0.41	164,410
Iowa	1,424,000	0.95	1,352,800
Illinois	7,633,000	0.89	6,793,370
Indiana	2,775,000	0.83	2,303,250
Massachusetts	1,800,000	0.16	288,000
Maryland	2,313,000	0.75	1,734,750
Maine	2,293,000	0.10	229,300
Michigan	30,411,000	0.21	6,386,310
Minnesota	11,855,000	NA	NA
Missouri	1,843,000	0.98	1,806,140
New Hampshire	2,028,000	0.14	283,920
New Jersey	3,852,000	0.23	885,960
New York	11,759,423	0.16	1,881,508
Ohio	9,323,000	0.83	7,738,090
Pennsylvania	31,193,000	0.41	12,789,130
Rhode Island	658,000	0.24	157,920
Vermont	2,372,000	0.07	166,040
West Virginia	1,182,000	0.66	780,120
Wisconsin	23,412,000	0.69	16,154,280
Total	151,934,423	0.41	62,235,998

Table 2. Annual Coniferous Evergreen Nursery Data for Selected U.S. States (USDA-NASS, 2004). Average values based on 2000 and 2003 nursery data reported. Nursery data reported for operations with annual sales greater than \$100,000.

State	Producers	Plants Sold	Sales
Alabama	56	2,104,500	8,236,500
California	123	5,610,000	64,606,500
Connecticut	26	2,843,000	25,080,500
Florida	144	2,649,500	20,389,500
Georgia	47	2,127,000	11,653,500
Illinois	82	665,500	19,063,500
Michigan	79	2,603,500	33,264,000
New Jersey	94	1,550,000	25,726,500
New York	52	530,000	9,866,000
North Carolina	125	1,607,000	19,776,000
Ohio	79	1,578,000	27,509,500
Oregon	161	10,601,500	97,569,500
Pennsylvania	96	1,093,500	27,986,500
South Carolina ¹	37	818,000	4,547,000
Tennessee	93	1,080,000	9,793,000
Texas	47	884,500	8,186,000
Virginia ²	32	708,000	9,535,000
Washington	39	637,000	7,298,500
Total	1,412	39,690,500	430,087,500

¹South Carolina only reported data for 2000

²Virginia only reported data for 2003

An attack by *S. noctilio* can cause tree mortality, in addition to a reduction in commodity value, which is an effect of the deposition of the phytotoxic mucus and the introduction of *A. areolatum* (Taylor, 1981). The damage caused by an attack could pose an economic threat to the U.S. softwood timber industry. The timber industry in the United States produces large quantities of timber; has timber processing mills throughout the country; and has annual sales of logs, lumber, pulpwood and veneer valued at nearly \$20 billion (LDAF, 2000; Prestemon *et al.*, 2005; USDA-USFS, 2001, 2003; USDC, 1999a, 1999b, 1999c) (Figure 12; Tables 6, 7, 8 and 9).

Sirex noctilio can successfully attack stressed trees (CPC, 2005; Neumann *et al.*, 1987). Common causes of tree stress include overcrowding, drought, nutrition deficiency, unsuitable site selection, damage from climate or other organisms, or suppression (Neumann *et al.*, 1987). From 1987 to 1989 southeastern Australia and southwestern Victoria, experienced severe outbreaks of *S. noctilio* in plantations of *Pinus radiata*; these outbreaks cost an estimated \$10 to 12 million in losses (Haugen *et al.*, 1990).

When evaluating the economic effects caused by the introduction of *S. noctilio* into three areas of the western United States (USDA, 1992), the best and worse case scenarios estimated losses between \$24 to 131 million. Of the five wood pests examined in this economic evaluation, *S. noctilio* had the greatest potential economic impact.

Sirex noctilio is a quarantine pest in the United States, Canada and Japan. Phytosanitary certificates and treatments will likely be necessary for the movement of pine logs from areas identified as infested with *S. noctilio* (CPC, 2005).

Table 3. 1996 United States Softwood Production in Thousand Cubic Feet (MCF) by Region (USDA-USFS, 2001).

United States Region	Saw Logs	Veneer Logs	Pulpwood	All Products
Northeast North Central	336,542	3,075	369,044	815,874
South	2,721,782	736,174	2,399,152	6,154,838
West	2,099,934	384,689	82,472	3,065,488
Total	5,158,258	1,123,938	2,850,668	10,036,200

Table 4. Value of Selected Softwood Commodities in the Northeastern and North Central States that Could be Affected by *S. noctilio* (LDAF, 2000; USDA-USFS, 2001, 2003; USDC, 1999a, 1999b, 1999c).

State	Softwood Logs and Bolts ¹	Softwood Lumber ^{1,3}	Softwood Veneer ^{1,4}	Softwood Pulpwood Production ²	Total
Connecticut	NA	NA	NA	60,778	60,778
Delaware	NA	NA	NA	1,042,405	1,042,405
District of Columbia	NA	NA	NA	NA	NA
Illinois	NA	NA	NA	85,282	85,282
Indiana	NA	2,985,000	NA	260,529	3,245,529
Iowa	NA	NA	NA	NA	NA
Maine	128,299,000	262,045,000	NA	29,551,781	419,895,781
Maryland	3,210,000	29,220,000	NA	2,144,664	34,574,664
Massachusetts	NA	10,809,000	NA	241,633	11,050,633
Michigan	2,659,000	24,857,000	NA	10,000,351	37,516,351
Minnesota	4,689,000	43,997,000	NA	11,742,430	60,428,430
Missouri	NA	2,891,000	NA	3,225	2,894,225
New Hampshire	5,368,000	110,054,000	NA	5,125,285	120,547,285
New Jersey	NA	NA	NA	119,913	119,913
New York	25,000,000 ⁵	40,000,000 ⁵	NA	6,622,404	28,840,404
Ohio	NA	5,406,000	NA	486,798	5,892,798
Pennsylvania	NA	9,254,000	NA	1,518,974	10,772,974
Rhode Island	NA	NA	NA	88,938	88,938
Vermont	NA	NA	NA	4,173,256	4,173,256
West Virginia	NA	6,311,000	NA	1,110,557	7,421,557
Wisconsin	NA	24,224,000	NA	17,698,846	41,922,846
Total	169,225,000	572,053,000	NA	92,078,047	790,574,047

¹Values in 1997 dollars

²Values based on the average 1996 Louisiana southern pine pulpwood price per cord adjusted to 1998 dollars.

³Refers to lumber that is not edge worked and not manufactured from purchased lumber (USDC, 1999b).

⁴Includes veneer backed with cloth, paper or another flexible material (USDC, 1999c).

⁵ NYS Forest Products Timber Estimate 2004 (Crawford, 2006)

Table 5. Value of Selected Softwood Commodities in the Southern States that could be Affected by *S. noctilio* (LDAF, 2000; USDA-USFS, 2001, 2003; USDC, 1999a, 1999b, 1999c).

State	Softwood Logs and Bolts ¹	Softwood Lumber ^{1,3}	Softwood Veneer ^{1,4}	Softwood Pulpwood Production ²	Total
Alabama	67,078,000	803,149,000	36,721,000	117,562,990	1,024,510,990
Arkansas	152,623,000	817,759,000	12,627,000	35,409,337	1,018,418,337
Florida	77,317,000	244,798,000	NA	81,277,478	403,392,478
Georgia	96,614,000	998,557,000	52,696,000	119,639,174	1,267,506,174
Kentucky	NA	14,057,000	NA	938,390	14,995,390
Louisiana	107,022,000	412,891,000	33,950,000	64,595,774	618,458,774
Mississippi	271,459,000	937,552,000	28,866,000	58,912,520	1,296,789,520
North Carolina	122,671,000	571,646,000	11,255,000	54,777,797	760,349,797
Oklahoma	NA	129,014,000	NA	8,145,651	137,159,651
South Carolina	38,898,000	532,022,000	NA	63,356,945	634,276,945
Tennessee	NA	6,046,000	NA	12,692,221	18,738,221
Texas	210,876,000	424,550,000	9,607,000	32,336,410	677,369,410
Virginia	10,642,000	234,115,000	6,310,000	31,883,749	282,950,749
Total	1,155,200,000	6,126,156,000	192,032,000	681,528,435	8,154,916,435

¹Values are in 1997 dollars.

²Values based on the average 1996 Louisiana southern pine pulpwood price per cord adjusted to 1998 dollars.

³Refers to lumber that is not edge worked and not manufactured from purchased lumber (USDC, 1999b).

⁴Includes veneer backed with cloth, paper or another flexible material (USDC, 1999c).

Table 6. Value of Selected Softwood Commodities in the Western States that Could be Affected by *S. noctilio* (LDAF, 2000; USDA-USFS, 2001, 2003; USDC, 1999a, 1999b, 1999c).

State	Softwood Logs and Bolts ¹	Softwood Lumber ^{1,3}	Softwood Veneer ^{1,4}	Softwood Pulpwood Production ²	Total
Arizona	NA	45,962,000	NA	339,814	46,301,814
California	154,140,000	1,758,190,000	60,169,000	NA	1,972,499,000
Colorado	NA	35,359,000	NA	NA	35,359,000
Idaho	245,711,000	823,895,000	31,398,000	4,833,801	1,105,837,801
Kansas	NA	NA	NA	NA	NA
Montana	115,308,000	509,193,000	NA	1,182,282	625,683,282
Nebraska	NA	NA	NA	NA	NA
New Mexico	NA	NA	NA	371,569	371,569
Nevada	NA	NA	NA	NA	NA
North Dakota	NA	NA	NA	NA	NA
Oregon	993,860,000	2,418,176,000	392,057,000	2,182,786	3,806,275,786
South Dakota	NA	NA	NA	NA	NA
Utah	NA	13,867,000	NA	NA	13,867,000
Washington	1,331,068,000	1,610,913,000	80,165,000	6,170,894	3,028,316,894
Wyoming	2,008,000	73,182,000	NA	NA	75,190,000
Total	2,842,095,000	7,288,737,000	563,789,000	15,081,146	10,709,702,146

¹Values are in 1997 dollars.

²Values based on the average 1996 Louisiana southern pine pulpwood price per cord adjusted to 1998 dollars.

³Refers to lumber that is not edge worked and not manufactured from purchased lumber (USDC, 1999b).

⁴Includes veneer backed with cloth, paper or another flexible material (USDC, 1999c).

The regional distribution of U.S. forests we used is based on the USFS classification system; this may affect their susceptibility to damage by *S. noctilio*. This USFS system divides the conterminous United States into eight regions (Figure 8) (USDA-USFS, 2005). The Northeastern and North Central States are covered by region R9 (eastern); the South is covered by region R8; and the Western States are divided into six regions (R1-R6).




Figure 8. USFS Regions

The Southern States produce 60 percent of the nation’s timber (USDA-USFS, 2003). This region produces more timber than any other country. Softwood (especially southern pine) forestry is a major source of revenue in the South (Table 3). Over the past 50 years, the South has surpassed the Western States as the nation’s leading producer of softwood timber and pulpwood (USDA-USFS, 2003) (Table 3) for the following reasons: technological advances in southern pine manufacturing and treatment; short rotation periods; and increased demand for strong pulp fiber (Adams, 1995; Helms, 1995; Tesch, 1995; USDA-USFS, 2003; Walker, 1995). The increase in softwood timber and pulpwood production results in an increased investment in southern pine production and advances in stand management and tree genetics (USDA-USFS, 2003). Models forecasting southern timber trends through the year 2040 have predicted: 1) softwood timber prices will increase; 2) there be an increase in softwood timber production; and 3) pine plantation areas (in the South) will increase by 67 percent (USDA-USFS, 2003).

Loblolly pine (*Pinus taeda*) is the most important pine species for timber and pulpwood production in the Southern States; it comprises over 50 percent of pine in this region (UFL, No Date; About Inc., 2005). Other southern pine species that *S. noctilio* can affect (besides loblolly pine) include longleaf pine, shortleaf pine, and slash pine (USDA-USFS, 1991) (Figure 6).

Overall, the West produces less softwood timber than the Southeast, and more than the Northeast and North Central States (Table 3). The total value of western softwood timber products analyzed in this assessment is the highest in the United States, (*i.e.* approximately \$11 billion annually in logs, lumber, veneer and pulp products) (Table 6); this may be due, in part, to the large quantity of high priced softwood timber, (*e.g.* Douglas fir, harvested in Oregon and Washington), and the rising timber prices, due to lower forestry investments (Helms, 1995; Tesch, 1995; Skog and Risbrudt, 1982; USDA-USFS, 2003).

Compared to the West and South, the Northeastern and North Central States have a lower density and distribution of pine hosts (Figures 5, 6 and 7). Consequently, the softwood timber industry in the Northeast and North Central States is not as large as the Southern or Western States (Tables 3, 4, 5 and 6). White pine is one of the major forest types in the Northeast and North Central States (Figure 7).

Given the current and future economic value of pine resources, and the uncertainty regarding what impact *S. noctilio* will have if and when it spreads from the current area of detection, it is prudent to protect pine resources by the most efficacious means available. The recommendation section of this document includes suggestions with regard to protection methodologies.

4.5 Risk Element #5: Environmental Impact

Impact categories include effects on ecosystem processes (e.g., increases fire risk due to feeding or disease transmission); impacts on the natural community composition (e.g., reduce biodiversity, affect native populations, and affect endangered or threatened species); and the impacts on the community structure (e.g., change density of a canopy layer, eliminate or create a canopy layer). Other impacts include those on human health, such as disease transmission or the production of allergens; and the stimulation of control programs, including toxic chemical pesticides or the introduction of a non-indigenous biological control agent.

Rating	Numerical Score	Explanation
High	3	Three or more of the above.
Medium	2	Two of the above.
Low	1	One of the above.
Nil	0	None of the above.

Rank: Medium

Sirex noctilio has caused significant damage to plantations of *Pinus radiata* D. Dom (Monterey pine) in the Southern Hemisphere (Madden, 1988; Carnegie *et al.*, 2005). *Pinus radiata* is listed as Endangered by the World Conservation Union (IUCN) Conifer Specialist Group, indicating that it is “facing a very high risk of extinction in the wild in the near future” (Earle, 2005). The native *P. radiata* occurs in three locations in central coastal California (Earle and Frankis, 1999); these forests have significant ecological and genetic resource value (Ciesla, 2003).

In Australia and other countries in the Southern Hemisphere where *S. noctilio* is a pest, the neotylenchid nematode, *Beddingia* (= *Delandenus*) *siricidicola*, is a biological control agent. This parasitic nematode was first isolated from *S. noctilio* in New Zealand in 1962 (Bedding and Akhurst, 1974). The APHIS *Sirex* Science Advisory panel recommends the initiation of a program to import and deploy *B. siricidicola* as quickly as possible (APHIS, 2006b).

Sirex noctilio attacks live pine trees, which results in rapid wilting of the crown needles and tree death (Eldrige and Taylor, 1989); however, in their native range siricids are considered to be of minor environmental importance (Smith and Schiff, 2002; Slippers *et al.*, 2003). *Sirex noctilio* is an “essentially secondary, opportunistic wood-boring pest,” that, in small numbers, may be

useful in killing unwanted trees; however, these populations must be maintained, or damage can be significant (Neumann *et al.*, 1987).

4.6 Cumulative Risk Element Score

Cumulative Risk Element Score	Risk Rating	Risk Score
5-7	Low	1
8-11	Medium	2
12-15 (Habitat Suitability = 3) + (Host Range = 1) + (Dispersal Potential = 3) + (Economic Impact = 3) + (Environmental Impact = 2) = 12	High	3

Rank: High

With regard to cumulative risk, *Sirex noctilio* scored **High**. It scored High for Habitat Suitability, Dispersal Potential and Economic Impact. Environmental Impact scored Medium for the U.S.—based on the potential impact of the woodwasp related to the endangered *Pinus radiata*, and the introduction of a non-native species of nematode, *Beddingia siricidicola*, for use as a biological control agent. *Sirex noctilio* scored Low with regard to Host Range due to its preference for species in the genus *Pinus*. These scores indicate that *S. noctilio* could pose a serious potential economic threat to the U.S. forestry industry.

V. Pathways of Introduction

5.1 Natural Spread

Based on survey trapping conducted in 2005, *S. noctilio* has been found in five counties in the United States and four counties in Canada (APHIS, 2006a; CFIA, 2006); a delimiting survey has not yet been completed. A grid trapping plan, extending 150 miles in radius from Oswego, New York, has been developed and will be deployed in 2006. *Sirex noctilio* is capable of dispersing large distances, 30 to 40 kilometers (18 to 25 miles), each year (Carnegie *et al.*, 2006); this demonstrated ability allows it to span areas of low density pine host plantings, such as those in New South Wales, Australia (Carnegie *et al.*, 2005). Since the extent of *S. noctilio* infestation is unknown, it is difficult to estimate its annual rate of spread, and the length of time until other areas of the United States are affected by this pest.




Figure 9. Counties where *S. noctilio* has been detected in New York, Pennsylvania and Ontario.

Due to the uncertainty of this insect’s biology, we utilized two estimates of spread. By using an 18 mile per year spread estimate, we identified the United States and Canadian counties where *S. noctilio* currently resides and the portions of Michigan, Ohio, Indiana, New Jersey, Delaware, Maryland, West Virginia, Virginia, Vermont, New Hampshire, Connecticut and Massachusetts that could be affected within ten years by natural spread (Figure 10).




Figure 10. Estimated Spread of *Sirex noctilio* over 10 years. The above figure utilizes an 18 mile/year spread rate in the United States and Canadian counties currently reporting detection.

By using a 25 mile/year spread estimate and the counties in the United States and Canada where *S. noctilio* are currently present, Michigan, Ohio, Indiana, New Jersey, Delaware, Maryland, West Virginia, Virginia, Vermont, New Hampshire, Connecticut, Rhode Island, Maine and Massachusetts could be affected within ten years by natural spread (Figure 11).




Figure 11. Estimated Spread of *Sirex noctilio* over 10 years. The above figure utilizes a 25 mile/year spread rate in the United States and Canadian counties currently reporting detection.

5.2 Artificial Spread

The following pine articles were considered at-risk for artificially spreading *S. noctilio* by USDA-APHIS-PPQ standards, which include all host materials, living, dead, cut, or fallen, including nursery stock; untreated round wood (logs); untreated sawn wood and untreated wood packaging; firewood; wood chips; stumps; Christmas trees; roots; and branches and debris.

Nursery Stock

Annual sales of coniferous evergreens for 2003 were valued at \$443 million, according to a USDA 17 state nursery survey (Table 2) (USDA-NASS, 2004). Several major production nurseries produce and distribute this stock.

Pine trees sold at garden centers and retail nurseries may have passed through several businesses before reaching the retail center. Some nurseries grow planting stock that is sold to other nurseries. Planting stock of pines is commonly sold in the range of 1 to 3 inches to 2 to 3 feet to other growers. These nurseries grow the trees to a larger size either in containers or in the field.

Christmas tree growers also purchase a large amount of pine seedlings. Two to three year old pine seedlings are commonly used for Christmas tree production (Koelling and Dornbush, 1992).

The majority of pine nursery stock over 36 inches in height is sold balled and burlapped, with a smaller percent sold in containers (Cinnamon, pers. comm., 2005). Wholesale growers in Ohio grow 5 to 7 feet balled and burlapped Austrian pine for distribution to retail garden centers, re-wholesalers, and landscaping companies (Storts, pers. comm., 2005). The number of trees per acre varies from 1,100 to 1,200. In Ohio, 5 to 7 feet tall pine trees are dug in the early spring during the months of February, March, and April, and then again in the fall from late September through October (Storts, pers. comm., 2005). The majority of trees are dug to order, and then placed on trucks and shipped within two days (Storts, pers. comm., 2005). Insecticides are only applied in the presence of pests; the trees are not sprayed on a routine calendar spray schedule (Storts, pers. comm., 2005). Trees from this wholesale nursery are shipped within a 14 state radius.

Sirex noctilio females prefer trees under stress. In nurseries, it is likely that trees will be well-managed, healthy and not under stress conditions; therefore, it is unlikely that pine trees in nurseries will attract *S. noctilio*, additionally healthy trees are more resistant to *S. noctilio* attack (Coutts and Dolezal, 1965). If a *S. noctilio* attack is successful, trees will be culled from nursery stock due to the presence of visual symptoms like needle browning and droop, which occurs within a few months of attack. Trees that exhibit these symptoms are unmarketable, and will not likely be shipped. Although *S. noctilio* attacks trees as small as one inch in diameter, it is much more common for trees larger than three inches to be attacked (Morgan and Stewart, 1966). Large pine nursery stock, either in pots, or balled and burlapped, that are over four inches in diameter at the base, will usually ship in the fall, and will not show the characteristic of browning and droop prior to shipment, if infestation by *S. noctilio* has just occurred.

Mugo pines typically have compact growth (5-20 ft tall), with small multi-trunk shrub/tree form and dense growth (U Conn, 2007). This compact form in nursery stock would likely be unsuitable and unattractive to ovipositing *S. noctilio* females due to small trunk size, limited accessibility for oviposition and the typical oviposition behavior (Morgan and Stewart, 1966), additionally, as outlined above, nursery stock would likely be healthy and not under stress conditions. There has been no known report of *S. noctilio* using *P. mugo* as a host in Europe or Australia. *Pinus mugo*, which have compact shrub like growth forms, the risk for artificial spread of *S. noctilio* through their movement would be **Low** to **Nil**.

The risk for artificial spread of *S. noctilio* through the movement of large nursery stock in the fall is greater than the risk of moving large nursery stock in the spring, as there are no visible symptoms of attack during the fall shipping season. The risk of artificial spread of *S. noctilio* through the movement of pine nursery stock from the areas of detection, and the bordering counties where no detection occurred, is **Medium** for large nursery stock during the fall, and **Low** for all pine nursery stock shipped during the spring.

Untreated Pine Roundwood (Logs)

In the 2004 New York Industrial Timber report (Crawford, 2005), the total estimated log production for all hardwoods and softwoods combined was 811 million board feet (MMbf), with

115 MMbf white pine (*P. strobis*) harvested and 45 MMbf red pine (*Pinus resinosa*) (Crawford, 2006). The majority (64 percent) of the timber harvested in New York remains in the state. Exported timber primarily ships to Canada (73 percent), Vermont and Pennsylvania (27 percent), with minor volumes of unreported logs going to multiple New England, Mid-Atlantic and Mid-West states. Of the exported softwood log timber harvested in New York, 96.1 MMbf was exported to Canada, and 12 MMbf was exported to Vermont and Pennsylvania. It is important to note that the New York industrial timber report only addresses logs and pulpwood, and does not address large stock wood products, such as utility poles and timbers used in the log home manufacturing industry.

We considered the potential risk for the movement of immature stages of *S. noctilio* in untreated pine logs to be **High** (Morgan and Stewart, 1966; Haugen *et al.*, 1990; USDA, 1992; Iede *et al.*, 1998). *Sirex noctilio* larvae can survive two or more years in wood, with a relatively low mortality rate if the moisture content of the wood is suitable (above 20 percent oven-dried weight) (Neumann *et al.*, 1987; USDA, 1992). Green pine logs are most attractive and susceptible to oviposition by *S. noctilio* 5 to 7 days after felling, with their attractiveness lasting up to three to four weeks (Taylor, 1981). The movement of infested material to uninfested areas without prior treatment is greatly discouraged (Iede *et al.*, 1998). The level of infestation by *S. noctilio* in the counties where detection has occurred and the effectiveness of visual inspection for log infestation and similar products are unknown. The risk associated with large stock wood products, such as untreated pine utility poles and logs or timbers used in log home manufacturing in untreated form, would be analogous to untreated logs. *Sirex noctilio* could survive within these stock wood products for extended periods of time and the effectiveness of visual detection on logs and poles is believed to be low, however visual detection in timbers and other wood products is probably higher (Crawford, 2006). Because of these characteristics, the risk of artificial spread of *S. noctilio* through the movement of these products from areas where detection has occurred and bordering counties to areas where no detection has occurred is **High**.

Untreated Sawn Wood (Green Pine Lumber) and Untreated Packing Material (Green Pine Packing Material)

Sirex noctilio larvae can survive in sawn and air dried wood and larval presence is difficult to detect if the lumber is rough cut (Fisher, 1955). Chandler (1959) reported other siricidae species emerging from non-kiln dried lumber inside buildings for up to three years after construction. In logs, the eggs and young larvae are susceptible to desiccation, if the wood is too dry. *Sirex noctilio* is not reported to lay eggs in very dry wood, but mature larvae can survive in wood with low moisture content (< 20 percent ODW) (Coutts and Dolezal, 1965). Adult *S. noctilio* emerging from very dry logs in insectaries are very small (Coutts and Dolezal, 1965), and the number of eggs per female are directly related to size (Newman *et al.*, 1987). Siricidae larvae can be moved in wood materials, *e.g.* dunnage, solid wood packing materials, and have been intercepted at U.S. ports-of-entry 149 times since 1985 (PIN, 2005). Of the 149 siricidae intercepted, 17 were identified as species of *S. noctilio*, while the remaining could not be identified below the family level.

Green pine lumber is typically utilized on a local basis, usually within a 50 mile radius of the mill, due to the lower value and limited utility of the product (Crawford, pers. comm. 2006). Solid wood packing material (SWPM) used for domestic applications will be untreated green

wood. McIntosh Box and Pallet Co. Inc. headquartered in East Syracuse, New York is a manufacturer of pallets and crating products with four of their five plants located in counties where *S. noctilio* has been detected or in bordering counties. McIntosh uses an estimated nine million board feet of wood per year, with 40 to 50 percent of the wood being softwood. Of the softwood utilized, more than 90 percent is imported from Canada as kiln dried “downfall” product, which is lumber that did not meet the qualifying grade for construction lumber. Kiln drying is a method utilized to kill pests, such as *S. noctilio*, in the wood (APHIS, 2005a). The remaining softwood material comes from New York and Pennsylvania, predominantly as kiln dried downfall product. The use of downfall products is a widespread practice among the SWPM industry in New York. McIntosh reported using untreated green white pine lumber in the manufacturing of a very small amount of specialty products (Huftalen, Pers. communication, 2006).

Due to the limited movement of green pine lumber, and the small amount of green pine material used in the manufacturing of SWPM from the counties where *S. noctilio* has been detected and the bordering counties in New York, we considered the potential risk for artificial spread of *S. noctilio* in untreated sawn wood (green pine lumber) and untreated packing material (green pine packing material) to be **Low**. If the area of *S. noctilio* infestation increases to include areas where untreated packing material utilizes greater amounts of untreated pine host material in the manufacturing of SWPM, the potential risk for the movement of *S. noctilio* by SWPM would increase proportionally, and the potential risk would be **High**.

Firewood (Fuelwood)

As indicated in the previous section, *S. noctilio* can survive and develop within wood for extended periods of time, it is difficult to detect larvae within the wood and it is possible to transport larvae if the material is moved from infested areas. Although there have been no specific papers discussing *S. noctilio* in firewood, the fact that it is able to survive in many other types of semi-dried wood products indicate it would be able to survive in firewood.

According to the 2002 FIA database, there is no indication of any pine being utilized as fuelwood in New York. The only fuelwood listed in New York was from hardwoods, with approximately six million cubic feet of roundwood product growing stock. The FIA collected this data on fuelwood by conducting residential telephone surveys of “individuals or groups of woodland owners who have harvested or allowed fuelwood to be harvested from their land” (Wharton, 2005). The general use and movement of firewood by private individuals on a local basis is probably not captured in the FIA database.

The risk associated with the artificial spread of *Sirex noctilio* through the transport of firewood is **Low** due to the limited production of pine firewood/fuelwood from NY. If the area of *S. noctilio* infestation increases to include areas where pine firewood is utilized in greater amounts, such as Massachusetts, New Hampshire, Maine, Indiana and Michigan (Figure 12), the potential risk for movement of *S. noctilio* by firewood would increase proportionately, and the potential risk would be **High**.




Figure 12. Cubic feet of pine fuelwood product reported by county in the Forest Inventory Analysis (FIA) database. (2002 Roundwood Growing Stock)

Wood Chips

The chipping of wood products into pieces 0.625 inch (5/8”) (or smaller) at the thickest point helped eliminate all life stages of the Asian Longhorned Beetle (ALB), *Anoplophora glabripennis*, and the Emerald Ash Borer (EAB), *Agrilus planipennis* (APHIS, 2003a; APHIS 2003b). Currently, chipping is an eradication treatment in the ALB and EAB quarantine areas. The larvae of ALB bore deep into the center of logs and are comparable in size to *S. noctilio* larvae. Because of the similarity in habitat and size, the chipping of pine wood products to a thickness of 0.625 inch or less would kill all stages of *S. noctilio*. We considered the risk of artificial spread of *S. noctilio* through the movement of pine wood chips 0.625 inch or less to be **Nil**.

Stumps

Processed stumps can create wood naval stores, which include products such as wood turpentine, wood rosin, dipentene, and natural pine oil (IPPC, 1995). Rosin products can be extracted from chipped stumps or through distillation (IPPC, 1994). Wood rosin is a small component of pine oleoresin production, constituting five percent of production worldwide, or 2,400 tons annually (IPPC, 1995). Wood naval stores production has declined to a low level in the United States in the past 60 years (IPPC, 1995).

Annual sales of rosin products from pine stumps are about 20 million pounds per year (Jacobs pers. comm., 2005). The Hercules Company is the only manufacturer of rosin from stumps in the world. They utilize nine to ten thousand gross tons of stumps per month for production (Jacobs pers. comm., 2005; PINOVA, 2005).

The wood rosin industry in the United States harvests stumps from the coastlines of North Carolina, South Carolina and Georgia (Jacobs pers. comm., 2005). The majority of harvested

stumps come from Florida and land 200 miles inland from the Gulf of Mexico in Georgia and Alabama (Jacobs pers. comm., 2005). After timber harvests, stumps from previous harvests are removed. Stumps must be at least 20 years old to produce quality rosin (Jacobs pers. comm., 2005). Harvested year-round, the longleaf and loblolly pine stumps produce naval stores. Because stumps harvested for naval stores are not removed from counties where *S. noctilio* has been detected, we did not consider this as a viable pathway for artificial movement.

In the counties where *S. noctilio* has been detected, and in the bordering counties, the harvesting of pine forest timber does not include the removal of stumps; this is the general practice in New York (Crawford, pers. comm. 2006).

Sirex noctilio will typically infest the stem portion (or bole) of the pine tree with over 90 percent of emergences recorded are from breast height upward to a diameter of three inches at the tree top (Morgan and Stewart, 1966). Green stumps are highly susceptible to attack from *S. noctilio* (Neumann and Minko, 1981). Because pine stumps are not removed from the forests in counties (and bordering counties) from where detection has occurred, *S. noctilio* preferentially attacks the upper stem portion of the tree. We considered the potential risk for artificial movement of *S. noctilio* by the movement of pine stumps to be **Low** to **Nil**. If *S. noctilio* were detected in areas where stumps were removed at the time of log harvest, the risk of artificial spread through the movement of stumps would probably be **Low**, as there may be low numbers of larvae found in the stumps. However, should stumps be utilized for fuelwood, and if processed (chipped) within a short period of time, the risk for artificial spread by stumps would be **Nil**.

Christmas Trees

The estimated proportion of pine Christmas trees sold in New York is 0.16, with an annual value of approximately \$1.9 million (Table 1). In New York Christmas tree farms, the estimated majority of newly planted trees are firs and spruce; pines are not as popular with consumers as in the past (Crawford, pers. Communication 2006). Only four of the fourteen tree varieties listed on the Christmas Tree Farmers Association of New York website (2006) were pines.

Sirex noctilio females prefer trees under stress (Neumann *et al.*, 1987). Christmas tree farm trees are healthy, well-managed and thrive under stress-less conditions. Pine trees infested with *S. noctilio* exhibit needle browning and droop within a few months following attack (Haugen *et al.*, 1990; Neumann *et al.*, 1987); these symptoms make the trees unmarketable and reduce the likelihood of shipment. It is unlikely that pine trees in Christmas tree farms will attract *S. noctilio*; healthy trees are able to resist attack (Couatts and Dolezal, 1965), and if an attack is successful, the trees will more than likely be culled during the cutting process. If a tree is attacked by *S. noctilio* in the early Fall, and then harvested prior to signs of attack, it is unlikely that the larvae will be able to complete the 1 to 2 year life-cycle before the tree is disposed of through various collection and recycling methods. Because there is a low proportion of pine Christmas trees produced in New York, a low likelihood of attack on well-managed, healthy trees, a low chance of survival of undetected larvae, and a rapid decline in the marketability of a tree following successful *S. noctilio* attack, we considered the potential risk for artificial spread of *S. noctilio* through the movement of pine Christmas trees from the areas where detection has occurred (and the bordering counties) to areas where no detection has occurred to be **Low**.

Roots

Sirex noctilio prefers to attack the bole of the pine tree, with over 90 percent of emergence reported from breast height and up to three inches in diameter at the tree top (Morgan and Stewart, 1966). The eggs and larvae infest the sapwood layer before moving deeper into the tree (Madden, 1988). There are no reports of *S. noctilio* larvae infesting the roots of pine trees. The roots of pine trees, like stumps, are not removed from the forests at the time of timber harvest. Consequently, we considered the potential risk for artificial spread of *S. noctilio* through the movement of pine roots from the areas where detection has occurred (and the bordering counties) to areas where no detection has occurred, to be **Nil**.

Branches and Debris

Sirex noctilio can emerge from pine branches, as small as one inch in diameter, on standing trees near the trunk. Low numbers are able to complete development in small diameter material, with *S. noctilio* development and emergence occurring when the diameter of the infested pine logs is three inches or greater (Morgan and Stewart, 1966). Larger green pine branches and logging residue are attractive and susceptible to ovipositing females (Neumann and Minko, 1981). Green pine logs are most attractive and susceptible to oviposition by *S. noctilio* 5 to 7 days after felling, with attractiveness lasting up to four weeks (Taylor, 1981). It is a common practice to leave the branches in the forest when pine trees are harvested in New York. These branches, in some instances, may be chipped on site for the utilization in power generating facilities (Crawford, pers. comm. 2006). Due to the low movement of pine branches and logging debris from forests, the reduced ability of *S. noctilio* to develop within smaller diameter material, and the potential risk for artificial spread of *S. noctilio* through the movement of pine branches from the areas where detection has occurred (and the bordering counties) to areas where no detection has occurred, we considered the artificial spread of *S. noctilio* through the movement of branches and debris to be **Nil**. If *S. noctilio* is detected in areas where pine branches and other logging debris are removed from the forest, there could be a **Low** potential risk for artificial spread through the movement of pine branches larger than one inch in diameter.

VI. Control Options

Eradication

In Pittwater, Tasmania (1952), an infestation of *S. noctilio* was discovered; however, attempts to eradicate this pest were unsuccessful (Taylor, 1981). The discovery of the pest in Victoria, Australia (1961) initiated the establishment of the National *Sirex* Fund and program in 1962. The program involved an extensive “search and destroy” effort against *S. noctilio*, in addition to multidisciplinary research into management of the pest (Taylor, 1981). The ability of *S. noctilio* to establish and maintain low population levels in moribund branches supported the belief that it would be difficult to eradicate it as a pest (Morgan and Stewart, 1966). The report of the *Sirex noctilio* Science Advisory Panel supported the conclusion that eradication was not a feasible option (APHIS, 2006b).

Chemical Control

The use of insecticides against the adult stage of *S. noctilio* is not feasible because of the adult's short life span the adult's tendency to not feed (making a contact insecticide necessary), the lack of known effective compounds; and the potential impacts on non-target organisms.

Cultural Control

Through good silvicultural practices, the effects of *S. noctilio* can be reduced and more effectively managed (Neumann *et al.*, 1987). Healthy stands that vigorously grow are less susceptible to attack from *S. noctilio* (Haugen *et al.*, 1990). The following recommended measures help to minimize *S. noctilio* outbreaks: 1) timely, selective thinning of forests to reduce overcrowding and tree stress, with increased importance placed on the removal of suppressed, deformed, or multi-stemmed trees and trees dying or diseased; limiting high pruning and non-commercial thinning activities during months when the wasp is active; good site selection, including adequate soil type, soil drainage, and avoidance of steep slopes; and minimizing injury to trees from fire (and other forestry treatments), and the rapid removal of trees damaged by natural events, such as wind, hail, lightning strikes or snow (Neumann *et al.*, 1987). These practices are similar to the management recommendations for the southern pine beetle, *Dendroctonus frontalis*, which promote the thinning of forests from below (to reduce competition) and the removal of high-hazard, damaged or weakened trees (Hyland, 1994).

Biological Control

There are several biological control options available for the management of *S. noctilio*, with the most widespread, well-recognized being the parasitic nematode, *B. siricidicola* (Bedding and Iede, 2005). Currently, *B. siricidicola* is utilized in the management of *S. noctilio* in New Zealand, Australia, South Africa, Brazil, Uruguay, Argentina, and Chile, saving millions of dollars in pine timber (Bedding and Iede, 2005). The biology and use of *B. siricidicola*, which has an insect parasitic stage and a free living mycetophagous stage, has been extensively studied and reported (Bedding, 1968; Bedding, 1972; Bedding and Akhurst, 1974; Akhurst, 1975; Bedding and Iede, 2005). The successful development and refinement of methods for mass rearing, long term storage and inoculation of *B. siricidicola* through the years are encouraging for management of *S. noctilio* (Bedding and Iede, 2005).

Seven North American species of wasps in the family Ibalidae, which are parasitic on Siricidae, have been identified; two of these species are *Ibalia anceps* and *I. leucospoides* found in the eastern United States (Smith and Schiff, 2002). In Australia, the rearing and release of several different parasitoids for *S. noctilio* management has been conducted, with varying effectiveness (Neumann and Minko, 1981). *Ibalia leucospoides*, reported to occur in New York (Smith and Schiff, 2002), has a life-cycle in near synchrony with its host, rapidly disperses long distances; it is the most effective parasitoid in Victoria (Neumann and Minko, 1981). The parasitic wasps, *Megarhyssa nortoni*, *Rhyssa hoferi*, and *Schlettererius cinctipes*, have also been recommended for release as biological control agents in Australia to manage *S. noctilio*; these species have been found on other species of *Sirex* in North America as well (USDA, 1992). Additional parasites of Siricids native to the United States include *Rhyssa howdenorum*, *R. lineolata*, and *R. alaskensis* (USDA, 1992).

VII. Risk Mitigation Options

To reduce the risk of artificial movement of *S. noctilio*, we recommend the following risk mitigation options:

Quarantine

No movement of untreated pine logs, untreated pine utility poles, untreated pine products used in the manufacturing of log homes, untreated sawn pine lumber, untreated pine firewood, and untreated pine branches and logging debris larger than 1 inch in diameter should be permitted from the counties where *S. noctilio* has been detected and the bordering counties.

Fumigation

For importing Monterey pine, *Pinus radiata*, and logs and lumber from Chile and New Zealand into the United States there is a code of federal regulations, Title 7 Chapter III Part 319.40-5, which outlines the mandatory steps and procedures necessary to allow importation. Logs and any regulated wood packing material to be used with the logs during shipment to the United States must be fumigated in accordance with Section 319.40-7 (f)(1) within 45 days following the date of felling, and prior to the arrival of the logs in the United States.

In the USDA-APHIS-PPQ Treatment Manual, the fumigation treatments listed for use on Siricidae (woodwasps) on wood products, including containers, are treatment schedules T404-b-1-1, and T404-b-1-2; the regulated wood packing material fumigation treatment is T404-e-1 (APHIS, 2005a).

Heat Treatment – Kiln Sterilization

In the USDA-APHIS-PPQ Treatment Manual, the kiln sterilization heat treatment listed for use on Siricidae (woodwasps) on wood products, including containers, is treatment schedule T404-b-4; the regulated wood packing material heat treatment schedule is T404-e-2 (APHIS, 2005a).

The heat requirement to kill *Sirex* listed in a Pest Risk Assessment on the importation of unprocessed logs and chips of eucalypt from Australia (USDA, 2003) is a core temperature of 65°C for two hours.

Drying wood to moisture content below 20 percent oven-dried weight reduces the ability of *A. areolatum* to survive and grow within the wood, and kills the eggs and young larvae of *S. noctilio* (Coutts and Dolezal, 1965).

Chipping

The chipping of wood products into pieces 0.625 inch (5/8”) at the thickest point or smaller eliminates all life stages of the Asian Longhorned Beetle (ALB), *Anoplophora glabripennis*, and the Emerald Ash Borer (EAB), *Agrilus planipennis* (APHIS, 2003a; APHIS, 2003b). Currently, chipping is an eradication treatment in the ALB and EAB quarantine areas.

Chemical Treatment of Immature Stages

Inorganic borate salt is a preservative treatment of wood against decaying fungi and wood destroying insects, such as the wood boring beetles in the families Lyctidae, Anobiidae, and Cerambycidae. Depending on the size of the material, borate salt can be applied through dipping or pressure treatment. Vacuum pressure treatment, followed by kiln drying to a moisture content below 18 percent, is currently being utilized by some log home manufacturers in New York. The combination of the two treatments may be effective in eliminating the immature stages of *S. noctilio* within the wood; however, testing and validation of treatment efficacy needs to be conducted.

Timing of Movement

Due to the limited trap catch and observation data currently available for *S. noctilio* in the United States, implementation of this option is under development. As additional data and information become available on the timing of adult emergence, this option may be utilized.

The movement of untreated pine logs, pine utility poles, and pine products used in the manufacturing of log homes during the periods of the year when adult emergence of *S. noctilio* is not likely to occur (15 October –June 15 in New York (*estimated*)), would be possible if all materials are treated or converted into a processed product (30 days) before adult emergence. This is not an option for firewood, as it is not a treated or processed product. Examples of treatment are: fumigation, heat treatment (kiln drying), and chipping. Examples of processed products are: pressure treated wood (pressure treated and chemically impregnated), creosote impregnated lumber, wood impregnated with creosote, and wood impregnated with lubricants (PPQ 578, 2005). Larvae of *S. noctilio* contained within the wood are killed prior to their emergence as adults by the treatments or processing.

The movement of untreated pine logs, pine utility poles, and large pine branches (especially if felled within 45 days) through counties where detection has occurred (and the bordering counties) during periods when adult *S. noctilio* are active (June 15- October 15 in New York estimated) are prohibited and/or regulated. During this period, materials could be attractive to ovipositing females, which could become attached to the wood, and be moved long distances.

The timing of movement risk mitigation option is similar to the regulations currently used on Pine Shoot Beetle, in which articles can or cannot move without treatment during certain periods of the year in relation to the risk presented by the biology of the pest. If *Sirex noctilio* is detected in other areas of the United States, the timing of movement would have to be adjusted to ensure that movement takes place only when adult emergence is not possible.

VIII. Conclusions

Sirex noctilio, a wood boring pest of pines, is a high risk invasive species with the potential to cause serious economic damage to nearly all regions in the continental United States where pines are grown. *Sirex noctilio* is capable of dispersing long distances, both naturally and through the artificial movement of infested materials, such as logs. Eradication of *S. noctilio* is not feasible; therefore, regulations and management are necessary to slow its movement and prevent the occurrence of outbreaks resulting in the loss of pine timber. Effective management of *S. noctilio* involves an integrated approach: restricting the movement of untreated materials during periods

of the year when risk of spread is greatest; monitoring infestation through aerial survey, ground survey and traps trees; good silvicultural practices; and using biological control agents, like *B. siricidicola*, when, and if, it is available.

IX. Recommendations

We recommend that strategies similar to those currently being used for the management of *S. noctilio* in Australia (Haugen *et al.*, 1990) and several other Southern Hemisphere countries, be implemented for the management of *S. noctilio* in the United States. These strategies are necessary to prevent the potentially serious economic damage observed when *S. noctilio* outbreaks occur.

To effectively implement the strategies for management, it is necessary to determine the extent and the current levels of *S. noctilio* infestation in the United States. This can be accomplished through grid trapping and aerial/ground surveys of suspected pine forests. It is necessary to prevent the artificial spread of *S. noctilio* in potentially infested materials through the treatment and regulated movement of materials. We recommend a restriction in the movement of untreated pine logs, including untreated pine utility poles, untreated pine logs and lumber used in the manufacturing of log houses, untreated green pine lumber (SWPM), firewood, and nursery stock larger than four inches in diameter at the base from the counties where *S. noctilio* has been detected (and the bordering counties). These materials are considered to pose the highest risk for the artificial movement of *S. noctilio* based on the current distribution.

To reduce the risk associated with the movement of these materials, we recommend utilizing risk mitigation options, such as fumigation, heat treatment, chipping and timing of the material movement. The promotion of education on effective silvicultural management practices is important in reducing a stand's susceptibility to *S. noctilio*. The overall success of the program for *S. noctilio* management may rely on the ability to utilize the parasitic nematode, *B. siricidicola*, as a biological control agent when, and if, the nematode is approved for release in the United States. In other countries, the use of *B. siricidicola* in the management of *S. noctilio* has dramatically increased the effectiveness of the program. However, it is important to realize that this is an ongoing program, and there is a continual need for monitoring *S. noctilio* populations, rearing re-isolation, and inoculation of the nematode. The goal of the *S. noctilio* management strategy is to prevent the occurrence of population outbreaks, and reduce the rate of its potential to spread.

X. Future Research Needs

10.1 Determine the Effects of Competitive Interactions between *S. noctilio*, Native Bark Beetles, and Native Siricids

Research should be conducted to determine the nature of the interaction between *S. noctilio*, native bark beetles, and native Siricidae species.

Research areas:

- Can *S. noctilio* compete with/displace native species?
- What effect(s) does competition have on *S. noctilio* reproduction, survival and host location?
- Will *S. noctilio* cause minimal forest damage in the United States, as it does in Europe and north Africa?
- Will it become a major forest pest as observed in the Southern Hemisphere?

10.2 Determine the Effects of Native Biological Control Agents on *S. noctilio* in North America

Research should be conducted to determine the ability and effectiveness of native Siricidae parasitoids to locate and utilize *S. noctilio* as a host. The large number of native parasitoids present in the United States may be useful in maintaining low populations of *S. noctilio*.

10.3 Information Needed on the Ability to Visually Detect *S. noctilio* and Periods of Adult Wasp Emergence

The ability to effectively detect *S. noctilio* presence in wood through visual inspection in forests and mills is unknown at this time. Better detection of infested materials early in the production chain would reduce the potential for artificial movement. Increased knowledge regarding the phenological development of *S. noctilio* in the United States would allow for better predictive modeling methods of adult wasp emergence. This information is useful in determining periods of the year when material movement would be possible without the risk of adult emergence.

XI. Authors & Reviewers

Authors:

Daniel Borchert: USDA-APHIS-PPQ-CPHST-PERAL

Glenn Fowler: USDA-APHIS-PPQ-CPHST-PERAL

Lisa Jackson: USDA-APHIS-PPQ-CPHST

Reviewers:

Lawrence Brown: USDA-APHIS-PPQ-CPHST-PERAL

James Smith: USDA-APHIS-PPQ-CPHST- PERAL

Melissa Fabiano: USDA-APHIS-PPQ-CPHST- PERAL

Leon Bunce: USDA-APHIS-PPQ-ER

Brian Kopper: USDA-APHIS-PPQ-ER

Sloane Crawford: NYS DEC

XII. Acknowledgements

This pest risk analysis (PRA) is based on a PRA process developed by Mike Firko and Joe Cavey, USDA-APHIS, for the evaluation of arthropod pests. The PRA process was expanded by Polly Lehtonen, APHIS, to evaluate pest plants (weeds), and was further modified for arthropod pests by Ken Lakin and Glenn Fowler (USDA-PPQ-APHIS-CPHST-PERAL).

Thanks to Charles Thayer for assistance in reference preparation.

XIII. References Cited

- About Inc. 2005. *Pinus taeda*: loblolly pine. <http://forestry.about.com/library/tree/bllobp.htm>
- Adams, D.L. 1995. The northern rocky mountain region. pp. 387-440. In Barrett, J.W. (ed). Regional silviculture of the United States, 3rd edition. John Wiley and Sons, Inc. New York. 643 pp.
- Akhurst, R.J. 1975. Cross-breeding to facilitate the identification of *Deladenus* spp., nematode parasites of woodwasps. *Nematologica* 21:267-272.
- APHIS. 2006a. Sirex Science Advisory Panel Report. January 9-10. Annapolis MD, <http://www.dec.state.ny.us/website/dlf/privland/forprot/health/panel.pdf>. Accessed 5/20/2006
- APHIS, 2006b. Emergency and Domestic Programs. *Sirex noctilio* updates http://www.aphis.usda.gov/ppq/ep/emerging_pests/sirex/updates.html. Accessed Date 4/17/2006.
- APHIS. 2005a. Treatment manual. http://www.aphis.usda.gov/ppq/manuals/port/pdf_files/Treatment_Chapters/05-05-T400-5.pdf Accessed Date 4/3/2006.
- APHIS. 2005b. Phytosanitary issues management. Processed products. PPQ form 578. http://www.aphis.usda.gov/ppq/pim/exports/processed_products.html Accessed date. 4/26/2006
- APHIS, 2003a. Asian longhorned beetle cooperative eradication program. Hudson County, New Jersey. Environmental assessment.
- APHIS, 2003b. Emerald ash borer cooperative eradication program. Lucas County Ohio. Environmental assessment. USDA-MRPBS-APHIS.
- Arnett, R. H., Jr., 1985. American Insects: A Handbook of the Insects of America North of Mexico. Van Nostrand Reinhold Co., New York. 850 pp.
- Backyardgardener. 2006. Africa plant hardiness zones. <http://www.backyardgardener.com/zone/africa.html>. Date accessed 4/3/2006.
- Bedding, R.A. 1968. *Deladenus wilsoni* n. sp. and *D. siricidicola* n. sp. (Neotylenchidae), entomophagous-mycetophagous nematodes parasitic in siricid woodwasps. *Nematologica*. 14: 515-525.
- Bedding, R.A. 1972. Biology of *Deladenus siricidicola* (Neotylenchidae) an entomophagous-mycetophagous nematodes parasitic in siricid woodwasps. *Nematologica*. 18: 482-493.
- Bedding, R.A., and R.J. Akhurst. 1974. Use of the nematode, *Deladenus siricidicola* in the biological control of *Sirex Noctilio* in Australia. *J. Aust. Ent. Soc.* 13:129-135

- Bedding, R.A. and E.T. Iede. 2005. Application of *Bedding siricidicola* for Sirex Woodwasp Control. Nematodes as Biocontrol Agents. CPC Publishing. Cambridge pp 385- 399.
- Benson, R. B. 1962. Holarctic sawflies (Hymenoptera: Symphyta). Bulletin of the British Museum (Natural History) Entomology. Vol. 12 No. 8 pp. 379-409.
- Benson, R.B. 1943. Studies in Siricidae especially in Europe and southern Asia (Hymenoptera: Symphyta). Bull. Ent Res. London. 34. 27-51.
- Borchert, D.M. and R.D. Magarey. 2005. A guide to the use of NAPPFAST. <http://www.nappfast.org/usermanual/nappfast-manual.pdf> 32.pp. Date accessed 7/5/2006.
- CFR. 2006. Code of Federal Regulations. 7 CFR 319.40-5 b.
- CFIA. 2006. *Sirex noctilio* reported detections in Ontario Canada. http://www.nappo.org/annualmtg/AnnualMtgHighlights/01_Tuesday/Greg%20Stubbings_NAPP_O_2006.pdf
- CPC. 2005. Crop Protection Compendium, 2005 ed. Wallingford, U.K.: CAB International [CD-ROM].
- Carnegie, A. J., Matsuki, M. Haugen, D.A., Hurley, B.P., Ahumada R., Klasmer, P., Sun, J., Iede, E.T., 2006. Predicting the potential distribution of *Sirex noctilio* (Hymenoptera: Siricidae), a significant exotic pest of Pinus plantations. Ann. For. Sci. 63 (2006) 119-128.
- Carnegie, A. J., R.H. Eldridge and D.G. Waterson. 2005. History and management of Sirex woodwasp in pine plantations in New South Wales, Australia. New Zealand Journal of Forestry Science 35(1): 3-24
- Chandler, L. 1959. Home Building Speed-Up Helps Make Horntail A Pest in Structural Timbers. Pest Control, June 1959.
- Chrystal, R.N. 1928. The Sirex Wood-Wasps and Their Importance in Forestry. Bull. Ent. Research Vol XIX Part 3: 219-245
- Ciesla, W.M. 2003. European Woodwasp: A Potential Threat to North America's Conifer Forests. Journal of Forestry, March 2003.
- Cinnamon, Mark. 2005. Nursery Program Manager, Illinois Department of Agriculture. Personal communication with Lisa Jackson APHIS. 10/05.
- Christmas Tree. 2006. Christmas tree farmers association of New York Inc. Website. <http://www.christmastreesny.org/resources/treetypes.asp> Date Accessed 4/24/2006.

Clark A.F. 1932. Insects Infesting *Pinus Radiata* in New Zealand. The N.Z. Journal of Science and Technology. Feb. 1932: 235-243.

Conifer Specialist Group 1998. *Pinus radiata*. In: IUCN 2006. 2006 IUCN Red List of Threatened Species. <www.iucnredlist.org>. Downloaded on **06 July 2006**.

Coutts, M.P. 1969a. The mechanism of pathogenicity of *Sirex noctilio* on *Pinus radiata*. I Effects of the symbiotic fungus *Amylosterum* sp. (Thelophoraceae). Aust. J. Biol. Sci. 22:915-924.

Coutts, M.P. 1969b. The mechanism of pathogenicity of *Sirex noctilio* on *Pinus radiata*. I Effects of *S. noctilio* mucus. Aust. J. Biol. Sci. 22:1153-1161.

Coutts, M.P. and J.E. Dolezal. 1965. *Sirex noctilio*- its associated fungus and some aspects of wood moisture content. Aust. For Res 1: 3-13.

Crawford, S. 2006a. NYS Department Environmental Conservation Division of Lumber Products. Email communication with Daniel Borchert APHIS. 5/11/2006

Crawford, S. 2006. Personal communication with Daniel Borchert APHIS. Syracuse New York.

Crawford, S. 2005. New York State Industrial Timber Harvest Production and Consumption Report-2004. NYS Department of Environmental Conservation Forest Products Utilization Program. Albany NY.

Dawson, I. 1991. Plant hardiness zones for Australia.
<http://www.anbg.gov.au.hort.research.html>. Date accessed 4/3/2006.

Earle, C.J. and M.P. Frankis. 1999. *Pinus radiata* D. Don.
<http://www.conifers.org/pi/pin/radiata.htm> Date accessed 4/3/2006.

IPPC. 1994. Chapter VII. Plant oleoresins. *In* International trade in non-wood forest products: An overview. FAO Corporate Document Repository. Forestry Department.
<http://www.fao.org/docrep/x5326e/x5326e0a.htm>.

IPPC. 1995. Chapter 1. Production trade and markets. *In* Gum naval stores: Turpentine and rosin from pine resin. FAO Corporate Document Repository. Forestry Department.
<http://www.fao.org/docrep/V6460E/v6460e06.htm>.

FIA, 2002. Forest Inventory Analysis, timber products output mapmaker version 1.0.
http://ncrs2.fs.fed.us/4801/fiadb/rpa_table/WebClass_rpa_table.ASP Date accessed 4/19/2006.

Fisher, R.C. 1955. Australian quarantine and the *Sirex* wood-wasp problem: Investigations in the United Kingdom. Timber Technology. Volume 63. No. 2195. September. pp. 480-483.

- Fong, L.K., and R.K. Crowden. 1976. Preliminary studies on the mucus secretion of the woodwasp *Sirex noctilio* F. I. Physico-chemical and biochemical properties. Aust. J. Biol. Sci. 29:21-32.
- Haugen, D.A., and E.R. Hoebeke. 2005. Pest Alert: Sirex woodwasp – *Sirex noctilio* F. (Hymenoptera: Siricidae). USDA Forest Service NA-PR-07-05.
- Haugen, D.A., R.A. Bedding, M.G. Underdown, F.G. Neumann. 1990. National Strategy for Control of Sirex Noctilio in Australia. Special Liftout Section No. 13.
- Helms, J.A. 1995. The California region. pp. 441-497. In Barrett, J.W. (ed). Regional silviculture of the United States, 3rd edition. John Wiley and Sons, Inc. New York. 643 pp.
- Helmsing, P. 2004. 2002 Ag Census. American Christmas Tree Journal. July: 56-57.
- Hoebeke, E.R., D.A. Haugen, and R.A. Haack 2005. *Sirex noctilio*: Discovery of a Palearctic siricid woodwasp in New York. Newsletter of the Michigan Entomological Society 50(1&2):24-25.
- Huftalen, R. 2006. President McIntosh Box and Pallet. New York. Personal communication with Daniel Borchert APHIS. 4/26/2006.
- Hyland, J. 1994. Southern pine beetle management. http://www.forestry.state.al.us/publication/TF_publications/forest_health/southern_pine_beetle_management.htm Date Accessed 4/26/2006.
- Iede, E.T., S.R.C. Penteado, and E.G. Schaitza. 1998. *Sirex noctilio* problem in Brazil: Detection, evaluation and control. In Proceedings, training and control of *Sirex noctilio* with natural enemies. 45-52. FHTET 98-13. Fort Collins, CO: USDA Forest Service, Forest Health Technology Enterprise Team.
- Jacobs, Susan. 2005. Global Logistics Team Leader, Pinova, A division of Hercules Incorporated.
- Kelley, S. 1998. Hardiness zone map of Europe. <http://www.uk.gardenweb.com/forums/zones/hze>. Date Accessed 7/5/2006.
- King, J.C. 1965. Some aspects of the biology of the fungal symbiont of *Sirex noctilio*. Aust. J. Bot., 1966, 25-30
- Koelling, M.R.; Hart, J.B. and Leefers, L. 1992. Christmas tree production in Michigan. Michigan State University, Agricultural Experiment Station, East Lansing, Michigan, USA. Special Report 61.

LDAF. 2000. Quarterly report of forest products data: 1955 to current report. Baton Rouge, LA: Office of Marketing. Louisiana Department of Agriculture and Forestry
<http://www.ldaf.state.la.us/divisions/forestry/reports/quarterreport/original-report.asp>

Liddle wonder. 2002. New Zealand plant hardiness zones.
<http://www.liddlewonder.co.nz/zones.php>. Date accessed 4/3/2006

Madden, J.L. 1974. Oviposition behaviour of the woodwasp *Sirex noctilio* F. Aust. J. Zool. 22. 341-351.

Madden, J.L. 1988. *Sirex* in Australia. In: Berryman, A.A. [Ed]. Dynamics of Forest Insect Populations: Patterns, Causes, Implications. pp. 407-429. Plenum Press. NY.

Madden, J.L. 1981. Egg and larval development in the woodwasp *Sirex noctilio* F. Aust. J. Zool. 29. 493-506.

Michigan Ag Connection. 2005. Michigan Christmas tree inventory results.
<http://www.michiganagconnection.com/story-state.cfm?Id=291&yr=2005>

Middlekauff, W.W. 1960. The Siricid wood wasps of California (Hymenoptera: Symphyta). Bulletin of the California Insect Survey. 59-75

Morgan, F.D. and N.C. Stewart. 1966. The biology and behavior of the woodwasp *Sirex noctilio* F. in New Zealand. Trans. roy. Soc. N. Z. Zool., Vol. 7 No 14. pp 195- 204.

Morgan, F.D. 1968. Bionomics of Siricidae. Ann. Rev Entomol. 13. 239-56.

NAPPO. 2006. Update on the *Sirex* woodwasp (*Sirex noctilio*) distribution in Canada- New records from Ontario. Date posted 3/24/2006. <http://www.pestalert.org/> Accessed date 4/17/2006.

Neumann, F.G., and G. Minko. 1981. The *Sirex* woodwasp in Australian radiata pine plantations. Aust. For., 1981, 44: 44-63.

Neumann, F.G., J.L. Morey and R.J. McKimm. 1987. The *Sirex* wasp in Victoria. Bull. 29. pp.1-50.

Olsen, R. 2005. Marketing Representative. Connecticut Department of Agriculture. Hartford, CT.

PIN. 2005. Port Interception Network. USDA-APHIS. Riverdale, Maryland.

Pinova, 2005. PinovaTM, products lines and processes.
<http://www.herc.com/pinova/products/html>.

Plantdeas.com. 2005. South and Central America plant hardiness zones.
<http://www.plantideas.com/zone/sazone.html>. Date accessed 4/3/2006.

Prestemon, J.; Pye, J.; Barbour, J.; Smith, G.R.; Ince, P.; Steppleton, C. and Xu, W. 2005. U.S. wood using mill locations. USDA Forest Service. Southern Research Station.

Rawlings G.B. and Wilson N.W. *Sirex Noctilio* as a beneficial and destructive insect to *Pinus radiata* in New Zealand

Rawlings, G.B. 1953. Rearing of *Sirex noctilio* and its parasite *Ibalia leucospoides*. N. Z. For. Ser. For. Res. Notes. 1 (8).

Simpson. R.F., and R.M. McQuilkin. 1976. Identification of volatiles from felled *Pinus radiata* and the electroantennograms they elicit from *Sirex noctilio*. Entomol. Exp. And Appl. 19: 205-213.

Sirex cyaneus, 2002. Blue horntail (woodwasp).
www.forestry.ubc.ca/fetch21/FRST308/lab7/sirex_cyaneus/horntail.html. Published on the internet, 1 p.

Sirex juvencus, 2002. *Sirex juvencus* L. (common *Sirex* or Bouvillon).
http://www.xylophages.com/html/fiche_sirex.html Published on the internet, 2 pp.

Skog, K. and Risbrudt, C. 1982. Trends in economic scarcity of U.S. timber commodities. USDA Forest Service. Forest Products Laboratory. Madison, Wisconsin. 24 pp.

Slippers, B., T.A. Coutinho, B.D. Wingfield and M.J. Wingfield. 2003. A review of the genus *Amylostereum* and its association with woodwasps. South African Journal of Science. 99 January/February 70-74.

Smith, D.R., N.M. Schiff. 2002. A Review of the Siricid woodwasps and their Ibaliid parasitoids (Hymenoptera:Siricidae, Ibaliiidae) in the Eastern United States, with emphasis on the Mid-Atlantic Region. Proc. Entomol. Soc. Wash. 104 (1), 2002, pp. 174-194.

Smith, J. W., 2004. Pathway assessment: wood-boring insects emergence from dead wood.

Spadberry, J.P., and A.A. Kirk. 1978. Aspects of the ecology of siricid woodwasps (Hymenoptera: Siricidae) in Europe, North Africa and Turkey with special reference to the biological control of *Sirex noctilio* F. in Australia. Bull. Ent. Res. 68, 341-359.

Spradbery, J.P. 1973. A comparative study of the phytotoxic effects of siricid woodwasps on conifers. Ann. Appl. Biol. 75, 309-320.

Spradbery, J.P. 1977. The oviposition biology of siricid woodwasps in Europe. Ecological Entomology (1977) 2, 225-230.

Storts, Dave. 2005. Production Manager for Container and Field. Berryhill Nurseries of Ohio, a Division of Monrovia Growers. Springfield, Ohio. Personal communication with Lisa Jackson APHIS. 10/05.

Taylor, K.L. 1981. The ecology of pests: some Australian case histories. The sirex woodwasp: ecology and control of an introduced forest insect. CSIRO Melbourne. pp. 231: 247.

Tesch, S.D. 1995. The pacific northwest region. pp. 499-558. In Barrett, J.W. (ed). Regional silviculture of the United States, 3rd edition. John Wiley and Sons, Inc. New York. 643 pp.

U Conn. 2007. *Pinus mugo* information page. University of Connecticut Plant Database. <http://www.hort.uconn.edu/plants/p/pinmug/pinmug1.html>

UFL. No Date. Florida forest trees: Loblolly pine (*Pinus taeda*). University of Florida. University of Florida Institute of Food and Agricultural Sciences. Florida 4-H Forest Ecology. <http://www.sfrc.ufl.edu/4h/index.html>

USDA. 1992. Pest Risk Assessment of the Importation of *Pinus radiata* and Douglas-fir Logs from New Zealand. Misc. Pub. No. 1508.

USDA-ARS. 1990. USDA plant hardiness zone map. USDA Agricultural Research Service. Miscellaneous Publication Number 1475. Washington, D.C.

USDA-ERS. 1990. Greenhouse/Nursery Statistical Bulletin. USDA Economic Research Service.

USDA-NASS. 2004. Nursery crops 2003 summary. USDA National Agricultural Statistics Service.

USDA-NASS. 2002. Census of Agriculture. National Agricultural Statistics Service. <http://www.usda.gov/nass/>

USDA-USFS. 1991. Forest Density. USDA United States Forest Service. Southern Forest Experiment Station, Forest Inventory and Analysis Research Unit.

USDA-USFS. 1998. Proceedings of a Conference: Training in the control of *Sirex noctilio* by the Use of Natural Enemies. Forestry Health Technology Enterprise Team. 98-13.

USDA-USFS. 2001. United States timber industry: an assessment of timber product output and use, 1996. Johnson, T.G. (ed.). USDA United States Forest Service. Southern Research Station. Asheville, N.C.

USDA-USFS. 2003a. Pest risk assessment of the importation into the United States of unprocessed logs and chips of eighteen eucalypt species from Australia. FPL GTR 137.

USDA-USFS. 2003b. Southern Forest Resource Assessment Final Technical Report. U.S. Forest Service Southern Research Station and Southern Region, U.S. Environmental Protection Agency,

U.S. Fish and Wildlife Service, the Tennessee Valley Authority and the National Association of State Foresters. USDA United States Forest Service.

<http://www.srs.fs.usda.gov/sustain/report/htm>

USDA-USFS. 2005. Regional areas of the forest service. USDA United States Forest Service. <http://www.fs.fed.us/contactus/regions.shtml>

USDC. 1999a. Logging. 1997 Economic Census. Manufacturing industry series. U.S. Department of Commerce. Economics and Statistics Administration. U.S. Census Bureau.

USDC. 1999b. Sawmills. 1997 Economic Census. Manufacturing industry series. U.S. Department of Commerce. Economics and Statistics Administration. U.S. Census Bureau.

USDC. 1999c. Softwood veneer and plywood manufacturing. 1997 Economic Census. Manufacturing industry series. U.S. Department of Commerce. Economics and Statistics Administration. U.S. Census Bureau.

Vasiliauskas, R., J. Stenlid, and I.M. Thomsen. 1998. Clonality and genetic variation in *Amylostereum areolatum* and *A. chailletii* from northern Europe. *New Phytologist* 139, 751-758.

Walker, L.C. 1995. The southern pine region. pp. 271-333. *In* Barrett, J.W. (ed). Regional silviculture of the United States, 3rd edition. John Wiley and Sons, Inc. New York. 643 pp.

Wharton, E. 2005. Forest resource use. Northeastern forest inventory and analysis. <http://www.fs.fed.us/ne/fia/studies/FRU/index.html#fuel> . Date accessed 5/11/2006.

Appendix B. Dicamba and Triclopyr Toxicity Data

Table B-1. Toxicity Data for Triclopyr (Triclopyr Triethylamine Salt and Triclopyr Acid).

Chemical	Test Species	Test Type	Toxicity Value
Triethylamine Salt	Northern Bobwhite (<i>Colinus virginianus</i>)	Acute LD ₅₀	NA
		Subacute LC ₅₀	11,622 ppm
		Chronic NOEL	NA
	Mallard (<i>Anas platyrhynchos</i>)	Acute LD ₅₀	2055 mg ai/kg
		Subacute LC ₅₀	>10,000 ppm
		Chronic NOEL	NA
	Rainbow Trout (<i>Onchorynchus mykiss</i>)	96 h LC ₅₀	240 ppm*
	Bluegill Sunfish (<i>Lepomis macrochirus</i>)	96 h LC ₅₀	471 ppm*
	Waterflea (<i>Daphnia magna</i>)	48 h EC ₅₀	1496 ppm
	Rat (<i>Rattus norvegicus</i>)	Acute LD ₅₀	1847 mg/kg
	Rat (<i>Rattus norvegicus</i>)	Subchronic 13 wk feeding study NOEL	5 mg/kg/day
	Rat (<i>Rattus norvegicus</i>)	Developmental NOEL Maternal toxicity NOEL	100 mg/kg
Rabbit (New Zealand White)	Developmental NOEL Maternal toxicity NOEL	30 mg/kg	
Triclopyr Acid	Northern Bobwhite (<i>Colinus virginianus</i>)	Acute LD ₅₀	NA
		Subacute LC ₅₀	2934 ppm
		Chronic NOEL	500 ppm
	Mallard (<i>Anas platyrhynchos</i>)	Acute LD ₅₀	1698 mg ai/kg
		Subacute LC ₅₀	5620 ppm
		Chronic NOEL	100 ppm
	Rainbow Trout (<i>Onchorynchus mykiss</i>)	96 h LC ₅₀	117 ppm
	Bluegill Sunfish (<i>Lepomis macrochirus</i>)	96 h LC ₅₀	148 ppm
	Waterflea (<i>Daphnia magna</i>)	48 h EC ₅₀	132.9 ppm
	Honeybee (<i>Apis mellifera</i>)	48 h Acute Contact LD ₅₀	> 100 ug/bee
Rat (<i>Rattus norvegicus</i>)	Acute LD ₅₀	630 mg/kg	

* formulated product; ai = active ingredient; NOEL = No Observable Effect Level; ppm = parts per million.
Source: EPA Office of Pesticide Programs. 1998. Reregistration eligibility decision: Triclopyr. EPA 738-R-98-011.

Table B–2. Toxicity Data for Dicamba (Dimethylamine Salt and Dicamba Acid).

Chemical	Test Species	Test Type	Toxicity Value
Dimethylamine Salt	Northern Bobwhite (<i>Colinus virginianus</i>)	Acute LD ₅₀	>2453 mg ai/kg
		Subacute LC ₅₀	>4533 ppm
		Chronic NOEL	NA
	Mallard (<i>Anas platyrhynchos</i>)	Acute LD ₅₀	NA
		Subacute LC ₅₀	>4533 ppm
Chronic NOEL		NA	
Rainbow Trout (<i>Onchorynchus mykiss</i>)	96 h LC ₅₀	1000 ppm	
Bluegill Sunfish (<i>Lepomis macrochirus</i>)	96 h LC ₅₀	1000 ppm	
Waterflea (<i>Daphnia magna</i>)	48 h EC ₅₀	1600 ppm	
Dicamba Acid	Northern Bobwhite (<i>Colinus virginianus</i>)	Acute LD ₅₀	188 mg ai/kg
		Subacute LC ₅₀	>10,000 ppm
		Chronic NOEL	1600 ppm
	Mallard (<i>Anas platyrhynchos</i>)	Acute LD ₅₀	1373 mg ai/kg
		Subacute LC ₅₀	>10,000 ppm
		Chronic NOEL	800 ppm
	Rainbow Trout (<i>Onchorynchus mykiss</i>)	96 h LC ₅₀	28 ppm
	Bluegill Sunfish (<i>Lepomis macrochirus</i>)	96 h LC ₅₀	135 ppm
	Waterflea (<i>Daphnia magna</i>)	48 h EC ₅₀	110 ppm
	Honeybee (<i>Apis mellifera</i>)	48 h Acute Contact LD ₅₀	90.65 µg/bee
	Rat (<i>Rattus norvegicus</i>)	Acute LD ₅₀	2,740 mg/kg/day
	Rat (<i>Rattus norvegicus</i>)	Subchronic 13 wk feeding study NOEL	500 mg ai/kg/day
Rat (<i>Rattus norvegicus</i>)	Developmental NOEL Maternal toxicity NOEL	>400 mg/kg/day 160 mg/kg/day	
Rabbit (New Zealand White)	Developmental NOEL Maternal toxicity NOEL	150 mg ai/kg/day 30 mg ai/kg/day	

ai = active ingredient; NOEL = No Observable Effect Level; ppm = parts per million.

Source: EPA Office of Pesticide Programs. 2005. Environmental Fate and Ecological Risk Assessment for the Reregistration of Dicamba and Dicamba Sodium, Potassium, Diglycoamine, Dimethylamine and Isopropylamine salts.