





Figure 2-1. Areas Closed Year-round to Bottom Trawling

NOTE: Very limited state-managed bottom trawling occurs in some of the depicted areas. Beam trawling for shrimp is allowed in southeast Alaska, Prince William Sound, and the Kodiak area, although effort is extremely low.












Figure 2-4. Alternative 2: Essential Fish Habitat for Late Juvenile and Adult Pacific Cod




Figure 2-5. Alternative 2: Essential Fish Habitat for Golden King Crab Eggs




Figure 2-6. Alternative 2: Essential Fish Habitat for Mature and Late Juvenile Golden King Crab




Figure 2-7. Alternative 2: Essential Fish Habitat for GOA Adult and Late Juvenile Weathervane Scallops





Figure 2-8. Alternative 2: Essential Fish Habitat for BSAI Adult and Late Juvenile Weathervane Scallops

Figure 2-9. Alternative 2: Essential Fish Habitat for Freshwater Salmon in Alaska



Source: EFH EA, June 1998






Figure 2-10. Alternative 3: Essential Fish Habitat for GOA Late Juvenile and Adult Pacific Ocean Perch








Figure 2-12. Alternative 3: Essential Fish Habitat for BSAI Late Juvenile and Adult Pacific Cod

Chapter 2 Final EFH EIS – April 2005




Figure 2-13. Alternative 3: Essential Fish Habitat for BSAI Golden King Crab




Figure 2-14. Alternative 3: Essential Fish Habitat for Weathervane Scallops

Chapter 2 Final EFH EIS – April 2005




Figure 2-15. Alternative 3: Essential Fish Habitat for Freshwater Chinook Salmon

Chapter 2 Final EFH EIS – April 2005




Figure 2-16. Alternative 3: Essential Fish Habitat for Marine Chinook Salmon