

Supplemental Table A

Incidence of non-neoplastic histopathological changes at 2 weeks in male and female B6C3F1 mouse liver by various compounds treatment

Treatment Group	Control		Oxazepam				PNT	ONT				Eug	Meug	Acet
Dose			125 ppm		2500 ppm		5000 ppm	1250 ppm		5000 ppm		75 mg/kg/day	75 mg/kg/day	6000 ppm
Sex	M	F	M	F	M	F	F	F	M	F	F	F	F	F
Total Animals	6	5	6	6	6	6	6	6	6	6	6	6	6	6
Infiltration Cellular, Mixed														
<i>minimal</i>	2	3	1	4	4	4	4	3	1	5	6	5	4	
<i>mild</i>												1		
Extramedullary hematopoiesis, granulocyte							1							
Granulomatous Inflammation														
<i>minimal</i>									1					
Hemorrhagic necrosis/ intraction							1							
Hypertrophy, centrilobular														
<i>minimal</i>			4											
<i>mild</i>					1	1								
<i>moderate</i>					2	2								
<i>marked</i>					3	3								
Necrosis, centrilobular														
<i>minimal</i>														1
Necrosis, zonal							1							
Necrosis, focal														
<i>minimal</i>	2						1							
<i>mild</i>														1
Single Cell Necrosis														
<i>minimal</i>					1									
<i>mild</i>							1							
Increased hepatocellular mitosis												1		

PNT: p-nitrotoluene, ONT: o-nitrotoluene, Acet: Acetaminophen, Eug: Eugenol, Meug: methyleugenol

M: Male, F: Female

