For more information

Contact the Oregon State University Extension forester who serves your county, or contact an Oregon or federal forester.

OSU Extension, Curry County 29390 Ellensburg (Hwy 101) Gold Beach, OR 97444 541-247-6672 or 1-800-356-3986 http://extension.oregonstate.edu/curry/

Oregon Department of Forestry http://egov.oregon.gov/ODF/

USDA Forest Service, Pacific Northwest Region http://www.fs.fed.us/r6/nr/fid/

Oregon Department of Agriculture http://egov.oregon.gov/ODA/PLANT/ path_sod_index.shtml

Informational websites about SOD:

OSU Extension Service http://extension.oregonstate.edu/ emergency/oak_death.php

California Oak Mortality Task Force http://nature.berkeley.edu/comtf/


USDA Agricultural and Plant Health Inspection Service (APHIS) http://www.aphis.usda.gov/plant_health/











You can help STOP the spread

People can spread *Phytophthora ramorum* across long distances by moving infected plants either purchased at a nursery or collected in the wild, or by moving infected wood, leaves, stems, or soil.

If you live, work, recreate, or travel in the quarantined portion of Curry County, Oregon (see map, over):

- Become familiar with the most recent regulations related to Sudden Oak Death (see websites, in "For more information").
- DO NOT collect and remove host plants or plant parts from the forest (see http:// www.aphis.usda.gov/plant_health/plant_ pest_info/pram/index.shtml).
- DO NOT collect or remove soil.
- Stay on established trails, and respect any trail closures.





A dead tanoak.

nurseries.

An infected rhodo-

dendron: note dead

blackened leaf stems

flower bud and

iust below bud.

Before leaving infested areas:

have used in infested areas.

undercarriage of your vehicle.

horses' hooves, and pets' paws.

bleach solution for cleaning.

Buy healthy plants from reputable

• Clean soil off shoes, mountain bikes,

• For maximum protection, use a 10%

• Clean and disinfect equipment (saws,

shovels, pruning equipment, etc.) you

• Wash soil off tires, wheel wells, and the

If you travel or work in any of the 14 quarantined counties in coastal California (see map, over), follow these same rules. Don't bring *Phytophthora*

ramorum into Oregon!

Oregon forests and plant nurseries — and the people who depend on them!

Help protect



Bleeding trunk on an infected tanoak.

EC 1608-E • June 2007

Where is Sudden Oak Death?



Area in Curry County, Oregon under state and federal quarantine regulation for Sudden Oak Death, as of June 2007.



California and Oregon coun ties with documented cases of Sudden Oak Death, as of June 2007.

© 2007 Oregon State University. This publication may be photocopied or reprinted in its entirety for noncommercial purposes. Produced and distributed in furtherance of the Acts of Congress of May 8 and June 30, 1914. Extension work is a cooperative program of Oregon State University, the U.S. Department of Agriculture, and Oregon counties. Oregon State University Extension Service offers educational programs, activities, and materials without discrimination based on age, color, disability, gender identity or expression, marital status, national origin, race, religion, sex, sexual orientation, or veteran's status. Oregon State University Extension Service is an Equal Opportunity Employer.

Important points to know **Sudden Oak Death**

Tanoak killed by Sudden Oak Death.

Sudden Oak Death is the common name for a disease caused by *Phytophthora ramorum*, a previously unknown and presumably nonnative pathogen. At this time, no one knows where the pathogen came from or how it was introduced into Oregon.

Phytophthora ramorum has killed more than 1 million oak and tanoak trees in 14 coastal counties in California and several hundred tanoaks in Curry County, Oregon. It also causes branch and shoot dieback and leaf spotting on a large number of woodland and nursery plant species.

The pathogen has a wide host range including tanoak, California black oak, Douglas-fir, grand fir, coast redwood, Pacific madrone, Pacific rhododendron, evergreen huckleberry, and many other tree and shrub species common in Oregon and Washington forests. Hosts in the nursery trade include varieties of rhododendron, camellia, and *Pieris*. A complete, current host list is at http://www.aphis.usda.gov/plant_health/plant_pest_info/pram/index.shtml

P. ramorum spreads naturally when clouds and rain move spores within forest canopies—from treetops to stems and shrubs below, or across landscapes from treetop to treetop. Artificial (human-assisted) spread occurs when people transport infected plants or plant parts or infested soil. The pathogen survives in infested plant material, litter, soil, and water.

State and federal inspectors survey forests and nurseries in Oregon regularly to detect the disease. Infected plants and adjacent host plants are destroyed; areas affected are monitored until the pathogen is no longer present.

State and federal quarantine regulations are in place to minimize the risk of new infections and

prevent human-assisted spread. Complete texts of these regulations (ORS 603-052-1230 and 7 CFR 301.92) are on the Oregon Department of Agriculture and the U.S. Department of Agriculture websites (see over).



Tanoak infection (left) is revealed in the inner bark, where dark cankers indicate areas killed by P. ramorum.



Infected Douglas-fir branch tip (above) and an infected huckleberry branch (left).