

# Ohio, 2005

## Primary and secondary syphilis

- CASES: Ohio reported 211 cases in 2005.
- RATES: Ohio ranked 24 among 50 states + DC + 3 territories: 1.8 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

## Male-female rates of primary and secondary syphilis

- In Ohio, the rate among males was 3.1 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 0.7 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

## Racial disparities of primary and secondary syphilis


In Ohio, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 7.1 among blacks, 1.6 among Hispanics, 0.6 among Asian/Pacific Islanders, and 3.5 among American Indians/Alaska Natives.

- The rate among blacks was 6.5 times that of whites.
- The rate among Hispanics was 1.5 times that of whites.
- The rate among American Indians/Alaska Natives was 3.2 times that of whites.

## Congenital syphilis

- CASES: Ohio reported 2 cases in 2005.
- RATES: Ohio ranked 28 among 28 states + 2 territories reporting at least one congenital syphilis case: 1.3 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Ohio, 1996-2005



## Total P&S Syphilis Rates by County, Ohio, 2005

