

## Guidance on Lisinopril; Hydrochlorothiazide

This guidance represents the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the Office of Generic Drugs.

**Active ingredient:** Lisinopril; Hydrochlorothiazide

**Form/Route:** Tablet/Oral

**Recommended studies:** 2 studies

1. Type of study: Fasting  
Design: Single-dose, two-treatment, two-period crossover *in-vivo*  
Strength: 25 mg/20 mg  
Subjects: Normal healthy males and females, general population  
Additional Comments: Female subjects enrolled in the BE studies should not be pregnant, and if applicable, should practice abstention or contraception during the study.

---

2. Type of study: Fed  
Design: Single-dose, two-treatment, two-period crossover *in-vivo*  
Strength: 25 mg/20 mg  
Subjects: Normal healthy males and females, general population  
Additional comments: Please see comment above.

---

**Analytes to measure:** Lisinopril and Hydrochlorothiazide in plasma.

**Bioequivalence based on (90% CI):** Lisinopril and Hydrochlorothiazide

**Waiver request of in-vivo testing:** 12.5/10 mg and 12.5 mg/20 mg based on (i) acceptable bioequivalence studies on the 25 mg/20 mg strength, (ii) proportional similarity of the formulations across all strengths, and (iii) acceptable in vitro dissolution testing of all strengths.

**Dissolution test method and sampling times:**

Please note that a **Dissolution Methods Database** is available to the public at the OGD website at <http://www.fda.gov/cder/ogd/index.htm>. Please find the dissolution information for this product at this website. Please conduct comparative dissolution testing on 12 dosage units each of all strengths of the test and reference products. Specifications will be determined upon review of the application.