

Draft Guidance on Topiramate

This draft guidance, once finalized, will represent the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the Office of Generic Drugs.

Active ingredient: Topiramate

Form/Route: Sprinkle Capsule /Oral

Recommended studies: 2 studies

1. Type of study: Fasting
Design: Single-dose, two-treatment, two-period crossover *in-vivo*
Strength: 25 mg
Subjects: Normal healthy males and females, general population
Additional Comments: Females must have a negative baseline pregnancy test within 24 hours prior to receiving the drug. Females should not be pregnant or lactating, and if applicable, should practice abstinence or contraception during the study.

2. Type of study: Fed
Design: Single-dose, two-treatment, two-period crossover *in-vivo*
Strength: 25 mg
Subjects: Normal healthy males and females, general population
Additional comments: Please see comments above.

Analytes to measure (in appropriate biological fluid): Topiramate in plasma.

Bioequivalence based on (90% CI): Topiramate

Waiver request of in-vivo testing: 15 mg based on (i) acceptable bioequivalence studies on the 25 mg strength, (ii) proportional similarity of the formulations across all strengths, and (iii) acceptable in vitro dissolution testing of all strengths.

Dissolution test method and sampling times:

Please note that a **Dissolution Methods Database** is available to the public at the OGD website at <http://www.fda.gov/cder/ogd/index.htm>. Please find the dissolution information for this product at this website. Please conduct comparative dissolution testing on 12 dosage units each of all strengths of the test and reference products. Specifications will be determined upon review of the application.