

Table 187. Persons 18 Years and Over With Selected Diseases and Conditions by Selected Characteristics: 2005

[In thousands (217,774 represents 217,774,000). Based on National Health Interview Survey, a sample survey of the civilian noninstitutionalized population; see Appendix III]

Selected characteristic	Total persons	Persons with selected diseases and conditions					
		Diabetes ^{1, 2}	Ulcers ¹	Kidney disease ^{3, 4}	Liver disease ³	Arthritis diagnosis ⁵	Chronic joint symptoms ⁶
Total^{7, 8}	217,774	16,186	15,104	3,791	2,965	46,941	58,863
SEX⁸							
Male	104,919	7,896	6,913	1,736	1,571	18,260	25,967
Female	112,855	8,290	8,190	2,055	1,395	28,681	32,896
AGE							
18 to 44 years old	110,431	2,640	4,787	931	738	8,624	17,161
45 to 64 years old	72,296	7,595	6,008	1,305	1,721	20,780	25,426
65 to 74 years old	18,446	3,421	2,100	675	321	8,609	8,313
75 years old and over	16,600	2,529	2,209	880	185	8,927	7,963
RACE							
Race alone ^{8, 9}	215,349	15,982	14,916	3,696	2,912	46,288	58,006
White	180,477	12,886	13,042	2,967	2,484	40,351	50,803
Black or African American	24,817	2,513	1,327	570	314	4,718	5,593
American Indian or Alaska Native	1,469	116	148	¹⁰ 38	¹⁰ 33	310	420
Asian	8,155	437	349	¹⁰ 104	¹⁰ 81	868	1,138
Native Hawaiian or other Pacific Islander	431	(B)	(B)	(B)	—	¹⁰ (B)	53
Two or more races ^{8, 11}	2,425	204	188	¹⁰ 95	¹⁰ 53	653	857
HISPANIC ORIGIN AND RACE^{8, 12}							
Hispanic or Latino	27,770	1,898	1,311	563	386	3,194	4,729
Mexican or Mexican American	17,163	1,242	745	351	213	1,677	2,631

— Represents zero. B Figure too small to meet statistical standards for reliability of a derived figure. ¹ Respondents were asked if they had ever been told by a health professional that they had an ulcer or diabetes. A person may be represented in more than one column. ² Excludes borderline diabetes. ³ Respondents were asked if they had been told in the last 12 months by a health professional that they had weak or failing kidneys or any kind of liver condition. ⁴ Excludes kidney stones, bladder infections, or incontinence. ⁵ Respondents were asked if they had ever been told by a health professional that they had some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia. ⁶ Respondents with joint symptoms that began more than 3 months ago prior to interview. Excludes back and neck. ⁷ Total includes other races not shown separately. ⁸ Estimates are age adjusted to the year 2000 standards using four age groups: 18–44 years, 45–64 years, 65–74 years, and 75 years and over. ⁹ Refers to persons who indicated only a single race group. ¹⁰ Figures do not meet standard of reliability or precision. ¹¹ Refers to all persons who indicated more than one race group. ¹² Persons of Hispanic or Latino origin may be of any race or combination of races.

Source: U.S. National Center for Health Statistics, *Vital and Health Statistics, Series 10, Number 232, Summary Health Statistics for U.S. Adults: National Health Interview Survey, 2005*; <<http://www.cdc.gov/nchs/data/series/sr10/sr10232.pdf>>.