

Section 2

Description of the Sample

This section describes the sample design and selection, the method of estimation, the sampling variability of the estimates, and the methodology of computing confidence intervals.

Domain of Study

The statistics in this report are estimates from a probability sample of unaudited Individual Income Tax Returns, Forms 1040, 1040A, and 1040EZ (including electronic returns) filed by U.S. citizens and residents during Calendar Year 2005.

All returns processed during 2005 were subjected to sampling except tentative and amended returns. Tentative returns were not subjected to sampling because the revised returns may have been sampled later, while amended returns were excluded because the original returns had already been subjected to sampling. A small percentage of returns were not identified as tentative or amended until after sampling. These returns, along with those that contained no income information, were excluded in calculating estimates. This resulted in a small difference between the population total (133,189,982 returns) reported in Table C and the estimated total of all returns (132,226,042) reported in other tables.

The estimates in this report are intended to represent all returns filed for Tax Year 2004. While most of the returns processed during Calendar Year 2005 were for Tax Year 2004, the remaining returns were mostly for prior years, and a few for non-

calendar years ending during 2005 and 2006. Returns for prior years were used in place of 2004 returns received and processed after December 31, 2005. This was done based on the assumption that the characteristics of returns due, but not yet processed, can best be represented by the returns for previous income years that were processed in 2005.

Sample Design and Selection

The sample design is a stratified probability sample, in which the population of tax returns is classified into subpopulations, called strata, and a sample is randomly selected independently from each stratum. Strata are defined by:

1. Nontaxable (including no alternative minimum tax) with adjusted gross income or expanded income of \$200,000 or more.
2. High combined business and farm total receipts of \$50,000,000 or more.
3. Presence or absence of special Forms or Schedules (Form 2555, Form 1116, Form 1040 Schedule C, and Form 1040 Schedule F).
4. Indexed positive or negative income. Sixty variables are used to derive positive and negative incomes. These positive and negative income classes are deflated using the Chain-Type Price Index for the Gross Domestic

Valerie Testa, and Jana Scali designed the sample and prepared the text and tables in this section under the direction of Yahia Ahmed, Chief, Mathematical Statistics Section, Statistical Computing Branch.

Product to represent a base year of 1991. (See footnote 1 for details.)

5. Potential usefulness of the return for tax policy modeling. Thirty-two variables are used to determine how useful the return is for tax modeling purposes.

Table C shows the population and sample count for each stratum after collapsing some strata with the same sampling rates. (See references 1 and 2 for details.) The sampling rates range from 0.05 percent to 100 percent.

Tax data processed to the IRS Individual Master File at the Enterprise Computing Center at Martinsburg during Calendar Year 2005 were used to assign each taxpayer's record to the appropriate stratum and to determine whether or not the record should be included in the sample. Records are selected for the sample either if they possess certain combinations of the four ending digits of the social security number, or if their ending five digits of an eleven-digit number generated by a mathematical transformation of the SSN is less than or equal to the stratum sampling rate times 100,000. (See reference 3 for details.)

Data Capture and Cleaning

Data capture for the SOI sample begins with the designation of a sample of administrative records. While the sample was being selected, the process was continually monitored for sample selection and data collection errors. In addition, a small subsample of returns was selected and independently reviewed, analyzed, and processed for a quality evaluation.

The administrative data and controlling information for each record designated for this sample was loaded onto an online database at the Cincinnati Submission Processing Center. Computer data for the selected administrative records were then used to identify inconsistencies, questionable values, and missing values as well as any additional variables that an editor needed to extract for each record. The editors use a hardcopy of the taxpayer's return to enter the required information onto the online system.

After the completion of service center review, data were further validated, tested, and balanced.

Adjustments and imputations for selected fields based on prior year data and other available information were used to make each record internally consistent. Finally, prior to publication, all statistics and tables were reviewed for accuracy and reasonableness in light of provisions of the tax law, taxpayer reporting variations and limitations, economic conditions, and comparability with other statistical series.

Some returns designated for the sample were not available for SOI processing because other areas of IRS needed the return at the same time. For Tax Year 2004, 0.03 percent of the sample returns were unavailable.

Method of Estimation

Weights were obtained by dividing the population count of returns in a stratum by the number of sample returns for that stratum. The weights were adjusted to correct for misclassified returns. These weights were applied to the sample data to produce all of the estimates in this report.

Sampling Variability and Confidence Intervals

The sample used in this study is one of a large number of samples that could have been selected using the same sample design. The estimates calculated from these different samples would vary. The standard error (SE) of an estimate is a measure of the variation among the estimates from the possible samples and, thus, is a measure of the precision with which an estimate from a particular sample approximates the average of the estimates calculated from all possible samples.

The standard error may be expressed as a percentage of the value being estimated. This ratio is called the coefficient of variation (CV). Tables 1.4 CV, 2.1 CV, and 3.3 CV contain estimated CV's for the estimates included in Tables 1.4, 2.1, and 3.3 of this report.

The sample estimate and an estimate of its standard error permit the construction of interval estimates with prescribed confidence that the interval includes the population value. If all possible samples were selected under essentially the same conditions and an estimate and its estimated standard error were calculated from each sample,

then:

1. About 68 percent of the intervals from one standard error below the estimate to one standard error above the estimate would include the population value. This is a 68 percent confidence interval.
2. About 95 percent of the intervals from two standard errors below the estimate to two standard errors above the estimate would include the population value. This is a 95 percent confidence interval.

For example, from Table 1.4, the estimate for State Income Tax Refunds, X , is \$22.736 billion, and its related coefficient of variation, $CV(X)$, is 0.86 percent. The standard error of the estimate, $SE(X)$, needed to construct the confidence interval estimate, is:

$$\begin{aligned} SE(X) &= X \cdot CV(X) \\ &= (\$22.736 \times 10^9) \cdot (0.0086) \\ &= \$0.196 \text{ billion} \end{aligned}$$

The p percent confidence interval is calculated using the formula:

$$X \pm z \cdot SE(X)$$

where z takes the value 1, 2, or 3 when p is 68, 95, or 99, respectively. Based on these data, the 68 percent confidence interval is from \$22.540 billion to \$22.932 billion, the 95 percent confidence interval is from \$22.344 billion to \$23.128 billion, and the 99 percent confidence interval is from \$22.148 billion to \$20.324 billion.

Table Presentation

Whenever a weighted frequency is less than 3, the estimate and its corresponding amount are combined or deleted in order to avoid disclosure of information for specific taxpayers. (The combined or deleted data, if any, are included in the corresponding column totals.) These combinations and deletions are indicated by a double asterisk (**). Estimates based on less than 10 sampled returns are considered to be unreliable. These estimates are noted by a single asterisk (*) to the

left of the data unless all of the sampled returns are selected with certainty (at the 100 percent rate).

In the tables, a dash (-) in place of a frequency or an amount indicates that either no returns in the population had the characteristic or the characteristic was so rare that it did not appear on any of the sampled returns.

Footnote

- [1] Indexing of positive and negative income is done by dividing each by the ratio of the Chain-Type Price Index for the Gross Domestic Product for the fourth quarter of 2003 to the fourth quarter of the base year of 1991. The indices were calculated using the Gross Domestic Product (GDP) Chain-type Price Index found in the table titles "Quantity and Price Indexes for Gross Domestic Product" released to the public on November 30, 2004 on the BEA web site (<http://www.bea.doc.gov/>).

References

- [1] Hostetter, S., Czajka, J. L., Schirm, A. L., and O'Connor, K. (1990), "Choosing the Appropriate Income Classifier for Economic Tax Modeling," in *Proceedings of the Section on Survey Research Methods*, American Statistical Association, 419-424.
- [2] Schirm, A. L., and Czajka, J. L. (1991), "Alternative Designs for a Cross-Sectional Sample of Individual Tax Returns: the Old and the New," *Proceedings of the Section on Survey Research Methods*, American Statistical Association, 163-168.
- [3] Harte, J.M. (1986), "Some Mathematical and Statistical Aspects of the Transformed Taxpayer Identification Number: A Sample Selection Tool Used at IRS," *Proceedings of the Section on Survey Research Methods*, American Statistical Association, 603-608.

