

Table 44. Tuberculosis Cases and Percentages in Persons Completing Therapy for Whom Therapy Was Indicated for One Year or Less: Reporting Areas, 1999–2003¹

Reporting Area	Year									
	1999		2000		2001		2002		2003	
	No. ²	(%) ³	No. ²	(%) ³	No. ²	(%) ³	No. ²	(%) ³	No. ²	(%) ³
United States	15,208	79.9	14,259	(80.9)	14,061	(80.8)	13,271	(81.4)	13,205	(81.4)
Alabama	260	(80.4)	263	(88.2)	224	(79.5)	203	(89.7)	235	(85.1)
Alaska	59	(88.1)	104	(90.4)	50	(88.0)	45	(91.1)	52	(90.4)
Arizona	231	(82.3)	228	(78.1)	254	(81.9)	225	(79.1)	260	(79.2)
Arkansas	156	(82.7)	176	(84.1)	138	(92.0)	114	(86.8)	116	(87.1)
California	3,183	(79.8)	2,931	(79.5)	2,969	(80.1)	2,820	(81.8)	2,876	(79.9)
Colorado	80	(88.8)	83	(92.8)	128	(93.0)	96	(84.4)	102	(93.1)
Connecticut	108	(78.7)	90	(68.9)	113	(70.8)	90	(70.0)	101	(73.3)
Delaware	30	(80.0)	23	(78.3)	30	(93.3)	23	(91.3)	28	(85.7)
District of Columbia	57	(84.2)	71	(80.3)	63	(85.7)	77	(97.4)	68	(91.2)
Florida	1,075	(84.0)	1,004	(82.7)	1,018	(80.8)	945	(81.6)	922	(85.4)
Georgia	593	(77.6)	599	(80.6)	507	(81.3)	464	(78.2)	484	(77.9)
Hawaii	168	(67.9)	119	(73.1)	138	(73.9)	137	(69.3)	99	(76.8)
Idaho	14	(71.4)	13	(53.8)	7	(71.4)	12	(58.3)	13	(84.6)
Illinois	700	(81.3)	636	(83.3)	636	(77.8)	583	(76.3)	567	(80.4)
Indiana	133	(79.7)	119	(89.9)	97	(92.8)	107	(96.3)	118	(95.8)
Iowa	54	(83.3)	38	(89.5)	36	(77.8)	33	(81.8)	36	(91.7)
Kansas	62	(80.6)	70	(82.9)	70	(72.9)	76	(80.3)	65	(76.9)
Kentucky	177	(80.2)	117	(88.0)	131	(87.8)	129	(85.3)	120	(85.0)
Louisiana	310	(77.4)	285	(77.5)	259	(74.5)	207	(81.6)	219	(79.9)
Maine	19	(94.7)	22	(72.7)	19	(94.7)	18	(77.8)	23	(69.6)
Maryland	258	(86.0)	255	(83.5)	235	(88.9)	265	(86.4)	242	(86.8)
Massachusetts	248	(80.6)	261	(83.9)	250	(77.6)	255	(76.9)	247	(76.5)
Michigan	293	(83.6)	248	(83.1)	275	(81.1)	263	(81.4)	211	(83.4)
Minnesota	184	(87.5)	172	(84.9)	221	(84.6)	220	(85.5)	201	(88.6)
Mississippi	187	(84.5)	153	(83.7)	137	(83.9)	113	(83.2)	112	(76.8)
Missouri	174	(79.3)	183	(81.4)	138	(84.8)	114	(86.0)	107	(71.0)
Montana	10	(90.0)	19	(89.5)	17	(94.1)	10	(90.0)	5	(100.0)
Nebraska	18	(55.6)	22	(63.6)	31	(83.9)	24	(66.7)	26	(69.2)
Nevada	81	(77.8)	91	(84.6)	87	(80.5)	71	(97.2)	96	(91.7)
New Hampshire	17	(94.1)	20	(80.0)	19	(89.5)	18	(88.9)	14	(85.7)
New Jersey	490	(71.8)	480	(74.2)	458	(75.5)	475	(78.1)	456	(80.0)
New Mexico	48	(87.5)	38	(71.1)	46	(82.6)	50	(78.0)	37	(86.5)
New York City	1,252	(78.8)	1,122	(83.9)	1,067	(84.9)	927	(85.5)	998	(85.6)
New York State ⁴	332	(75.6)	358	(79.6)	375	(81.6)	318	(76.4)	301	(81.1)
North Carolina	431	(88.6)	391	(90.3)	340	(88.5)	369	(90.5)	324	(89.2)
North Dakota	7	(57.1)	5	(100.0)	5	(100.0)	6	(100.0)	4	(25.0)
Ohio	262	(71.4)	298	(73.8)	258	(79.1)	219	(80.4)	194	(88.7)
Oklahoma	180	(73.9)	128	(82.8)	159	(80.5)	169	(84.6)	137	(69.3)
Oregon	114	(78.1)	110	(80.9)	113	(87.6)	104	(86.5)	99	(81.8)
Pennsylvania	370	(76.2)	337	(74.2)	280	(76.4)	306	(73.9)	290	(79.3)
Rhode Island	44	(68.2)	43	(74.4)	60	(73.3)	45	(68.9)	40	(90.0)
South Carolina	264	(69.7)	245	(76.3)	227	(85.0)	222	(80.2)	234	(85.5)
South Dakota	17	(76.5)	12	(75.0)	12	(66.7)	9	(88.9)	15	(73.3)
Tennessee	319	(84.0)	322	(80.4)	265	(84.2)	276	(83.7)	238	(79.8)
Texas	1,441	(79.8)	1,323	(78.9)	1,443	(75.8)	1,361	(75.6)	1,411	(74.7)
Utah	34	(73.5)	41	(73.2)	33	(81.8)	29	(96.6)	34	(97.1)
Vermont	2	(50.0)	2	(100.0)	7	(100.0)	8	(87.5)	8	(100.0)
Virginia	292	(85.6)	254	(86.2)	272	(80.9)	289	(86.5)	304	(84.9)
Washington	235	(81.3)	233	(82.0)	241	(84.6)	236	(83.5)	235	(80.9)
West Virginia	36	(69.4)	24	(50.0)	25	(64.0)	26	(65.4)	17	(70.6)
Wisconsin	96	(86.5)	75	(78.7)	75	(81.3)	67	(82.1)	61	(86.9)
Wyoming	3	(100.0)	3	(100.0)	3	(100.0)	3	(100.0)	3	(100.0)
American Samoa ⁵
Fed. States of Micronesia ⁵
Guam ⁵	65	(53.8)	47	(66.0)	56	(37.5)	79	(51.9)	58	(87.9)
Marshall Islands ⁵
N. Mariana Islands ⁵	62	(82.3)	72	(83.3)	50	(88.0)	51	(68.6)	42	(78.6)
Puerto Rico ⁵	148	(82.4)	128	(87.5)	92	(92.4)	93	(79.6)	90	(70.0)
Republic of Palau ⁵	0	(0.0)	0	(0.0)	0	(0.0)	0	(0.0)	7	(57.1)
U.S. Virgin Islands ⁵

¹Most recent year for which data are available.

²Total cases for which therapy less than 1 year indicated in persons alive at diagnosis, with an initial regimen of one or more drugs prescribed, who did not die during therapy.

³Percentage of total cases in persons who completed therapy for whom therapy less than 1 year was indicated.

⁴Excludes New York City.

⁵Not included in U.S. totals.

See Technical Notes for description of Completion of Therapy calculation (page 9).