

**Table 41. Tuberculosis Cases and Percentages by Completion of Tuberculosis Therapy (COT): Reporting Areas, 2004<sup>1</sup>**

Reporting Area	Total Cases	Therapy ≤1 Year Indicated <sup>2</sup>		Therapy >1 Year Indicated <sup>3</sup>		All Drug Therapy		
		No. <sup>4</sup>	COT ≤1 Year(%)	COT(%)	No. <sup>4</sup>	COT(%)	No. <sup>4</sup>	COT(%)
<b>United States</b>	<b>14,502</b>	<b>12802</b>	<b>(82.3)</b>	<b>(91.0)</b>	<b>335</b>	<b>(77.9)</b>	<b>13137</b>	<b>(90.7)</b>
Alabama	211	179	(89.9)	(96.6)	8	(100.0)	187	(96.8)
Alaska	43	38	(84.2)	(100.0)	0	...	38	(100.0)
Arizona	272	246	(74.8)	(85.4)	5	(80.0)	251	(85.3)
Arkansas	133	114	(88.6)	(93.0)	3	(66.7)	117	(92.3)
California	2,992	2639	(81.5)	(92.0)	79	(75.9)	2718	(91.6)
Colorado	128	116	(96.6)	(98.3)	4	(100.0)	120	(98.3)
Connecticut	101	85	(77.6)	(95.3)	3	(100.0)	88	(95.5)
Delaware	33	29	(93.1)	(93.1)	0	...	29	(93.1)
District of Columbia	81	66	(93.9)	(98.5)	1	(100.0)	67	(98.5)
Florida	1,074	928	(85.9)	(94.6)	18	(83.3)	946	(94.4)
Georgia	537	458	(83.2)	(91.7)	11	(90.9)	469	(91.7)
Hawaii	116	108	(79.6)	(88.9)	1	(100.0)	109	(89.0)
Idaho	11	10	(80.0)	(80.0)	0	...	10	(80.0)
Illinois	567	505	(79.6)	(93.3)	8	(62.5)	513	(92.8)
Indiana	128	111	(93.7)	(98.2)	0	...	111	(98.2)
Iowa	47	47	(76.6)	(91.5)	0	...	47	(91.5)
Kansas	62	57	(84.2)	(89.5)	1	(100.0)	58	(89.7)
Kentucky	127	110	(87.3)	(98.2)	2	(100.0)	112	(98.2)
Louisiana	248	231	--	--	1	--	232	--
Maine	20	18	(66.7)	(94.4)	0	...	18	(94.4)
Maryland	313	279	(90.7)	(97.1)	6	(100.0)	285	(97.2)
Massachusetts	283	256	(76.6)	(93.4)	5	(100.0)	261	(93.5)
Michigan	272	239	(87.0)	(93.7)	6	(83.3)	245	(93.5)
Minnesota	199	187	(90.9)	(95.7)	5	(80.0)	192	(95.3)
Mississippi	119	101	(83.2)	(94.1)	5	(80.0)	106	(93.4)
Missouri	127	111	(82.0)	(89.2)	1	(100.0)	112	(89.3)
Montana	15	14	(92.9)	(100.0)	0	...	14	(100.0)
Nebraska	39	37	(86.5)	(94.6)	0	...	37	(94.6)
Nevada	95	88	(90.9)	(93.2)	2	(100.0)	90	(93.3)
New Hampshire	24	21	(100.0)	(100.0)	2	(100.0)	23	(100.0)
New Jersey	482	429	(82.3)	(92.8)	14	(71.4)	443	(92.1)
New Mexico	42	33	(84.8)	(93.9)	2	(50.0)	35	(91.4)
New York State <sup>5</sup>	323	293	(86.3)	(97.3)	10	(80.0)	303	(96.7)
New York City	1,037	901	(84.5)	(90.9)	40	(62.5)	941	(89.7)
North Carolina	381	332	(92.5)	(96.7)	10	(80.0)	342	(96.2)
North Dakota	4	4	(100.0)	(100.0)	0	...	4	(100.0)
Ohio	219	190	(87.9)	(92.6)	3	(100.0)	193	(92.7)
Oklahoma	178	164	(80.5)	(95.1)	2	(100.0)	166	(95.2)
Oregon	106	97	(89.7)	(92.8)	0	...	97	(92.8)
Pennsylvania	328	280	--	--	8	--	288	--
Rhode Island	51	49	(85.7)	(91.8)	0	...	49	(91.8)
South Carolina	233	213	(82.6)	(90.6)	1	(100.0)	214	(90.7)
South Dakota	11	9	(44.4)	(88.9)	0	...	9	(88.9)
Tennessee	278	237	(85.7)	(93.2)	2	(100.0)	239	(93.3)
Texas	1,672	1472	--	--	49	--	1521	--
Utah	36	32	(96.9)	(100.0)	2	(100.0)	34	(100.0)
Vermont	6	6	(66.7)	(83.3)	0	...	6	(83.3)
Virginia	329	292	(88.4)	(92.1)	7	(85.7)	299	(92.0)
Washington	245	226	(88.1)	(95.6)	7	(85.7)	233	(95.3)
West Virginia	24	21	(81.0)	(100.0)	1	(100.0)	22	(100.0)
Wisconsin	95	89	(79.8)	(92.1)	0	...	89	(92.1)
Wyoming	5	5	(100.0)	(100.0)	0	...	5	(100.0)
American Samoa <sup>6</sup>	...	...	...	...	...	...	...	...
Fed. States of Micronesia <sup>6</sup>	...	...	...	...	...	...	...	...
Guam <sup>6</sup>	51	45	(84.4)	(86.7)	0	...	45	(86.7)
Marshall Islands <sup>6</sup>	...	...	...	...	...	...	...	...
N. Mariana Islands <sup>6</sup>	...	...	...	...	...	...	...	...
Puerto Rico <sup>6</sup>	123	89	(79.8)	(92.1)	1	(100.0)	90	(92.2)
Republic of Palau <sup>6</sup>	...	...	...	...	...	...	...	...
U.S. Virgin Islands <sup>6</sup>	...	...	...	...	...	...	...	...

<sup>1</sup>Most recent year for which data are available.

<sup>2</sup>Initial isolate susceptible to rifampin (n=9,699) or susceptibility unknown (n=157); culture negative (n=2,418); culture status unknown (n=656); age unknown (n=1).

<sup>3</sup>Initial isolate rifampin resistant, or pediatric patients (aged <15) with meningeal, bone, joint, or miliary disease.

<sup>4</sup>Number of cases in persons alive at diagnosis, with an initial regimen of one or more drugs prescribed, who did not die during therapy. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information reported for ≥90% of cases.

<sup>5</sup>Excludes New York City.

<sup>6</sup>Not included in U.S. totals. Data for 2006 are interim as of April 6, 2007.

**Note:** Ellipses indicate data not available. See Technical Notes for description of Completion of Therapy calculation (page 9).