Leading Causes of Death by Age Group, Hispanic Males- United States, 2004*

| | Age Group in Years | | | | | | | | | | | | | |
|------|--|--|---|-------------------------------------|------------------------------------|------------------------------------|------------------------------------|------------------------------------|--|-----------------------------------|--|--|--|--|
| Rank | 1-4 | 5-9 | 10-14 | 15-19 | 20-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65 + | 65-74 | 75-84 | 85 + | All Ages |
| 1 | Unintentional injuries 34.8% | Unintentional injuries 37.4% | Unintentional injuries 36.3% | Unintentional injuries 44.5% | Unintentional injuries 43.7% | Unintentional injuries 38.1% | Unintentional injuries 24.3% | Heart disease 18.6% | Cancer 27.3% | Heart disease 31.1% | Cancer 29.4% | Heart disease 31.1% | Heart disease 35.2% | Heart disease 21.9% |
| 2 | Birth defects 15.3% | Cancer 21.2% | Cancer 15.7% | Homicide 26.8% | Homicide 25.1% | Homicide 18.6% | Heart disease 12.0% | Cancer 18.5% | Heart disease 25.7% | Cancer 24.2% | Heart disease 28.7% | Cancer 24.5% | Cancer 15.2% | Cancer 19.0% |
| 3 | Homicide 8.4% | Birth defects 8.8% | Homicide 7.3% | Suicide 10.2% | Suicide 11.7% | Suicide 10.3% | Cancer 10.1% | Unintentional injuries 11.5% | Diabetes 6.5% | Stroke 6.6% | Diabetes 6.7% | Stroke 6.7% | Stroke 7.8% | Unintentional injuries 11.4% |
| 4 | Cancer 5.4% | Homicide 4.4% | Birth defects 6.5% | Cancer 5.7% | Cancer 4.2% | Cancer 5.9% | HIV disease 8.7% | Chronic liver disease 9.8% | Chronic liver disease 6.0% | Diabetes 5.6% | Stroke 5.7% | Diabetes 5.8% | Influenza & pneumonia 5.3% | Stroke 4.7% |
| 5 | Heart disease | Septicemia 2.2% | Suicide 6.3% | Heart disease 2.0% | Heart disease 2.5% | Heart disease 5.4% | Homicide 6.4% | HIV disease 5.3% | Unintentional injuries 5.1% | respiratory | Chronic lower respiratory diseases 3.4% | Chronic lower respiratory diseases 4.3% | Chronic lower respiratory diseases 5.2% | Diabetes 4.2% |
| 6 | Septicemia 2.6% | (6)Heart disease 1.8% | Heart disease 2.6% | Birth defects | (6)Birth defects 0.8% | HIV disease 3.7% | Chronic liver disease 6.3% | Stroke 4.2% | Stroke 4.9% | Influenza & pneumonia 3.2% | Chronic liver disease 3.0% | Influenza & pneumonia 3.2% | Alzheimer's disease 3.8% | Homicide 4.1% |
| 7 | Influenza & pneumonia 1.7% | (6)Influenza & pneumonia 1.8% | Influenza & pneumonia 1.3% | (7)Influenza & pneumonia 0.5% | (6)Legal intervention 0.8% | Stroke | Suicide 5.9% | Diabetes 3.8% | Kidney disease 1.9% | Kidney disease 2.3% | Unintentional injuries 2.6% | Kidney disease 2.3% | Diabetes 3.3% | Chronic liver disease 3.5% |
| 8 | Chronic lower respiratory diseases 1.3% | (8)Chronic lower respiratory diseases 1.1% | (8)Stroke 1.0% (8)Chronic | (7)Legal intervention 0.5% | HIV disease | Chronic liver disease 1.3% | Stroke 3.2% | Suicide 2.9% | HIV disease | Unintentional injuries 2.2% | Kidney disease 2.2% | Unintentional injuries 2.1% | Kidney disease 2.3% | Suicide 2.7% |
| 9 | Perinatal conditions 1.1% | (8)Perinatal conditions 1.1% | inatal tions % (8)Kidney diseases 1.0% (8)Kidney disease 1.0% cal & al care cations | ., . | Influenza & pneumonia 0.6% | Diabetes 0.8% | Diabetes 1.9% | Viral hepatitis 2.4% | Chronic lower respiratory diseases 1.6% | Alzheimer's disease 1.9% | Influenza & pneumonia 2.0% | Alzheimer's disease 1.9% | Unintentional injuries 1.9% | Chronic lower respiratory diseases 2.4% |
| 10 | Anemias 0.6% Stroke 0.6% | Benign neoplasms 0.7% Medical & surgical care complications 0.7% | | | Stroke 0.5% | Influenza & pneumonia 0.7% | Viral hepatitis 1.4% | Homicide 2.3% | Influenza & pneumonia 1.3% | Chronic liver disease 1.7% | Septicemia 1.4% | Chronic liver disease 1.5% | (10)Septi- cemia 1.5% (10)Hyper- tension 1.5% | Perinatal conditions 2.2% |

*Percentages represent total deaths in the age group due to the cause indicated. Numbers in parentheses indicate tied rankings. The white, black, American Indian/Alaska Native, and Asian/Pacific Islander race groups include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Some terms have been shortened from those used in the National Vital Statistics Report. See the next page for a listing of the shortened terms used in the table and their full, unabridged equivalents used in the report. To learn more, visit *Mortality Tables* at http://www.cdc.gov/nchs/datawh/statab/unpubd/mortabs.htm (HHS, CDC, NCHS).

Short and Full Terms for Leading Causes of Death

Some terms in the leading causes of death table have been shortened from those used in the National Vital Statistics Report. Below is a listing of the shortened terms used in the table and their full, unabridged equivalents used in the report.

| SHORT TERMS | FULL TERMS |
|---------------------------------------|--|
| Aortic aneurism | Aortic aneurism and dissection |
| Benign neoplasms | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior |
| Birth defects | Congenital malformations, deformations and chromosomal abnormalities |
| Bronchitis | Acute bronchitis and bronchiolitis |
| Cancer | Malignant neoplasms |
| Chronic liver disease | Chronic liver disease and cirrhosis |
| Diabetes | Diabetes mellitus |
| Gallbladder disorders | Cholethiasis and other disorders of gallbladder |
| Heart disease | Diseases of heart |
| HIV disease | Human immunodeficiency virus (HIV) disease |
| Homicide | Assault (homicide) |
| Hypertension | Essential (primary) hypertension and hypertensive renal disease |
| Kidney disease | Nephritis, nephrotic syndrome and nephrosis |
| Kidney infection | Infections of kidney |
| Medical & surgical care complications | Complications of medical and surgical care |
| Perinatal conditions | Certain conditions originating in the perinatal period |
| Pneumonitis | Pneumonitis due to solids and liquids |
| Pregnancy complications | Pregnancy, childbirth and the puerperium |
| Stroke | Cerebrovascular diseases |
| Suicide | Intentional self-harm |
| Unintentional injuries | Accidents (unintentional injuries) |
| | |