

Notes on literature.

Microbiology + Chemistry  
Genetics!

J. Lederberg  
Columbia University  
Yale University

Coleman, R. CRAS 216:616 1943 Action des rayons X  
sur la fréquence d'une mutation bactérienne.

S<sup>-</sup> to S<sup>+</sup>

Spont.  $5 \times 10^{-8}$

$\approx 5$  mins (~~70%~~  $\rho S = 1$ ) (75 000  $\mu$ !!!)  $60 \times 10^{-8}$

Cooper KE + D Woodman, JPB 58:75-84 (1946) The diffusion of antiseptics  
through gargles...  
Dept Law Med  
Dr. Bristol

$$m' = M_0 e^{\left(\frac{-x^2}{4Dt}\right)}$$

$x = \text{distance}$

$$c_{\text{conc}}(x) = m'$$



$D = \text{const.}$

$$\frac{dc}{dt} = D \frac{\partial^2 c}{\partial x^2}$$

Witkin, E.M., PNAS 34(3): 59-68 (1946) indicated differences in sensitivity to radiation in *Escherichia coli*.

u.v. - GE Hg lamp 2537 Å. Irradiated on petri plates. Colony counts at 24 hours.

B,  $5 \times 10^7$ , irradiated  $\bar{e}$  1000  $\text{ergs}/\text{mm}^2$ . At 24h. (nutrient agar) 4 colonies developed. One was propagated as B/r and proved to have a different resistance. All proved to be more resistant.



No other levels of resistance were found.

B/r is also X-ray resistant

at  $pS = 2$ , there are breaks in the killing curves of B only partially repl. by B/r.

lag B/r in broth is less; m.g.t. 19 mins. At 50  $\text{ergs}$ ,  $pS$  of B is 1; of B/r is 0. However, after 3 hours, the cells of B are elongated and undivided, of B/r  $\rightarrow$  100 cells.

A second irradiation of 700  $\text{ergs}$  will reduce each B/r microcolony leaving a representative but kill each undivided long cell of B. The effectiveness of the technique in mixtures of B and B/r indicate that the long cells behave like individual bacteria in sens. to radiation. With large samples, surviving colonies are tested for resistance by a test dose & elongation phenomenon. Delbrück analysis induced mutations are not detected.  $B \rightarrow B/r$   $10^{-5}$  generations.

The curve for B/r is a multiple hit curve.



Lurdegen, CC, PNAS, 32: 68-70 (1946) A new gene theory and an explanation of the phenomenon of dominance to ~~the~~ Mendelian segregation of the cytoplasm

chromosome = place of attachment for cytoplasm.

contaminated recessive = chromosome<sup>-</sup> cytoplasm<sup>+</sup>

Adsorption of cytoplasm on certain recessive loci.

$$F \times f \rightarrow \begin{matrix} F \\ f \\ f^{cont.} \end{matrix}$$

$$f \times f^{cont} \rightarrow 1:1 \text{ in most cases.}$$

Hooley!

Ferguson, T.B. + S.O. Thorne, Jr., J Pharm 86: 288-63 (1946) The effect of some amine compounds on the growth + respiration of E. coli. Dulse.

ATCC 6522 SG.

Amines:

1. 3-amino
- 2a 5-amino, 1,2,3,4 tetrahydro
- 2 2-dimethyl-7-amino
3. 5-amino
- 4 2,7-diamino
- 5 2,8-diamino.

Riboflavin had no effect.

Effect on oxidation of various substrates (glycose, pyr, lact, aspar, olive) is in different order (1,4,2,5,3) from growth (1...5)

% inhibition increases  $\bar{c}$  pH.

Caspe, S. + G. Cameron, ~~AE~~ JCCP 27:43-52 (1946) Effect of a  
respiratory enzyme system + creatine upon the growth of cells in vitro

diphthase ( ~~ADP~~ ) (FAD)

At  $10^{-6}$  ~~has~~ elicited response in tissue culture

do. creatine 50 mg%. only when unfitted.

together, synerg.

Hesley, A.D. J. Bact. 38: 563-78 (1939) Factors limiting bacterial growth.  
VII Respiration & growth properties of coli surviving sublethal temperatures

Waddell, Agnes H., Edinburgh Math. Notes, #35 Dec. 1945 Curves formed  
by colonies of microorganisms growing on a plane surface.

Mathematical analysis of outlines of conjoined colonies of sectors.

Wmslow-CE-A, G.R.B. 9:259-74 (1934)

Fells. 15 The role of certain ions on bacterial physiology. *A. reevis*. *Abstr. Bact.* 7: 33, 87, 133 (1923).

X

distilled water as good as NaCl for *E. coli*.

Whistler + OR Brock. J Bact 15: 235-43 (1927) The stability of various spp. of bacteria in aqueous suspensions.

Slant growth suspended in H<sub>2</sub>O & incubated 18-20 hours at 37°.

E coli highly resistant even when carefully washed. (high conc ca 10<sup>9</sup>)

10<sup>-3</sup> broth protects B. cereus from ~~being~~ death in saline.

N. rapid effects (1-2h.).

• 0.0145 M NaCl best medium for stability -

7.725 is toxic.

Only 5-10% killed in H<sub>2</sub>O in 9h.

20-40 x 10<sup>6</sup> conc.

85% = 8.5g/l = ca. 2 N.



Sherris, J. M. + H. B. Naylor, *Ageing & reproduction and the viability of young bacterial cells at low temperatures.* J. Bact 43:749 (1942)

Effects of certain mild agents (cold, low saline etc. are) greater on "young cells." During lag, bacteria become sensitive just before active reproduction.

A 4-hour *E. coli* culture at 37° grad. cooled to 1° C. (15 min.) Samples were warmed gradually & suddenly killed. As a control, a 24 hr. culture in 1% peptone was semi-treated over a period of 3 hours.

The "young" cultures were held at 1° for periods up to 36 days & responded to cold shocks by being killed & <sup>in</sup> lag in returning growth at 37°. *S. luteus* cells did age.

When held at 1° "young cells" die more rapidly.

Days held.	<u>Y.</u> x 10 <sup>6</sup>	<u>Viable cells/ml. Mature.</u>
0	8.6	
2	1.47	650
4	.49	460
7	.125	440
14	.004	192
24	400	95
36	72	43
42	—	39
51	—	16
62	—	10

Nelson, F. E. J. Bact 48:473-7 (1944) Factors which influence the growth of heat treated ~~and~~ bacteria.

Basal -  $\text{NH}_4$ ,  $\text{KPO}_4$  glucose agar + peptone - typtone used most.

Heat E. coli 55° 8 min.

Medium.	Counts (dupl.) $\times 10^3$	
Minimal	.46	.32
.01% typtone	.74	.39
.04%	1.0	.89
.2%	3.0	4.6
.5%	6.5	16.0
+ .01% thio glyc.	14.0	25.0
+ .01% typtone. better		

Unheated organisms were essentially same in all plates.

I.

45:395-403 (1943)

Iowa State College  
Ames, Iowa.

# Temperature

Ceuran, H.R. + F.R. Evans, J Bact 34: 179 - 1937

The importance of enrichments in the cultivation of bacterial spores previously exposed to lethal agents.

*B. subtilis*, *C. haereus*, + *S. lactis* - ATCC  
 "CC" *E. coli*

"Nutrient agar" gave much lower plate counts when treated cultures were tested than were obtained if supplemented variously, e.g.

"1 drop of st. defibrinated cow's blood per plate"

.3cc 10% glucose.

These supplements had no effect on untreated cultures.

Temp - 98° how long?

H<sub>2</sub>O<sub>2</sub> .05%

Details not stated

Y. Ex. deleterious, if anything.

Spores germinated on the NA but later did not respond to supplement.

E. coli 18 hours culture.

	Untreated 5000	U-V	Δ.
N.A.	57	20	27
" + blood	57	65	102
" + glucose	60	45	105
" yeast	61	25	27
Muesli agar	61	38	189
Tamaleptic + milk pdi.	54	69	237.

This can be investigated.

Hansen, P.A. *Arch. f. Microbiol.* 5:99-122 (1933) The growth of  
thermophilic bacteria.

Temperature - tolerance

Williams, F. T. J Bact 32: 589-97 (1936)

Attempts to increase the heat resistance of bacterial spores.

*Vaccinia stevensii*. Peptone - beef extract - sugar

Temperature - tolerance by bact.

Edwards, OF + LF Kettger, J. Bact 34: 489 - 1937

The relation of certain respiratory enzymes to the maximum growth temperatures of bacteria.

M.G.T. measured by observation in liquid + solid tubes in a variety of organisms. Solid or liquid had no effect.

A statistical correlation was found, among different strains, between temperature of destruction of enzyme activity (cytochrome oxidase, catalase and succinic dehydrogenase).

E.g. °C.	M.G.T.	Cytochrome Oxidase	Catalase	Succ. dehy.
<i>B. mycoides</i>	40	41	41	40
"Thermophilus"	76	65	67	59
	1	2	3	4

A correlation of  $.8466 = R_{1,234}$  was found for these items.

"Indophenol" oxidase activity gave best correlation.

$$r_{12} = .8431 \quad r_{13} = .8451 \quad r_{14} = .7737$$

Qualitative tests: intact cells

- (2) - CN sensitivity, ~~indophenol~~ p-phenylenediamine oxidation
- (3)  $H_2O_2$
- (4) Thunberg. Met Blue.

Endospores guarded.

Dunn, M.S., et al., JBC 156:703 - 713 (1944)

XVIII. The amino acid requirements of *Leucostoeus macleodensis*.

Standard curves found for arg, cyst, glut, hist, isoleu, leuc, lys, meth, P.A., pro, trypt, tyr + val.

Alanine, Hopsol, norl, & norv, were non-essential or auxiliary.

In medium "C", P.A. was required, 150+ / tube giving near ex. prod.

XIX The determination of lysine in protein hydrolyzates by a microbiological method.

Shaulman, S., HSDrum + L. B. Rubin, JBL 151:511- (1943)

The microbiological analyses of 7 amino acids in *L. casei*.

72-hour acclimation.

*S. aureus*: 30r tube for 1/2 max-growth.

Medium of Hettler + Peterson PSEB 152: 76 (1943).

50 mg in



HISTIDINE; ASSAY

Reun, M.S., et al. JBC 159: 653

Histidine by Luccanostoe

TRYPTOPHANE

Substrate utilization  
and synthesis.

*L. arabinosus*

Wright, L.P. and Steggs, H.R. JBC 159: 611- 1945

Tryptophane utilization and synthesis by strains of *L. arabinosus*

PYRIDOXINE + CO<sub>2</sub>

Amino Acid Assay.

Jeyman, C.M. et al JBC 162:173-4 (1946)  
pyridoxine in lactiae. bacteria.

On the function of  
Setler.

Amino acid requirements modified by CO<sub>2</sub>.

CO<sub>2</sub> + pyridoxine removes requirement for P<sub>A</sub>, Tyr, Arg in L. californicus  
(16r)  
and Aspartic in S. faecalis

Texas.

THREONINE essay  
S. FAECALIS amino acid analysis

Greenhut, I. T., BS Schweigert & CA Elvejlum,  
SBC 162: 69-76

The amino acid requirements of *S. faecalis* and the use of  
this organism for the determination of the in. natural products.

Ileuc, thr, gl, asp, lys, val, isole, meth, arg, hist, ser, trypt,  
and cyst required

Alan, try, OA, glyc stimulatory.

Differ  $\bar{c}$  Anell and Secciard who did not require meth, val, hist  
and isole, and that alan was

Purines, biotin, panth, B<sub>2</sub>, B<sub>6</sub>, nic, + folic  
Glucose, citrate, Mg, Fe, Na, Mn

Response to dl is not linear. Unnatural isomer (~~dl~~ l(+)? inactive

2-5 hour hydrolysis  $\bar{c}$  2N HCl, autocl. gave satisf. recovery

ATC 8043

Atkins, P & J L Ward, BSEP, 26:120 - 1945  
Effects of analogues of vitamin K.

The antibacterial

Wobley, DW PSEBM 60:225-1945 Observations on the antimicrobial  
activity of 2,3-dichloro-1,4-naphthoquinone & its mesal bystanis  
R.

Shive, W. + J. Tharow, JBC 162:451-462 (1946.)  
 Biochemical transformations...  
 I Hypoxanthine.

dl hydroxyaspartic acid is inh. to E coli, reversed by glutamic acid or by aspartic acid. (c.) pantothenic acid raises anti-bacterial index.

An E coli strain initially non-prototrophic was adapted by serial transfer for use in these expts (!!). (Reisolated?).

Antibact index ca 10-15. index in E coli. In *Sarcospora* 60-100.


II tried in coli. similar, but index 100-200.

At low levels of I, 1r prot = 10r aspart in reversal. do β-alanine: Hypoxanthineless effluvia. Panto. increases antibact index from 3-20. e.g. other g-t. had no effect.

At higher (I) glut. decreases in activity. β-alanine, valer, succ, + fumaric ineffective. Isoserine had no effect at 1mg/cc!

Interpts off. of panto as indicating shift of limiting nutrient from β-alanine synthetic to another one. Interpts glut. effect as panto... aspect by transamin.

II ~~SA~~ pabr. 463 -  
 also



II reversed completely by methionine.

Series of antib. indices made with addition of different substrates. 1. Methionine 2. adenine 3. . . ?

SA: pabr

3000 nonmeth.  
 10000 meth.  
 20000 panto.

Presumably II is ineffective only at a certain locus of pabr action


Medinaveitia, J. Biol. Chem. 139:85-91 (1945). Antibacterial substances related to pantothenic acid.

"pantamides". Reference vials. P.T.:  $P-NHCH_2CH_2SO_3Na$ .

L. casei used.

pant-hydroxide was active, but not highly so:  $P-NHNH_2$ . No other act.

Also, pantoyl - N-2 aminoethyl - (p-aminophenyl) - 1 sulfone.



Not covered by pant; " by pantothen.

Therapeutic activity, in rats & Spyzogones.


Martin, AR + FL Rose, substances related to pant.

39:91-

1945. Antibacterial sub-

(overlap Wyrstal; Green, Johnson + Pauli).

	2.	3.	5.	6.
1.		Cl		
2.	Cl			15
3.		L		16
4.		Me		17
5.	Me			18
6.		HO		19
7.		MeO		20
8.	MeO			21
9.		EtO		22
10.		NH <sub>2</sub>		23 MeO
11.		COOH		24.
12.		MeS		
13.		EtS		
14.		MeSO <sub>2</sub>		
		EtSO <sub>2</sub>		
		Cl	Cl	
		Cl		Cl



2.	3.	5.	6.
	<del>SO<sub>2</sub>Et</del>		<del>Me</del>
	Cl	Cl	<del>Cl</del>
	Cl		Cl
	Cl		NH <sub>2</sub>
	Cl		NHAc
	Bz	Bz	
	Me		Me
	MeO		MeO
		Me	
	MeO		Me.



"S. pyogenes; Wrigley's tooth. + blood.


I:  $\frac{1}{27}$  eff. as SA. |  $\frac{2}{5} + \frac{4}{5}$  anti SA.

S. mut.

4-amino isophthalic

4-(4'-amino benzamide) benzoic ac.

& Et. 4-amino benzoate



Sl. anti SA activity

McLwain H. Biochem. J. 39:329-33 (1945) Biochemical characterization of actions of chemotherapeutic agents. S. lack of gross displacement of pantothenate and pabate from microorganisms by pantooyctaurine & Sulphamida-mide.

Sheep. hemolyticus. limiting pantothenate medium  $\rightarrow$  pantothenate poor cells. No all ex part in heavy part medium growth removed by mic washing.

Suspensions contg 15-60 mg (dry) of cells in 2-5 ml  $^{32}P$   $PO_4^-$  part determined by digesting + Proteus growth.

With cells (lg. batches) exposed to SA. No release of anti SA occurred as exposure to buffer, saline or SA.

Pnt. content of bugs grown in initially  $2 \times 10^6$  was

30 mmol/g (dry) Growth for shorter periods - more part, the contemporary level being important. The cells inactivate part.

Cells up to 700 mmol/g were obtained

No part was liberated on exposure to part-taurine of the poor part cells. No did washing. plasma vials. part inactivation.

In part vials cells, part stable at R.T. was released into saline at  $37^\circ$ . The quantity remaining being ca that of part poor. Large inc part-taurine had no effect on quantity removed.

The amt of SA-antigen present is not altered by large amt of SA.

\* It is suggested that although part + part functions in resting bacteria these activities, when the resp. substances are incorporated are not influenced by SA + PT but the reactions involved are the as simulations of the substrates. These are stably bound.

Therefore expect a lag in action for detection of ~~part~~ substrates.

McIlwain, H + DE Hughes, *Biochem. J.* 39:133-139 (1945). 3. Relations  
ships between metabolic and growth inhibition by paralothenate analogues  
: their structural and sp. specificity.

Assay  $\bar{c}$  Proteus.

Several analogues tested for (1) effects on growth, reversed ~~by~~  
by P<sub>th</sub>

Some comp. ind. growth but not P<sub>th</sub> inactivator:  
bis-nordesoxy paralothenate. ~~These~~ These were not reversed by  
paralothenate.

All analogues which competed  $\bar{c}$  P<sub>th</sub>, inhibited the  
inactivation of prot.

order of activity of different analogues ~~is~~

& of prot T. in different species is the same for growth &  
prot metabolism.

McIlwain, H., *Biochem. J.* 39:279- (1945) 4. Time-relationships between metabolic and growth inhibition by pantoyltaurine.

1. put + streptococci  $\rightarrow$  slow inactivation of put at uniform rate.
2. not occur at 0°.
3. Inhibited by pantoyltaurine immediately.
4. Growth inhibition has lag ca. 1 hour.; recovery also lags.
5. Reversible on washing & removal of put. occurs very quickly.

$\therefore$  assumes action of ~~put~~ PT is to inhibit the synthesis of a put derivative essential for growth, which can be produced in excess.

Field, J.B., EG Linsen, J. Spero, and KP Feiler, JBC 156: 725-737 (1944)


Studies on the ~~hemorrhagic~~ hemorrhagic sweet clover disease.  
XIV. Hypertension and hyperproteinemia induced by methyl xanthines and its  
effect on the action of 3-3'-methylenebis-4-hydroxycoumarin).

Caffeine, theobromine + theophylline stimulate liver production of  
prothrombin + fibrinogen, reversing dicoumarol.

NICOTINIC AC. analogues (Acetylpyridine)

Woolley, D. W. JBC 162:179-80 (1946) Reversal by trypt of the biological effects of 3-acetylpyridine.

Tryptophane was as effective as nic in reversing effect of 3-AP on mice (pellagra!).



Rodriguez.

RIBOFLAVIN, analogues

*L. casei*

Sarett, H.P. JBC 162:87-97 (1946) The effect of riboflavin analogues upon the ~~use~~ utilization of riboflavin and FAD by *L. casei*

Review: isoriboflavin has  $< .5\%$  activity of  $B_6$  for *L. casei*  
inhibits subgrowth at low  $B_6$

Shows: in presence of suboptimal  $B_6$  or FAD, stimulates ac. prod.

Deaminopyrimidine competitively inhibits utilization of  $B_6$ .

Lumiflavin competes  $\bar{c}$  low  $B_6$ , stimulates  $\bar{c}$  high.  
inhibits FAD utilization at lower concn.

*L. casei* is alkali-treated peptone, or Casamino (Tandy + Dickson)


main effects on  $B_6$  enzymes, and not on  $B_6 \rightarrow$  FAD reaction

ANTIBIOTIC: Buttercup Juice

Baer, Harold, M. Holden and BC Seegal, JBC 162(1):65-68 1946

The nature of the antibacterial agent from *Anemone pulsatilla*.

~~Anemonin~~ ANEMONIN obtained, a polymer of proto-A.



Activity measured against *E. coli*, *Staph.* and *Candida albicans*.

Acetylcyclic ac., nor vinylcyclic had no antibacterial effect.



Kimball, R.F., *Genetics* 24:49-58 (1939). A delayed change of phenotype following a change of genotype in *Paramecium aurelia*.


Following induction there is a delay in the expression of change of mating type that may occur.

Frederick, C.C. + G., Genetics 24:1-7 (1939) Non-random crossing over in the 2d chromosome of Neurospora crassa.

See L. '36. Genetics 32: 243-56.

9 chromosomes.  
= 38.7

kuitorlore, peach, tuft + thuffy.



1. Excess of 2 strand exchanges. Deficiency of multiple exchanges.

Jeweries & Tamer, J Bact 49:383- 1945.

The inheritance of environmentally induced characters in bacteria.  
Graded cone.

(Selection favoring wild type in mixed cultures in absence adaptive agent.)

Inoculate mass populations into Agar.

Changes of  
critical  
conc.

NaCl - from 3 to 8%
CuSO <sub>4</sub> - 1:4000 to 1:800
HgCl <sub>2</sub> - 1:300,000 to 1:50,000

∴ use 6% salt agar


Horowitz, N JBC 162:413 1946.

The isolation & identification of a natural precursor of choline.

$\text{CH}_3\text{-NH-CH}_2\text{CH}_2\text{OH}$  isolated from 47904, active on 34486

Appears only after 7 days. more conc. in mold than medium.

47904 must synthesize type. choline. methylation of diMeEtOH $\text{NH}_2$  also affected.



Fries, Nils. Svensk Botanisk Tidskrift, 39: 270-8 (1945)  
Two X-Ray induced auxo-heterotrophies.

*Ophiostoma* (*Ceratostomella*) *multiauriculatum*.

wild type requires: B<sub>1</sub> + B<sub>6</sub>. Mutants for Biotin (225) and  
pab (617) obtained by X-Ray. Isolated by special selection technique.

Ark. för Botanik, 32: 1-9 (1945) über Röntgen-induzierte  
physiologische Mutationen bei *Ophiostoma multiauriculatum*.

50 kv. 2-3 ma. 100 m. Plated irradiated spore suspensions onto minimal  
"Fries agar" + B<sub>1</sub> + B<sub>6</sub>. Mutants "deutlich schlechteres Wachstum abweisen"  
wurden. Von den die auswachsenden Ansporenyzelien wurden deshalb  
nur solche isoliert, die sich in dieser ~~von~~ Beziehung von - des Meistens -  
normalen Myzelien unterscheiden.

1. Temporary radiation effects (back mutations?)
2. Morphologicals.
3. Mutants.

527 isolated. 30 mutants - 6 biochemicals.  
None from ~~the~~ unirradiated material.

- # 225 Biotin
- 358. Reduced S. (parathiotroph - cysteine etc. or 4 valent S. (SO<sub>3</sub><sup>=</sup>)
- 446 Parathiotroph - can use <sup>not</sup> tetravalent S.
- 460 - ~~yes~~ Oracil
- 513 Adenine? low activity
- 617 pab.
- 848 Guanine.

Naturw 30: 44/5 - 1942. Adenine als Wachstumsfaktor  
für *Ophiostoma ulmi* (Bresinans) Kaurf.  
Requires only B<sub>6</sub>.

Nature, No. 3947: 757 (June 23 1945) X-ray induced mutations in the physiology of *Ophiostoma*.

*O. multiauratum*. strains mentioned above.

Parathiotyphs in crosses lost ability to reduce tetravalent S. (#358). Other features identical as 1 gene in crosses.

Needed large quantities of adenine.

Uracil-less used cytidine or cytidylate. but not cytosine (lib 129P).

Nature #3847: 105 July 24, 1943.

Vitamin B<sub>1</sub>, Vitamin B<sub>6</sub> + Biotin as growth substances for some *Ascomycetes*.

*Ophiostoma*:

	Needed	Stimulate
<i>O. piceae</i>	Pyr	—
<i>steroceras</i>	P, S, C	Biotin
<i>coeruleum</i>	Pyr	B <sub>6</sub> "
<i>quercus</i>	Pyr	" "
<i>pinus</i>	Pyr, Biotin	B <sub>1</sub>
<i>ulmi</i>	B <sub>6</sub>	Pyr
<i>fagi</i>	B <sub>6</sub>	Biotin
<i>pilliferum</i>	B <sub>6</sub>	Biotin
<i>multiauratum</i>	B <sub>1</sub> + B <sub>6</sub>	—

"Artificial symbiosis" tested + worked. (Heterocaryon?)

Nitrate needs biotin  $\bar{e}$  NH<sub>4</sub> for N; respirable  $\bar{e}$  NO<sub>3</sub> + acid!

Hollander, A. Effect of long uv & short visible radiation on E. coli  
J. Bact 46: 531-11 1943.

Saline = NaCl 3g RCl .2g CaCl<sub>2</sub> .2g / 100 ml H<sub>2</sub>O. Protected by hyp barth  
somewhat.

1. Growth delaying effect before app. lethality (plate counts)
  2. Survival in saline: (incubation).  
control survived quite well 10 hours. (98%).  
irradiated died much more rapidly
- Longer wavelengths much less efficient ( $10^5$  energy eq.).

Wickscham 145

8 ascospores/ascus. after copulation. Relatively anaerobic. Bottom fermentation 3 pellets.

Under slide conditions, hyphae are found. (rel. anaerobic). Nucleus visible in terminal hyphae, ca. 8-10 $\mu$ , particularly anaerobically.

glucose, maltose & sucrose rapidly fermented. Also melibiose.

Not galactose or lactose

Sporulation did not occur from hyphae, or was diminished temporarily.

Trypan blue in agar leads to dark pigm. in aggl. phase (abt. from normal). Growth rapid 30-37°. Colonies develop slowly - 4-6 days. Copulation occurs readily at 20-33°. Ascus ruptures before completing development.

Wickscham, L.S., & Eugénie Duprat.

J. Bact 50: 597- 1945.

A remarkable fermenting yeast, *Pichyosporichia nycos vesatilis*, n.s.,



Lwoff, A. + A. Audenaert, Ann Inst Pasteur — ? 1941.


Sur une mutation de *Thiopylla lwoffii* apte a se développer dans les milieux à l'acide succinique.

pp 1-2 missing  
Typical strain will not utilize succinate.  
Rarely mutations appear, influenced by succ. from S- to S+. In presence of EtOH S- outgrows S+. S+ → S- not found. Rate S- to S+  $\approx 10^{-8}$ .

70:51- 1944. *Recherches enzymologiques sur les mutations bactériennes.*

Succinoxidase is present in both strains. *Arabobacteriacae*. is decarbox. spontaneously but not rapidly enough for growth.

Hydroxy fumaric acid studied (enol form of




Rate of decarboxylation studied. Rapid at first as ~~by~~ ~~cost~~. S+, but slows down to spont. rate (almost as rapid)

Prove there is an enzyme present? in S-. which is not present in S+?

Mitchell, H K and M B Houlahan, ASB 33:31- 1946.

*Neurospora* CV. A temperature sensitive *Neurospora* mutant.

51602. At 31° or above, requires riboflavin absolutely.  
S-shaped response curve .1-2.5  $\mu$ g. At high B<sub>2</sub>, growth curves  
like wild, at low levels, bimodal temperature response.



Grows on 20 ml

at higher temperatures,  $\bar{c}$  a small B<sub>2</sub> supplement,  
(ca. 3 r) full wt. can eventually be obtained (200 hours =  
8 days.) containing full B<sub>2</sub> content by L. casei.

For B<sub>2</sub> determ., autoclave cultures in medium & analyze filtrate. F. vial

ca 6-9 r/100 mg. Mutant grows intermittently, rising up + syn-  
thesizing vitamins. Not tested as *Neurospora*.

Inhibited by leucichrome; reversed by B<sub>2</sub>. R<sub>50</sub> = 1.2-2.5.

Some unlabeled in tissue extracts.

*Neurospora* may contain a doubly functioning set of genes for different temperatures.

Abb 4A x 21a.

Tatum, E. L. + T. T. Bell.

A. J. B. 33(18): 15-21 (1946)

Neurospora<sup>44</sup>. Biosynthesis of thiamin.

		Distance from center
1090 (sitophila).	45 asci	23
9185	24 "	8.3
18558	8 "	0
17084.	33 "	35

No interspecific heterozygosis.

3 day growth, some / 125 ml flask.

18558 requires thiazole  
9185 intact thiamine

When growth is limiting thiamin, accumulation of pyrimidine was established by 18558 (tested on 17084, + Phycomyces). Analogues of thiazole had activity very similar to Phycomyces, except that 5th ethyl may have ca. 1% activity of B<sub>1</sub> for 18558.

2-methyl derivative was also app. active

Factor S did not influence 9185 response.

17084, 1090 (and 56501), require both pyr and thz. Mixture has same activity as thiamin. Filtrates have a 9185 active component, which loses activity on sulfite treatment. It is also active for 18558 and Phycomyces. Not active for 17084.


299 as low B<sub>6</sub> responds only to B<sub>1</sub> or pyr + thz.

Wodley's conclusions on pyritiamin not confirmed. 17084 and 1090 cannot pyritiamin for pyrimidine.

A thiamin metabolism error may exist in 1090 + 10084.

These strains have a higher requirement.

i.e.



Carrel, A. *Pr. Am. Phil. Soc.* 68: 129-32 (1929) The nutritional properties of malignant cells.

Neurology

Kellogg, W.N., et al S 103:49. 1946.  
LogS.

Special conditioning in

## RADIATION: Cathode

Wychoff, RWS + T.M. Rivers, ~~JEM~~ JEM 51: 921- 1930.

The effect of cathode rays upon certain bacteria.

$1.5 \times 10^5$  volts

The absorption of a single electron will kill a cell.

Concluded that only .008 of the incident electrons are absorbed from phantom expts.

" Only 85% of the single hits were effective, but when death occurred, a single hit sufficed ..

(data from dose response curve, and calculated absorption by the bacteria.)

[How can this be compared to the production of rays by radioactive P, etc?]

RADIATION: u-v

Hollaender, A + RM Duggar, J. Biol 36:17 1938

The effects of sublethal doses of monochromatic u-v radiation on the growth properties of bacteria.

2653 Å



Reles, N.H., *Genetics* 28: 398- 1943. Comparative studies of the cytogenetical effects of neutrons and X-Rays.

Hollander, W.F. Gen. 28:76-1943 Abst. A possible case of directed mutation in the pigeon.

$PT^1$  = sexlinked Almond  $a^{br}$  → mosaic of brown and  $a^{bl}$ .

Al  $\cdot a^+$  → mosaic = black between beams.

do. Al. - (homozygous ♀♀).

Evidence that Al →  $a^+$ , etc. If so, mutation is directed by the other allele. (rather than somatic loss or crossing over).

Sonneborn, T.M. do.:90 Development and inheritance of serological characters in variety 1 of *P. aurelia*.

Stork P has antigen; 60 lacks it. Single dominant gene.

P x 60 → some homozygotes which retain antigen 4-8 fissions (cytoplasmic lag).

Aa x aa → slowly developing antigen detectable only after several fissions + increasing to standard level.

Anti-A kills most of  
Anti-60 kills most numbers, but some resistants arise.  
Lost within a few fissions unless continued exposure to serum.  
Some lines then retain their resistance (275 qmcr.); others lose it more rapidly. Lost as endomixis or fertilization in 9 fissions (Dauer modification!!)


Note: bar is dominant.

Sutton, E. Genetics 28: 97- 1943 Bar eye in *D. melanogaster*: a cytological analysis of some mutations and reverse mutations.

Sum: Hemizygous  $\sigma^7$ , and Bar deficiency have no phenotypic effect.

The Bar effect is produced by interaction with other loci, which may mutate... The Bar effect may be destroyed by mutation of one of the two interacting loci, as well as by separation of these loci through chromosomal rearrangement.

Reversals:



a) deficiency of one duplication, incl. 15F - 16A.

b) inv.  $16A_7 - 17A$ .

c) no detect. change

d) def.  $16A_1 A_2$

e) del  $16A$ .

f) inv - long from 4A to  $16A_2$

g) " -  $16A_7$  to C.

Similar effects in double Bar.

Oeshov, S.L., Acta Path, Microbiol Scand, 22:523 (1945) Investigation  
of the permeability of yeast cells.

White  
Woolley  
Hutner

Geol. D., J.G.P 29: 219 (1946) *P. Steolyta Eupymest.*

P.4.

Strong, L.C. XII. Yale JBM 18: 145-155 (1946)

The effects of selection toward resistance.

1. Meth induces ap. tumors in homozygotes; likewise in heterozygotes; particularly in strains selected for resistance to local tumor formation.

An increased mutation rate is also postulated.

Mice here strongly selected for resistance.

1. In one subline, no change [biotype - pure line?]

2. In 4 lines, a decrease, but accompanied by an increase in mutation rate to susceptibility.

Owen, FV, J Agr Res. 71:423 1945 Cytoplasmically inherited  
male sterility in sugar beets.

Asteiger G. Hereditas 30: 213-16 (1944) Inefficient sterility  
for the induction of ~~sterility~~ homozygotes.

Alleles costs 3da. 1 meter.



## Auxanography

Hayes, W. JPB 57: 457-466 (1945) The effect of agar depth in the plate method for the assay of penicillin.

200000/ml opt. For penicillin, agar depths less than 5-6 mm. give sharply increasing size zones of inhibition, varying  $\propto$  concentration.

The assay value increases at agar depths considerably greater than the apparent radius of diffusion.

8.8 cm plates require 50 ml for 8 mm. agar, which is required for uniform results.

Selmann, FE + H Wobser, Verh. Schw. Naturf. Ges., 120: 181-2 1940.  
Verschwinden embryonaler Zellkerne v. Testisfix nach Collicinbehandlung.

6:20.74 (1910) Method of counting

Smith, Suter MC; A.S. Med Techn  
bacteria....

Beyerinck, H.W. Archives Néerlandaises des Sciences, 23: 367-72 (1887)

J'Anaxanographie, ou la méthode de l'hyalodiffusion dans la gélatine appliquée aux recherches microbiologiques.

Add liquid supplement to the surface of an agar or gelatin pour plate to allow requirements. e.g. yeast & phosphate (yeast is more resistant than most bugs to killing under such conditions). Also, double diffusion zones for cardioline giving "une figure lenticulaire opaque de couleur jaunâtre." Glucose + agarose, etc. as to source. Inhibitors also easily demonstrable. Also suggests drying the plate.

Points out that optimal conc. to not have to be known. Used large plates for multiple effects.

Furth, J. + M.L. Boon, AAAS Research  
Conference on Cancer, 1944, 129-138.

The time and site of origin of the leukemic  
cell.

Malignant cells determined by bioassay - intravenous adm. to rec. an.

→ 1 cell needed for transmission.

1. Young leukemic mice do not harbor ~~by~~ neodymphocytes.
2. Some neodymphocytes can be found before clinical leukemia.
3. Thymectomy reduces incidence leukemia. (ca 60 to 10%). Do. undefining. Splenectomy is effect. Does not influence transmissibility. <sup>They</sup> may have a general effect in inhibiting tumor growth.
4. Undefining reduced incidence from 65 to 10%. Also interferes w. transmission. May have leukemic cells by bioassay & evidence of leukemia. Rarely in bone marrow; probably not typical site.
5. Hecht leukemogenesis. a. X-Radiation increases. b. Used +, hybrids which do not develop spont. Overlaps in 70-100 days. contains neodymphocytes a short time before leukemia develops.

Earle, W R, AAAS Career 1944.139.

A summary of certain data on the production  
of malignancy in vitro

Oedel, Zf H.K. Pausch. J Bact 51: 791-2 (1946) The  
biotin requirements of *Neisseria sicca*

only biotin required opt. .0001  $\mu$ /ml

Reyes - Teodoro, R. + M.N. Michaelson, J Bact 51:569s (1946)  
Recovery of vitamins from cultures of acetone, butyl alc. bacteria.  
Synth. medium.

75-80% recovery. 15-20% in medium.

acid hydrolysis or papain-diastase are best methods.



Kleinberger Nobel J Hyg 44:99 1945

Jbl Bact I ~~1/10~~ ~~1/10~~ 1240 (miscultures)

JID 54:313.

~~Haldenester 3. Bact 13:111~~ symbiosis

J Bact 30:301

Green, FE + EUMykon J (D) 4/2: 525-36 (1938).  
42: 545-

Feldstein, HC + ML Snyder, J Bact 42: 653-64 (1941) The inhibition of the spreading growth of Proteus and other bacteria to permit the isolation of associated streptococci.

a) Fry's technique of pouring layered plate  
1. Prevent spreading with a top layer

6% NaCl inhibits spreading but not growth markedly.  
(probably cuts diffusivity of water - as ind. by dye)  
(probably not a good idea)

Hydric inhibits spreading at  $10^{-4}$  but growth as well.

alcohol 5% inh. spreading but not growth.


[Detting has proteus phages.]

Fry - BJEP 15: 456-7 (1932)

Burrows, W. J. D. 54:135- 1939 The nutritive requirements  
of the Salmonellas.

Many strains require tryptophane. Carabe + mid. Tryptoph. conc.  
does not affect rate, or final growth, but only lag. Replaceable by  
lysine in one strain. Tryptophane assay increased after growth.

$N_2S$  response similar:



$N$  variations affected both rate + amount. Glucose was all or none.  
not lag.

$NH_4SO_4$   
NaCl  
 $KH_2PO_4$   
glucose

high  
resp. rates ~~lower~~  $\hat{=}$  low tryptophane -

(selection?)

Demere, M. CSH 9: 145 (1949) *Crustaceans in Zoology*.  
see Demere 1935.

Plough, H.H. CSH. 9:127 (1941) Spontaneous mutability in *Drosophila*

Goldschmidt, R. Biol Zentr. 49: 437-48 (1929) Experimentelle Mutation  
und Problem der sog. Parallelinduktion. Vers. an *Drosophila*

By heat-treatment of larvae, phenotypic sooty which is sooty were found.

"simultaneous somatic + germinal mutations," favored. !

Bluhm, A. Biol Zbl. 48: 641-8 (1928) Einige fragende Worte zum Mutationis-  
griff. (Hansson, basit)

see Bauer. —



Delbruck, M. Biol Rev. 21:30 - 1946. (Bacterial viruses or bacteriophages)

Winge, O. CR Carlsberg 24:79-95 (1944) on segregation and mutation in yeast.

~~S. cerevisiae~~ *S. cerevisiae* - only 1/2 spores survive. (lethal?)

*S. uvarum* - (single spore form) probably varying segregants

Ditteusen, E. CR Calsbeeg 24: 31-37 (1944) A case of simple segregation  
in *Pachomyces italicus*.

1:1 segregation of a morphological gene (L.) long down - short cell type.

Spore lines are of two types & when they sporulate, they bud true (particularly  
ll). LL sporulate only rarely. Hybridization attempted L x l &  
yielded substantially the P<sub>1</sub>, again segregating 1:1. L x L rare; l x l freq.


Twombly, G.H., + D. Meese, *Cancer Research* 6: 82- (1946) The growth of mammalian tumors in fertile eggs. Is a fertile ovum produced?

Rebbecca R39, Bagmouseca 755 + the RC mouse ca. were grown in fertile hen's eggs.


Tumor producing activity could not certainly be dissociated from viable cells.

Dawson, R. Alkaloid formation in plants. Zoology Colloquium 3/6/46.

Tobacco alkaloids:



Nor-nicotine is demethylated nicotine.  
nic + normic = fairly constant in various strains



Also N-methyl anabasine  
Nicotyrine is a 1'-2"ene - nicotine.

Pyridyl common; side group varies. A similar series in cissampelos, cactes alkaloids.

Accumulation of nicotine in leaves is not modified by most procedures on leaves.

Grafting tomato top to tobacco roots  $\rightarrow$  nicotine containing leaves + fruit.

Tobacco/tomato  $\rightarrow$  no alkaloid

Solomonson, U. V., Chem. Rev. 37: 481- 1946. Synthesized Esters  
gives the relations between their structure and their activity.

Res. Labs  
Hoffman La-Roche Inc  
Nutley 10, N.J.

Thaugnot, G. Rev. Cytol et Cytophysiol. Vig. 5:169-264 (1941)  
Substances mitochondriales et cellules végétales

Shemin, D. JBC 162:297-307 (1946) The biological conversion of L-serine to glycine.


Benzyl ac. and labelled comp. injected into rats, guinea pigs.

$N^{15}$  in hippuric ac. determined + comp'd  $\bar{c}$  that in the labelled injectum.

The dilution factor was lowest for glycine (2.8, 2.4 resp.) and v.

high for <sup>al-</sup> glutamic (1500, 450...)  $NH_3 \rightarrow$  400, 20 resp. in the two spp. d-serine was rel. ineffective. l-serine was 5.5, 3.9.

l-glutamic is 45, 10.



Ratio of  $\frac{N^*}{C^*}$  in hippo glycine

demonstrates the direct conversion and

eliminates ethanotamine. Nor is  $\begin{array}{c} COOH \\ | \\ CHNH_2 \\ | \\ COOH \end{array}$  the intermediate, unless

reversible deamination. N-benzoylglycine  $\nrightarrow$  hippuric.

Probably no reversible deamination  $\bar{c}$  glycine...



Leiria, SE, Genetics 30:84-1945. Mutations of bacterial viruses affecting their host range.

Coli B. Viruses  $\alpha$ ,  $\nu$ .

$B/\alpha$ ,  ~~$B/\nu$~~  readily obtained. Also  $B/\alpha\nu$ . Also  $B/\alpha$ , etc. morph. variants.

$B/\nu$  more difficult.

$\nu + B/\nu \rightarrow 10^{-5}$  to  $10^{-7}$  clear plaques. A new virus, active on  $B/\nu$  can be isolated.  $\nu'$ . It can be obtained from single plaque isolates.

No virus active on  $B/\alpha$  found. But  $\alpha \rightarrow \alpha'$  active on  $B/\alpha_2$ , not active on  $B/\alpha_1$ .

$\nu' \rightarrow$  a smaller plaque count on  $B/\nu$  than  $B$  (.2 to .6). This is not due to  $\nu' \rightarrow \nu$ . After absorption by  $B/\nu$ , the plating efficiency does not vary. It is likely that  $\nu'$  is less readily absorbed by  $B/\nu$  than by  $B$ .  $\nu'$  interferes  $\bar{c}$   $\nu$ . (Self-interference also likely).

$\alpha'$  is identical  $\bar{c}$   $\alpha$  on  $B$ . Plating efficiency .3-.7 on  $B/\alpha_2$ . Absorption is lower. Delbruck analysis,  $\bar{c}$  complication of bacterial mutation to resistance & sp. multiplication. Fluctuation  $\rightarrow$  conclusion of mutation. Some cultures had a mutant population  $\bar{c}$  smallest burst size indicating mutation in cell.

Serologic identity of  $\alpha$  &  $\alpha'$ ;  $\nu$  &  $\nu'$  is tableted. Bact. resistance independent:  $B/\alpha$  sens. to  $\nu'$ .

$B/\alpha_1 \rightarrow B/\alpha_1\nu'$  but was sens. to  $\alpha$

mutant can be obtained from  $B/\alpha_1\nu' \rightarrow B/\alpha_1\nu'\alpha$  resist. to  $\alpha, \nu, \nu'$

McDowell, -

Genetic factors - High incidence in CSB. Incidence related to "amt. of inheritance" of leukemic strain. Genes vs. cytoplasmic elements.

f, heterozygotes: differences in reciprocal hybrids. Maternal effect?!

Variability in f, - isolates. f, x p, (n). Low incidence (to 1/4.) Still problems of segregation due to imperfect penetrance + masking of phenotype. Binding tests essential. (Test of genotype)

RR x rr

↓  
Rr. 1:1 ratio in progeny expected for monogenic inheritance.

Stoli = Little-Stones. "S"

resistant

Why backcross rather than inbred??

(Ask for reports) CSB. (1 generation = 100%??) (Selection??)

RR x rr

↓

S x c

Rr x rr

↓

X sc

X So

Rr, rr

test by x n!!

↓ Test progeny by mating to S ♀. Variability in backcross

~~FF~~ F1s genetically uniform, reduced incidence. ∴ non genetic determin.

all crosses to high strains CSB. Nursing CSB ♀♀ inhibits leukemogenesis.

Planned as high uniformity as possible.

7 each ♂ x 10 ♀

D used as B albino.

heterozygote between families.

Effect on 1/2 or homozygotes.

age or litter no?

p1 RR x rr



f1 Rr x Rr



f2 Rr, rr.

test. the progeny of these.

x rr. Some lines should have no leaks.  
Some up to 50% leaks.

Variability found between ♂♂. is 1-2

2 = 5 differ in 3 genes on pigment. 2 correlated = leaks.  
transmission of a longevity factor from ♂♂. non sp. leaks

but had a much influence as leakiness...

These affect greatest in ♂♂. Also ♂♂ - fighting; cystitis; these  
improved competition + improved cystitis.

- Age of mother at parturition. (Stoli) Young → higher incidence.

50 families are not adequate for multivariate analysis.

Test # of genes??

Effect of nursing greater on hybrids. (Sex-linked testis)

Young removed as born... divided between 3 strains of nurses.

No mice got 1st milk (everything fostered). 4/6-1s.

1. Reciprocal hybrids still vary. S-nursing protects in both directions except in final % leukemia.

⊖ B nurse, the cytoplasmic effect is much greater, and affects final rate.

Freese, HC + JW Gowen, *Genetics*, 27:212 - (1942) Analysis of data  
on X-ray induced visible <sup>gene</sup> mutations in *D. melanogaster*.

Timofeef-Resovsky's data indicate no significant detection of mutation,  
or mutability of any allele in the w series.

Hauffmann, BP, *Genetics* 27:537- 1942. Reversion from rough to wild type in *D. melan.*

Sex-linked recessive. Deactivated at low temps.  $rst^3$  flies are a mosaic of smooth + rough facets, rough part. in  $\sigma^7$ . Associated  $\bar{c}$  along inversion from  $rst$  to the right of bobbed. left bush is in 3C2-3C4 region.  $rst^2$  is allelic (see Zurenberg 1937).

$Rst \sigma^7$  In (1)  $rst^3, rst^3 carbb \bar{c}$  4000<sub>2</sub> X-rays and X Y females  $\rightarrow$  revertants, which were sterile (heterozygous hemizygous for inversion).

Then radiated  $\sigma^7$  x  $brst^3$   $\sigma^7$ . 21,104 F,  $9\%$  examined.

171 were  $Rst$  phenotypically. 72 analyzed. 25 sterile 7 lost  
23  $rst \bar{c}$  poor expression; 17 revertants. (ca. 4%).

16 had karyos in proximal heterochromatin of the  $brst^3$  X chrom.  
4 were revertants; 2 also transloc. 7 unip. transl. 2 could be maintained as ~~hetero~~ homozygotes & were  $\sigma^7$  fertile. After two years some  $rst$  flies appeared again (cytological modifications).

There exist some data that new arrangements have weak spots.

Other genes tested. No reversion of forked or pearl found.

Gruneberg, H. *J. Genetics* 34:169-89 1937 The position effect proved  
by a spontaneous reversion of the X-chromosome in *D. melan.*



Zeffers, AB+CS Stone  
1937.

Reverse Mutations & the position effect. Gen 24: 73  
The  $w^{m5}$  and its desc. U. Tex. 4032: 190-200

Seidie, et al., Gen. 24:88-1939 Reversal of lethal factors.

Olewe, (P., PNAS 26:452-4 (1946) A reversion to wild type assoc.  
= crossing over in D. melan.

<sup>(lz<sup>1</sup>)</sup>  
Glossy and Spectacle (lz<sup>3</sup>) are sexlinked, recessive, alleles of lz,  
are in ~~the~~ the dl-49 inversion.

lz<sup>1</sup> Bx / lz<sup>3</sup> f ♀♀ x lz<sup>3</sup> Bx ♂♂, 11/5584 2857 ♀♀  
were wild type + dominant to lz<sup>1</sup> or lz<sup>3</sup>. The inversion was not lost.

Ten of the offspring were Bx. ∴ the crossing over occurs  
~~between~~ in the inversion, and has been shown to be between v and lz.

The complementary type was not picked up. The only compound  
which reverts is lz<sup>1</sup> lz<sup>3</sup>

Roblin, Richard O., Chem Rev. 38(2): 255-377 (1946).  
Metabolite Antagonists. ✓

Chemotherapy, American Cyanamid Co., Stamford Res. Labs., Ct.


Fosdick, L.S., et al., JACS 68:840-1946 pressorhormone contg.

nuclear Cl and F.

mil.

Synthesis.

p F-styrene



Lertan, A. SACS 68:835 - 1946. The microbiological synthesis of riboflavin - a theory concerning its inhibition.

decomposition of  $B_2$  increased by addn of Fe (.18-.36 mM/l)  
do. decreased production by *C. acetobutylicum*. Traces of catalase +  
 $N_2 \rightarrow 2O_4$  mic. yield.  $H_2O_2$  unchanged.

Fatajot, R. Rev. Can. Biol., 5:9-47 (1946) L'effet biologique  
primaire des radiations et la structure des microorganismes.

R✓

Wahl, R., Ann Inst Pasteur 72:73-80 (1946) Influence de la composition du milieu sur la bactériophagie.

B<sub>1</sub>, Ca needed by some strains. Causes multiplication & lysis.



Raayer, M + R. Satajet, Ann Inst Pasteur 72: 89 - 1946. Accumulation du nombre de bactériophages en présence de bactéries stériles par irradiation.

*S. paratyphus* Y6R; phage C16. X-Rays 33kV 30mA.

8 - 16000 r/min.  $10^9$  cells irradiated + given doses of 150000 - 400000 r (p5 = 12, 32 resp!!) Tested for ability to form colonies + for titre of added phage.

Non-irradiated cells from ~~5~~  $11 \times 10^3$  to  $146 \times 10^6$  in 6 h. Irradiated ~~from~~ to  $800 \times 10^3$ . There was no increase in irradiated bacteria.

after 4 h. in culture, irradiated bacteria did not support phage.

1 single c.d. / 200 bacteria would allow phage multipl. found.

Increase in phage about same at 400000 as 100000 r.

Expl. on basis of growth giving giant forms.

Woolley, D.W. JBC 163:481- 1946. Reversal of the action of  
phenyl pantothenate by certain amino acids.

Sp. requiring ~~pp~~ ~~ppant~~ are not reversibly by  $\phi$ mit. Sp. synth.  
prot are not protected by it from  $\phi$ mit. H.C. reversed  $\phi$ mit. Amino  
acids which were active were histidine, glut, prol, glyc + asp.  
S. cerevisiae. Similar results in L. casei

Keithwood, S + PH Phillips. JOC 163: 251 (1942). (Geanti-  
riocitol effect of  $\alpha$ -hexachlorocyclohexane.

S. curvicae.

Insecticidal.

Carlson, J.G. *Biol Bull* 90:109- 1946. Polytamni viscosity changes in different regions of the grasshopper mandible during mastication.

Whitaker W.L. PSEBM 61:420- 1946 Postalvein ligations and  
the celiac fistula in the rat.

Grant Mills Ann Arbor

Dumrese, M PNAS 32:36- 1946.

B/1. (called B in this paper). Ca.  $5 \times 10^8$  phage / plate.

u.v. - GE lamp at 92 cm. = 4.2 ergs/sec. Exposed on plate

X-Ray 180kv 25ma 20sd 1/m.

24hr bacteria <sup>!!!</sup> concentrated to give  $10^9$ /cc.

(time spent from "phaging" ???) Irradiated 0 - 4 min.  
to lysis?

(Distinct increase in 4 hours from 0 to 295 of mutations in unrad. ctrl.)  
somewhat greater  $\bar{c}$  u.v.

after 2 hours, increase of 10x in controls

1 min ir.	4.4
2 min	2.2
4 min	1.6.


mutation rate increases until 1-2 div., falls to normal by the 13th div  
(6 hours). Killing not given.

Rubos, RJ + BD Davis JEM 83: 409 - 1946. Factors influencing  
the growth of *H. baileyi* in jejunal media.

Oleic acid (water sol) facilitating surface growth.  
Sera albumin

Ammonia and - citrate - yes.

Meulien, V.  
Z.R. 1:548 1941.



~~21658~~  
21913  
4637

McLowan Clin MJ 48: 305 '41. Mutations Theory Cancer



6, BC Science + Culture 7: 299-1141. Regarding wound hormones.

1. hemolysates of *Toxopneustes*

2. irradiated tumor cells - M. Helf

Pelczar, M.J. + J.R. Porter, Arch. Biochem. 2: 323-329 + 3.

The Nutrition of *Proteus morganii* Amino Acid + Growth Factor Req.

T/O) essentially  $pH 7.2-7.4 \pm NaOH$ .

Cysteine  $M/10^4$

Pant 1 r/ml  
Nic 1 r/ml



(intact)

nicotinic ac. or amide eq. - effective.

Inf. before, medium ca. 2x as dense as synthetic. ( $\bar{c}$  amac.)

cysteine or methionine is only essential amino ac. cysteine better. Others a.g. have little effect.

of aqueous soln. animal materials have a stimulating effect.  
Norvaline, norleucine + all-threonine are inhibitory but reversed by other amino acids.

Purines + pyrimidines had no effect.

Not B's. : B<sub>1</sub>, B<sub>2</sub>, B<sub>6</sub>, choline, betain, foli, pab, inos, panthoi, glutamine...  
all tried  $\bar{s}$  effect.

Try Vitamin C, fat soluble, K, etc.

Bach, Med, State U. Iowa, Iowa City.

Meyers, F.P. + J.R. Porter, J Bact 50: 323-31 (1945) The nutrition of *Proteus morganii*: sulphur requirements.

Basal:

NH <sub>4</sub> Cl	1.	Glucose	5g
NH <sub>4</sub> SO <sub>4</sub>	1	Cystine	24mg
NaCl	1	Pnt	1mg
KH <sub>2</sub> PO <sub>4</sub>	1	Nic	1mg.
K <sub>2</sub> HPO <sub>4</sub>	1		
MgSO <sub>4</sub>	1		
<del>H<sub>2</sub>S</del>			
H <sub>2</sub> O	1l.		

Other >- compounds (cystine 4+).

lanthionine	3+	
Methionine	2+	(variable)
Na <sub>2</sub> S	2+	
<u>Cysteine</u>	variable	!!
homocysteine	2+ var.	

Porter + Meyers. Arch Biochem 8: 169-176 (1945) Amino acid relationships in the nutrition of *P. morganii*.

Altoth allolthreonine increased by 20 am. eq.  
 norvaline by leucine, meth. valine.  
 norleucine (l, d, all) methionine. (leucine 11/150)

Stokes, J L + H Guinness, J Bact 51:570 1946.

*Theca campyriteni* microorganism

abstr.

Finley, H.E. Morehouse College, Atlanta Ga. Brooklyn.  
6(108): 31- 1946.



(R)

Patterns of sexual reproductive cycles in *Hydractis*.



Wyso, O. PSEBM. 48:122-1941. The nature of SA inhibition:

See Elvehjem.



$$\frac{1}{v_i} = \frac{1}{V_0} \left( K_s + \frac{K_s}{K_i} (I) \right) \frac{1}{(S)} + \frac{1}{V_0}$$

then  $\frac{1}{v_i} \propto \frac{1}{S}$

$$\frac{1}{v_i} = k_s \left( 1 + \frac{(I)}{K_i} \right) \cdot \frac{1}{(S)} + \frac{K_s}{V_0}$$

$$\frac{K_s}{V_0} = k_s$$

$$\delta = \frac{1}{V_0}$$

$$k_s = \frac{R_s}{V_0}$$

$$\frac{k_s}{\delta} = K_s.$$

Lewis  
Diener

(Kraysi)

Dubbs

Mellan

Bowen

Shuman + Wang.

Lindgren

Genetics of Patti. Organ.

JID 71:



Jennison, H W + S P Wadsworth J Bact 39: 389-97 (1940) Evaluation of the errors involved in estimating bacterial numbers by the plating method.

Regnier et S. Lambros. Bull Sci Pharmacol  
(do.)

Perry, CA + Epstein. AJCP-T.S. 3: 70-1 (1931). ~~Problems~~ the use of double-poured ~~plates~~ blood plates in the examination of throat swabs cultured for hemolytic streptococci.

Belcher, J. Brit. Real Pflanzen 26:221-49 1939.

Alteration of formation in *Chl. spp.*

*C. variabilis*  
*paradoxa*

Braarud, T. + Bigotthe, F., - BA 7:2826

*C. sp.*  
1 cell =  $2.98 \times 10^{-12}$  g N;  $.98 \times 10^{-12}$  P

Kelso, O. JGP 14:315-37 1931.

\* Harvey Ann Bot 23 181 1909

\* Strehlow, ZBot 21:625-92 1929 *C. paradoxa* x *botryodes*

Kaesi-Wilhelm, best; Berlin

Moeuws, F. Biol Zentrbl. 60: 597-626 (1940). Über Mutationen der Sexualkeime bei *Chlamydomonas*.

~~70°~~ 75°C. 15m. → rate mutations of .3%  
6000r → .002%

60: 143-166 1940. Homoser.

ke *Monostroma*. 60: 225-38 (1940). Über Zygosen-Kopulationen

*M. vittoriae* Kopulationen of gametes → zygote. In 2-3 weeks → sporophyte → 32 haploid zoospores  
ouch!

60: 484-498 (1940) *Polydum granulatum*

~~Whitford~~ Whitford, LA.  
4.) West Port Diesel.

Freshwater algae of No Carolina. (Ohio St  
*C. fenestrata* found: new form

Pitau, K. Z. ind Abst. J. 79: 317-19 (1941). Stat. Moeuws work prob 10<sup>-6</sup>

Comman, I. Bot Gaz. 104: 50-62 (1942). Coleicine

*Chlamydomonas pseudococcus* - resistant to ~0.15%

\*# Moeuws, Z. ind Abst. J. 78: 418 1940 distofutelle. Zoon in

Krogg's Zygote generation by indirect. 10-14d / generation

Leber, L.F. + Muñoz, J.M. (1938) Ethyl Alcohol metabolism in animal tissues. *Biochem J.* 32: 299-307.

"The action of kidney was especially marked in a rat which had previously received alcohol orally for a month."

fasting 2h. diminishes ~~the~~ GETH in liver.

Alcohol tolerant animals have liver with - GETH = 8, at upper range of normal variation.

pyruvic acid stimulated alcohol disappearance, especially in fasted animals (undoubtedly a H acceptor).

Alcohol disappears more rapidly in intact tolerant animal, site of difference might be kidney?

Abdelkhalik, E. et al. (1914). *J. Physiol. Ch.* (90: 369-387).

+ Bassani, E. Studie über das Verhalten des Bluteserums gegenüber Dextrose, Lävulose u. Galaktose vor und nach erfolgter parenteraler Zufuhr dieser Zuckersorten.

Usually, no optical changes noted in any serum tested. So. with serum of fete or amino acids & or peccines.

\* Wildermuth, F. Weitere Untersuchungen über das Verhalten des Bluteserums gegenüber Kohlenzucker vor u. nach erfolgter parenteraler Zufuhr dieses Disaccharids. Versuche ~~an~~ an Kaninchen. 23/24 rabbits responded  
388-418.

The adapted rabbits showed no polarimeter activity on lactose or galactose. " Ein vorläufiger Versuch, durch Verfütterung von Milch eine Änderung des erwähnten Resultates herbeizuführen, war bis jetzt ohne Erfolg. Es wurden noch Versuche mit parenteraler Zufuhr von Milchzucker in Angriff genommen, um festzustellen, ob hier ganz spezifisch spezifische Reaktionen vorliegen."

Used 10 cc 10% sugar. Activity found within 24h.

(1 cc serum ( $n_D$  =  $-0.28^\circ$   $\rightarrow$   $+0.25^\circ$  initially  $\rightarrow$   $+0.16$  at 23h.)

L. Sugriven  
Vesuvius an Hindu.      similar effects with Sans animals.

3.  
P. Ausent.

It is has since been apparent that LA-22 is actually  
genetically a <sup>stable,</sup> single mutant although <sup>it was</sup> isolated in two steps,  
a single genetic

does not revert, and has a complete mutation.



Röhmann, F. (1917) *Bioch. Z.* 84:382 - Über die durch parenterale  
Rohrzuckerinjektionen "hervorgebrachten" Fermente des Pfortaderumms  
von trächtigen Kaninchen.

In repeating earlier work, found adaptive serum sucrose to be  
quite regular. Studied gravid animals to determine relation with  
lactogenesis. Regularly found sucrose in 7-10 days & sucrose disappears  
from urine.

v. 57:380 (1913) 61:464 (1914); 72:26 (1915).

Memmer, R.H.A., (1906-7) On the presence of lactase in the  
intestine of animals and on the adaptation of the intestine to  
lactose. J. Physiol. 35:20-31.

For lactose metabolism:

JBC 81:541- (1959)

80:33-36.

see also

JGP 19:879 Lactose synthesis in mammary gland.

JPhys. 71:342

Colby. Disposal of intravenous lactose in rabbit

1 gm. adm. Unfermentable sugars returned to urine in 3h.

> 75% accounted for by the urine as non-ferm. red. sugars

Insulin had no effect. Urine resulted in only slightly delayed  
removal. No blood lactose found.

Walteris J. rabbit in woman  
confinement

Lactosuria seen during

Plummer did not find adaptation to lactase  
young animals contain lactase which is lost in later life

does not accept Weirland's conclusions on presence of amylase in  
adapted fowl intestine

Potter, O.R. + Klug, H.L. (1947) Dietary alteration of enzyme activity in rat liver. *Arch. Biochem.* 12: 241-248.

High fat diet did not increase citric acid relative activity of liver, ~~not any part of~~ fat fed liver showed marked decreases in octanoin oxidase when lysed. Succinylase  $\downarrow$  in high fat + high carbohydrate animals.


what is SBC in pures.

Lighthbody HD + Klemmair A (1939) Variations produced by food differences in the concentration of arginase in the livers of white rats. JBC 129:71-78.

High protein diets caused a) increase in size  
b) increase in relative arginase conc.

Gelatin augmentation caused b)  $\bar{s}$  a).

Warburg, O. & Christian, W. (1942) Isolation & Kristallisation des Gärungsfermentes Endose. *Beih. Z.* 310: 384-421.



Determined spectrophotometrically at 240m $\mu$  in .5 cm cell,  
 $\epsilon$  3 ml M/300 also combined  $\epsilon$  3 $\mu$ .

Half saturated  $\epsilon$  MgSO $_4$  in phosphate buffer at  $2.8 \times 10^{-3}$   $\mu$ M 6.74  
 $\text{HCO}_3$   $6.1 \times 10^{-4}$  7.34.

3 hypotheses for F inhibition:

1. binds ~~to~~ Mg.
2. displaces substrate from enzyme Mg
3. a MgF compound displaces Mg. 3- affirmed.

When the product:  $(\text{Mg})(\text{PO}_4)(\text{F}^2)$  has same value, inhibition is same.  $\epsilon$  Mg  $> 4/100$ , i.e. inhibition was noted.

6 for 50% inhibition,  $3.2 \times 10^{-12} (\frac{4}{2})^4$

Arsenate replaces phosphate. Pyrophosphate cannot, but is itself inhibitory.  $\text{Zn}$ , a few compounds were also inhibitory.  
 Carboxylase is inhibited by fluoride at higher conc;  $\text{PCy}$  had no effect.

Wilson, W. J. (1910) Variation among bacteria. Brit Med. J. (2), 1909-1910

Understood selection vs. slow fermentation.

see Adams  
"Principles of Pathology"  
1908. I: 104.  
and J. Exp. Med. 4: 349 (1895)

is intermediate coli-typhi related:

Prompt (< 2 da) fermentation of lactose at 22°. Negligible >> 1<sup>hr</sup>  
at 37. See also J.P.B. 14:1 (1909) re dulcitol. Showed  
no agglutinin associated with the lactase. lactase diff. test

at 37, MHL, Mal and Glu fermented & gas

- I. The utilization of lactose by *Escherichia coli-mutabile*. Deere, C.J., Dulaney, Anna D., and Michelson, I.D. J. Bact. 31: 625-633 (1936).

White form of Ecm uses very little lactose (determined as reducing sugar with Cu) before the red forms appear.  $\text{NH}_3$  production indicates that amino acids are used as C source if lactose is unavailable

- II. The lactase activity of *Escherichia coli-mutabile*. ib. 37: 355-363 (1939).

Used Shaffer-Somogyi (JBC 100:695-713 '33) method, with Reagent # 50 and 15 minutes heating. Thymol used to sterilize heavy cell suspensions (req. 1 hr.) Dry cells prepared after Morrison & Hisey (JBC 117: 693-706). Substrate was 50 ml  $\frac{1}{2}\%$  lactose in 1% acacia an M/10 P buffer 7.0-7.2.

Dried cells suspended in 25 ml 2% acacia in .2M P buffer, 10-20 mg thymol added and incub. 37 1-1 $\frac{1}{2}$  h. 25 cc. 1% lactose added, and samples taken for analysis. .01% Cu used to stop enzyme action. Activity expressed as  $u = 2.5$  mg lactose split / 12 h/ mg.

Lac $\nearrow$  grown on lactose had activity ca 2.8 if grown on lactose; 0.2 on plain agar, 0.1 on glucose. Lac- had activity of 1.0 on lactose, etc. on others. No difference whether dried or not. These values characterize the Lac- itself, as no Lac $\nearrow$  were seen at this interval, on Endo's agar.

- III On the activation of the lactase of *Escherichia coli-mutabile*. Deere, C.J. J. Bact. 37:473-483.

"Earlier experiments led us to believe that the antiseptics employed "activated" the lactase which was present, but inactive, in living growing cultures of the non-lactose-fermenting (white) form." Later found that drying would also activate lactase while only partially inhibiting glycolysis, so that  $Q_{O_2}$  might increase

Garrett white: /plain agar:	Wet:	Lac 11.7	Dry: 30.7	
		Glu 139	91.7	
	/Lac	Wet:	Lac 19	72.6
			Glu 136	132
			-- 9	
Red: /plain		Lac 19.2	42.3	
		Glu 117	88.9	
Red:/Lac		Lac 128	1.8	This prep. was obviously overdried. but may have been too acid.
		Glu --	1.9	
		-- 7		

Ex tracts of dried cells contained demonstrable lactase.

No valid test was made of the possibility of lactase activation in Lac $\nearrow$ , but he concluded that adaptation was based upon increased permeability rather than increased enzyme.



Papacostas G + J. Gaté - Les associations microbiennes :  
Leurs applications thérapeutiques .  
Devient mix culture phenomena

W. Harris, Anna Harris 1951 Degeneration and regeneration  
of antibiotic-producing strains of *Streptomyces griseus*  
(Kranzberg) Waksman + Henrici. M.S. Thesis U of W.

Yeast glucose agar Y. Ex. 10 Glu 5  $K_2HPO_4$  Agar 15  
tap water

Maltose (or starch) Spor. Agar (pH 6.8-7)

Maltose 10

Tryptone 5

$K_2HPO_4$  .5

NaCl .5

$FeSO_4$  .1

Agar 20

H<sub>2</sub>O

more stable. Sporogenesis restored in this medium.

*S. gressia* *reflexus* + *indica*


B13

B21	Glucose g.	10	20	<del>15</del> <del>20</del>	
	(NH <sub>4</sub> ) <sub>2</sub> HPO <sub>4</sub>	4			
	CaCl <sub>2</sub>	.4			
	K <sub>2</sub> HPO <sub>4</sub>	2	2		Lee Dulany et al. Mycologia 1949
	MgSO <sub>4</sub> · 7	1	1		Savage ) Bart 57:429
	NaCl	5			
	FeSO <sub>4</sub> · 7 mg	20	10		Carvajal Mycologia 1948:7
	ZnSO <sub>4</sub> · 7	10	10		Kelmer ) Bart 56:157 57:73
	seawater		Ca 5		
	pH 7		Mn 5		
	Sod lact		8		
	NH <sub>4</sub> NO <sub>3</sub>		2.5		Walsman: Streptomyces
	CaCO <sub>3</sub>		1		

Spore suspensions:

Serial potato dextrose agar. 7-10 days 30°. 5ml H<sub>2</sub>O, gently shake. suspensions shaken with 10g. "glow beads". (480 s.p.u. 2 min) diluted with Neurolog OT to give 1:1000. Filter aseptically through cotton.

Summary of



UV - p.s.  
45 cm  
Stirling

Williams Smith, H., (1948) Investigations on the typing of staphylococci by means of bacteriophage I. The origin and nature of lysogenic strains. J. Hyg. 46. 74-81.

A number of coagulase +,  $\phi 420^s$ , strains were studied. Many were mutually lysogenic. 7/23 were lysogenic for the other 16., and sometimes mutually. None of these  $420^s$  types were  $\lambda$  for other strains. Presence of  $\lambda$  did not necessarily confer sero-resistance. Very few resistant were non-lysogenic.

Williams Smith, H., (1948) II. The significance of lysogenic strains in staphylococcal type designation. J. Hyg. 46: 82-89.

a) Mixture of  $a(\lambda_1) + b(\lambda_2)$  led to the production of new phage types,  $c(a, \lambda_1, \lambda_2)$ . A genetic classification was attempted with limited success. Much of the resistance pattern depends on the  $\lambda$  carried.

Cowles, P. B. (1931) J. Bact 22: 119-123. The recovery of bacteriophage from filtrates derived from heated spore suspensions.

1. *B. anthracis*. Reduced  $\lambda$ . Filtrates from cultures heated to  $90^{\circ}$  10 min. were  $\lambda$ ;  $95^{\circ}$  survivors were not, at least from isolated colonies.
2. *B. megatherium* 899 (de Jong) Spores survived  $90^{\circ}$ , and "all colonies ... showed ... bacteriophage".
3. *B. subtilis* (d'Helle) survived  $90^{\circ}$  10 min. or  $100^{\circ}$  5 min. Some, but not all, of the spores carried  $\lambda$ .  
 $75^{\circ}$  10 min. inactivated all the free phage used.

Regards as evidence against spont. generation of  $\phi$ .

Flu, P.C., (1938). Etude sur le bacteriophage du *Bacterium megatherium*. Ann inst Past 60, 610-632.

From summary: Used de Jong's 899 as lysogenic; 338 as indicator.

a) found less phage than bacteria, in contrast to Wollmans

b) very young cultures carry phage also, but saline destroys the phage and prevents its filtrability.

Wollman, E. and Wollman, E., (1938) Recherches sur le phenomene de Twort-d'Herelle. V. (Bacteriophage ou autolyse heredo-contagieuse). Ann inst Past 60, 13-57.

lysogen superior. have rel. low titre

phage ca = bacteria argue that phage particles exist as such  
in bacteria  
phage survival at division

not compatible i parasitism L'existence de "phages"  
de la fraction lysogene et la production de novo des particules corporelles  
bacteriophages paraissent demontree l'origine endogene de  
ceux-ci

Phage.  
Summary.  
Burnet, F.M. & McKie, M. (1929). Observations on a permanently lysogenic strain of *B. enteritidis* Gaertner. AJMS 6:276-284.

Lysogenicity determined by growing test strain with indicator, heating to 56 for 30 mins to kill bacteria and plating on indicator for plaques. Titters of  $10^7$  -  $10^8$  often obtained in most isolates; others showed  $10^3$ - $10^4$ .

Repeated washing continued to liberate phage. After almost exhaustive washing with saline, distilled water liberated additional large quantities of phage. Lysis by other phages diminished the yield.

Lysogenicity was found to be permanent. "The permanence of the lysogenic character makes it necessary to assume the presence of bacteriophage or its anlage in every cell of the culture, i.e., it is part of the hereditary constitution of the strain.

Rough enteritidis produces the phage although it will lyse only smooth cultures of other organisms.

A mucoid resistant variant of the enteritidis to phage 13 was found to be lysogenic of 13 as well as for gallinarum. The mucoid strain was unstable and gave off rough and smooth colonies.

ib. Type differences amongst staphylococcal bacteriophages. 6:21-31. 4 phages found for a white coccus "SF". Some resistant variants were aureus pigmented, but nonpathogenic. (Among the phages was C-C'- see induced lysogenicity.)  
/B is C-resistant.

Burnet 1932 JPB 55:851

A B C D N phage types from BD (groups B and D)

A: hole at margin, filled center

B: smaller, layers, uniform.

serol. uniform.

serol. heterogeneous

About 50% para B → A type only.

see Burnet 1930a

JPB 33:647

enteritidis → B most usually

typhimurium → A, D, N.

A+B are specific for smooth!

C is SR

gallicum

D, N are SR or R.

rough strains may often produce S phages.

BTM strain (enteritidis?) → phage S<sub>1</sub> (A phage) This is specific for smooth BD (evidently no action on para A).

A phage from para A did not attack any out sanguis and 1 enteritidis.  
B<sub>1</sub> (antigenic value?) role of I?

supports common origin of enteritidis, and para B with later divergence of somatic antigen (does not refer to 'common XII component').

Argues ecol. advantage of symbiosis

(over):



para C  
highly pathogenic  
to mouse!

superstifer - Weischild VI - VII

"European" superstifer 5/8 tyrogene for smooth or rough sang.

Other rarely tyrogene for super, but did not on typhi suis.

typhi suis (F12) best indicator.

para C  $\Rightarrow$  only FT2

most others (e.g. Thompson) also  $\Rightarrow$  second R phase

2 serological and resistance types: H (Weischild) +  
S (superstifer)

Range of action not clear e.g. interaction not tested

Burnet + Fresh (1936.) 14:27-38.

Culture	x-resistance						Absorption by heat-killed cells	
	A	B	C	C'	D	Au1	C	C'
SF	+	+	+	+	+	+	+++	++
SF/C	+	+	-	-	+	+	-	-
SF/C'	+	-	-	-	+	+	-	-

SF and SF/C are serologically identical, SF/C' distinct.

If SF is spread fairly heavily on dense C, no loss of colonies, but SF/C found.

SF + stated C, then excess C'.

Explosive production of C grown on SF cultures, infected with a few particles  
Do. single bursts, 80-150 per burst, in 10-90 mins.

C' appeared in older cultures of SF/C, reaching a peak of 50%.

SF/C/Au1 remained lysogenic; SF/C could not be disinfectant by

anti C serum. SF/C colonies were noted in the center of C' plaques

SF/C/B did not liberate C' mutants.

Estimates 10-20% contacts to become lysogenic.

See). d'Herelle, F & Rakietin, TL. (1934) JID 54, 313.

Bruce White, P. (1937) Lysogenic strains of *V. cholerae* and  
the influence of lysogeny on double phage activity. *J. Nat'l Bur. of  
Stand.* 44:276-278.

Phage LL $\phi$  acts weakly on certain strains. Addition of lysogeny  
(egg white 1:25) enhances action to give more active filtrates.

(Bacteria)  
LL-resistant strains of agglutinable *V. cholerae* are invariably  
 $\lambda$ -infected with it. Most existing lysates are therefore probably contaminated  
with it.

~~Some~~ Chinese strains were sensitive could be made lysogenic  
El Tor and other vibrios ~~to~~ were omitted  $\lambda^+$  or  $\lambda^s$ .

On agar, no lysis was seen with LL $\phi$  on Rough vibrios, but  
the phage multiplied and became lysogenic. "blockade mini-  
mity" interpretation:

cf. Doorebos

Fester, L.B. (1945) A bacteriophage for *Pseudomonas pyocyanea*.  
↓ *Bact* 50: 301-303.

Evans, A.C. (1940) The potency of nascent streptococcus bacteriophage B. J Bact 39: 597-604.

phage as released from lysing bacteria more active. Lysis?

(1942) Technique for the determination of the sensitivity of a strain of streptococcus to bacteriophage of type A, B, C, & D. J Bact 44: 207-~~208~~ 209.

Phage references:

CRSB.

Lomstedt

125:846 ~~126:~~ 127:962 128:379  
129:151,267 130:602,144

φ · X · 174

138:497

See also

JPB 58:259

J Biol 54:313

Proc Soc 48:359 (format 4)

Geldmeester, E. (1941) Z. Balet. (I), 147: 417- ~~4~~

~~Robertson~~ d'Heulle, F. & Robertson, T. L. (1934) J. I. D. 54: 313

Quelen, A. (1948) Lyse bacterienne par un filtrat bacteriophageique  
sans multiplication des corpuscles. Ann. IP 75: 472-484

C16 - lysis & plaque formation on paratyphoid Y6R

on coli 36, however, conc. phage reaches a sterile area, but when  
spread, no plaques are formed, only a granular growth.

It is not regenerated from coli 36. (Sumet). Is readily adsorbed.  
I show by mixing cultures to eliminate adsorbed phage. Cells are lysed  
by microscopic examination in liquid medium.

Title of C16 does not increase on coli 36, but does on dys.

Considers possibility of "lysin". Shows same behavior when grown on other  
hosts. ~~Host bacteria do not lyse~~ coli 36. Phage autolysis inhibits  
lysis. Lysin agent is removed by adsorption with sensitive Y6R.  
bacteria

Does not show numerical relationships of adsorbed to bacteria  
killed.



Eldemester, E., & Milfeld, I. (1941) Beitrag zum Bakteriophagenproblem.  
Z. Bakt. (I) Orig., 147: 417-437.

Most intestinal contents carry phages (77% on dys., 7% on  
para B; 5% on S. typhi.) The latter are more often found in Salmon.  
convalescents

Refer to earlier work Z. B. 91:12 (1923)

" dass in den lysoresistenten Kulturen immer einige wenige  
lysosensible Keime vorhanden sind, welche zur Entwicklung von Phagen  
ausreichen. Experimentelle Beweise für diese Annahme sind jedoch  
bisher nicht erbracht worden." Manywigs colonies of coli 88 tested.

Believes in growth without bacterial destruction. Diss. sequitur.

Tested  $\lambda$  by filtration of suspensions.  
32/50 (64%) of a variety of *Salmonella* strains tested were  $\lambda+$ , usually  
best for homologous types. S. typhi, Para B, dysenteriae, para C, *Fluorensis*

11/30 (37%) of dys. tested were  $\lambda+$  (9E, 1Y, 1Shiga, 1Fluorensis,  
usually for homologous type.

5/16 cholera  $\lambda+$ , specific for vibrio.

Coli  $\lambda$  usually active on dysenteriae.

Believes in activation of latent  $\lambda$  rather than infection  $\bar{c}$  with  $\lambda$ . Opposes  
virus theory.

Animal cultures can be temporarily  $\lambda-$ .

d'Herelle, F., + Kalmieten, T.L. (1934) J.I.D. 54:313-344.

Mutations as governing bacterial characters and ecologic reactions.  
also book.

Reduced lysogenicity. [See Malone, A.H., and Sakari, M., Studies on Asiatic Cholera. Indian Medical Research Memoirs #14, Calcutta 1930: Thebent + Spirals I.

*S. enteritidis*, ATCC Oany<sub>27</sub>, 404. stated to be  $\lambda^-$ . Lysogenicity was induced by addition of a lysin  $\phi$ . Activity of  $\lambda$  became attenuated by daily transfer over several months. Some cultures became partially sensitive, especially after 150 transfers. [S.E. not isolated?]

With  $\lambda_1+$ ,  $\lambda_2$  could be added.

Some of the symbiotic "mutants" are mentioned.

Nicolle, P., Grabar, L, + Sibert, P. (1946) AIP 12: 81~~4~~-88.

Fréquence de la lysogénéité et moindre fréquence apparente de la lyso-sensibilité parmi les bacilles paratyphiques B.

31 tested for  $\lambda$  on ~~3~~ *Shigella* indicators. strain 12, and to 1 + 9.

26 were  $\lambda+$  (71%) With one exception,  $\lambda+$  were resistant to  $\lambda_I$ ,  $\lambda-$  were sensitive. The exception was on old very rough culture.  
↓  
2 exceptions.  $\lambda$  from strain 1 and strain 9 shown to be different, serologically & in host range.

Bordet, J. + Bordet, P. (1946) Bactériophagie et variabilité  
microbienne. AIP 72: 161-173; 321-334.

S( $\lambda$ -)  $\rightarrow$  R( $\lambda$ +), especially in <sup>absence</sup> ~~presence~~ of Ca.

"excès de calcium entrave l'apparition du type R producteur de principe".

Complete Ca deficiency (oxalate 20 drops 2.5% / 5ml). also prevents the change.

Tests for the  $\lambda$  involve brief heating culture. [May have been resorbed!]

See Hadley 1924 J.I.D. *Pyocyanus*  $\lambda$ ]

Lisbonne's bact. at 37° has a metallic sheen, "gleineuse" at 10-12.  
cells capsulated & metachromatic material (toluidine blue).

Change does not require Ca. Cold bacteria have not produced

$\lambda$ , reappears in 24h. at 37.

Lisbonne's *indes Stigma lysogeni*. antiserum does not remove  $\lambda$   
although phage is inhibited. Lysis by  $\lambda$  is inhibited by oxalate,  
but cells are not decolimated.

Write for strains ]

Fisk, Roy, T. (1942) Studies on staphylococci. I. Occurrence of bacteriophage caucis among strains of *Staphylococcus aureus*.  
J. Inf. Dis. 71: 152-160.

Took a 4mm loopful over an area of 1x6 cm. Spotted loopful likewise used in both directions; not always seen reciprocally. Incubated 5h. at 37°, then at room temperature. Used zephiran 1:50,000 - 1:150,000 to sterilize lysates. [used milk agar for chromogenesis: 30cc strains milk + 70cc 15% agar, mixed after autoclaving.]

With 45<sup>2</sup> combinations, 43 phages lysis was found.

No lysogenic combinations were found in coagulase-negative, albus strains. Ultimately found that 19/43 = 44% of coagulase positive strains carry  $\lambda$ . Considerable specificity found. Reciprocal lysogenesis was not observed here. But sequences such as:

69 → 47 → 44 → 68 → 49  
                    ↖          ↗          ↖          ↗  
                    77                    77

24 groups of  $\lambda$  noted. None active on albus.

5 frankly lytic cultures were found.

II. Identification of *Staphylococcus aureus* strains by means of bacteriophage. 71: 161-165.

showed that staph. from related series give some responses to a series of 27  $\lambda$  isolates as follows.

See Amer. J. Hyg. 40, 232-238 (1944) for III.

Thomas, R.C. (1948) Ohio J. Sci. 48(3):102-106. A method for removing transmissible lysis from secondary cultures of bacteria. L. Ohio Ag Expt Sta - Wooster).

Exposure of lytic cells to nucleic acid from various sources gave colonies reacting with original lysis. Saw lysogenic (?) bacteria with 2/9% NA in H<sub>2</sub>O. R. temp 1-12h. Poured plates and tested colonies.

Science 88:56-57 (1958). Transmissible lysis in water extracts of seeds.

90599  
PS

Phytopath. 30: 602-611 (1940) Additional facts regarding bacteriophage lytic to *Agrobacterium stewartii*.

Phage strain resistant corn. Typical phage reaction. "Transmissible in seeds".

NR

Mckie, M. (1934) The lysogenicity of coliform bacilli. H.J.E.B.M.S.  
12: 169-175.

82 coliforms and 9 atypicals tested for lysogenicity by testing filtrates.  
> 31% gave phages in the primary filtrate, and in several cases there  
were two or more phages. (52 & from 37 &+). Rough Flexner VR dysentery  
was most susceptible. (38 & active). 13 were active on rough

<sup>398R</sup>  
gallinarum.

15/52 were weak and lost on passage

28 on Flexner VR

3 as coli KR, weak on Flex VR

3 on 398R, — on VR

3 specific S' & on 398S; shiga S and YS.

Complex cross-resistance

Dunbar, James M. (1948) Bacteriophage typing of untypable  
*Salmonella typhi* organisms. *Nature* 162:851. (Nov. 27)

Many cultures are contaminated with an "anti S" phage, rather "rough".

When reduced, "agglutinations" are characteristic and ... type to  
I and IV & ... and highly specific Type II S phage. Growth in anti S  
serum is used to type those previously untypable strains.

These contaminated bacteria are "interfered with" by specific phages.

"Central Pathological Laboratory  
M.E.L.F."



Taylor, H.E., (1949) Additive effects of certain transforming agents from some variants of pneumococcus. *J. Exp. Med.* 89:399-424.

Small scale (1500 ml) preparations of TP described. Bovine Serum Albumin is necessary factor.

Strains: A66 (SIII)  
R36A (R) from D39 SII. Never reverts and readily transformed.  
ER Extremely rough from R36A. Stays in aggregates.  
SIII-1 ← SIII  $\xleftarrow[\text{TP}]{\text{ALL}}$  R36A.  
SIII-2 " " .

ER can revert to R, especially in liquid medium. Stable on agar on shallow layers.  
When SIII TP is added, R is regularly formed. BSA needed for regular effect.

RTP activity only from SIII and R36A bacteria. ER DNA and other NAs inactive.  
In view of parallel  $\bar{c}$  S transformations, the ER  $\rightarrow$  R effect is regarded as an induced change, not selection.

anti R prevents ER  $\rightarrow$  R. Thus it can be shown that ER  $\rightarrow$  S with SIII. "like other morphological mutants obtained from R36A, ER is 'incompetent' to undergo direct transformation into the SIII condition.

ER  $\rightarrow$  R  $\rightarrow$  S was obtained in one tube by adding ~~5~~<sup>3</sup>/<sub>4</sub> anti R after 5<sup>3</sup>/<sub>4</sub> h. and using SIII TP. ~~or TP from~~ R36A TP gave only R.

type-specific antiseria inhibit transformation of R36A →  $ST4^+$   
but is essential for  $ST4^-$

$\text{SIII-N}$  (normal) - 1 and - 2 differ in amount of  $\text{III}$  substance.

anti- $\text{III}$  enzyme makes - 1 and - 2 cultures rough. ~~Does not~~ also less effective on  $\text{III-N}$ .

$\text{III-1}$  requires very little antibody for agglutination. Does also agglutinated by R. No quelling. Not mucoid.

$\text{III-2}$  mucoid, quelling but less  $\text{III}$  than  $\text{III-N}$ . Not virulent.

TP from  $\text{III-1}$  and  $\text{III-2}$  transform R36A to comparable S type. and ER to R.

Roughs obtained from  $\text{III-1}$  and  $\text{III-2}$  were transformable to  $\text{III-N}$ .

When mixtures of  $\text{SIII-1}$  and  $\text{SIII-2}$  were applied together,  $\text{III-N}$  bacteria were found as well as the -1 and -2 types.

$R \xrightarrow{1} \text{III-1} \xrightarrow{N} \text{III-N}$ .

$R \rightarrow \text{III-2} \nrightarrow \text{III-N}$ .

$R \rightarrow \text{III-N} \nrightarrow \text{III-1}$   
 $\nrightarrow \text{III-2}$

Does not believe this goes through R as mediate.


TP from  $\text{SIII-N} (\leftarrow -1 \leftarrow R)$  shows no signs of inducing  $\text{SIII-1}$  from R. They show no signs of the intermediate stage.

$R \rightarrow \text{III-1} \xrightarrow{\text{TP III-2}} \text{III-2}$   
 $\xrightarrow{\text{TP III-2}} \text{III-N}$

Summation may or may not take place

No statement whether the  $\text{III-N}$  type prepared by summation is "heterozygous".

TP1  
TP2  
TPN  
TPR



Does not III-N from summation contain both transforming principles? [Evidence that intertransformations do not go through R?]

Austrian, R., and MacLeod, C.M. (1949) J. Exp. Med. 89:451-460  
Acquisition of M protein by pneumococci through transformation reactions.

I - SVI }  
III - A66 } used. { I -  
                  }                  { III - 3M

The "Dawson Rough" seems to correspond to Taylor's ER.

When <sup>or-36A</sup> II - R36NC } (II; 2'M) was transformed with  
III - A66 TP, III 2'M was obtained.

do,  $\in$  TPI transformation.

Dawson<sup>ER</sup> Roughs were obtained from R36NC.

Some of these were transformed to III 3M.  
from cells which <sup>obtained to</sup> still had some 2'M (serologically detectable) <sup>III 2'M</sup>. These may arise

This dequiformation does not take place so regularly. Griffith Roughs not tested for TP.

In vivo: ER + vaccine I <sup>2/10</sup>  
                  + vaccine III <sup>2/10</sup>

Concomitant acquisition  
of M3 protein noted in  
one case each.

↓  
R  
↓  
II.

Byatt, Pamela H., Jaun, G. J. & Salle, A. J. (1948) Variation in pigment production in *Staphylococcus aureus*.

Extracts of chromogenic *S. aureus* (strains??) ~~did~~ transformed white strains to colored. Transformed strains retained bac - character.

Burnet, FM + McKie, M. (1929) Type differences amongst  
Staphylococcal bacteriophages. Aust. J. EBMS. 6: 21-21.

SF: MR - Lact + gel - .


Phage B gave three kinds of SF/B: opaque white; colorless +  
translucent; frankly aureus. 1B was also resistant to C.

SF/B was non-lysozyme, but after being kept on agar for some  
weeks gave rise to papillae some of which were of the chalky white  
type, others frankly aureus. Either in this way, or directly  
... SF/B ... the aureus type of SF/B could be obtained.

Goldstein, Abram (1944) The mechanism of enzyme-inhibitor substrate reactions. J Gen Physiol. 27:529-580

Non-competitive.

E = total enzyme  
+ = free



(1)  $K_I = \frac{(E_f)(I_f)}{(EI)} = \frac{(E_f)(I - EI)}{(EI)}$        $E = E_f + EI$   
 $= aE + EI$

(2)  $I = K_I \frac{(1-a)}{a} + (1-a) E$       Let  $I' = \frac{I}{K_I}$ ;  $E' = \frac{E}{K_I}$   
 = "specific concentrations"

(3)  $I' = \frac{1-a}{a} + (1-a) E'$  (2ml B)

(free) (combined)

Zone A:  $I' = \frac{1-a}{a}$  (i.e.  $I \approx I_f$ )  
 $E < \frac{K_s}{10}$

Zone B:  $I' \neq I_f \neq EI$

Zone C:  $I' = (1-a) E'$  ( $I \approx EI$ )



$a = \frac{v}{v_{max}}$

$v = k_D(ES)$   
 $v_{max} = k_D(E)$

(3b) (4A) and  $S' = \frac{a}{1-a} + a E'_s$

Most enzyme systems operate in zone A., i.e.  $S' = \frac{a}{1-a}$  (MM equation)

They prefer to plot  $\frac{v}{v_{max}} / \log_{10} S$ . Consider  $1.1 \times 10^{-3}$ ,  $1.25 \times 10^{-3}$ ,  $1.7 \times 10^{-3}$  as good fits for  $K_s$ .

The zone B equation is fitted as follows:

$\frac{S}{a} \neq K_s \frac{1}{1-a} + E$  and  $\frac{I}{1-a} = K_I \frac{1}{a} + E$



$$\frac{V_{\max}}{v} = 1 + \left[ K_s + \frac{I}{K_I} \right] \frac{1}{S}$$

For  $I=0$ ,  $\frac{V_{\max}}{v} = 2$  when  $\frac{K_s}{S} = 1$ . ✓

otherwise, for a given, constant activity:

$$\frac{K_s}{S} + \frac{I}{SK_I} = C$$

$$C = \frac{1}{S} K_s + \frac{I}{S} \cdot \frac{1}{K_I}$$

$$SC = K_s + \frac{I}{K_I}$$

$$Sa = 1 + \frac{I}{K_s K_I}$$

$$aS - bI = 1.$$

# Competitive equilibrium.

$$\frac{E_f I_f}{(EI)} = K_I$$

$$\frac{E + S_f}{(ES)} = K_S$$

$$\frac{(ES)}{E} = a. \quad ES = aE.$$

$$E = ES + EI + E_f.$$

$$\frac{EI + E_f}{E} = 1 - a$$

$$\begin{aligned} EI &= (1-a)E - E_f \\ &= (1-a)E - \frac{K_S a E}{S - aE} \end{aligned}$$

$$I' = \left[ (S' - aE'_S) \left( \frac{1-a}{a} \right) - 1 \right] + \left[ 1 - a \left( 1 + \frac{1}{S' - aE'_S} \right) \right] E'_I$$

Free combined (= (EI)')

If  $I_f \approx I$   
or if  $EI \approx I$

$$I' = (S' - aE'_S) \left( \frac{1-a}{a} \right) - 1$$

$6 A_I B_S$

$$I' = \left[ 1 - a \left( 1 + \frac{1}{S' - aE'_S} \right) \right] E'_I$$

He finds  $\frac{I'}{S'} = \frac{1-a}{a}$  i.e. for  $a = 1/2, \frac{I}{S} = \frac{K_I}{K_S}$ .

$$\frac{1-a}{I'} = \frac{a}{S'}$$

$$\frac{\frac{EI}{E}}{I'} = \frac{\frac{ES}{E}}{S'} \quad \text{and} \quad \frac{\frac{EI}{I}}{ES} = \frac{K_S}{K_I}$$

Hoder, F. + Akano, R., *Z. Vermutl.* 85:423- (1935)

Foley, G.E. and Schwachman, H. (1950) <sup>Rev. 1112</sup> ~~Journal~~  
4: 141-149 Some observations on streptomycin-dependent  
strain of *Staphylococcus aureus*. RR

Bawden, F.C., Kassarjian, B., and Nixon, H.L. (1950) The mechanical  
transmission and reproduction of *Rhizostoma paracumbale* virus.  
JGM 4: 210-219.

Fleming, A., Younker, A., Kramer, I.R.H., & Hughes, V.H. (1950) The  
morphology and motility of *Proteus vulgaris* and other organisms re cultured in  
the presence of penicillin. JGM 4: 257-269.

RR

Eriksen, K.R. (1949) Studies on the mode of origin of penicillin resistant staphylococci. Acta path 26: 267-279.  
From Univ Inst General Path. Copenhagen.

Broth is various P inoculated with varying amounts ( $10^{-1}$  to  $10^{-6}$ ) of a 24 hr. broth culture. Later plated loopful (ca. 0.02 ml) on ~~and~~ agar. With large inocula, secondary growth is found up to  $1/4$  ou/ml; with initial bacteria of  $10^{-3}$ , no sec. gr., but eventually comes up.

"Demerec is not correct and that the resistant bacteria appear only after contact with penicillin for some ~~time~~ length of time."

Reasoning?? Note that with ca  $1/8$  ou/ml and perhaps  $10^{-5}$  ml, any secondary growth was delayed 24-48 hours.

In 6 ~~days~~<sup>tests</sup>, it appeared only after 6 days. "In these cases where the secondary growth appears at such a late juncture, presumably it can be taken ~~that~~ for granted that the growth does not originate from resistant bacteria present in the original culture."

(Some confusion about isolation of pure resistant cultures in testing for stability.)

Found variance in mutant numbers only in 3 ml cultures, not in 15 ml cultures.

# Treatment of recombination in texts since 1948


1950 Clifton Introduction to the bacterias pp 73-75

"Possibilities of recombination of genes by other than sexual mechanisms may exist, and our original definition of bacteria as 'apparently sexless' organisms is still valid." Fair statement of expts. T+L 1947

1949 Burrows et al. p. 184 passing reference  
extensive ~~study~~ for general analysis of variation 12 1947.

Stolzer, B.A.D. (1949) Measurement of rate of mutation of flagellin gene  
phase is relevant to typical *S. typhimurium*. J. Hyg. 47: 398-413.

[ Dept. of Biology & Microbiology, University of Texas at Austin, Austin, Texas ]

Stagnant + culture in used, especially water is, to keep  $\mu$  stable.  
occasional mixed strains are found. Some non-viable strains (<2%)  
were found. Some populations at mutational equilibrium were noted.  
Rate of  $3.5 \times 10^{-4}$  / generation found by D. Vsted   
pharmaceutical. p. 405

KR

Klebsiella - Nobel, E. (1941) *Klebsiella* the significance of the  
ability of *K. typhimurium*. J. Hyg. 11: 495-505.

Layers filterable, possibly, + the ability of *K. typhimurium* to  
infect the mouse possible.

Stern, C. 1936 Somatic crossing over and segregation in *Drosophila melanogaster*  
*Genetics* 21: 625-730.

Minute flies (M/m) show m spots. Originally interpreted as elimination of M-carrying (deficient chromosome). By use of  $\theta$ -translocators, it was shown that the M phenotype (not merely deficiency, covered by duplication) was necessary for spotting. bobbed (bb) spots not found: interpreted as partial elimination.

Autosomal M also cause X-mosaics (~~sn~~  $sn^3$  (singid))


However, the Bld ~~is~~ Minute causes X-spots, but not III-spots!!!!

Effect of autosomal Mw on Notch<sup>8</sup> cf. was studied:

$N^8/y \text{ } \text{f} \times sn^3; Mw/+ \text{ } \text{m}$  Among females:  
 $N^8/sn^3$  9/280  $y/sn^3$  15/381 No difference.  
 $sn$  setae (elim?  $N^8$ )  $\downarrow$  2  $y$  spots  
 2  $sn^3$  spots

No  $ysn$  spots.  $\therefore$  2-strand and no reduction.  $(y/sn)$  suggesting segregation of X-chromosome.

$ysn/++$  flies  $\rightarrow$  110  $y sn$  43  $y$  7  $sn$  spots.  $y$  and  $sn$  simply somatic crossing-over as well as segregation. But no  $y-sn$  twin spots were found, ruling out two-strand crossing over. Equatite reduction is ruled out by absence of  $ysn-sn-y(+)$  triple spots.



Region of crossing-over varies with spot size (developmental stage). Crossing-over to the right of  $o$  in  $ysn$  spots supported by expts. with  $\theta$  translocators. Segregation is probably nearly always equatorial.



bb fails to show segregation in +/bb flies. Assumption of phenotypic masking seemed unlikely.  $\therefore$  Crossing-over to the right of bb considered very rare.

Determined X-ploidy of spots by color of 5-6th abd. segments.  
Most spots in females were XX by color.

### Autosomal mosaic

Under influence of autosomal M.

Secondary Sources:

1. Sorsby "Clinical Genetics"; pp/ 337-40; 313-15
2. Kallmann and Sander 1947. in Hoch & Knight, "Epilepsy". Chap. 3
3. Neel 1947 Medicine 26:115. at 123-125

Acc (3): 25-30% of propositi have family history (5-6x as frequent in parents sibs and children of propositi). monozygotic twin correlation 70%. Quotes Lennox extensively on cerebral dysrhythmia. In 24% of families both parents showed dysr. Obvious complexity.

(2) Examples in animals; also audiogenic seizures. *Lennox:* From Conrad: (incidence figures) %

gen. pop.	childr.	sibs	neph&nieces	dizyg. twins	monozyg cotwins
.3	6.3	4	1.2	3.1	66.6

concordance in twins:

	diz	monoz	
idipath.	4.3	86.3	Thus even sympt. epilepsy has a genetic component. Index twins were restricted to severe hospital cases.
symptom.	0	12.5	

also found consanguinity correlations with mental deficiency, but not with schizophrenia.

From Lennox:

dysrhythmia

general pop	.10
epileptics	.9
par and sibs	.6

in twins, 85% show concordance of encephalo. if monozyg; 5% if dizyg.

1. Similar to 2, but emphasizes consanguinity correl. with psychopathy.

Conclusions: inheritance not simple (probably several different mechanisms). Certainly a very large genetic component in severe cases, from Conrad's twin studies. Most frequent suggestion is dominant with low penetrance, but high incidence of dysrhythmia in both parents of propositi (Lennox) suggests recessive factors also.


(Lennox '47 is Res Pub Ass Res nerv ment dis 26:11)

CC: Dr. Javid

1954  
1/2/54

copied  
MAY 17 1985

### Conjugation in yeast.

Fowell 1951 emphasizes dicauson: mating of cells gives  from which either haploid or diploid or dicauston (i.e.  $\rightarrow$  + and - haploid)


buds may be generated. Tools care to remove profusion buds.

paired 250+/- cells; 30 zygotes formed. 50% zyg.  $\rightarrow$

only haploid. Other zygotes  $\rightarrow$  only 2n. "An investigation of sporulation revealed that nuclear fusion apparently always occurs in zyg. formed by this proc." Renard 1946 also suggest dic.

Also discussed by Gaimann 16; (Pulhennond 25. bot Rev 1940 6:1) Comp. Morph. of Fungi 1928.

Winge: Roberts 1948. Unsuccessful crossing: spores may give haploid cells "before fusing"

W & L figure  spores. But W'35 also shows substantially complete copulation and diploid buds.  $\therefore$  some variation.

But note analogy of Fowell's dic. i conjugation formation.

Karnada, H. Zbl. Bakt. I 118: 304-16 (1930)

S. para B + G+ soil bact → frequent antigenic variations  
in Salmonella → enteritidis; breslau.

JPB 35:851

19 Burnett 1932 Lysogen.

Palauten 69. J Bact 34:285

Andrews 7. Pr. Roy Soc Med 33 Dec 39]

3 Kueger Physiol Rev 16:129

18 Burnett Arch Exp Biol 6:277

8/3

Delbrück JGP 23:443 Adsorption no.

Ext. lysis → loss of virus.

22,365 -

temperature same as for cell divisions

Receptors: 63 - Lurie + Friedel JEM 59:213 ✓

See Burnett 9. AJEM 15:227

J Immun 46:281.

(leave out glucose in virus media)

Tryptose 2% glucose .1% NaCl 1% - pH 7

AD Hickey.

$\frac{1}{8}$  [ .6% agar  
mixture ]

.5 ml phage

2 ml (2-24. bact.  $10^8$  / ml)

3 > virus later; .5 ml mixture + 3.5 ml .7% agar

pour on plate =

back up!!!

Freundzel, J. + Z. Szymanowski, CRSB 117:543-546 (1934)

Recherches sur la Paragglutination: Différenciation des antigènes H et O.

They had shown that P. exhibits a different serological specificity from the "agglutinin composite de Schützge". But the R strains do contain an antigen related to the preceding strain.

~~This~~ paragglutinable strains are homogeneous + repeated re-isolation indicates that the modification is heritable. Only some E. coli are capable of paraggl.

coli-typhoid paragglutination.

The P. coli absorb H-<sup>agglutinin</sup> antigen from anti-typhoid sera. The original coli does not. anti-H was removed by absorption on Stanley. There was little further agglutinin absorption. However, there was still considerable aggl. of coli. ∴ Paraggl. coli has all H antigens, and a fraction of the O of typhi. anti-P coli serum has a low titer on heated typhi. Typhi phages do not lyse (P) coli.

2. Balat (I, 121:448-451 (1931) Paragglutination des Bacc.

Bang mit Typhusserum. —

Zirconi - ctd.

Using para A and ~~the~~ triple, (P) is also obtained with cross-reactivity, but very little in para B.

Could not transform steps.

Relates paracyclization to the

ps. transformation

Smith WE, J Bact. 47:417-418 (1944)

Wahlen + Almader JID 65:147-55 (1957)



Appleby, J. C. J Bact 38:641-51 (1939) Cytology and methods  
of reproduction of two cocci and the possible relation of these organisms  
to a spore forming rod.  
~~Appleby~~

Cocci appeared in a culture of the bacillus.

11

Agri Bact Dept, Univ Reading England

Sex in Bacteria. Literature:

J. Bact 50

Nuclei - El. Micro.

(R)

Bayler, M.B., MO Appleman, OH Sears + GL Clark, J Bact 50: 249-56 (1945)  
Chem. + Agronomy Illinois

Some morphological characteristics of nucleole fact as shown by the electron microscope II. [ See Soil Sci Soc Am. Pt. 7: 269-71 (1942) ]

4-5 granules/cell untreated +  $\bar{e}$  .02%  $N_2HCO_3$   $\frac{1}{4}$  hrs. Attempts  
at staining w.g. M <sup>15 min.</sup> saline left mottled cells. (several transparencies; corres-  
ponding to nuclei? After  $N_2HCO_3$  saline did not remove granules.  
acetone removed granules. also  $HNO_3$ ,  $HCl$

Krayci, B. J Biol 49:475- 1945. A study of ... factors... in ... of ...

low pH n.g.

zones are not found until sugar + glycolytic products are added + also the autogenous comp.

" healthy cells, facing starvation, aerobic ... "

Geo; Green HC J Biol 35:261

U. d. d. 24. 1943

Knausi, G. + S. Mudd J. Bact 45: 347-57 (1943)

Enzell.

The internal structure of certain bacteria.

Apparent <sup>DR</sup> nucleic ac. material in granular form is 5. S. Stenococcus.

Most diploid cells contain 2 granules each.

R.R. Mellon, J. Bact. 10: 481-501 (1925) Studies in Microbic Heredity I Observations on a primitive form of sexuality (zygospore formation) in the colon-styphoid group.

*B. coli* (Nx) In patient benign mucrotropin appeared as filamentous form & "many very large coccus like forms were encountered developing from the filaments."

Broth, peptone-veal - 5% NaCl broth + 1% Na<sub>2</sub> glycerophosphate at pH 6.8 autoclaved; ppt. ~~redissolved~~ filtered + reautoclaved. Ppt redissolved in alcohol. Single cell isolate inoculated into broth 37° 72h. Then at R.T.; streaked out on Endo. (with broth - glyf base pH 8) was incubated at 37° 18-24 hours, periphery of colonies were fungoid & zygospore formation.

"no attempt has been made to study the fate of these spore like bodies".

Similar forms were found in smaller cells.

No convincing evidence of origin from > 1 cell.

Mystic on sexuality + variability  
Does not understand basis of relationship.

Assumes that cell-fusion has taken place. Criticizes Almqvist.

"unless it necessary... to rule out the purely symbiotic influence of the accompanying strain."


10: 579-88 (1925)

Lindgren, CC + Ralph R Mellan, Nuclear phenomena suggesting  
 a sexual mechanism for the tubule bristles.  
 Proc Soc 30:110 - 1932.

Mellan et al Proc Soc 30:80 1932

AFB → fettablegonidia → var AF diplococci →  
 tetradiplococcus → diploths → actastgonidia → R+bc  
 → Stbc

Acetocarium pupus.



Tetranomus



Mauhiel, JG. Contribution à l'étude de la variation en micologie.  
logie. Th. doc. de nat., Nancy 307p. 1932.  
: from Annales Biologiques ✓

temporary variations in pigment in prodigiosus:

La aut. My. in pigm prod.



Brown, F.M. + H.M. Heffron, Science 49:198-200 (1929). Mendelism among bacteria?


yellow "bacillus" *B. lutyus* (Brown)  
 G+ young cultures "sph. refractile bodies" in old cult

a) yeast like organisms associated "both, while annoying at the time have been found to be merely phases in the life history..."  
 b) loss of color.

a) filament formation in old cultures

b) sudden toxin: e.g. Pb.: 8 rods fuse into a mass, staining intensely  $\bar{c}$  fuchsin "symplesm."

Mass ~~then~~ divides into 4 sph. non-staining bodies. Each of these  $\rightarrow$  tetrad  $\rightarrow$  16 "cocci" On transfer to new medium, cocci divide and  $\rightarrow$  rod form.



As white strains, old cultures, or symplesm formation  $\rightarrow$  both yellow + white colonies. Each, be it true in daily transfer.

Single cell isolations of each made of 2 wks. After 11 transfers, "substrains" that showed no change of color from the 1st. single cell isolations were taken and considered to be pure strains of that color. Biologically identical.

Both cultures were mixed. On transfer, almost entirely white. (Fountain Valley School, Colorado Springs, Colo);

isolated symplasmas from mixed cultures on Pb-untreated agar slant  
grown in broth  
and plated out.  $\rightarrow 362:138$  w:y. (8:3)

$\rightarrow w+y \quad w > y.$


In one instance  $\rightarrow$  all white.

V. nuy: single cell isolation after mixed culture???)

1. Pure strains stable
2. Mixed cultures  $\rightarrow$  unstable white colonies.

Assume that there is a diploid segregation in  $F_1$   $\bar{e}$  same name:

1) Should have studied the progeny for variance.



Kowen + Zmick J Bact 44:551- (1942) A factor  
sexual fusion in bacteria

1. yellow + wh. strains of *Phytonoma steudtii*.

a) Look for heterozygous colonies, resulting colonies from  
mixture. Also mated by R, S. No recombination found.


4/20/22 mixed colonies found. Stable on replating  
(this is not unexpected in view of a small % of sticking together).

2. Recombination of characters (haploid)

509 - large, smooth, mucoid, wh. 400 small, rough, non muc.,  
yellow.

Mutations observed in parentals, as frequently, as in mixed  
culture.

↓  
app. recombination.



Data not analysed.  
> 200 000 colonies examined.  
no literature.

See J Bact 33.

Erskay, L + W. Hesselbrock, J Bact 49: 233 - . 1945. Some observations  
on the filterability of *M. tularensis*.

Filter ca 300-350  $\mu$  demonstrated by infectivity + sedimentation.

Imięński A J Bact 52:49:1-5 (1945) On the structure of anaerobic  
bar

Hollande, Arch Protistool. 83:465-608 (1934) Contrib. à l'étude  
cytologique des microbes (Coccidies...)

Diener, L. J. *Bact* 50: 441-458. Morphology + Nature of the  
Pleuropneumonia group of organisms.

(R)

Altme - Weber, E., *Jal J Bact* 50: 291-5 (1945) The effect of  
incompletely inh. conc. of penicillin on *E coli*.  
Dept Lab, Jewish Hosp.  
Brooklyn, N.Y.

Nutrient broth:  
75 units/ml *Stamm* → "bipolar" dephthroids<sup>≠</sup>

Ab. at 300/ml

at 100, mycelium

150 "zygospores"  
200 early small cells.

Ade, P.A. J Bact 51(6): 699-701 (1946) Mutation in certain  
phytopathogenic bacteria induced by acenaphthene.

16 Path  
4E Subclg

*Phytophthora michiganensis* + *Erwinia carotovora*  
acenaphthene saturated nutrient broth. 2 vols 28°.

by *P. mich* "a sudden + complete mutation" → only a wh. shiny  
smooth type of colony. Neither intermediate nor typical forms  
were found after a certain time.

*E. carot* → several types - perid. grayish compact flat colony



Ramchandani, J. C. Ann. Bot. 44: 975-987 (1930) *Serratia*  
in *Bacteria*. III *B. violaceus*

color variations.

· Geib. 40: 2, 43: 579.

→ wh. mutant + reversion.

Hort. EC PRS 389:468 (1917). Morph St. in the life history  
of bacteria.

Breeding? (Centricity.)

Stewart FH J Herp 27:379-95 (1928) The life cycle of *Paramecium*,  
alternate asexual and asexual phases.

Rosen HR Mycologia 20: 251-75 (1928) Variations within a  
bacterial sp - I Morphologic Variations Aick.

"Gurney-Dixon, S. "The transmutation of bacteria" 1919.

"B. mesentericus?" Very particles attached to flagellae were  
seen.  
interpreted as germinis.


No direct evidence of vitality. Filtrates did show culture.

∴

not relevant.

Beebe, J.M. J. Bact 42: 193-223 (1941) The morphology and cytology of *Myxococcus xanthus*. m.s.  
Univ. Ariz. Tucson. (R)

Describes nuclei  $\bar{c}$  2 chromosomes and autogamous fuses before sporulation. Meiosis not observed.



(SUNSON.) *Bubamixobacterium* !!

Nyberg, C. Acta Soc Med Fennicae 12: 1-18 (1930)  
des Bacillus myzoides.

Zur Biologie

Broadhurst, J. *et J. Bact* 27: 48 (1934) SAB.  
*Zygote phoresis in bacteria.*

[Find "Lommel" 1926; Wygodtshchikoff + Mamulowa 1930  
Roux + Terain 1890.]

# Classification of Literature.

## 1. Growth

- a general
- b growth factors
- c " " - analogues
- d antibiotics
- e regeneration - see sp. organs
- f genetics.

## 2. Genetics

- a transmission
- b gene acting as gene; induced mutation.
- c action
- d biochemical, in microorganisms
- e other, "
- f adaptive enzymes.



JPB  
JCP  
JLEM  
JEM  
J Col Res  
J. O. Ch.  
J. Ph. Ch.  
Faraday Soc.  
J. Russ.  
J. Hyg.  
J. Chem Soc.  
Exp Ph + Th.

Enzymologia  
Advances

Biol Rev.  
Zool Rev.  
Q. R. Biol

"Flux" in "specific" proteins not established.

Is order specific???

Is order maintained in derivat.??

Are <sup>sp.</sup> no. proteins made by enzymes??

Spont. reactivation.

"Specificity" - <sup>+ substrate:</sup> enzymatic -  
immunologic  
[genetic].  
How else.

Boils J 39(5):) 1945.

$$B \xrightarrow{m} B/r$$

$$r \xrightarrow{m} r'$$

$$r' \propto B, B/r$$

$$Br \xrightarrow{m} Brd.$$

$$d \notin Brd$$

$$r' \propto Brd.$$

$\therefore$  this mutational resistance is specific.

$$Bd, \xrightarrow{m} Bd, r'$$

$$r', r \notin Bd, r'$$

$$d, \propto Bd, r'$$

!!!

$$Bd, r' \xrightarrow{m} Bd, r'd$$

resistant to  $d, r, r'$ .

The mutant viruses are all active in original host!

$$B \xrightarrow{m} Bd_1.$$

$B \xrightarrow{m} Bd_2$ . small colony mutant on nutrient agar!

$$T1 = d$$

$$T2 = r$$

Some Aspects of the Nitrogen Metabolism of a Lysogenic  
Strain of Bacillus megaterium

The total nitrogen of the infected and uninfected cells was determined by the semi-micro Kjeldahl technique. The uninfected cells were found to contain a larger amount of total nitrogen than the infected cells. It was found that the desoxyribonucleic acid (DNA) content of the infected cells was slightly higher than that of uninfected cells. The presence of the virus in lysogenic cells in the immature form is believed to be the explanation for the slightly larger amount of DNA in the infected cells.

The technique of Feldman and Gunsalus was used to study the activity of the transaminases of B. megaterium. Pyridoxal-PO<sub>4</sub> was required as a coenzyme and a number of amino acids could serve as amino donors.

The effect of sodium azide, sodium fluoride and iodoacetate on growth and virus production was studied. NaF had little or no effect in the concentrations used. Sodium azide and iodoacetate depressed growth and virus formation. The inclusion of ATP in the medium, along with the inhibitor, produced inconclusive results.

Studies with N<sup>15</sup>-ammonium carbonate showed that after a 30 min. exposure the amount of N<sup>15</sup> taken up by both strains was the same. A study of N<sup>15</sup> distribution in amino acids, purines and pyrimidines was done also.

A complex amino acid medium was developed; it supported better growth of the lysogenic strain than nutrient agar, the amount of virus produced was significantly less. The addition of asparagine or adenine to the amino acid medium increased virus formation.

The two strains of B. megaterium were grown on synthetic media containing purines, pyrimidines and nucleotides as the sole sources of nitrogen. The uninfected cells showed good growth on these media, but the growth of the lysogenic strain was only fair. The lag phase could be shortened appreciably with larger inocula, i. e., direct transfer. Little or no ammonia was liberated, and there was little change in pH over a 48 hour growth period. Attempts to isolate and identify intermediate products of metabolism by chemical methods, paper chromatography and UV irradiation were unsuccessful.

References

- Feldman, L. I., and Gunsalus, I. C. 1950 The occurrence of a wide variety of transaminases in bacteria. *J. Biol. Chem.*, 187, 821,830.
- Lwoff, A., and Gutman, A. 1950 Recherches sur un Bacillus megaterium lysogene. *Ann. inst. Pasteur*, 78, 711-739.

4/52

Rhoades *ijap*.  
"Kuntze's ... 1950"

①  $I_j i_j$  shows no effect.

② Give  $i_j i_j$  segregants

which are stupid. ③ Stupid segregants  $\times I_j \rightarrow$  stupid

$I_j i_j F_2$ . ④ Stupid  $F_2 \times \bar{I}_j \bar{i}_j \rightarrow$  stupid  $I_j I_j$ . (genetic marker.)

1.  $\therefore$  Pleistid abnormality is inherited at least two generations in presence of  $I_j$ . Further selfing of  $I_j I_j \bar{i}_j$ .

② Virus is brought in from  $i_j I_j$  stock. This virus has no effect in presence of  $I_j$  but can be propagated in presence of  $\bar{i}_j$ .

See Jenkins MT J Her. 15: 467-472.

Notes that green  $\bar{I}_j i_j$  plants show "conditioning"

In summer-grown plants,  $F_1$  plants are pure green. In out-of-season plants (req. 4 mos for maturity) white-stripping is seen: intracellular competition. Lectus modifies *ijap* pleistids in same sense as others.

Other genes do not behave in the same way. (1948)

Dojap.  $ij/ij$  are striped.

$ij \sigma \times Ij \rightarrow$  normal  $F_1$ .  $ij \text{♀} \times Ij \rightarrow$  white and striped  
as well as green  $F_1$ .

$F_2$   
3:1  $Ij:ij$

$ij$  plastids are smaller as well as chlorotic. "Both types of plastids were found in certain green cells".

Striped  $F_1$  ( $ij \text{♀} \times Ij \sigma$ )  $\times$  unstriped  $Ij$   $\rightarrow$  plants  $1/2$  should be  $Ij Ij$ .

Occasionally all progeny of a backcross ear (white sector) were white seedling, though  $1/2$  were  $Ij Ij$ . Concludes that mutant plastids retain their individuality.

(Persistence of striping in  $Ij Ij$  striped plants?)

Later, glossy-1 was used to mark  $Ij$  to prove homozygous condition. Normal sized plastids were paler in cells adjacent to white tissue. Also proportion of white offspring less than expected from proportion of maternal white tissue. Direct effect of  $ij$  on cytoplasm supposed: indecisive whether the permanent changes are in cytoplasm or plastid: plastid or plasmagene mutation? Segregation as case of Rennie; case is best evidence for cyto factor

- x Aschner, M. 1946 The symbiosis of Lucanosipoda aegyptia Mcq. Bull. Soc. Foued Entom., 30, 1-6.
- x Borghese, E. 1947 Il ciclo del batterio simbiote di Blattella germanica (L.). Mycopathologia 4. (18 pp)
- Borghese, E. 1948 Osservazioni sul batterio simbiote della Blattella germanica L. Monitore Zool. Italiano 56 (Suppl.), 252-253.
- x Bramstedt, F. 1948 Über die verdauungsphysiologie der aphiden. Zeit. f. Naturforschung, 3b, 14-24.
- ✓ Buchner, P. 1947 Symbiosis in Oliarius. Nature, 160, 264. *ref. Mahdihassan 1947*
- Buchner, P. 1949 Symbiose der Tiere mit pflanzlichen Mikroorganismen. Sammlung Göschen, 1128, 128 pp.
- Csáky, T. Z., and Tóth, L. 1948 Enzymatic breakdown of nitrogen compounds by the nitrogen fixing bacteria of insects. Experientia, 4: 4 pp.
- ✓ Hier, H. T. 1947 Intracellular bacteroids in the cockroach (Periplaneta americana Linn.). J. Bact., 53, 173-189.
- ✓ Glaser, R. W. 1946 The intracellular bacteria of the cockroach in relation to symbiosis. J. Parasitol., 32, 483-489.
- Goetsch, W. 1946 Darm-Symbionten als Eiweißquelle und Vitamin-spender. Österreichische Zool. Zeit., 1, 58-86.
- Goetsch, W. 1946 Vitamin "T" ein neuartiger Wirkstoff. Österreichische Zoologische Zeitschrift, 1, 49-57.
- x Gubler, H. W. 1947 Versuche zur Zuchtung intracellulärer Insekten-symbionten. Inaugural-Dissertation, Univ. Zürich. 34 pp.
- x Keller, H. 1950 Die Kulture der intrazellulären Symbionten von Periplaneta orientalis. Z. f. Naturforschung, 5b, 269-272.
- x Koch, A. 1948 Begriff und Bedeutung der Symbiose. Mikrokosmos, 38, 4 pp.
- ✓ Koch, A. 1948 Wege und Ziele der experimentellen Symbioserforschung. Naturwiss. Rundschau, No. 4, October, 1948, pp 166-171.
- x Koch, A. 1949 Die Bakteriensymbiose der Küchenschaben. Mikrokosmos, 38, 6 pp.
- ✓ Koch, A. 1950 Fünfzig Jahre Erforschung der Insektensymbiosen. Die Naturwissenschaften, 37, 313-317.
- MAHDIHASSAN, S. 1946 The micro-organisms in Melophagus ovinus. Current Sci., 15, 166-167.

- ↓ Mahdihassan, S. 1946 Bacterial origin of some insect pigments. *Nature*, 158, 58-59.
- Mahdihassan, S. 1947 *Cicadella viridis*, Its symbiotes and their function. *Curr. Sci.*, 16, 58-59.
- ✓ Mahdihassan, S. 1947 Specificity of bacterial symbiosis in Aphrophorinae. *Proc. Ind. Acad. Sci.* 25, 155-162.
- Mahdihassan, S. 1947 Bacterial symbiosis in a *Margarodes* sp. *Current Science*, 16, 379-380.
- + Mahdihassan, S. 1947 Bacterial symbiosis in *Aphis rumicis*. *Acta Entomologica Musei Nationalis Pragae*, 25, 123-126.
- Mahdihassan, S. 1948 Bacterial origin of muscle pigment in a Cicada. *Rev. de Entomologia*, 19, 585-586.
- \* Peklo, J. 1946 Symbiosis of azotobacter with insects. *Nature*, 158, 795-796.
- \* Rau, A. 1943 Symbiose und Symbiontenerwerb bei den mesbraciden (Homoptera-Cicadina) *Zeit. Morphol. Okol. Tiere*, 39, 369-522.
- Schanderl, H. 1949 Unsere derzeitigen Kenntnisse über die Hefen. *Biologia Generalis*, 19, 130-162.
- ↖ Schanderl, H., Lauf, G., Becker, H. 1949 Studien über die Mycetom- und Larmsymbionten der Aphiden. *Zeit. f. Naturforschung*, 4b, 50-53.
- \ Tóth, L. 1946 The biological fixation of atmospheric nitrogen. Monographs in Nat. Sci., No. V, Hungarian Museum of Nat. Sci. 116 pp.
- Tóth, L. 1950 Protein metabolism and nitrogen fixation by means of microorganisms living in symbiosis with insects. Eighth International Congress of Entomology, 303-306.

LATIN 18

ag



Test H213 for partial segregation, heterozygosity of Lac.

Couie, D.B.; Roberts, R.B.; Roberts J.Z. (1979). JCCP 34: 243-257. potassium metabolism in *Escherichia coli*. I. Permeability to sodium and potassium ions.

$\text{Na}^+$  reaches equilibrium rapidly between water space of cells and environment.  
 $\text{K}^+$  concentrated: 2-15 mg/ml K bound in cell; also diffusible K in equilibrium. "After initial equilibrium there is a further slow uptake of K in resting cells suspended in a medium with no energy source. This appears to be due to the residual metabolism of the cells."

When glucose is added, K is taken up at a minimum rate of  $1 \mu\text{g(K)}/\text{min}/\mu\text{l cells}$ . Bound K (low K medium for growth) is not readily lost. Free K is lost upon washing. In metabolism, cells exchange K rapidly (5%/min.) but membrane must be highly permeable.

$2.3 \pm 0.3$  atoms K taken up per mole glucose.

iodoacetate inhibited K-exchange but not P-loss. DNP prevented K turnover. Azide inhibited P uptake. Excess  $\text{PO}_4$  partially. Attempts to isolate K compounds failed. K was released by suspending cells  
a) in NaCl pH 9 2)  $\text{Et}_2\text{O}$ ; water 3) freezing + thawing; 4) ext. 50%  $\text{EtOH}$ .  
Implied that K-compounds are extremely unstable & destroyed when extracted. Uptake with D-1-P accelerated.

See Lebowitz & Kaperminty.

Potassium metabolism in *Escherichia coli* II Metabolism in the  
presence of carbohydrates & their metabolic derivatives JCCP 34: 259-291.  
Roberts, Robert 1.2., + co. ii

Kl behaved like K<sub>and</sub> could be used as a tracer.

K-uptake unaffected by UV or hyperinhibition.

## MATSUURA'S SPIRAL THEORY OF CROSSING OVER

- Beadle, G.W. 1938.  
Further studies of asynaptic maize. *Cytologia* 4:269-287.
- Belling, John 1933.  
Crossing over and gene rearrangement in flowering plants. *Genetics* 18:388-413.
- Huskins, C. Leonard and Stanley Smith 1934.  
Chromosome division and pairing in Fritillaria meleagris: The mechanism of meiosis. *J. Genet.* 28(3): 397-406.
- and Jane D. Spier 1934.  
The segregation of heteromorphous homologous chromosomes in pollen mother cells of Triticum vulgare. *Cytologia* 5:269-277.
- and G.B. Wilson 1938.  
Probable causes of the changes in direction of the major spiral in Trillium erectum L. *Ann. Bot. N.S.* 2:281-291.
- and H.B. Mewcombe 1941.  
An analysis of chiasma pairs showing chromatid interference in Trillium erectum L. *Genetics* 26:101-127.
- Morgan, L.V. 1933.  
A closed X chromosome in Drosophila melanogaster. *Genetics* 18:250-283.
- Matsuura, H. 1937.  
Chromosome studies on Trillium Kantschaticum. III. The mode of chromosome disjunction at the first meiotic metaphase of the P.M.C. *Cytologia* 8:142-177.
- 1940.  
----- XII. The mechanism of crossing over. *Cytologia* 10:390-405.
- 1938.  
----- VII. Additional evidence for the Neo-two-plane theory of univalent constitution. *Cytologia* 9:78-87.
- 1941.  
----- XIV. Primary and secondary chiasmata. *Cytologia* 11:380-387.
- 1944.  
----- and its allies. XVII. A study of chromosome interlocking in T. Tschonoskii Maxim. *Cytologia* 13:369-379.
- 1949.  
----- XXII. Critical evidence of the spiral theory of crossing over. *Chromosoma* 3:432-439.

## CHROMOSOME STRUCTURE (GENERAL)

- Belling, J. 1928. The ultimate chromomeres of *Lilium* and *Aloe* with regard to the number of genes. *Univ. Cal. Pub. Bot.* 14:307-318.
- Caspersson, T. and J. Schultz. 1940. Ribonucleic acids in both nucleus and cytoplasm, and the function of the nucleolus. *Proc. Nat. Acad. Sci.* 26:507-515.
- Darlington, C. D. 1939. Misdivision and the genetics of the centromere. *J. Genet.* 37:341-364.
- Darlington, C. D. and L. Lacour. 1941. The detection of inert genes. *Jour. Hered.* 32:115-121.
- Gates, R. Ruggles. 1942. Nucleoli and related nuclear structures. *Bot. Rev.* 8:337-409.
- Heitz, E. 1931. Die Ursache der gesetzmässigen Zahl, Lage, Form und Grosse pflanzlicher Nucleolen. *Planta* 12:775-844.
- Heitz, E. 1935. Chromosomenstruktur und Gene. *Zeit. ind. Abstam. Vererb.* 70:402-447.
- Huskins, C. I. 1941. Structural differentiation of the nucleus. in: *The Structure of Protoplasm.* (ed. W. Seifriz) 109-126.
- Lewis, I. M. 1941. The cytology of bacteria. *Bact. Rev.* 5:181-230.
- McClintock, B. 1934. The relation of a particular chromosomal element to the development of the nucleoli in *Zea mays*. *Zeit. Zellf. und Mikr. Anat.* 21:294-328.
- Morgan, D. T. J. 1943. The formation of chromocenters in the interkinetic nuclei of maize. *J. Hered.* 34:194-198.
- Nebel, B. P. 1939. Chromosome structure. *Bot. Rev.* 5:563-626.
- Rhoades, M. M. 1940. Studies of a telocentric chromosome in maize with reference to the stability of its centromere. *Genetics* 25:483-520.
- Robinow, C. F. 1944. Nuclear apparatus and cell structure of rod-shaped bacteria. in: *The Bacterial Cell* (R. J. Dubos), 355-377.
- Schultz, J. and T. Caspersson. 1940. The genetic control of nucleolar composition. *Proc. Nat. Acad. Sci.* 26:515-523.
- Straub, J. 1933. Chromosomenstruktur. *Naturwiss.* 31:97-108.
- White, M. J. D. Nucleus, chromosomes and genes. 1942. in: *Cytology and Cell Physiology* (ed. C. Bourne) 139-159.

Halpern, S.

on the nature of adaptive enzymes

Genetics (2): 363-367 1938

Stenfeld, L. and F. Saunders, J. Bact. 36: 53-56 (1938)

The fermentation of mucic acid by some intestinal bacteria.

+ : aerobacter, coli, para B, typhus, enteritidis

- : typhi, paratyphi, cholera-suis, dysenteriae.

---

Knopfmacher, H.P. + A.J. Salle, J. Gen. Physiol. 24: 377-397 (1941)

Studies on the lactase of E. coli.

Hessley + Benfante.

① China-Blue - Rosolic Acid Indicator medium.

Toluene supposedly inhibits oxidation but not hydrolysis. after Reaction.

No activity in autolysates.

Deere et al 1936. - lactose is not removed from broth by lac-

Measured lactase by increase in total reducing power caused by toluene or thymol-treated cells. Thymol study is 1 hour.

Substrate: .5% lactose in 1% acacia + .1M Phosphate at 7.0-7.2.

Samples dried by vacuum distillation. Dried cells (20-30 mg.), suspended in 25 cc 2% acacia, 10-20 mg thymol & incubated. After 1 hr, 25 cc 1% lac added. Dil. c .01%  $\text{CuSO}_4$  to stop enzyme action.

Activities: small activity noted in unadapted cells! .1-.2% hydr/mg cells.


This increased to about 4.5%

No specific statements on no-cell controls in lactose-acacia system.

Acacia might be hydrolyzed! 12 hour incubation period. No statement on contamination! [ 10mg thymol / 50 cc. ] Dried + Non Dried cells had similar activity.

Porter, R.R. (1948) *Acta Biochem.* 2(2):105-112. The unreactive amino groups of proteins.

Only <sup>19</sup> of the 32  $\epsilon\text{NH}_2$  (lysine) of  $\beta$ -lactoglobulin react with



(FDNB) unless deactivated. All can be acetylated.



W 327.

~~Mal S<sub>M</sub> + T L B<sub>1</sub> -~~ x

M<sub>1</sub> + M<sub>3</sub> - S<sub>M</sub> + T - L - B<sub>1</sub> -

x S<sub>M</sub> - M<sub>1</sub> - B - M - H.

S<sub>M</sub> - M<sub>1</sub> - M<sub>3</sub> + B M T L B<sub>1</sub> ...

S<sub>M</sub> + M<sub>1</sub> + M<sub>3</sub> -

S <sub>M</sub>	M <sub>1</sub>	M <sub>3</sub>	Glu	Mal
-	-	+	+	-
-	-	-	-	-
+	-	+	+	?
+	-	-	<del>-</del>	?
+	+	-	-	+
+	+	+	+	+
-	+	-	-	-
-	+	+	+	+

If suppressor affects M<sub>1</sub> -

S<sub>M</sub> + M<sub>1</sub> - M<sub>3</sub> + and S<sub>M</sub> + M<sub>1</sub> - M<sub>3</sub> -

have to be identified

from +++ and - (++) (wild types).  
Glu + Mal +

Need progeny tests of

- ① Measure " $K_m$ " of adaptation and compare  $\bar{c} K_m$  for the enzyme.
- ② Determine u.v. absorption spectrum of ONPG + substrate (i.e.  $\beta$ -galactosidase) for spectro-photometric evidence of complex formation. Do. enzyme + ONPG in presence of inhibitor - Mg $\cdot$ F $\cdot$ PO $_4$  (?)

If  $S_M \rightarrow Mal_3^-$  in  $S_M + M_1 - M_3 +$ .

$S_M \pm M_1$

Wild types

vs.  $S_M + M_1 - M_3 +$

$S_M + M_1 +$   
Cross segregants  $\bar{c}$

wild type and look for Mal-segregants.

If  $S_M \rightarrow Mal_3^-$  in  $S_M + M_1 - M_3 -$  [blue-Mal+], must be distinguished from  $S_M \pm M_1 + M_3 -$ . Take  $M_3 +$  papillae and cross  $\bar{c}$  wild type...

blue-Mal+ is index of  $S_M + M_3 -$ .

Cross W108-Mal+ - blue+ :  $S_M + Mal_3 + Mal_1 + \times S_M - Mal_3 + Mal_1 -$

and look for Mal segregation. If red, that + type.

- Beasley, J. O., 1940. The Origin of the American Tetraploid Gossypium Species. Am. Nat. 74, 285.
1942. Meiotic Chromosome Behaviour in Species Hybrids, Haploids, and Induced Polyploids of Gossypium. Genetics, 27, 25.
- Harland, S. C., 1932a. The Genetics of Gossypium. Bib. Genet. 9, 107.
- 1932b. The Genetics of Cotton, VI The Inheritance of Chlorophyll Deficiency in New World Cottons. J. Genet. 25, 271
- 1936a. The Genetical Conception of the Species. Biol. Rev. 11, 82.
- 1936b. Some Notes on Cotton in Colombia. Trop. Agr. T'dad, 13, 31.
1939. The Genetics of Cotton. Jonathan Cape.
- 1940a. Taxonomic Relationships in the Genus Gossypium. J. Wash. Acad. Sci. 30, 426.
- 1940b. New Polyploids in Cotton by the Use of Colchicine. Trop. Agr. T'dad. 17. 53.
- Hutchinson, J. B., 1932. The Genetics of Cotton, VII 'Crumpled': A New Dom. in Asiatic Cottons, Produced by Complementary Factors. J. Genet. 25, 281.
1938. The Distribution of Gossypium and the Evolution of the Commercial Cottons. 1st Cotton Conf. Ind. Cent. Cotton Cttee., Bombay
- 1939a. The Classification and Evolution of Cotton. Trop. Agr. T'dad. 16, 82.
- 1939b. The Relationships of Gossypium raimondii Ulb. Ibid. 16, 271.
- 1943a. A Note on Gossypium brevilanatum Hoch. Ibid. 20, 4.
- 1943b. The Cottons of Jamaica. Ibid. 20, 56.
- 1943c. Notes on the Native Cottons of Trinidad. Ibid. 20, 235.
- 1943d. The Effect of Environment on Hair Characters and Spinning Value in Sea Island Cotton. J. Text. Inst. 34, T61.
1944. The Cottons of Puerto Rico. Puerto Rico Agricultural Journal, 28, 35.
1946. On the Occurrence and Significance of Deleterious Genes in Cotton. J. Genet, 47, 279.
- (In press). The Genus Gossypoides Skovsted.
- Hutchinson, J. B. and Panse, V. G., 1936. The Introduction of Improved Strains of Crop Plants in Central India and Rajputana. Ind. J. Agric. Sci. 6, 397.
- Hutchinson, J. B. and Ghose, R. I. M., 1937a. Studies in Crop Ecology I. The Composition of the Cotton Crops of Central India and Rajputana. Ibid. 7, 1
- 1937b. The Classification of the Cottons of Asia and Africa. Ibid. 7, 233.
- Hutchinson, J. B. and Govande, G. K., 1938. Cotton Botany and the Spinning Value and Hair Properties of Cotton Lint. Ibid. 8, 21.
- Hutchinson, J. B. and Bhola Nath. 1938. A Note on the Occurrence of Chlorophyll Deficiency in Gossypium arboreum. Ibid. 8, 425
- Hutchinson, J. B., Panse, V. G., Apte, N.S. and Pugh, B.M., 1938. Studies in Plant Breeding Technique. III. Crop Analysis and Varietal Improvement in Malvi Fower (Andropogon sorghum). Ibid. 8, 131.
- Hutchinson, J. B. and Silow, R. A. 1939. Gene Symbols for Use in Cotton Genetics. J. Hered. 30, 461.

- Silow, R. A., 1941. The Comparative Genetics of Gossypium anomalum and the Cultivated Asiatic Cottons. J. Genet. 42, 259.
- 1944a. The Inheritance of Lint Colour in the Asiatic Cottons, Ibid. 46, 78.
- 1944b. The Genetics of Species Development in the Old World Cottons. Ibid. 46, 62.
1946. Evidence of Chromosome Homology and Gene Homology in the Amphidiploid New World Cottons. Ibid. 47, 213.
- Silow, R. A. and Yu, C. P., 1941. Anthocyanin Pattern in Asiatic Cottons. J. Genet. 43, 250.
- Skovsted, A. 1933. Cytological Studies in Cotton. I. The Mitosis and the Meiosis in Diploid and Triploid Asiatic Cotton. Ann. Bot. 47, 227.
1934. Cytological Studies in Cotton. II. Two Interspecific Hybrids between Asiatic and New World Cottons. J. Genet. 28, 407.
- 1935a. Cytological Studies in Cotton. III. A Hybrid between Gossypium davidsonii Kell. and G. Sturtii F. Muell. Ibid. 30, 397.
- 1935b. Some New Interspecific Hybrids in the Genus Gossypium L. Ibid. 30, 447.
- 1935c. Chromosome Numbers in the Malvaceae I. Ibid. 31, 263.
1937. Cytological Studies in Cotton. IV. Chromosome Conjugation in Interspecific Hybrids. Ibid. 34, 97.
- Stephens, S. G., 1942. Colchicine-Produced Polyploids in Gossypium. I. An Autotetraploid Asiatic Cotton and Certain of its Hybrids with Wild Diploid Species. J. Genet. 44, 272.
- 1944a. The Genetic Organization of Leaf Shape Development in Gossypium. Ibid. 46, 28.
- 1944b. Cotton Growing in the West Indies during the Eighteenth and Nineteenth Centuries. Trop. Agr. Trinidad. 21, 23.
- 1944c. Phenogenetic Evidence for the Amphidiploid Origin of New World Cottons. Nature 153, 53.
- 1944d. Note on the Meiosis of a Triple Species Hybrid in Gossypium. Ibid. 153, 82.
- 1944e. Colchicine-Produced Polyploids in Gossypium. II. Old World Triploid Hybrids. J. Genet. 46, 303.
- 1944f. The Modifier Concept. A Developmental Analysis of Leaf Shape 'Modification' in New World Cottons. Ibid. 46, 331.
- 1944g. Canalization of Gene Action in the Gossypium Leaf Shape System and its Bearing on Certain Evolutionary Systems. Ibid. 46, 345.
1946. The Genetics of 'Gorky'. I. The New World Alleles and their Possible Role as an Interspecific Isolating Mechanism. Ibid. 47, 150.

Publicazioni della Stazione Zoologica di Napoli vol 22 suppl. 1950, June

Relazioni tenute al convegno su GLI AGENTI MUTAGENI 27-31 maggio 1949

1. ~~W.~~ Auerbach, Ch. (Edin) Possible differences between the effects of chemical and physical mutagens.  
1-21
2. C.D. Darlington (London): Physical and chemical breakage of chromosomes  
22-31
3. E. Haeflorn (Zurich): Erfahrungen mit Phenol-Behandlung von Drosophila-Gonaden  
32-49 in vitro  
\*\*
5. H.E.-Taylor (Paris): Biological significance of the transforming principles of Pneumococcus  
~~50-54~~ 65-77  
~~65-77~~
6. R. Latarjet (Paris): Induction d'une mutation spécifique chez une bactérie par des cancérogènes hydrosolubles. *W.P. Bau Hoi + CA Elias*  
65-78-93
4. \*\*B. Ephrussi (Paris): Induction par l'acridavine d'une mutation spécifique chez la levure  
50-64
7. N. Visconti (Milan): Le mécanisme d'action létale de la moutarde azotée sur Bacterium coli  
di Madame 94-113
8. M. Vogt (Neustadt im Schwarzwald): Urethane-induced mutations in Drosophila  
1154-124
9. E. Battaglia (Pisa): Nuove sostanze inducenti frammentazione cromosomica  
125-157
10. F.D'Amato (Pisa): The chromosome breaking activity of chemicals as studied by the Allium Cepa test  
158-170
11. A. Buzzati-Traverso (Pavia): Perspectives of research on mutagens (A discussion with the participants in the Symposium)  
171-186

Bibliografia generale 187

Handwritten notes:

Handwritten notes:

Handwritten notes:

Porter and Taylor

J. Neurophys. 8


(1945)

Interruption of tubanes + pain.

post tibial nerve stim., neuromuscle tib. ant. response spinal cat.

Stim. n. at each respiration. (artificial). Pain produced by acid in other

nerve fields. Response increased. No response to conc. reflex stimuli.



Weinstein, EA + M B Bender, *Devel. Neuro Psychol.* 50:34-42 (1973)

Integrated facial patterns elicited by stimulation of the brain stem.



Bender, MB + EA Weinstein, Functional Representations in the  
oculomotor and trochlear nuclei. ~~49~~

Arch. Neurol. Psychol. 49: 98-106 (1943)

~~Am. Nat.~~ Timofeïf - Resovsky, *Am. Nat.* 63: 118-24 (1929) The effect  
of  $\beta$ -rays in producing somatic gene variations of a definite locus in different  
directions in D. m.

Jeq, et al. P.R.S. B123:1

Juria, S. CRAS 209:604 (1939)

Remere,

Gen. 22:469-

Lea,

Action of Radiation on living cells.

1946.

Bauer, K. H. Arch. Klin. Chir. 152: 278-83 (1928). Fortschritt d.  
Vererbungslehre u. Geschlechtsfrage.

Tumorsomatei mutationi.

Trypanosome Refs.

PSEBM 67: 77, '48

J. Pharm 82: 137-151 '44

Arch. Protist. 68: 187, 1929

J.I.D. 81: 139, '47

Z. Bakt. I, 53: 303, '10

Trans. Roy, Soc. Trop Med Hyg. 22: 85, '28



## CYTOLOGICAL TECHNIQUE

- Belling, J. 1930. The use of the microscope. McGraw-Hill Co., New York.
- Coleman, L. C. 1938. Preparation of leucobasic Fuchsin for use in the Feulgen reaction. Stain Tech. 13: 123-124.
- Conn, H. J. 1940. Biological stains, 4th ed. Biotech. Pub., Geneva, New York.
- Darlington, C. D. and L. F. LaCour. 1942. The handling of chromosomes. Unwin Bros., London.
- Hillary, B. B. 1939. Use of the Feulgen reaction in cytology.  
I. Effect of fixatives on the reaction. Bot. Gaz. 101: 276-300.
- \_\_\_\_\_ . 1940. Use of the Feulgen reaction in cytology.  
II. New techniques and special applications. Bot. Gaz. 102: 225-235.
- Johansen, D. A. 1940. Plant microtechnique. McGraw-Hill Co., New York.
- LaCour, L. F. 1937. Improvements in plant cytological technique. Bot. Rev. 5: 251-258.
- Lee, Bolles. 1937. The microtome's vade-mecum (ed.) J. B. Catenby and T. S. Painter 10th ed. Churchill, London.
- Smith, Luther. 1947. The acetocarmine smear technic. Stain Tech. 22: 17-31.

Huggins, C. & Smith, D.R. (1947) *Acromogenic substrates. III. p-nitrophenyl sulfate as a substrate for the activity of phenol sulfatase activity.* JBC 170: 391-398.

$\text{DN}(\text{CH}_3)_2$  (DMA) 47 ml +  $\text{CS}_2$  50 ml are mixed in a 500 ml suction flask ~~and~~ in ice bath in hood. Add 9.1 ml 1%  $\text{CO}_2\text{SO}_3\text{H}$  dropwise. Add 13.9 g p-nitrophenol rapidly. Stir one hour & let stand overnight.

Add 100 ml .4 M KOH  $\rightarrow$  yellow crystals. Stir thoroughly. Evaporate  $\text{CS}_2$  at  $80^\circ$  *in vacuo*. Recrystallize crude product 3-4 x in 80% EtOH. [Method from J. Ch. S. 1:684 (1926)].

Formed activity measurable in 10 hours.  
opt. at pH 6.12 in acetate 1/2.  
 $K_m = 7 \times 10^{-5} \text{ M}$ . from talca diastase.

Dept Surgery, UChicago.

BA - for serum adapted enzymes.  
Indices

Enzymes

Lactase + Lactose

Adaptation

serum (~~not~~)

18

8838 "protectini" R. Abdalalden. Munch. Med. Wschr. 88:726

5415 Localization of lactase in yeast cell See

Hjörbäck + Vasselund. Z. physiol. ch.

277: 171-180 (1943). T. cremoris

fermented but did not hydrolyse !?

Über die Faktoren für  
und die Lokalisation  
der Enzyme in den Hefezellen

17 13310 arguari. See JBC 147:99-108 (1943).

481 { Conyell + Christman JBC 150: 143-154 (1943).

16447 } Utilization of lactose by the fermenting rat.

16 632  
4676

15



than adequate for rate of fermentation, but faster fermentation of lactose than the hexoses not explained.

"In cell-free extracts, toluene treated, or acetone dried cells, glucose fermentation becomes the fastest so that either the enzymes necessary for the direct fermentation are more labile, or the different rates are due to some structural factor such as a differential permeability to lactose."

No No Fr

25g MM<sub>1</sub>'s in 150ml H<sub>2</sub>O

add 2 ml conc H<sub>2</sub>SO<sub>4</sub>

milk, 5g MM<sub>1</sub>'s in 7.5 ml H<sub>2</sub>O

2 ml H<sub>2</sub>O in 150 ml H<sub>2</sub>O

24-48h 37°

fermentation

50g glucose 6h

30 ml H<sub>2</sub>O (3.0g) = 4.78 ml H<sub>2</sub>O by density

etc in soluble bottle

add 2/16.5 ml H<sub>2</sub>O in 150 ml H<sub>2</sub>O +  
4.78 ml H<sub>2</sub>O in 150 ml H<sub>2</sub>O

A bibliography of Neurospora

A. Biochemical mutants.

1. Beadle, GW & JL Tatum, PNAS 27:499-508. Genetic control of biochemical reactions in Neurospora (( (pab), (pyr), (tz1) 1941
2. Tatum & Beadle PNAS 28:234-43 1942 Do.: An aminobenzoic acid class mutant. (1533, 5359)
3. Tatum & Beadle, Growth S4: 27-30 1942. The relation of genetics to growth factors and hormones.
4. Bonner, D., Tatum & Beadle, Arch Bioch 3:71-81 1943. : A mutant strain requiring isoleucine and valine. 15117.
5. Seb AM & NH Horowitz, JBC 154:129-39 1944. The Ornithine cycle in Neurospora and its genetic control
6. Horowitz & Beadle JBC 150:325-33 1943 A microbiological method for the determination of choline by use of a mutant of Neurospora. 34466
7. Loring HS & JG Pierce, JBC 153:51-59 1944. Pyrimidine nucleosides and nucleotides as growth factors for mutant strains of Neurospora.
8. Horowitz, JBC 154:141-9 1944. The d-aminic acid oxidase of Neurospora.
9. Ryan and E Brand, JBC 154:161-75 1944. A method for the determination of Leucine in protein hydrolysates and in feedstuffs by the use of a Neurospora mutant.
10. Regnery, DC JBC 154:151-60 1944. A Leucineless mutant strain of N.c.
11. Tatum Bonner & Beadle, Arch Bioch 3:477-8 1943 Anthranilic acid and the biosynthesis of Indole and Tryptophan by N.
12. Tatum & Bonner, JBC 151:349 1943 Synthesis of Tryptophan from indole and serine by Neurospora.
13. Bonner & R Dorland, AB 2:451-62 1943. Experiments on the application of N. sitophila to the assay of Pyridoxin in tomato plants.
14. Stokes, JL JW Foster & CR Woodward, jr. AB 2:335-45 1943. Synthesis of pyridoxin by a pyridoxinless X-ray mutant of N. sitophila
15. Stokes, A Larsen, CR Woodward, & Foster, JBC 150:17-24 1943 A neurospora assay for pyrioxine
16. Wyss, O, VG Lilly & LH Leonian, S 29:18-9 1944 The effect of pH on the availability of p-aminobenzoic acid to N. crassa
17. Ryan Beadle & Tatum AJB 30:784-88 1943. The tube method of measuring the growth of Neurospora.
18. Ryan, Tatum & AC Giese, JOCP 23:83-94 1944. The four-carbon respiratory system and the growth of the mold Neurospora.
19. Doerman, AB: A lysineless strain of Neurospora and its inhibition by arginine.

B. Genetic and cytological.

1. Sengar, CL & ED Dodge, J Agr Res 34:1019-22 1947 Life histories and heterothallicism of the red bread mold fungi of the *Aspergillus nitropila* group.
2. Dodge, JAR 25:289-305 1937 Nuclear phenomena associated with heterothallicism and homothallicism in the ascomycete *Neurospora*.
3. Dodge, JAR 25:1-14 1936. Production of fertile hybrids in the ascomycete *Neurospora*.
4. Dodge, B Terry 67:167-75 1940. Unequal division of spores and crossing over in the fungi.
5. Dodge B Terry 68:75-91 1942 Heterocaryotic vigor in *Neurospora*.
6. Ryan B Terry 71:698-11 1943. Crossing over and unequal division of spores in fungi.
7. Dodge Mycologia 34:302-7 1942. Conjugated nuclear division in the fungi.
8. Beal, & VL Coe, Gen 23:281-303 1944 Heterocaryosis in *N.C.*
- 9.

cross #	camping from	3A	3B	3C	51
1335		++	CL	CL	++
"	( 211 )	-	CL	++	-
"	( 145 )	-	+	-	(+)
"	( 1339 )	-	(CL)	CL	-
211		-	CL	++	-
145		-	+	-	(CL)
1339		-	(+)	CL	-
1394		EL	CL	CL	++
"	( 145 )	-	+	-	(+)
"	( 211 )	-	CL	++	-
284		CL	++	-	+
"	( 145 )	-	+	-	+
145		-	+	-	CL
"	( 1394 )	-	+	-	+
1371		CL	CL	CL	+

similar results in other groups.



In addition to references cited in my American Journal of Veterinary Research, Vol. VIII, 1947 paper, I found the following possibly useful references in my file.

Martin, Gustav J. Acridine Antiseptics. Medicine 23: No. 1, 1944.

Glassman, H. H. Surface active agents and their application in bacteriology, Bact. Rev. 12: No. 2, 1948.

x Brodie, J. and Shepherd, A. Further observations on the differential inhibition of coliform bacilli and rough variants of intestinal pathogens. J. Gen. Microbiol. 3: 74-79, 1949.

x Brodie, J. Observations on the differential inhibition of coliform bacilli and rough variants of intestinal pathogens. J. Gen. Microbiol. 2: 1-7, 1948.

Caselitz, F. F. Ein Beitrag zum S-R-Formenwechsel der Salmonella Bakterien. Zeitschr. f. Immunitat, 109: 149-158. 1952.

Dolkner, W.A.L., van der Meer, C. and Scholten, R. Th. The electrophoretic behaviour in relation to the changes of the antigenic structure observed in the smooth-rough variation. Antonie van Leeuwenhoek, Jour. Microbiol. and Serol. 8: 59-70. 1942.

Zagrodski, E. Nouvelle methode de differenciation des souches R et S du bacille de Bang au moyen de l'agglutination non specifique. Iadonesci veter 18: 23, 1939. (cited in Bull. Inst. Past. 37: 1259.

Bennet, F.M. (1927) B.J.E.P. 8:121-129. The relationships between  
 heat-stable agglutinogens and sensitivity to bacteriophage in the  
 Salmonella group.

I	enteritidis	(I), IX, XII
(II)	typhi	IX, XII, VI
(III)	pulchrum	IX, XII
(IV)	deby	I, IV, XII
(V)	para B	IV, XII
(VI)	stanley	IV, V, XII
(VII)	reading	IV, XII
(VIII)	typhimurium	(I) IV (V) XII

1, + 12

Phages ~~1, 11, 13 + 18~~ attack typical smooth only of (I), (II) + (III).

Phages 11, 13 + 18 regularly attacked only typical roughs of (II) + (III).

Special "r" strains were attacked only by phage 8, which attacked  
 all but a few variants, of (I).

~~18~~ 1R/13 was "smooth", and lysed by 12; it carried 1  
 lysogenically. 1S/12 were "rough" and sensitive to 11, 13 + 18.

¶ 8 attached S forms of ~~①, ②~~, ①, ②, ③, <sup>⑤</sup>, but not ⑥, ④, ~~⑦~~  
para A, super-tifer., R forms of 1, 2, 3, ⑤, ⑥, ⑦, para A.

¶ 1 + ¶ 12 attached typical S forms only of ①, ② + ③ + ④. no R.

¶ 11 " most R forms regardless of type. do. 13 + 18.

¶ 20 " R + S forms of ①, ②, ③ para A

¶ 21 R + S ①, ②, ③ + decy S.

Conclusion: Sphages probably associated with the factor  
now recognized as IX. R are cosmopolitan, as are the  
serological behaviors.

Burnet 1929b. Further obs. - Kymant.

φ 8 eq. active on R + S of gallinacum. No serological difference

detectable between S + S/8, or R + R/8. R/8 did not absorb φ 8.

R + S sera showed little cross-reaction. R was obtained with φ 1.

1929a. Classified phages:

A	B	C	D
1, 12, 33	8, 18, 28, 31, 34, 38	20, 25, 32, 35	11, 13

Testing on variants obtained S phase.

A are S φ.

18, 35, 11 + 13 are R only.

8, 34 are indifferent to R/S. Other φ are more active on R than S.

32 + 38 : 32 gallinacum R or S, 38 R only.

gallinacum S/12 were variably "rough" if really resistant, but frequently reacted with both R + S sera. Various colonial types noted.

The mucoids which were found were hypogenic → smooth sensitivities. sensitive to R φ.

All /H were rough. S/8 → smooth; correlated with resistance to φ 8.

Smooth mutants could be recovered from rough strains. Reversibility may be associated with a slight O-egg lectinogen content. (titre ca 80)

R-S-R → ... could take place.

Gumet 1930) Bacteriophage activity and the antigenic structure of  
 salina. J.P.B. 33:647-664.

Table 4. *S. gallinarum*:

A B C D D'

Discussion of mixture patterns in terms of "change" planes

For some phages, ~~the~~ susceptibility & specificity are uniform in R + S phases.

It is possible that different directions of modification of the O-substance

are responsible. In Staph, sensitivity is more closely correlated c

serology:

Phages

<u>Antigen</u>	1	2	3	4	
ABC	S	S	S	S	SF
BC	R	S	S	S	SF/1
<del>ABC</del> AC	S	R	R	S	SF/2
ABC.	S	S	R	S	SF/3

Table 4. *S. gallinarum*.

Cells.	A	B	C	D	D'	Reaction
	12	39	40 8	18 38 25	35	13
398S	S	S	S S ±	R S S	R	S
398S/8	S	S	S R R R	S S	R	S
398S/25	R	R	<del>R</del> <del>R</del> <del>R</del> ±	R R	R	S
398S/39	S	R	R R ±	R S S	R	S
398R = S/12	R	R	S		→	R
398R/8	R	R	R R R R	S S	S	R
398R/35	R	R	S S S	R R	R	R
398R/13	R	R	S S S	R R	R	R

Note: R are R to A, B., S to C + D / 8 is C<sub>R</sub> D<sub>S</sub> / 13 or 35 is C<sub>S</sub> D<sub>R</sub>

Burnet, + McKee, (1930) Bacteriophage reactions of flexner dysentery strains. SBB 33:637-646.

4 groups of phages. ~~A~~ - smooth only.  
others - most roughs, some smooths.  
antigenic types characteristically different.

Groups C + D are homologous with the Salmonella phages active on rough gallinarum.




Burnet + McKie JPB. 36: 299-306; 307-318 (1933).

The classification of dysentery-coli bacteriophages. <sup>I + II.</sup> resistance patterns  
& serology. Some phages may act equally on dysentery R, coli +  
Salmonella.

pH (ca<sup>2+</sup>)

Oxalate		4.76
Barbiturate		3.98
Benzoate		4.20
Citrate		2.06 4.11 4.11
Formate		3.75
Lactate		3.85
Malate		3.40 5.05
Nitrate		3.4
Oxalate		1.19 4.21
Phosphonate		2.12 7.21 12.32
Phthalate		2.89 5.41
Succinate		4.19 5.51
Sulfurous		1.76 7.20
Tartrate		3.02 4.54



Sulfurous.

Oxalate

Absorption of p20 by W578 and W811.

532

4/20/49

Assay on E. coli B to avoid confusion with  $\lambda$  action

# Temperature sensitive resume

W:	31	1	108
	35	20	110
	40	3	124
	42	56	125
	43	58	138
	44	60	137
	45	71	178
	47	78	179
	48	88	200
	65	87	242
	67	83	259
	72	42	305
	74		
	76		

tested for  
P.S. by E.L.

W305 maybe faster at 37 than at 40.  
W110 - at 31 ++ at 40. W42 maybe  
similar.

Lactulose

ca 1:12 of p. 467-468 NBS "Sugar".

100g lactose in <sup>ca 75g</sup> 500ml H<sub>2</sub>O add Ca(OH)<sub>2</sub> at 55° kept several days.  
Concentrate in vacuo to wt of 125g. Dilute residue with 125ml MeOH  
and evaporate to dryness several times. Remove small amount, ca 75g by  
filtration & wash with 400 ml MeOH. Concentrate filtrate to a syrup.  
Dilute to 500 ml H<sub>2</sub>O + 100g to remove color. Dilute to 200ml and  
filtrate & wash.

5 ml sample. in 200 ml Erlenmeyer. Add 5 ml .1N iodine from burette  
Add 7.5 ml .1N NaOH ~~to remove~~ dispense. Repeat for 6 times. (3ml I<sub>2</sub>)  
Acidify with 10 ml N HCl + back titrate with N/10 Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> standard.  
For 2.5 g, Br = 1/20 iodine titration of 5 ml sample. Add 26 ml 20% KI  
equivalent. Add 15g CaCl<sub>2</sub> liquor. Add benzoin dispense &  
mechanical stirring. Remove residue 2 mg, 20g 16%  
per equivalent Br. H<sub>2</sub>O to filtrate to remove 2 mg. Separate  
filtrate to 125 ml to remove H<sub>2</sub>O.

(b) Selected Bibliography

BRINK

1. The physiology of pollen. Amer. J. Bot. 11: 213-228, 283-294, 351-364, 417-436. 1924.
2. Segregation for the waxy character in maize pollen and differential development of the male gametophyte. Amer. J. Bot. 11: 465-469. 1924. (with J. H. MacGillivray).
3. Mendelian ratios and the gametophyte generation in angiosperms. Genetics 10: 359-394. 1925.
4. Dynamics of the waxy gene in maize. I. The carbohydrate reserves in endosperm and pollen. Genetics 11: 163-199. 1926. (with F. A. Abegg).
5. Differential action of the sugary gene in maize on two alternative classes of male gametophytes. Genetics 12: 348-378. 1927. (with C. R. Burdhan).
6. The occurrence of semi-sterility in maize. J. Heredity 18: 266-270. 1927.
7. Dynamics of the waxy gene in maize. II. The nature of waxy starch. Biochem. J. 22: 1349-1361. 1928.
8. Studies on the physiology of a gene. Quart. Rev. Biol. 4: 520-543. 1929.
9. The association of semisterile-1 in maize with two linkage groups. Genetics 16: 595-628. 1931. (with D. C. Cooper).
10. A structural change in the chromosomes of maize leading to chain formation. Amer. Nat. 66: 310-322. 1932. (with D. C. Cooper).

11. A strain of maize homozygous for segmental interchanges involving both ends of the P-Br chromosome. Proc. Nat. Acad. Sci. 18: 441-447. 1932. (with D. C. Cooper).
12. The coumarin content of Melilotus dentata. Science 86: 41-42. 1937. (with W. L. Roberts).
13. Chromosome homology in races of maize from different geographical regions. Amer. Nat. 71: 582-587. 1937. (with D. C. Cooper).
14. Somatoplastic sterility in Medicago sativa. Science 90: 545-546. 1939. (with D. C. Cooper).
15. Translocations in maize involving chromosome 3. Genetics 25: 299-309. 1940. (with E. G. Anderson).
16. Double fertilization and development of the seed in angiosperms. Bot. Gaz. 102: 1-25. 1940. (with D. C. Cooper).
17. Incomplete seed failure as a result of somatoplastic sterility. Genetics 26: 487-505. 1941. (with D. C. Cooper).
18. A hybrid between Hordeum jubatum and Secale cereale reared from an artificially cultivated embryo. J. Heredity 35: 66-75. 1944. (with D. C. Cooper and L. E. Ausherman).
19. The antipodals in relation to abnormal endosperm behavior in Hordeum jubatum x Secale cereale hybrid seeds. Genetics 29: 391-406. 1943. (with D. C. Cooper).
20. Seed collapse following matings between diploid and tetraploid races of Lycopersicon pimpinellifolium. Genetics 30: 376-401. 1945. (with D. C. Cooper).
21. The endosperm in seed development. Bot. Rev. 13: 423-477, 479-541. 1947. (with D. C. Cooper).
22. Growth in vitro of immature Hordeum embryos. Science 106: 547-549. 1947. (with Nancy Kent).

23. The endosperm-embryo relationship in an autogenous species, Trigonotis officinale. Bot. Gaz. 111: 139-153. 1949. (with D. C. Cooper).
24. The mutagenic action of mustard gas on Zea mays. J. Heredity 41: 232-238. 1950. (with P. B. Gibson and M. A. Stahmann).
25. The stimulative effect of Hordeum endosperms on the growth of immature plant embryos in vitro. Amer. J. Bot. 38: 253-256. 1951. (with Nancy Kent Ziebur).
26. The relation between light variegated and medium variegated pericarp in maize. Genetics 37: 519-544. 1952. (with R. A. Milan).
27. Very light variegated pericarp in maize. Genetics 39: 724-740. 1954.
28. The relation between Modulator and Activator in maize. Proc. Nat. Acad.



Science 76:916  
Arthur S. Thomas Jr. 3:193/104.

Hawaii - Morph. Inc. Rate of Growth of Plants;

24.

Dobell, C. QJMS 52: 121-138 (1908).

(I).

QJMS 53: 55-76

II

QJMS. 58: 375-506

---

Pijper, S. Nfi Med Rec 23: 222 1925 *Agaricacephalus*

Stewart F.H. Segregation + Autogamy in bacteria  
Allard + Son, London.

Badian, J. *Autothelminthol.* 7: 409-418 (1950)

Sp: *autogary*

1  
Allan, Appleby + Wolf *J. Natl. Paras. II* 100: 3-16. 1939.

$$\sim p \cdot \sim q$$

$$p \vee q = \sim(\sim p \cdot \sim q)$$

$$\sim(p \vee q) = \sim p \cdot \sim q$$

$$\sim(p \cdot q) = \sim p \vee \sim q$$

Föhlis + Smith J Agr R 23:401-432 (1923)

*Conjugatum isogonatum*.

Holman - in Jordan + Falls New Knowledge.

III Int Congress Microbiol N. Y. 1939-40.

V Int Bot Congress 193 Sect B

Life Cycles -

Föhnio

Mem. Nat Acad Sci. 16, 1921



Hollande + Hollande, Besch. Zool exp. ges. 72:445-1184.

Dennis, L. J. *Bent* 48: 125.