

CONNECTICUT VALLEY BRANCH
SOCIETY OF AMERICAN BACTERIOLOGISTS
ANNUAL MEETING

Saturday - 2,00 P.M. - December 7, 1946

Auditorium of Brady Memorial Hall, Yale
University, New Haven, Connecticut

2,00 P.M. Business Meeting

Dr. Paul R. Burkholder, President, presiding.

2,30 P.M. Scientific Program

GENETIC EVIDENCE FOR SEX IN BACTERIA

JOSHUA LEDERBERG, Fellow of Childs Fund for Medical
Research and E. L. TATUM, Professor of Microbiology,
Osborn Botanical Laboratory, Yale University,
New Haven, Connecticut,

A QUANTITATIVE METHOD FOR THE DETERMINATION OF THE
FUNGISTATIC ACTIVITY OF ANTISEPTIC POWDERS

MARION B. SHERWOOD, Research Bacteriologist,
Wellcome Research Laboratory, Tuckahoe, N.Y.

THERAPY OF TRYPANOSOMIASIS

BENJAMIN ARNOLD RUBIN, Dept. of Bacteriology,
Yale University, New Haven, Conn.

VARIATION IN TISSUE SPECIFICITY OF THE ROUS CHICKEN
SARCOMA VIRUS FOLLOWING GROWTH OF THE TUMOR IN THE
MAMMALIAN EYE.

EDWARD W. SHRIGLEY, Assistant Professor of Bacteriology,
Department of Bacteriology, Yale University, School of
Medicine, New Haven, Conn.

THE ISOLATION OF PARACOCCIDIODES BRASILIENSIS FROM A
CASE OF SOUTH AMERICAN BLASTOMYCOSIS

ROSALIE FERGUSON (by invitation) and MARGARET F.
UPTON, Ph.D., Bacteriologist, St. Lukes Hospital,
New York 25, N.Y.

Connecticut Valley Branch, Society of American Bacteriologists

Yale University School of Medicine, December 7, 1946.

Genetic evidence for sex in bacteria. Joshua Lederberg, Jane Coffin Childs
Fellow in Medical Research, and E+L+Tatum, Professor of Microbiology, Yale U.

In the short time allotted, it will be impossible to discuss in any detail
all the experiments that have led us to the conclusion that there is a sexual
phase in the life history of a strain of *Escherichia coli* with which we have
been working. It seems best to present a single experiment in detail, which
most forcefully illustrates the kind of evidence that has been accumulated.
The general thesis of these experiments ~~was~~ is the same as that which was pro-
posed by Sherman and Wing 10 years ago, that if there is a sexual phase in
bacteria, it should be possible to show that when mixed cultures containing
organisms differing in various characters are studied, new types will be found
which are characterized by recombinations of those characters. The strains used
by these authors were too variable for them to draw any conclusions as to the
occurrence of sexual fusions, and other authors have had similar difficulties.

Several types of characters have been used in our investigations.
~~XXXXXXXXXXXX~~ They have all been obtained as mutations in a single
strain of *E. coli* (strain K-12). The spontaneous variability of each of these
characters has been studied intensively and it can be anticipated categorically
that none of the phenomena which I am about to describe can be accounted for
on this basis.

Most of the characters used are nutritional requirements, i.e. the need
for a specific growth factor, including such compounds as biotin, thiamine,
threonine, leucine, and other vitamins and amino acids. They are sharply de-
terminable, since optimal growth can be obtained in the presence of the
specific requirement, whereas in its absence growth does not take place.
The requirements ^{were obtained} ~~are found to originate~~ by mutation, one at each mutational
step. By ultraviolet irradiation of a strain which is already a biochemical
mut. . . multiple mutants can be obtained, and these have been very useful

Table.

in these studies. Other characters which have been used are resistance to a specific bacteriophage, and the inability to ferment lactose (as indicated on Eosin-Methylene Blue-Lactose plates such as these.)

Character recombination in *E. coli* can be detected by the appearance of wild types in mixed cultures of biochemical mutants. Such wild types have never been found in the individual cultures. ~~The wild types~~ wild types can be detected readily since they form colonies in minimal agar medium, whereas the original mutants are unable to proliferate in the absence of their specific requirements. Thus although only about 1 cell in 10 million in these mixed cultures is a wild type, they can be readily detected by plating about 500 million cells after careful washing, into a minimal agar plate.

If these wild types are the result of recombination of characters, ~~such these~~ colonies should also show recombination of other character differences present in the 'parental strains'. This has been found to be the case.

~~XXXX~~ By plating Y-53, which is $B_1^+ T^- L^- B_2^- Lac^- T_1^+$ and Y-10 into agar containing thiamin, but lacking biotin, methionine, threonine or leucine, the colonies which appear should represent those cells which have recombined with respect to B.M, T and L, as indicated on the board. One should also expect to find that the other 3 characters should recombine also, namely B_1^- ; Lac^- and T_1^+ . There are 8 possible recombinations of these characters, as indicated, and the ones which are part of the parental complexes are encircled. All 8 have been recovered, as can be demonstrated by this plate, and these tubes. The figures on the board represent the relative frequencies with which they have been obtained. The disparity in the numbers is of course, of great interest; some evidence has been obtained that this disparity may be explicable on the basis of the same kind of factorial linkage that characterize ^{higher} organisms.

Evidence has been obtained that in mixed cultures containing more than 2 types, recombination takes place between only 2 types of cell at a time, strengthening the hypothesis that these data are ^{likely} to be explained by a cell fusion and the orderly segregation of genes.

SEGREGATION OF BACTERIAL CHARACTERS: B₁ T₁ and Lac

.. .. .

-	-	+	+	+	R	+		
B	M	T	L	B ₁	T ₁	Lac	(Y-40)	
<hr/>								
B	M	T	L	B ₁	T ₁	Lac	(Y-53)	
+	+	-	-	-	S	-		

Progeny:

<u>B</u>	<u>M</u>	<u>T</u>	<u>L</u>	B ₁	T ₁	Lac		
					+	R	-	100	100
					+	R	+	55	55
					+	S	-	50	50
					+	S	+	4	4
100.					-	R	-	17	850
					-	R	+	9	450
					-	S	-	13	650
40.					-	S	+	1	50

B₁ - and B₁ + are from separate series. Actually, the B₁- types as a whole are about 10X as frequent as B₁+. With this correction, the last column gives the absolute proportions.

Tentative Linkage Map

