

TABLE 1  
 THE 20 MOST FREQUENTLY REPORTED SALMONELLA SEROTYPES  
 FROM HUMAN SOURCES REPORTED TO CDC IN 1999 AND FROM  
 CLINICAL AND NON-CLINICAL NONHUMAN SOURCES REPORTED TO CDC AND NVSL IN 1999

HUMAN 1999				CLINICAL NONHUMAN 1999				NON-CLINICAL NONHUMAN 1999			
RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT
1	TYPHIMURIUM *	8051	24.6	1	TYPHIMURIUM *	1865	34.9	1	TYPHIMURIUM *	2334	19.6
2	ENTERITIDIS	5343	16.3	2	HEIDELBERG	330	6.2	2	HEIDELBERG	1860	15.6
3	NEWPORT	2618	8.0	3	AGONA	233	4.4	3	KENTUCKY	942	7.9
4	HEIDELBERG	1816	5.5	4	CHOLERAESUIS **	207	3.9	4	ENTERITIDIS	568	4.8
5	MUENCHEN	1332	4.1	5	NEWPORT	183	3.4	5	MONTEVIDEO	533	4.5
6	JAVIANA	1197	3.7	6	ANATUM	178	3.3	6	DERBY	471	3.9
7	MONTEVIDEO	851	2.6	7	DERBY	174	3.3	7	SENFTENBERG	466	3.9
8	ORANIENBURG	616	1.9	8	SENFTENBERG	159	3.0	8	HADAR	453	3.8
9	THOMPSON	602	1.8	9	MUENSTER	154	2.9	9	MUENSTER	295	2.5
10	INFANTIS	596	1.8	10	MONTEVIDEO	144	2.7	10	ANATUM	263	2.2
11	BRAENDERUP	529	1.6	11	DUBLIN	129	2.4	11	INFANTIS	243	2.0
12	AGONA	528	1.6	12	KENTUCKY	86	1.6	12	AGONA	202	1.7
13	HADAR	516	1.6	13	MBANDAKA	76	1.4	13	THOMPSON	198	1.7
14	SAINTPAUL	472	1.4	14	BREDENEY	72	1.3	14	CHOLERAESUIS **	189	1.6
15	TYPHI	352	1.1	15	INFANTIS	67	1.3	15	OHIO	185	1.6
16	JAVA	314	1.0	16	CERRO	64	1.2	16	READING	168	1.4
17	POONA	249	0.8	17	MUENCHEN	61	1.1	17	BRANDENBURG	163	1.4
18	MISSISSIPPI	248	0.8	18	THOMPSON	55	1.0	18	MUENCHEN	150	1.3
19	MBANDAKA	231	0.7	19	MELEAGRIDIS	53	1.0	19	MBANDAKA	144	1.2
20	DERBY	174	0.5	20	ORANIENBURG	49	0.9	20	ORION	124	1.0
	SUB TOTAL	26635	81.2		SUB TOTAL	4339	81.3		SUB TOTAL	9951	83.4
	TOTAL	32782			TOTAL	5337			TOTAL	11931	

\* TYPHIMURIUM INCLUDES VAR. COPENHAGEN  
 \*\* CHOLERAESUIS INCLUDES VAR. KUNZENDORF