

Table 33. Tuberculosis Cases and Percentages by Noninjecting Drug Use¹: 60 Reporting Areas, 2004

Reporting Area	Total Cases	Cases with Information on Noninjecting Drug Use		Cases Reporting Noninjecting Drug Use	
		No.	(%)	No.	(%)
United States	14,517	14,114	(97.2)	1,038	(7.4)
Alabama	211	211	(100.0)	12	(5.7)
Alaska	43	39	(90.7)	4	(10.3)
Arizona	272	258	(94.9)	31	(12.0)
Arkansas	132	132	(100.0)	1	(0.8)
California	2,989	2,901	(97.1)	169	(5.8)
Colorado	127	126	(99.2)	6	(4.8)
Connecticut	101	93	(92.1)	5	(5.4)
Delaware	32	32	(100.0)	0	(0.0)
District of Columbia	81	81	(100.0)	4	(4.9)
Florida	1,076	1,060	(98.5)	163	(15.4)
Georgia	536	510	(95.1)	52	(10.2)
Hawaii	116	79	(68.1)	—	—
Idaho	11	8	(72.7)	—	—
Illinois	569	501	(88.0)	40	(8.0)
Indiana	128	128	(100.0)	14	(10.9)
Iowa	47	47	(100.0)	2	(4.3)
Kansas	62	62	(100.0)	9	(14.5)
Kentucky	127	126	(99.2)	9	(7.1)
Louisiana	249	230	(92.4)	31	(13.5)
Maine	20	20	(100.0)	0	(0.0)
Maryland	314	310	(98.7)	14	(4.5)
Massachusetts	283	277	(97.9)	15	(5.4)
Michigan	273	265	(97.1)	14	(5.3)
Minnesota	199	198	(99.5)	4	(2.0)
Mississippi	119	119	(100.0)	10	(8.4)
Missouri	127	122	(96.1)	12	(9.8)
Montana	15	15	(100.0)	0	(0.0)
Nebraska	39	39	(100.0)	0	(0.0)
Nevada	95	91	(95.8)	5	(5.5)
New Hampshire	24	24	(100.0)	2	(8.3)
New Jersey	482	482	(100.0)	23	(4.8)
New Mexico	42	30	(71.4)	—	—
New York State ²	324	319	(98.5)	9	(2.8)
New York City	1,039	1,015	(97.7)	76	(7.5)
North Carolina	382	378	(99.0)	52	(13.8)
North Dakota	4	4	(100.0)	1	(25.0)
Ohio	219	218	(99.5)	25	(11.5)
Oklahoma	178	178	(100.0)	12	(6.7)
Oregon	106	106	(100.0)	4	(3.8)
Pennsylvania	327	325	(99.4)	10	(3.1)
Rhode Island	51	51	(100.0)	1	(2.0)
South Carolina	234	228	(97.4)	38	(16.7)
South Dakota	11	11	(100.0)	0	(0.0)
Tennessee	279	275	(98.6)	29	(10.5)
Texas	1,683	1,663	(98.8)	88	(5.3)
Utah	36	36	(100.0)	2	(5.6)
Vermont	6	6	(100.0)	0	(0.0)
Virginia	329	326	(99.1)	4	(1.2)
Washington	244	239	(98.0)	25	(10.5)
West Virginia	24	23	(95.8)	0	(0.0)
Wisconsin	95	92	(96.8)	9	(9.8)
Wyoming	5	5	(100.0)	0	(0.0)
American Samoa ³	3	3	(100.0)	0	(0.0)
Fed. States of Micronesia ³	8	8	(100.0)	1	(12.5)
Guam ³	51	50	(98.0)	0	(0.0)
Marshall Islands ³	41	39	(95.1)	0	(0.0)
N. Mariana Islands ³	55	53	(96.4)	1	(1.9)
Puerto Rico ³	123	121	(98.4)	33	(27.3)
Republic of Palau ³	5	5	(100.0)	0	(0.0)
U.S. Virgin Islands ³

¹Noninjecting drug use within past 12 months of TB diagnosis. Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for $\geq 75\%$ of cases.

²Excludes New York City.

³Not included in U.S. totals.

Note: Ellipses indicate data not available.