

TABLE A14.—Experiments concerning the effect of cigarette smoke on pulmonary surfactant and surface tension

Author, year, country, reference	System	Method	Results																				
Miller and Bondurant, 1962, U.S.A. (165)	Rat lung extracts	Cigarette smoke: (1) Applied to extract. (2) Exposure of rats.	(1) Exposure to cigarette smoke was associated with decreased surface tension in lung extract. (2) Surface tension of rats (lung extracts) exposed to cigarette smoke was decreased as compared with those not exposed.																				
Cook and Webb, 1966, U.S.A. (57)	40 subjects undergoing bronchoscopy: 14 normal 7 nonsmokers with pulmonary disease 19 smokers with and without pulmonary disease.		<p style="text-align: center;">Surface tension values of surfactant</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th>20 percent area</th> <th>100 percent area</th> <th>Stability index (reflects surfactant activity)</th> <th>† Values significantly different from values of normals at p<0.02 level.</th> </tr> </thead> <tbody> <tr> <td>Normal</td> <td>6.5</td> <td>60.0</td> <td>1.61</td> <td></td> </tr> <tr> <td>Pulmonary patients</td> <td>†17.0</td> <td>150.0</td> <td>1.00</td> <td></td> </tr> <tr> <td>Chronic smokers ..</td> <td>15.7</td> <td>51.0</td> <td>1.04</td> <td></td> </tr> </tbody> </table>		20 percent area	100 percent area	Stability index (reflects surfactant activity)	† Values significantly different from values of normals at p<0.02 level.	Normal	6.5	60.0	1.61		Pulmonary patients	†17.0	150.0	1.00		Chronic smokers ..	15.7	51.0	1.04	
	20 percent area	100 percent area	Stability index (reflects surfactant activity)	† Values significantly different from values of normals at p<0.02 level.																			
Normal	6.5	60.0	1.61																				
Pulmonary patients	†17.0	150.0	1.00																				
Chronic smokers ..	15.7	51.0	1.04																				
Giammona, 1967, U.S.A. (94)	<i>In vitro:</i> Surfactant material induced from dogs and rats. <i>In vivo:</i> Dogs, cats, and guinea pigs.	Exposed to cigarette smoke for 3 hours/day for up to 3 weeks.	<i>In vitro:</i> Exposure to cigarette smoke was associated with a significant decrease in maximal surface tension. <i>In vivo:</i> Dogs and cats (exposed for 1 week)—no significant change. Guinea pigs (exposed for 3 weeks)—significant decrease in maximal surface tension.																				
Webb, et al., 1967, U.S.A. (224)	Bronchial washing, from dog lungs.	Direct exposure to cigarette smoke.	<p style="text-align: center;">Surface tension values of surfactant</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th>Number</th> <th>20 percent area</th> <th>100 percent area</th> <th>Stability index</th> </tr> </thead> <tbody> <tr> <td>Control</td> <td>11</td> <td>7.1</td> <td>60.7</td> <td>1.60</td> </tr> <tr> <td>Smoke</td> <td>10</td> <td>18.7</td> <td>45.8</td> <td>0.84</td> </tr> </tbody> </table> <p style="text-align: center;">(p<0.002) (p<0.002)</p>		Number	20 percent area	100 percent area	Stability index	Control	11	7.1	60.7	1.60	Smoke	10	18.7	45.8	0.84					
	Number	20 percent area	100 percent area	Stability index																			
Control	11	7.1	60.7	1.60																			
Smoke	10	18.7	45.8	0.84																			

TABLE A15.—Studies concerning the relationship of smoking to infectious respiratory disease in humans
(Actual number of cases shown in parentheses)
SM = Smokers NS = Nonsmokers

Author, year, country, reference	Number and type of population	Data collection	Results				Comments			
			NS	SM	Cases	Controls				
Mills, 1960, U.S.A. (167).	118 male and female patients with pneumonia and 472 healthy individuals from "random" sample.	Hospital Interview.	Mean age		49.6	49.6	The author stated that there was a significant difference in tobacco usage between the two groups.			
			NS		16.25	26.21				
			Cigarettes only		63.56	52.33				
			Mixed		21.19	22.46				
Lowe, 1956, England (157).	520 male and 185 female tuberculosis patients and 419 male and 249 female control outpatients.	Interview by trained social worker.	Males		Females		Cigarette smokers include pipe smokers. The author noted a significant deficiency of non- and light smokers and an excess of heavy smokers among the cases			
			NS	2.5	8.1	Cases		Controls	Cases	Controls
			Cigarettes/day: 1-9	9.2	12.9	37.3		51.4	20.5	25.7
			10-19	38.1	35.6	30.8		20.5		
			20-29	29.4	27.4					
			30-39	11.3	9.3				11.4	2.4
			>40	9.4	6.7					
Dowling, et al., 1957, U.S.A. (72).	Individuals exposed to "infectious cold agent" and placebo.	Interview and medical examination.	Exposed to placebo		Exposed to infectious agent		No statistically significant differences noted.			
			Percent developing "cold"		Percent developing "cold"					
			NS	111	10	328		34		
SM	78	14	249	36						

TABLE A15.—Studies concerning the relationship of smoking to infectious respiratory disease in humans (cont.)

(Actual number of cases shown in parentheses)
SM = Smokers NS = Nonsmokers

Author, year, country, reference	Number and type of population	Data collection	Results			Comments	
Boake, 1968, U.S.A. (33).	Parents of 59 families.	Interview		<i>Person-years</i>	<i>Number of respiratory illnesses/ person-years</i>	No statistically significant differences noted.	
			NS (24)	120	524		5.2
			Cigarettes/day: 1-10 (19)	99	529		5.3
			11-20 (25)	108	486		4.5
			>20 (19)	99	424		4.3
Pipe, cigar (14)	72	304	4.2				
Shah et al., 1969, India (205).	Tuberculosis institute employees.	Survey, X-ray, and interview.		<i>Tuberculous by X-ray</i>	<i>Normal or nontuberculous</i>	† Numbers in parentheses represent figures "expected" by use of 2 x 2 contingency table. Tuberculous employees were found to have significantly fewer nonsmokers and more smokers.	
			NS †10 (10.7)	178 (168.3)			
SM 36 (26.3)	215 (224.7)						

TABLE A15.—Studies concerning the relationship of smoking to infectious respiratory disease in humans (cont.)

(Actual number of cases shown in parentheses)

SM = Smokers NS = Nonsmokers

Author, year, country, reference	Number and type of population	Data collection	Results			Comments	
Brown et al., 1961, Australia (4).	306 male and female tuberculosis patients, 221 male and female outpatients.	Interview	<i>Smoking habits prior to diagnosis</i>			Data presented only on Queensland sample. The authors noted that the significant difference between the patients and controls was not present when the groups were matched for alcohol intake.	
				<i>Tuberculous patients (percent)</i>	<i>Controls (percent)</i>		
			NS	9.1	19.0		
			Cigarettes/day: 1-9	10.5	15.4		
			10-19	34.3	19.5		
			20-29	26.3	26.8		
			30-39	7.2	5.4		
>40	6.2	9.1					
Pipes	5.9	4.6					
Haynes et al., 1966, U.S.A. (108).	191 male prep school students.	Interview	<i>Average number of respiratory illnesses/10 students (adjusted for age)</i>				
				<i>All respiratory episodes</i>	<i>All severe respiratory episodes</i>		<i>All severe lower or combined respiratory episodes</i>
			NS (99)	11.1	1.6		0.36
			SM (92)	20.2	6.7		3.34
Parnell et al., 1966, Canada (181).	47 smoking-nonsmoker pairs of student nurses and parents' occupational class.	Interview and health service records.	<i>Median number of illnesses/student</i>			The authors noted that these differences were statistically significant. † Particularly tracheitis, bronchitis, and pneumonia.	
				<i>All respiratory diseases†</i>	<i>All other illnesses</i>		
			NS (47)	2.08	2.99		
SM (47)	2.54	5.00					

TABLE A15.—Studies concerning the relationship of smoking to infectious respiratory disease in humans (cont.)

(Actual number of cases shown in parentheses)
 SM = Smokers NS = Nonsmokers

Author, year, country, reference	Number and type of population	Data collection	Results	Comments		
Peters et al., 1967, U.S.A. (183).	1,496 Harvard and 370 Radcliffe students.	Medical history, chart review, and questionnaire.	Number of visits to student health unit for respiratory illness/student (common colds, pharyngitis, bronchitis, laryngitis, pneumonia—not allergic rhinitis)		† p<0.001.	
				Harvard		Radcliffe
			NS	1.44 (771)		1.44 (193)
			SM	12.27 (726)		2.27 (177)
			<2 years smoked	2.00		
			3-4	2.30		
>5	2.50					
Finklea et al., 1969, U.S.A. (83).	1,811 male college students.	Questionnaire prior to A ₂ /HK/68 epidemic and follow-up on morbidity.	Heavy smokers—21 percent more clinical illnesses than nonsmokers; 20 percent more requiring bed rest than nonsmokers Light smokers—10 percent more clinical illnesses than nonsmokers; 7 percent more requiring bed rest than nonsmokers.	The authors also noted that: (a) Smokers exhibited serologic evidence of increased subclinical A ₂ /HK/68 infection. (b) There was no difference in the vaccination status between smokers and nonsmokers.		

TABLE A16.—Complications developing in the postoperative period in patients undergoing abdominal operations

Men over 20					
Group	Cases	Percent chest clear	Percent bronchitis	Percent broncho-pneumonia and atelectasis	Percent total complication rate
Smokers	300	41.7	53.0	5.3	58.3
Light Smokers	180	68.4	27.7	3.9	31.6
Nonsmokers	66	92.5	6.0	1.5	7.5
Women over 20					
Smokers	23	39.1	43.5	17.4	60.9
Light Smokers	62	77.5	20.9	1.6	22.5
Nonsmokers	518	88.8	8.1	3.1	11.2

SOURCE: Morton, H. J. V. (173)

TABLE A17.—Arterial oxygen saturation before and after operation

Arterial oxygen saturation (percentage)					
Group	Case number	Before operation	Day 1	Day 2	Day 3
Nonsmokers	1	94	93	94	..
	2	94	93	94	..
	3	96	93	94	..
	4	95	90	94	..
	5	94	90	93	..
Smokers	6	95	91	89	91
	7	92	89	81	89
	8	91	89	85	89
	9	93	91	88	92
	10	90	87	88	92

SOURCE: Morton, A. (172).

Chapter 4
Cancer

Source: 1971 Report, Chapter 4, pages 231 - 384.

Contents

	<i>Page</i>
Introduction	263
Lung Cancer	265
Epidemiological Studies	266
Prospective Studies	266
Retrospective Studies	266
Lung Cancer Trends in Other Countries	270
Histology of Lung Tumors	272
Lung Cancer Relationships in Women	277
Lung Cancer, the Urban Factor, and Air Pollution....	278
Lung Cancer and Occupational Hazards	282
Uranium Mining	282
Other Occupations.....	282
Nickel.....	282
Asbestos.....	283
Arsenic.....	283
Chromium.....	283
Pathological Studies.....	284
Pulmonary Carcinogenesis	284
General Aspects of Carcinogenesis	284
Polynuclear Aromatic Hydrocarbons	290
Nitrosamine Compounds.....	290
Pesticides and Fungicides.....	292
Radioactive Isotopes	292
Inhibitors of Ciliary Movement	293
Experimental Studies	293
Skin Painting and Subcutaneous Injection....	293
Tissue and Organ Culture	293
Tracheobronchial Implantation	
and Instillation	294
Inhalation	294
Reduction in Tumorigenicity	301
Summary and Conclusions	302
Cancer of the Larynx	303
Epidemiological Studies	303
Pathological Study	306
Experimental Study	307
Summary and Conclusions	307
Oral Cancer	310
Epidemiological Studies	311
Experimental Studies	314
Summary and Conclusions	315

	<i>Page</i>
Cancer of the Esophagus	315
Epidemiological Studies	315
Pathological Study	318
Experimental Studies	318
Summary and Conclusions	319
Cancer of the Urinary Bladder and Kidney	319
Epidemiological Studies (Bladder)	319
Epidemiological Studies (Kidney)	322
Experimental Studies	322
Summary and Conclusions	325
Cancer of the Pancreas	325
Summary and Conclusions	325
References	325

FIGURES

1. Lung cancer, Finland and Norway	271
2. Percent of smoking dogs with tumors	300
3. Percent of lung lobes with tumors in smoking dogs	300
4. Effects of chronic cigarette smoke inhalation on the hamster larynx	310

LIST OF TABLES

1. Lung cancer mortality ratios	267
2. Lung cancer mortality ratios for males by duration of cigarette smoking	270
A3. Outline of methods used in retrospective studies of smoking in relation to lung cancer	349
A4. Group characteristics in retrospective studies on lung cancer and tobacco use	355
5. Annual means of total lung cancer mortality and sex ratios for selected periods in Finland and Norway	272
6. Epidemiologic and pathologic investigations concern- ing smoking and histology of lung cancer	273
A7. Grouping of pulmonary carcinomas	360
8. Tumor prevalence among males and females 35-69 years of age, by type of tumor and smoking category	276
9. Epidemiologic investigations concerning the relation- ship of lung cancer to smoking, air pollution, and urban or rural residence	279
10. Pathologic and cytologic findings in the tracheo- bronchial tree of smokers and nonsmokers	285

LIST OF TABLES (Continued)

(A indicates tables located in appendix at end of chapter)

	<i>Page</i>
11. Identified or suspected tumorigenic agents in cigarette smoke	291
A12. Autopsy studies concerning the presence of radioactivity in the lungs of smokers	361
A13. Experiments concerning the effects of the skin painting or subcutaneous injection of cigarette smoke condensate or its constituents upon animals	363
A14. Experiments concerning the effect of cigarette smoke or its constituents on tissue and organ cultures	369
A15. Experiments concerning the effect of the instillation or implantation of cigarette smoke or its constituents into the tracheobronchial tree of animals	372
A16. Experiments concerning the effect of the inhalation of cigarette smoke or its constituents upon the respiratory tract of animals	375
17. Data on pedigreed male beagle dogs of groups F, L, H, h and N	296
18. Summary of principal cause of death (days No. 57 through No. 875) in dogs of groups F, L, H, h and N	297
19. Data on dogs with lung tumors indicating type of tumor and lobe in which the tumor was found	298
20. Laryngeal cancer mortality ratios—prospective studies	304
A21. Outline of retrospective studies of tobacco use and cancer of the larynx	380
A22. Summary of results of retrospective studies of tobacco use and cancer of the larynx	384
A23. Number and percent distribution by relative frequency of atypical nuclei among true vocal cord cells, of men classified by smoking category	385
A24. Number and percent distribution, by highest number of cell rows in the basal layer of the true vocal cord, of men classified by smoking category	386
25. Deposition of ¹⁴ C-labeled smoke particles in particular regions of the respiratory tract	308
26. Classification of the five registered stages of epithelial changes at the larynx	309
27. Oral cancer mortality ratios—prospective studies	312
A28. Outline of retrospective studies of tobacco use and cancer of the oral cavity	387
A28a. Summary of results of retrospective studies of smoking by type and oral cancer of the detailed sites	394
	261

LIST OF TABLES (Continued)

(A indicates tables located in appendix at end of chapter)

	<i>Page</i>
A29. Experimental studies concerning oral carcinogenesis	397
30. Esophageal cancer mortality ratios—prospective studies	316
A31. Summary of methods used in retrospective studies of tobacco use and cancer of the esophagus	401
A31a. Summary of results of retrospective studies of tobacco use and cancer of the esophagus	404
A32. Atypical nuclei in basal cells of epithelium of esophagus of males, by smoking habits and age	405
A33. Atypical nuclei in basal cells of epithelium of esophagus of males, by amount of smoking and age	406
34. Kidney and urinary bladder cancer—prospective studies	320
A35. Summary of methods used in retrospective studies of smoking and cancer of the bladder	407
A35a. Summary of results of retrospective studies of smoking and cancer of the bladder	409
36. Pancreatic cancer mortality ratios—prospective studies	324

INTRODUCTION

During the early years of this century, a number of pathologists and clinicians reported a dramatic increase in the incidence of lung cancer. Autopsy studies and studies of lung cancer death rates revealed a significant increase beginning prior to World War I and continuing during the ensuing years. This epidemic of lung cancer continues to the present day, with nearly 60,000 deaths expected from this disease in the United States during 1970.

Beginning in the 1920's, a number of reports appeared which suggested a relationship between lung cancer and tobacco smoking (4, 203, 278). Since that time, many clinical and epidemiological studies have been published which confirm this relationship. The 1964 Report (291) contains a thorough review and analysis of the data available at that time as well as an excellent discussion of the considerations necessary for their evaluation.

Major epidemiological studies have demonstrated that smokers have greatly increased risks of dying from lung cancer compared to nonsmokers. An increased risk of lung cancer has been found for every type of smoking habit investigated, but two characteristics of the risk are particularly evident: The risk is much greater for cigarette smokers than for smokers of pipes and cigars, and among cigarette smokers a dose relationship exists. That is, the more one smokes, as measured by total pack-years of smoking, present level of smoking, degree of inhalation, or age at start of smoking, the greater is the risk. It has also been shown that the risk of lung cancer among ex-smokers decreases with time almost to the level of nonsmokers; the time required is dependent on the degree of exposure prior to cessation.

Pathologists have found that the squamous cell or epidermoid form of lung cancer is the most prevalent one in cigarette smoking populations and that this form accounts for a major portion of the rise in lung cancer deaths (154). Such studies have also indicated a lower prevalence among smokers for oat-cell and adenocarcinomas of the lung than for the squamous form, but in most studies a higher frequency of these tumors is found among smokers than among nonsmokers.

Smoking has been implicated in the development of other types of cancer in humans. Among these is cancer of the larynx. A num-

ber of epidemiological studies have demonstrated increased mortality rates for laryngeal cancer in smokers, particularly cigarette smokers, compared with nonsmokers. Autopsy studies have revealed that a clear dose-relationship exists between smoking and the development of cellular changes in the larynx, including carcinoma *in situ*.

Cancers of the mouth and oropharynx have been found to be more common among users of all types of tobacco than among abstainers. Although smoking is a definite risk factor in the development of malignant lesions of the oral cavity and pharynx, its relative contribution in conjunction with other factors such as poor nutrition and alcohol consumption has not been fully clarified.

Similarly, although smokers are more likely to develop carcinoma of the esophagus than nonsmokers, the relative additional contribution of smoking in conjunction with nutritional factors and alcohol consumption requires clarification.

Smokers have been found to be more at risk for the development of cancer of the urinary bladder than are nonsmokers, and there is evidence to suggest that some smoking-induced abnormal metabolic product or abnormal concentration of a metabolic product may be responsible for this increased risk. In addition, cancer of the kidney is apparently more common in smokers than in nonsmokers, but the epidemiologic evidence for this relationship is not as definite as for bladder cancer.

Epidemiological studies have indicated an association between smoking and cancer of the pancreas. The significance of this relationship is unclear at this time.

Experimental studies have demonstrated the carcinogenicity of the condensate of tobacco smoke, or "tar." This material, when painted on the skin of animals, leads to the development of squamous cell tumors of the skin. Researchers have shown that this condensate contains substances known as carcinogens, capable of inducing cancers. Among these carcinogens are several chemicals which have been identified as tumor initiators, that is, compounds which initiate changes in target cells and also tumor promoters, or compounds which promote the neoplastic development of initiated cells. Other, as yet unidentified, factors are presumably also involved because the sum of the carcinogenic effects of the known agents does not equal that of cigarette smoke condensate.

Numerous experiments have been performed in which whole cigarette smoke, filtered smoke, or certain constituents of smoke, such as the "tar," are administered by varying methods to animals or to tissue and cell cultures in order to investigate the neoplastic-inducing properties of cigarette smoke. Particular difficulty has been encountered in experiments which have attempted to deliver

whole cigarette smoke to the larynx and into the lungs of experimental animals. This has resulted in the use of other methods such as the implanting of pellets containing suspected carcinogens and the instilling into the trachea of suspected carcinogens as such, or adsorbed onto fine inert particulate matter as a carrier. The difficulty with the inhalation studies has been twofold. First, the animals, particularly the smaller species such as the rat, frequently die from the acute toxic effects of the nicotine and carbon monoxide in the tobacco smoke. Second, the upper respiratory tract of experimental animals, particularly the nose, is much different from analogous human structures, resulting in a more efficient filtration of smoke in the upper respiratory tract. Nevertheless, in rodents and canines, progressive changes apparently indicative of ultimate neoplastic transformation have been identified in the respiratory tract.

Recently, two studies in different species and in different target organs have been reported concerning the development of early invasive cancer following the prolonged inhalation of cigarette smoke. Auerbach and his coworkers (11) trained dogs to inhale cigarette smoke through a tracheostoma. After approximately 29 months of daily exposure, these investigators found a number of cancers of the lung.

Dontenwill (76) in the second of these two studies, exposed hamsters to the passive inhalation of cigarette smoke over varying and prolonged periods of time. He observed the development of pre-malignant changes and, ultimately, invasive squamous cell cancer of the larynx.

LUNG CANCER

Cancer of the lung in the United States accounted for 45,383 deaths among males and 9,024 deaths among females in 1967 (289). It is presently estimated that approximately 60,000 people will die of lung cancer during 1970.

The alarming epidemic of lung cancer is a relatively recent phenomenon. Death rates for lung cancer (ICD Codes 162, 163) rose from 5.6 (per 100,000 resident population per year) in 1939 to 27.5 in 1967 (289, 290). This rapid increase followed the increased use of cigarettes among the United States population. The increase has occurred principally among males, although more recently females have shown a similar rising pattern.

The converging evidence for the conclusion that cigarette smoking is the major cause of lung cancer is derived from varied types of research including epidemiological, pathological, and laboratory investigations.

EPIDEMIOLOGICAL STUDIES

Numerous epidemiological studies, both retrospective and prospective, have been carried out in different parts of the world to investigate the relationship between smoking and cancer of the lung. These studies are outlined in tables 1, 2, A3, and A4.

Prospective Studies

The major prospective studies concerning the relationship of smoking and lung cancer are presented in table 1. In all, these investigations have studied more than a million persons from a number of different populations for up to 10 years. These studies show increased lung cancer mortality ratios for cigarette smokers of all amounts ranging from 7.61 to 14.20 among male smokers as compared to nonsmoking males. The one major prospective study of female cigarette smokers reveals an overall mortality ratio of 2.20 (118).

Also uniformly present in these studies is a dose-related increase in the mortality from lung cancer with increasing amounts of cigarettes smoked per day. Other measures of exposure show similar trends. Hammond (118) reported increased mortality ratios associated with increased inhalation (table 1) as well as with increased duration of smoking (table 2).

Ex-smokers show significantly lower lung cancer death rates than continuing smokers. In their study of more than 40,000 British physicians, Doll and Hill (74, 75) noted a decrease in lung cancer mortality rates with increasing time since smoking stopped (table 1). During the past 20 years, half of all the physicians in Britain who used to smoke cigarettes have stopped smoking. While the death rates from lung cancer rose by 7 percent among all men from England and Wales during the period from 1953-57 through 1961-65, the rates for male doctors of the same ages fell by 38 percent (96).

Pipe and cigar smokers have been shown in the prospective studies to have lung cancer mortality rates higher than those of nonsmokers, although these are generally substantially lower than those of cigarette smokers (table 1).

Retrospective Studies

More than 30 retrospective (case-control) studies have been reported concerning the relationship of smoking and lung cancer. These studies are outlined in tables A3 and A4. Table A4 presents the percent of nonsmokers and of heavy smokers among both cases and controls as well as the relative risk ratios for all smokers.

TABLE 1.—Lung cancer mortality ratios
(Actual number of deaths shown in parentheses)¹
SM = Smokers. NS = Nonsmokers.

Prospective studies									
Author, year, country, reference	Number and type of population	collection Data	Follow-up years	Number of deaths	Regular cigarette smoking only (cigarettes/day)	Pipe cigar	Inhalation	Exsmokers	Comments
Hammond and Horn, 1968, U.S.A. (140).	187,783 white males in 9 States ages 50-69.	Questionnaire and interview.	3½	448	SM . 443	Pipe	No data	Bronchogenic	341/448 deaths with microscopic proof. Includes those regular cigarette smokers who also smoked pipes and cigars. † With or without microscopic proof.
					NS . 15	NS ... 1.00 (15)	SM ... 2.67 (18)	(Excluding adenocarcinoma)	
					<10 ... 8.00 (24)	Cigar		Never smoked ... 1.00	
					10-20 ... 10.50 (84)			Previously <1 pack/day	
					>20 ... 23.40 (117)	NS ... 1.00 (15)		Continuing ... 16.94	
					All ... †10.73 (397)	SM ... 1.00 (7)		Duration } <1 year ... 16.60	
								of } 1-10 years ... 10.44	
								cessation } >10 years ... 1.51	
								Previously >1 pack/day	
								Continuing ... 46.21	
								Duration } <1 year ... 58.23	
								of } 1-10 years ... 22.82	
								cessation } >10 years ... 17.79	
Doll and Hill, 1964, Great Britain (74).	Approximately 41,000 male British physicians	Questionnaire and followup of death certificate.	10	212	NS ... 1.00 (3)	Pipe and Cigar Grams/day	No data	Cigarette smokers	
				SM . 209	1-14 ... 8.14 (22)	NS ... 1.00 (3)		NS ... 1.00 (3)	
				NS . 8	15-24 ... 19.86 (53)			Continuing ... 18.29 (124)	
					>25 ... 32.43 (57)	1-14 ... 6.00 (12)		Duration } <5 years ... 9.57 (5)	
						15-24 ... 6.43 (6)		of } 5-9 years ... 7.00 (7)	
						>25 ... 13.71 (3)		cessation } 10-20 years ... 2.67 (3)	
								>20 years ... 2.71 (2)	
Beat, 1966, 1966, Canada (21).	Approximately 78,000 male Canadian veterans.	Questionnaire and followup of death certificate.	6	331	NS ... 1.00 (7)	Pipe	No data	Ex-smokers of cigarettes only	† Refers to current cigarette smokers only.
				†SM . 324	<10 ... 10.00 (57)	NS ... 1.00 (7)		NS ... 1.00 (7)	
				NS . 7	10-20 ... 16.41 (204)	SM ... 4.35 (18)		Ex-smokers of cigarettes only ... 6.06 (18)	
					>20 ... 17.31 (53)	Cigar			
					All ... 14.20 (245)	NS ... 1.00 (7)			
						SM ... 2.94 (2)			

TABLE 1.—Lung cancer mortality ratios (cont.)

(Actual number of deaths shown in parentheses)¹

SM = Smokers. NS = Nonsmokers.

Prospective studies									
Author, year, country, reference	Number and type of population	Data collection	Follow-up years	Number of deaths	Regular cigarette smoking only (cigarettes/day)	Pipe cigar	Inhalation	Exsmokers	Comments
Kahn (Dorn), 1966, U.S.A. (139).	U.S. male veterans 2,265,674 person years.	Questionnaire and followup of death certificate.	8½	1,256					
				SM .1,178	NS 1.00 (78)	NS 1.00 (78)	No data		
				NS . . 78	1-9 5.49 (46)	SM 1.84 (17)		NS 1.00 (78)	
					10-20 9.91 (303)			Number of cigarettes/day:	
					21-39 17.41 (815)	NS 1.00 (78)		1-9 0.95 (4)	
					>39 23.93 (82)	SM 1.59 (6)		10-20 3.48 (89)	
					All 12.14 (749)			21-39 9.38 (57)	
								>39 8.24 (19)	
Hammond, 1966, U.S.A. (118).	440,558 males 562,671 females 35-84 years of age in 25 States.	Interviews by ACS volunteers.	4	Males	Current cigarettes	Pipe		Males	
					1,159	NS 1.00 (49)	NS 1.00 (49)		ICD code
					SM .1,110	SM 2.24 (21)	Slight 8.42 (120)		162 only.
					NS . . 49	NS 1.00 (49)	Cigar	Moderate 11.45 (811)	
					Females	1-9 4.60 (26)	NS 1.00 (49)	Deep 14.31 (141)	
					183	10-19 7.48 (82)	SM 1.85 (22)	Females	
					SM . . 81	20-39 13.14 (381)	Pipe and cigar	NS 1.00 (102)	
					NS . . 102	>40 16.61 (82)	NS 1.00 (49)	Slight 1.78 (25)	
						All 9.20 (719)	SM 0.90 (11)	Moderate } 3.70 (45)	
						Females		Deep }	
						NS 1.00 (102)			
						1-19 1.06 (20)			
						>20 4.76 (50)			
		All 2.20 (81)							

TABLE 1.—Lung cancer mortality ratios (cont.)

(Actual number of deaths shown in parentheses)¹

SM = Smokers. NS = Nonsmokers.

Prospective studies									
Author, year, country, reference	Number and type of population	Data collection	Follow-up years	Number of deaths	Regular cigarette smoking only (cigarettes/day)	Pipe cigar	Inhalation	Exsmokers	Comments
Buell et al., 1967, U.S.A. (49).	69,868 American Legionnaires 35-75 years of age and older.	Questionnaire and followup of death certificate.	3	304	NS 1.00 <20 2.30 20 ... 3.50 >20 4.90				
Hirayama, 1967, Japan (125).	265,118 male and female adults 40 years of age and older.	Trained PHS nurse interview and followup of death certificate.	1½	43 SM . 40	NS 1.00 (3) 1-24 ... 2.69 (29) >25 5.68 (5)				Preliminary report.
Weir and Dunn, 1970, U.S.A. (508).	68,153 males in various occupations in California.	Questionnaire and followup of death certificate.	5-8	888	NS 1.00 ±10 8.72 ±20 9.05 >30 9.66 All 7.61				NS include pipe and cigar smokers SM include ex-smokers.

¹ Unless otherwise specified, disparities between the total number of deaths and the sum of the individual smoking categories are due to the exclusion of either occasional, miscellaneous, mixed, or exsmokers.

TABLE 2.—Lung cancer mortality ratios for males
by duration of cigarette smoking
(Actual number of deaths are shown in parentheses)

Age began cigarette smoking	35-54	55-59	70-84	35-84
25 or older	2.77 (5)	3.39 (12)	3.38 (3)	3.21 (20)
20-24	5.83 (31)	11.11 (72)	12.11 (7)	9.72 (110)
15-19	8.71 (112)	13.06 (176)	19.37 (27)	12.81 (315)
<15	12.80 (36)	16.81 (57)	16.76 (9)	15.10 (101)

SOURCE: Hammond, E. C. (118).

These smoker-nonsmoker risk ratios range from 1.2 to 36.0 for males and from 0.2 to 5.3 for females.

Although not presented in tabular form, the data concerning lung cancer and pipe or cigar smoking are similar to those found by the prospective studies mentioned above. However, a study by Abelin and Gsell (1) conducted on a rural Swiss population noted that an increased risk of lung cancer was present among heavy cigar and pipe smokers (as well as cigarette smokers) to a greater degree than previously reported. The authors suggest that their findings might be due to differences in either the amount smoked or the carcinogenicity of Swiss and German cigars. The difference might also be explained by the greater use and more frequent inhalation of small cigars in Switzerland as compared to other countries where large cigars are more commonly smoked but rarely inhaled. Kreyberg (154), in a review of 887 cases of lung cancer in Norway, noted that pipe smokers showed an increased risk of lung cancer, although this risk was substantially lower than that for cigarette smokers.

LUNG CANCER TRENDS IN OTHER COUNTRIES

Several studies of particular interest are those in which the changing mortality from lung cancer has been investigated in countries in which cigarette smoking has become popular and widespread only in recent years. In those countries where accurate statistics for lung cancer mortality are available for both the pre-smoking and post-smoking periods, long-term trends can be studied in some detail.

Two such studies have dealt with lung cancer mortality trends in Iceland. Dungal (83) noted in 1950 that lung cancer was a rare disease in Iceland and felt that this rarity could be explained by the relatively late onset of heavy tobacco smoking in the Icelandic population when compared to that of Great Britain and Finland. He observed that the annual per capita consumption of tobacco did not reach one pound in Iceland until 1945, while Great Britain and Finland passed that amount before 1920. In 1967, Thorarinsson, et al. (276) noted a sharp rise in the incidence of lung cancer in Ice-




FIGURE 1.—Lung cancer, Finland and Norway.
SOURCE: Kreyberg, L. (154).

land after 1950 and found a correlation between that increase and the increasing sale of cigarettes in that country.

Kreyberg (154) analyzed the lung cancer death rates of both Norway and Finland in relation to the use of tobacco in those two countries over the past 100 years. Figure 1 shows the substantial difference in lung cancer mortality between the two countries. Kreyberg observed that cigarettes came into use in Norway in 1886 while the Finnish population (more closely allied to Russia socio-economically) was consuming more than 100 million cigarettes per year during the decade of the 1880's. Cigarettes remained scarce in Norway until after World War I, and this 30-year lag in consump-

TABLE 5.—Annual means of total lung cancer mortality and sex ratios for selected periods in Finland and Norway

Year	Finland		Norway	
	Males	Females	Males	Females
1936-38	192	33	34	30
Sex ratio	5.8 : 1		1.1 : 1	
1963-66	1,319	121	355	79
Sex ratio	10.9 : 1		4.5 : 1	

SOURCE: Kreyberg, L. (154).

tion behind that of Finland is reflected in a similar lag in total lung cancer mortality and sex ratios (table 5).

HISTOLOGY OF LUNG TUMORS

A number of investigators have focused their interest upon the relationship of cigarette smoking to the varied histology of lung tumors. The major histological types of lung cancer include squamous cell (epidermoid) carcinoma, small and large cell anaplastic carcinomas, adenocarcinoma (including bronchiolar and alveolar types), and undifferentiated carcinoma (153). A review of these studies (table 6) indicates a closer relationship between cigarette smoking and epidermoid carcinoma than between cigarette smoking and adenocarcinoma (42, 113).

The work of Kreyberg (153) in Norway, over the past 20 years, provides evidence of a specific histologic relationship. This investigator noted that a clearer association is obtained if the various types of pulmonary carcinomas are grouped. Table A7 presents his groupings of the specific histologic types. Using this classification as a basis for analysis of lung cancer sex-ratios in Norway, Kreyberg has observed that Group I carcinomas are significantly more frequent among males while Group II carcinomas show an approximately equal distribution among males and females. The author considers the recent rise in lung cancer in Norway to be a reflection of the increased prevalence of Group I carcinomas. Table 8 presents a summary of Kreyberg's investigation concerning 793 male and female cases of lung cancer. Among both males and females, the risk ratio among smokers is substantially higher for Group I types than for those of Group II. However, adenocarcinoma among males shows a risk ratio of 2.9, signifying a relationship with smoking. Kreyberg attributes the lower rates noted among females to their significantly lower consumption of tobacco in all forms.

TABLE 6.—Epidemiologic and pathologic investigations concerning smoking and the histology of lung cancer¹
(Actual number of cases shown in parentheses)

Author, year, country, reference	Number of persons and case selection method	Results			Comments	
Wynder and Graham, 1950, U.S.A. (316).	644 autopsies on males with confirmed lung cancer.	<i>Percent cases by histologic type and smoking history</i>			The percentage of chain smokers in the general population (7.6) was significantly less than among the patients with adenocarcinoma. The authors refrained from making any definite conclusions due to the insufficient number of cases.	
		<i>All lung cancers other than adenocarcinoma (695)</i>				
		Nonsmokers	1.3	Adenocarcinoma (59)		10.3
		Light cigarette smokers	2.3			7.7
		Moderate	10.1			15.4
		Heavy	35.2			38.5
Excessive	30.9		10.3			
Chain	20.3		18.7			
Doll and Hill, 1962, England (73).	916 male and 79 female cases with histologically confirmed lung cancer.	<i>Percent patients with lung cancer by average amount smoked daily over 10 years</i>			No statistically significant difference was found between the amounts smoked by the patients in the different histological groups. Number of proven adenocarcinomas too small for conclusions.	
		<i>Males</i>				
			<i>Epidermoid (475)</i>	<i>Oat-cell or anaplastic (93)</i>		<i>Adenocarcinoma (35)</i>
		Nonsmokers	0.2 (1)	0.7 (2)		6.1 (2)
		Smokers:				
		<6 cigarettes/day ..	2.9 (14)	3.9 (12)		6.1 (2)
		5-14	35.6 (169)	36.3 (110)		21.2 (7)
		15-25	36.8 (176)	34.7 (105)		48.5 (16)
		>25	24.4 (116)	24.4 (74)		18.2 (6)
		<i>Females</i>				
			<i>Epidermoid (18)</i>	<i>Oat-cell or anaplastic (38)</i>		<i>Adenocarcinoma (10)</i>
		Nonsmokers	61.1 (11)	31.6 (12)		50.0 (5)
		Smokers:				
<6 cigarettes/day ..	5.6 (1)	15.8 (6)	20.0 (2)			
5-14	22.2 (4)	23.7 (9)	10.0 (1)			
15-25	5.6 (1)	18.4 (7)	...			
>25	5.6 (1)	10.5 (4)	20.0 (2)			
				Males—106 unclassified tumors. Females—13 unclassified tumors.		

TABLE 6. *Epidemiologic and pathologic investigations concerning smoking and the histology of lung cancer¹ (cont.)*
(Actual number of cases shown in parentheses)

Author, year, country, reference	Number of persons and case selection method	Results	Comments				
Breslow et al., 1954, U.S.A. (42).	493 male and 25 female cases with histologically proven lung cancer. 518 age and sex-matched controls.	<i>Percent of patients with specific lung cancers by tobacco usage during the 20 years prior to study</i>					
		<i>All lung cancers other than adenocarcinoma</i>					
			Adenocarcinoma (47%)	Controls (518)	Nonsmokers include pipe and cigar smokers only. The authors conclude that cigarette smoking appears to affect the development of epithelial carcinoma more than that of adenocarcinoma.		
Nonsmokers	5.9	13.0					
Cigarette smokers	94.1	87.0					
Schwartz et al., 1957, France (247).	430 male and female cases with histologically confirmed lung cancer. 4 matched control groups.	<i>Percent of smokers by histologic type and smoking history</i>					
			Epidermoid	Anaplastic	Unknown type	Cylindrical	† Difference significant at $p \leq 0.05$ level.
		Cases	96.0	97.0	96.0	100.0	
Controls	79.0†	83.0†	79.0†	96.0			
Haenszel et al., 1958, U.S.A. (119).	158 female cases of lung cancer.	<i>Relative risk for specified tumors (smokers/nonsmokers)</i>			134 cases with final histological determination. † Difference from unity significant at $p \leq 0.01$.		
		Adjusted for age and occupation,	Group I (Kryberg)			Adenocarcinoma	
			3.0†	1.19			
Haenszel and Shlmskin, 1962, U.S.A. (112).	2,191 male cases of lung cancer with adequate histologic data.	<i>Standardized mortality ratios</i>			Cases obtained from a 10 percent sample of lung cancer deaths in U.S.A. during 1968. The authors noted an absence of important differentials by histologic type.		
		<i>Epidermoid and undifferentiated carcinomas</i>				Adenocarcinoma	
		White males total	100			100	
		Never smoked	6			18	
		Ex-smokers	34			46	
		<1 pack/day	123			116	
		>1 pack/day	499			467	

TABLE 6.—Epidemiologic and pathologic investigations concerning smoking and the histology of lung cancer¹ (cont.)

(Actual number of cases shown in parentheses)

Author, year, country, reference	Number of persons and case selection method	Results				Comments	
Cohen and Hossain, 1966, U.S.A. (58).	417 male and female cases of lung cancer with histologic diagnosis 1939-63 at one hospital.	<i>Percent cases by histologic type and smoking history (number of smokers)</i>				The authors also noted that: 1. Adenocarcinomas were 2½-3 times more common in women 2. Only 1 percent of Kreyberg Group I cases were nonsmokers.	
			<i>Squamous</i>	<i>Undifferentiated</i>	<i>Adenocarcinoma</i>		<i>Alveolar</i>
		Nonsmokers	1.0 (3)	10.0 (17)	23.0 (8)		20.0 (1)
		Smokers	89.0 (183)	90.0 (146)	60.0 (20)	...	
Ashley and Davies, 1967, England (6).	442 male and female cases of histologically diagnosed lung cancer.	<i>Percent cases by histologic type and smoking history</i>				The authors noted that cigarette smoking appears to be as strongly related to adenocarcinoma as to the other 2 types. Ashley's data on total number of cigarette smokers are inconsistent with his breakdown of smokers into groups based on number of cigarettes smoked per day.	
			<i>Undifferentiated</i>	<i>Squamous</i>	<i>Adenocarcinoma</i>		
		Nonsmokers	2.8 (4)	2.5 (6)	3.4 (2)		
		Pipe	9.9 (14)	9.9 (24)	1.7 (1)		
		Cigarette	87.3 (124)	87.6 (211)	94.9 (56)		
		<10/day	14.1 (20)	22.4 (54)	22.0 (13)		
		10-20	33.8 (48)	41.5 (100)	33.9 (20)		
		21-30	12.0 (17)	21.6 (52)	16.9 (10)		
31-40	14.1 (20)	12.9 (31)	8.5 (5)				
>40	7.1 (10)	6.2 (15)	5.1 (3)				
Ormos et al., 1969, Hungary (204).	118 male and female cases of histologically proven lung cancer with adequate smoking information.	<i>Percent cases by histologic type and smoking history</i>				The author noted that the small number of cases allows for no definite conclusions.	
			<i>Group I</i>	<i>Group II and large cell carcinomas</i>			
		Nonsmokers	21.0 (18)	36.0 (9)			
	Smokers	79.0 (68)	64.0 (16)				

¹ Data obtained from patient interview and other sources.