

TABLE 8

NUMBER OF REPORTED SHIGELLA ISOLATES BY REGION
FOR 1986-2000

REGION	YEAR														TOTAL	
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999		2000
NEW ENGLAND	729	645	651	546	646	1757	555	520	414	586	373	486	366	851	385	9510
MID-ATLANTIC	1263	2331	1179	1310	1595	1263	1071	1215	1562	1758	2240	1874	1739	750	1726	22876
EAST NORTH CENTRAL	1550	3816	3597	2673	2096	1643	2154	2816	1970	2105	1190	1457	1580	1853	1334	31834
WEST NORTH CENTRAL	330	671	1151	1359	630	403	589	828	1368	1391	699	494	623	806	2064	13406
SOUTH ATLANTIC	686	1425	2562	2568	1894	1565	1727	2860	4790	2181	1912	1136	1275	534	1171	28286
EAST SOUTH CENTRAL	60	219	683	482	799	915	305	768	1023	802	334	233	272	223	207	7325
WEST SOUTH CENTRAL	1611	1671	2535	1408	1832	1409	1453	2943	1771	1944	1225	1731	2427	1688	1549	27197
MOUNTAIN	1137	1174	1692	1161	1655	2248	1174	1348	1436	2713	1665	1238	764	773	868	21046
PACIFIC	4906	4812	8286	4465	6159	5302	5895	6213	4449	5850	4433	3665	3439	2606	3428	73908
TOTAL	12272	16764	22336	15972	17306	16505	14923	19511	18783	19330	14071	12314	12485	10084	12732	235388

TABLE 8A
 NUMBER OF REPORTED S. SONNEI ISOLATES BY REGION
 FOR 1986-2000

REGION	YEAR													TOTAL		
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998		1999	2000
NEW ENGLAND	596	491	481	358	448	1573	435	428	302	456	264	341	248	739	299	7459
MID-ATLANTIC	772	2030	883	1015	1321	1020	880	985	1325	1527	2048	1664	1470	547	1536	19023
EAST NORTH CENTRAL	1123	3279	2685	2061	1443	1176	1671	2440	1707	1773	896	1242	1354	1592	1232	25674
WEST NORTH CENTRAL	147	503	870	1084	446	255	478	698	1166	1281	575	405	538	696	1971	11113
SOUTH ATLANTIC	462	1099	2155	2268	1535	1210	1451	2280	2695	1966	1772	999	1121	396	1003	22412
EAST SOUTH CENTRAL	52	136	523	319	513	587	152	341	649	511	232	211	262	218	196	4902
WEST SOUTH CENTRAL	1161	1293	2103	1022	1415	1072	1268	2765	1595	1721	1085	1514	2239	1449	1286	22988
MOUNTAIN	571	470	921	522	1124	1601	722	824	920	2180	1115	695	374	405	506	12950
PACIFIC	2421	1981	4407	1593	2871	2240	3049	3578	2087	3396	2275	1736	1781	1321	2105	36841
TOTAL	7305	11282	15028	10242	11116	10734	10106	14339	12446	14811	10262	8807	9387	7363	10134	163362

TABLE 8B

NUMBER OF REPORTED S. FLEXNERI ISOLATES BY REGION
FOR 1986-2000

REGION	YEAR													TOTAL		
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998		1999	2000
NEW ENGLAND	93	109	127	150	122	94	107	92	106	115	94	123	102	99	74	1607
MID-ATLANTIC	341	278	257	269	254	227	177	211	213	206	179	188	247	176	154	3377
EAST NORTH CENTRAL	314	361	422	439	465	339	330	287	238	289	267	185	191	223	91	4441
WEST NORTH CENTRAL	168	134	156	168	125	115	77	67	121	71	105	82	79	95	70	1633
SOUTH ATLANTIC	116	154	143	101	156	124	137	173	343	196	122	120	136	127	135	2283
EAST SOUTH CENTRAL	6	9	8	9	10	8	2	4	18	7	7	20	9	5	11	133
WEST SOUTH CENTRAL	380	328	376	333	367	280	170	136	141	188	113	184	146	185	131	3458
MOUNTAIN	496	556	503	472	381	491	371	382	353	464	441	484	352	338	312	6396
PACIFIC	1653	1831	2346	1741	2151	2034	1879	1709	1568	1483	1376	1187	945	777	754	23434
TOTAL	3567	3760	4338	3682	4031	3712	3250	3061	3101	3019	2704	2573	2207	2025	1732	46762

TABLE 8C
 NUMBER OF REPORTED S. BOYDII ISOLATES BY REGION
 FOR 1986-2000

REGION	YEAR													TOTAL		
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998		1999	2000
NEW ENGLAND	6	14	7	16	17	12	4	.	5	11	6	14	6	8	7	133
MID-ATLANTIC	16	16	21	17	10	10	6	11	15	12	5	13	10	12	16	190
EAST NORTH CENTRAL	37	28	64	42	29	20	16	30	19	31	19	23	25	23	6	412
WEST NORTH CENTRAL	1	2	6	5	5	7	3	.	7	6	3	4	4	6	7	66
SOUTH ATLANTIC	6	7	5	10	8	8	5	10	27	7	10	7	9	9	14	142
EAST SOUTH CENTRAL	1	2	2	1	1	.	1	.	.	.	2	10
WEST SOUTH CENTRAL	54	36	39	37	35	44	11	33	24	27	19	27	17	18	13	434
MOUNTAIN	43	59	52	48	47	30	14	14	18	32	91	41	22	23	38	572
PACIFIC	209	192	276	290	246	181	164	123	92	103	120	124	115	59	73	2367
TOTAL	373	356	472	466	398	312	224	221	207	229	275	253	208	158	174	4326

TABLE 8D

NUMBER OF REPORTED *S. DYSENTERIAE* ISOLATES BY REGION
FOR 1986-2000

REGION	YEAR														TOTAL	
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999		2000
NEW ENGLAND	4	4	12	9	8	6	6	.	.	3	6	6	10	3	2	79
MID-ATLANTIC	11	7	18	9	9	6	8	2	6	6	6	8	7	2	11	116
EAST NORTH CENTRAL	7	27	21	23	15	11	21	4	6	11	8	2	9	9	3	177
WEST NORTH CENTRAL	3	8	8	7	3	1	6	1	2	1	2	2	1	2	.	47
SOUTH ATLANTIC	2	4	4	2	5	6	4	4	5	2	8	5	6	2	4	63
EAST SOUTH CENTRAL	1	1	2	.	2	.	1	1	1	.	.	2	1	.	.	12
WEST SOUTH CENTRAL	13	12	17	15	14	10	2	9	10	5	7	6	9	1	3	133
MOUNTAIN	9	10	8	25	7	12	5	12	9	6	18	12	16	7	5	161
PACIFIC	68	159	121	126	118	93	73	72	55	56	48	36	28	23	28	1104
TOTAL	118	232	211	216	181	145	126	105	94	90	103	79	87	49	56	1892

TABLE 8E
 NUMBER OF REPORTED SHIGELLA ISOLATES WITH UNKNOWN SEROGROUP
 BY REGION FOR 1986-2000

REGION	YEAR																TOTAL
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000		
NEW ENGLAND	30	27	24	13	51	72	3	.	1	1	3	2	.	2	3	232	
MID-ATLANTIC	123	.	.	.	1	.	.	6	3	7	2	1	5	13	9	170	
EAST NORTH CENTRAL	69	121	405	108	144	97	116	55	.	1	.	5	1	6	2	1130	
WEST NORTH CENTRAL	11	24	111	95	51	25	25	62	72	32	14	1	1	7	16	547	
SOUTH ATLANTIC	100	161	255	187	190	217	130	393	1720	10	.	5	3	.	15	3386	
EAST SOUTH CENTRAL	.	71	148	153	273	320	149	422	355	284	93	2268	
WEST SOUTH CENTRAL	3	2	.	1	1	3	2	.	1	3	1	.	16	35	116	184	
MOUNTAIN	18	79	208	94	96	114	62	116	136	31	.	6	.	.	7	967	
PACIFIC	555	649	1136	715	773	754	730	731	647	812	614	582	570	426	468	10162	
TOTAL	909	1134	2287	1366	1580	1602	1217	1785	2935	1181	727	602	596	489	636	19046	